







# Nursing: Mental Health and Community Concepts - 2e

*The 2nd edition is currently in progress and will be fully released by June 30, 2025. The book link will not be changed.*

*ERNSTMAYER & CHRISTMAN - OPEN RESOURCES FOR NURSING (OPEN RN)*

WISTECH OPEN



*Nursing: Mental Health and Community Concepts - 2e Copyright © 2025 by WisTech Open is licensed under a Creative Commons Attribution 4.0 International License, except where otherwise noted.*

# Contents

Introduction	1
Preface	4
Standards and Conceptual Approach	xi

## Part I. Chapter 1 Foundational Mental Health Concepts

1.1 Introduction	21
1.2 Mental Health and Mental Illness	23
1.3 Introduction to Trauma-Informed Care	49
1.4 Stigma	53
1.5 Boundaries	57
1.6 Establishing Safety	61
1.7 Psychiatric-Mental Health Nursing	78
1.8 Learning Activities	83
I Glossary	85

## Part II. Chapter 2 Therapeutic Communication and the Nurse-Client Relationship

2.1 Introduction	91
2.2 Basic Concepts of Communication	92
2.3 Therapeutic Communication	100
2.4 Motivational Interviewing	119
2.5 Teletherapy and Telehealth	124
2.6 Learning Activities	127
II Glossary	132

## Part III. Chapter 3 Stress, Coping, and Crisis Intervention

3.1 Introduction	137
3.2 Stress	138
3.3 Coping	156
3.4 Defense Mechanisms	164
3.5 Crisis and Crisis Intervention	170
3.6 Applying the Nursing Process to Stress and Coping	186
3.7 Learning Activities	190
III Glossary	192

## Part IV. Chapter 4 Application of the Nursing Process to Mental Health Care

4.1 Introduction	197
4.2 Applying the Nursing Process	199
4.3 Assessment	201
4.4 Diagnosis	252
4.5 Outcome Identification	264
4.6 Planning	268
4.7 Implementation	275
4.8 Evaluation	290
4.9 NCLEX Next Generation	293
4.10 Spotlight Application	298
4.11 Learning Activities	300
IV Glossary	302

## Part V. Chapter 5 Legal and Ethical Considerations in Mental Health Care

5.1 Introduction	315
5.2 Ethical Principles	316
5.3 Standards of Care	332
5.4 Laws, Torts, Malpractice, and Disciplinary Actions	338
5.5 Client Rights	354
5.6 Learning Activities	370
V Glossary	372

## Part VI. Chapter 6 Psychotropic Medications

6.1 Introduction	379
6.2 Review of the Nervous System	381
6.3 Antidepressants	391
6.4 Mood Stabilizer - Lithium	402
6.5 Antianxiety Medications	405
6.6 Antipsychotics	411
6.7 Stimulants	421
6.8 Psychoactive Substances and Medications to Treat Substance Use and Withdrawal	425
6.9 Learning Activities	427
VI Glossary	429

## Part VII. Chapter 7 Depressive Disorders

7.1 Introduction	435
7.2 Causes of Depression	437

7.3 Types of Depression	451
7.4 Treatments for Depression	470
7.5 Applying the Nursing Process to Depressive Disorders	488
7.6 Spotlight Application	522
7.7 Learning Activities	525
VII Glossary	531

## Part VIII. Chapter 8 Bipolar Disorders

8.1 Introduction	537
8.2 Basic Concepts of Bipolar Disorders	539
8.3 Treatments for Bipolar Disorders	548
8.4 Applying the Nursing Process to Bipolar Disorders	559
8.5 Spotlight Application	598
8.6 Learning Activities	599
VIII Glossary	603

## Part IX. Chapter 9 Anxiety Disorders

9.1 Introduction	607
9.2 Basic Concepts	609
9.3 Anxiety - Related Disorders	614
9.4 Treatments for Anxiety	626
9.5 Obsessive - Compulsive Disorder	635
9.6 Post-Traumatic Stress Disorder	644
9.7 Applying the Nursing Process to Anxiety Disorders	657
9.8 Spotlight Application	682
9.9 Learning Activities	685
IX Glossary	687

## Part X. Chapter 10 Personality Disorders

10.1 Introduction	691
10.2 Basic Concepts	692
10.3 Treatment for Personality Disorders	713
10.4 Applying the Nursing Process to Personality Disorders	718
10.5 Spotlight Application	748
10.6 Learning Activities	750
X Glossary	762

## Part XI. Chapter 11 Thought Disorders

11.1 Introduction	767
11.2 Psychosis and Delirium	768
11.3 Schizophrenia	777
11.4 Applying the Nursing Process to Schizophrenia	798
11.5 Spotlight Application	846
11.6 Learning Activities	848
XI Glossary	853

## Part XII. Chapter 12 Childhood and Adolescence Disorders

12.1 Introduction	859
12.2 Common Disorders and Disabilities in Children and Adolescents	861
12.3 Psychological and Behavioral Therapy	914
12.4 Autism Spectrum Disorder	927

12.5 Applying the Nursing Process to Mental Health Disorders in Children and Adolescents	936
12.6 Spotlight Application	967
12.7 Learning Activities	968
XII Glossary	970

## Part XIII. Chapter 13 Eating Disorders

13.1 Introduction	975
13.2 Basic Concepts	976
13.3 Treatment for Eating Disorders	995
13.4 Applying the Nursing Process to Eating Disorders	1000
13.5 Spotlight Application	1032
13.6 Learning Activities	1035
XIII Glossary	1038

## Part XIV. Chapter 14 Substance Use Disorders

14.1 Introduction	1043
14.2 Substances: Use, Intoxication, and Overdose	1046
14.3 Withdrawal Management/Detoxification	1085
14.4 Substance-Related and Other Addictive Disorders	1098
14.5 Neurobiology of Substance Use Disorders	1107
14.6 Addiction Cycle	1117
14.7 Treatment of Substance Use Disorders	1126
14.8 Prevention of Substance Use Disorders	1157
14.9 Applying the Nursing Process to Substance Use Disorders	1165
14.10 Spotlight Application	1191
14.11 Learning Activities	1193



XIV Glossary	1196
--------------	------

## Part XV. Chapter 15 Trauma, Abuse, and Violence

15.1 Introduction	1203
15.2 Adverse Childhood Experiences	1205
15.3 Trauma-Informed Care	1215
15.4 Abuse and Neglect	1226
15.5 Intimate Partner Violence	1243
15.6 Workplace Violence	1255
15.7 Applying the Nursing Process to Clients Experiencing Trauma, Abuse, or Violence	1269
15.8 Spotlight Activity	1272
15.9 Learning Activities	1276
XV Glossary	1278

## Part XVI. Chapter 16 Community Assessment

16.1 Introduction	1283
16.2 Community Health Concepts	1286
16.3 Applying the Nursing Process to Community Health	1307
16.4 Learning Activities	1335
XVI Glossary	1337

## Part XVII. Chapter 17 Vulnerable Populations

17.1 Introduction	1343
17.2 Vulnerable Populations	1344
17.3 Spotlight Application	1376

17.4 Learning Activities	1384
XVII Glossary	1386

## Part XVIII. Chapter 18 Environmental Health and Emergency Preparedness

18.1 Introduction	1389
18.2 Environmental Health	1391
18.3 Emergency Preparedness, Response, and Recovery	1398
18.4 Nurse Roles in Emergency Response	1423
18.5 Spotlight Application	1429
18.6 Learning Activities	1431
XVIII Glossary	1433

## Part XIX. Answer Key

Chapter 1	1437
Chapter 2	1438
Chapter 3	1439
Chapter 4	1440
Chapter 5	1441
Chapter 6	1442
Chapter 7	1443
Chapter 8	1444
Chapter 9	1445
Chapter 10	1446
Chapter 11	1448
Chapter 12	1449
Chapter 13	1450

Chapter 14	1452
Chapter 15	1453
Chapter 16	1454
Chapter 17	1455
Chapter 18	1456



# Introduction

This is the second edition of the *Nursing Mental Health & Community Concepts* OER textbook that has been developed specifically for prelicensure nursing students. Content is based on the Wisconsin Technical College System (WTCS) statewide nursing curriculum for the Nursing Mental Health & Community Concepts course (543-114), the 2023 NCLEX-RN Test Plan,<sup>1</sup> the American Psychiatric Nurses Association Education Council's *Crosswalk Toolkit: Defining and Using Psychiatric-Mental Health Nursing Skills in Undergraduate Nursing Education*,<sup>2</sup> and the Wisconsin Nurse Practice Act.<sup>3</sup>

This book discusses mental health and community health concepts while emphasizing stress management techniques, healthy coping strategies, referrals to community resources, and other preventative interventions. Nursing care for individuals with specific mental health and substance use disorders is examined, and the nurse's role in community health needs assessments and caring for vulnerable populations is introduced.

Overall updates to the second edition were made based on feedback received from a WTCS faculty workgroup and include the following:

- Learning objectives were updated based on revised WTCS state curriculum.
- Content including DSM-5 criteria and definitions was updated with

1. NCSBN. (n.d.). *Test plans*. <https://www.nclex.com/test-plans.page>
2. American Psychiatric Nurses Association Education Council, Undergraduate Branch. (2016). *Crosswalk toolkit: Defining and using psychiatric-mental health nursing skills in undergraduate nursing education*. <https://www.apna.org/resources/undergraduate-education-toolkit/>
3. Wisconsin State Legislature. (2024). *Chapter 6: Standards of practice for registered nurses and licensed practical nurses*. Board of Nursing. <https://docs.legis.wisconsin.gov/statutes/statutes/441>

content from DSM-5-TR.

- Content based on the ANA Code of Ethics was updated to reflect the 2025 edition.
- Hyperlinks to supplementary information were updated.
- Nursing process sections were standardized across all chapters based on content originally included in Chapter 4 (“Application of the Nursing Process to Mental Health Care”). The NCSBN Clinical Judgment Measurement Model steps were included in parentheses next to the steps of the nursing process (e.g., Recognize Cues, Analyze Cues, Generate Solutions, Take Action, and Evaluate Outcomes). The “Assessment” sections were standardized and include Mental Status Examination, Psychosocial Assessment, a sample PQRSTU assessment with client responses for assessing the reason they are seeking health care, Suicide and Self Injury Screening, Cultural Assessment, Spiritual Assessment with sample client responses, and Lifespan Considerations. The “Implementation” sections include tables summarizing nursing interventions based on categories of the APNA Implementation Standard and common physiological concerns. Communication tips were also included with rationale.
- New Next Generation-style case studies were added to every chapter to help students develop clinical judgment and prepare for the NCLEX.
- Several supplementary YouTube videos were added to enhance student comprehension of the content.

Here is a [document](#) summarizing the updates made to the second edition.

This e book is free with [CC BY 4.0](#) licensing and can be viewed online or downloaded in multiple formats for offline use. It is part of the Open RN<sup>©</sup> Nursing OER textbook series, originally funded by a \$2.5 million Open Textbook Pilot grant from the Department of Education with continued funding by WisTech Open. This book was written by subject matter experts and is based on reliable scholarly research and evidence. Read about other Open RN open educational resources available on the [WisTech Open](#) website.

The following YouTube video provides a quick overview of how to navigate the online version.



*One or more interactive elements has been excluded from this version of the text. You can view them online here: <https://wtcs.pressbooks.pub/nursingmhcc/?p=4#oembed-1>*

## Preface

## Editors

Kimberly Ernstmeyer, MSN, RN, CNE, CHSE, APNP-BC

Dr. Elizabeth Christman, DNP, RN, CNE

## Graphics Editor

Nic Ashman, MLIS, Librarian, Chippewa Valley Technical College

## Developing Authors and Faculty Workgroup

The second edition of this textbook was developed by the editors based on feedback from a workgroup consisting of faculty and industry members:

- Dawn Barone, MSN, RN, Chippewa Valley Technical College
- Kellea Ewen, MSN, RN, Lakeshore Technical College
- Jennifer Hinz, BSN, RN, Manitowoc Public School District
- Kimberly Keith, MSN, RN, Midstate Technical College
- Angela Landry, MSN, RN, Waukesha County Technical College
- Kortnie McIlquham, BS, CBT-C, Chippewa Valley High School
- Krista Polomis, MSN, RN, CNE, Nicolet College

The first edition was created by developing authors by remixing existing open educational resources and developing content based on evidence-based sources and experience in the mental health field:

- Dawn Barone, MSN, RN, Chippewa Valley Technical College
- Dr. Elizabeth Christman, DNP, RN, CNE
- Sue Dzubay, MSN, RN, CHSE, RN-BC, Chippewa Valley Technical College
- Kimberly Ernstmeyer, MSN, RN, CNE, CHSE, APNP-BC, Chippewa Valley Technical College
- Jeanne Green, EdS, MSN, RN, CNE, CHEP, RN-BC, Chippewa Valley Technical College
- Deb Johnson-Schuh, MSN, RN, CNE, Mid-State Technical College



- Rorey Pritchard, EdS, MEd, MSN, RN-BC, CNOR(E), CNE, Senior RN Clinical Educator, Allevant Solutions, LLC
- Dr. Miriam Sward, DNP, PMHNP, APNP, Psychiatric Mental Health Nurse Practitioner
- Alicia Tays, MS, LPC, Counselor

## Contributors

Contributors who assisted in creating the second edition of this textbook include the following:

- Kelly Carpenter, MLIS, Moraine Park Technical College
- Jane Flesher, MST, Proofreader, Chippewa Valley Technical College
- Vince Mussehl, MLIS, WisTech Open Project Director, Chippewa Valley Technical College

## Peer Reviewers

Peer reviewers of the first edition include the following:

- Dr. Caryn Aleo, PhD, RN, CCRN, CEN, CNE, CNEcl, Pasco-Hernando State College
- Dawn Barone, MSN, RN, Chippewa Valley Technical College
- Lisa Bechard, MSN, RN, Mid-State Technical College
- Shanleigh Bechard, MS, LPC, Beloit Area Community Health Center
- Ginger Becker, Nursing Student, Portland Community College
- Nancy Bonard, MSN, RN-BC, St. Joseph's College of Maine
- Dr. Joan Buckley, PhD, RN, Nassau Community College
- Kathleen Capone, MS, RN, CNE, EdD, Utica College
- Lorraine Chiappetta, MSN, RN, CNE, Emeritus Washtenaw Community College
- Travis Christman, MSN, RN, Hospital Sister's Health System
- Maria Darris, MSN, RN, St. Louis Community College
- Tamara Davis, MSN, RN, Chippewa Valley Technical College
- Dr. Andrea Dobogai, DNP, RN, Moraine Park Technical College

- Dr. Judith Dornbach, DNP, MSN, RN, NEA-BC, Southwestern Oregon Community College
- Jessica Dwork, MSN-Ed, RN, Maricopa Community Colleges-Phoenix Collège
- Dr. Terri J. Farmer, PhD, PMHNP-BC, University of Arizona College of Nursing
- Dr. Rachael Farrell, EdD, MSN, CNE Howard Community College, Columbia, MD
- Dr. Vivienne Friday, EdD, MSN, RN, CNE, MSOL Goodwin University
- Jocelyn E. Goodwin, MSN, RN, Paradise Valley Community College
- Kerry L. Hamm, MSN, RN, Lakeland University
- Julia Harelstad, MSN, RN, Mayo Clinic Health System, Eau Claire WI
- Deborah Harmon, BSN, RN, AdventHealth Zephyrhills/Dade City
- Melissa Hauge, MSN, RN, Madison College
- Camille Hernandez, MSN, APRN-Rx, ACNP/FNP, Hawaii Community College – Palamanui
- Katherine Howard, MS, RN-BC, CNE, Middlesex College New Jersey
- Susan Jepsen, MSN, RN, CNE, Lansing Community College
- Eric Johnson, MSN, RN, Chippewa Valley Technical College
- Dr. Kathi L. Johnson, DNP, MS, RN-BC, CNE, Howard Community College
- Jenna Julson, MSN, RN, NPD-RN, Mayo Clinic Health System
- Dr. Andrew D. Kehl, DNP, MSN, MPH, APRN, RN, Vermont Technical College
- Kathryn Kieran, MSN, PMHNP-BC, MGH Institute of Health Professions
- Lindsay Kuhlman, BSN, RN, HSHS Sacred Heart
- Kathy L. Loppnow, MSN, RN, WTCS Health Science Education Director (retired)
- Dawn M. Lyon, MSN, RN, Saint Clair County Community College
- Dr. Erin Micale-Sexton, DNP, MSN-RN, CNL, Lansing Community College, Madonna University, Arizona College
- Sara K. Mitchell, MSN, RN, Fox Valley Technical College
- Dr. Jamie Murphy, PhD, RN, SUNY Delhi
- Kelly Nelson, MSN, RN, Mid-State Technical College
- Tram Nguyen, Nursing Student, Portland Community College

- Dr. Tennille O'Connor, DNP, RN, CNE, College of Central Florida
- Andrea Olson, MSN, RN, Trempealeau County Health Care Center
- Amanda Olson-Komisar, CSW, Social Worker; Pupil Services Teacher Assistant, Pepin Area Schools
- Dr. Grace Paul, DNP, MPhil, RN, CNE, Glendale Community College
- Krista Polomis, MSN, RN, CNE, Nicolet College
- Mary A. Pomietlo, MSN, RN, CNE, University of Wisconsin – Eau Claire
- Cassandra Porter, MSN, RN, Lake Land College
- Rachel Potaczek, BSN, RN, Chippewa County Department of Public Health
- Rorey Pritchard, EdS, MSN, RN-BC, NPD, CNOR(E), CNE, Allevant Solutions, LLC
- Dr. Debbie Rickeard, DNP, MSN, BScN, BA, RN, CNE, CCRN, University of Windsor
- Dr. Argie Rivera, DNP, MAN, PMHNP-BC, Polara Health, Grand Canyon University, Cara Behavioral Health
- Kathleen S. Rizzo, MSN, RN, St Louis Community College at Forest Park
- Ann K. Rosemeyer, MSN, RN, Chippewa Valley Technical College
- Brenda Scheurer, MS, PMP, Eau Claire City-County Health Department
- Cynthia Schroder, MSN, RN, FNP, Yavapai College
- Katherine A. Sell, MSN, RN, IBCLC, CNE, University of Wisconsin – Eau Claire
- Alexis Smith, MScN, RN, Western University
- Justine Sparrgrove, MSN, RN, Southwest Wisconsin Technical College
- Morgan Stock, MSN-Ed, RN, CNE, University of Arizona
- Dr. Miriam A. Sward, DNP, PMHNP, APNP, Northlakes Community Clinic
- Maria E. Thomas, MSN, MS, RN, CNE, Yavapai College
- Jane Trainis, MS, PMHCNS-BC, CNE, Community College of Baltimore County
- Jennie E. Ver Steeg, MLIS, AHIP, Mercy College of Health Sciences
- Emily Vergenz, MSN, RN, Lakeshore Technical College
- Devon Rice Weaver, MSN, RN, Clatsop Community College
- Dr. Nancy Whitehead, PhD, APNP, RN, Milwaukee Area Technical College

# Licensing/Terms of Use

This textbook is licensed under a Creative Commons Attribution 4.0 International (CC-BY) license unless otherwise indicated, which means that you are free to:

- SHARE – copy and redistribute the material in any medium or format
- ADAPT – remix, transform, and build upon the material for any purpose, even commercially

The licensor cannot revoke these freedoms as long as you follow the license terms.

- Attribution: You must give appropriate credit, provide a link to the license, and indicate if any changes were made. You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.
- No Additional Restrictions: You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits.
- Notice: You do not have to comply with the license for elements of the material in the public domain or where your use is permitted by an applicable exception or limitation.
- No Warranties Are Given: The license may not give you all of the permissions necessary for your intended use. For example, other rights such as publicity, privacy, or moral rights may limit how you use the material.

## Report Adoptions or Corrections

Please let us know if you would like to get involved in creating OER or suggest a correction by going to the WisTech Open website and using this [form](#). We appreciate your feedback to help improve and advocate for WisTech Open and Open RN resources!

# Attribution

Some of the content for this textbook was adapted from the following open educational resources. For specific reference information about what was used and/or changed in this adaptation, please refer to the footnotes at the bottom of each page of the book.

- [Nursing Fundamentals, 2e](#) by [Chippewa Valley Technical College](#) is licensed under [CC BY 4.0](#)
- [Nursing Management and Professional Concepts, 2e](#) by [Chippewa Valley Technical College](#) is licensed under [CC BY 4.0](#)
- [Nursing Pharmacology, 2e](#) by [Chippewa Valley Technical College](#) is licensed under [CC BY 4.0](#)
- [mhGAP Intervention Guide – Version 2.0](#) by [World Health Organization](#) is licensed under [CC BY-NC-SA 3.0 IGO](#)
- [Culture and Psychology](#) by Worthy, Lavigne, and Romero is licensed under [CC BY-NC-SA 4.0](#)
- [Human Relations](#) by [LibreTexts](#) is licensed under [CC BY-NC-SA 4.0](#)
- [Anatomy and Physiology](#) by [OpenStax](#) is licensed under [CC BY 4.0](#). Access for free at <https://openstax.org/books/anatomy-and-physiology/pages/1-introduction>
- [Nursing Care at the End of Life](#) by Lowey is licensed under [CC BY-NC-SA 4.0](#)
- [Doing What Matters in Times of Stress: An Illustrated Guide](#) by [World Health Organization](#) is licensed under [CC BY-NC-SA 3.0 IGO](#)
- [StatPearls](#) by Orenstein & Lewis is licensed under [CC BY 4.0](#)
- [Psychology 2e](#) by [OpenStax](#) is licensed under [CC BY 4.0](#). Access for free at <https://openstax.org/books/psychology-2e/pages/1-introduction>
- [StatPearls](#) by Jabbari & Rouster is licensed under [CC BY 4.0](#)
- [StatPearls](#) by Sheffler, Reddy, and Pillarisetty is licensed under [CC BY 4.0](#)
- [StatPearls](#) by Dhaliwal and Gupta is licensed under [CC BY 4.0](#)
- [StatPearls](#) by Shin and Saadabadi is licensed under [CC BY 4.0](#)
- [StatPearls](#) by Wilson and Tripp is licensed under [CC BY 4.0](#)
- [StatPearls](#) by Chokhawala and Stevens is licensed under [CC BY 4.0](#)
- [StatPearls](#) by Ahuja and Abdijadid is licensed under [CC BY 4.0](#)

- [StatPearls](#) by Padda and Derian is licensed under [CC BY 4.0](#)
- [Foundations of Addiction Studies](#) by Flori and Trytek is licensed under [CC BY-NC-SA 4.0](#)
- [StatPearls](#) by Vasan and Kumar is licensed under [CC BY 4.0](#)
- [StatPearls](#) by Kang, Galuska, & Ghassemzadeh is licensed under [CC BY 4.0](#)
- [Action Steps Using ACEs and Trauma-Informed Care: A Resilience Model](#) by Laurie Leitch is licensed under [CC BY 4.0](#)
- [StatPearls](#) by Kisling and Das is licensed under [CC BY 4.0](#)
- [Community Tool Box](#) by [Center for Community Health and Development](#) at the University of Kansas is licensed under [CC BY NC SA 3.0](#)
- [Daily New Confirmed COVID-19 Deaths Per Million People](#) by [Our World in Data](#) is licensed under [CC BY 4.0](#)
- [The Emerging Neurobiology of Bipolar Disorder](#) by Harrison, Geddes, & Tunbridge is licensed under [CC BY 4.0](#)
- [StatPearls](#) by Fisher, Hany, and Doerr is licensed under [CC BY 4.0](#)
- [StatPearls](#) by Lewis and O'Day is licensed under [CC BY 4.0](#)
- [StatPearls](#) by Vasan and Padhy is licensed under [CC BY 4.0](#)
- [The Scholarship of Writing in Nursing Education: 1st Canadian Edition](#) by Lapum, St-Amant, Hughes, Tan, Bogdan, Dimaranan, Frantzke, and Savicevic is licensed under [CC BY-SA 4.0](#)

Suggested attribution statement for this book: Ernstmeyer, K., & Christman, E. (Eds.). (2025). [Nursing Mental Health and Community Concepts, 2e](#) by [Chippewa Valley Technical College](#) is licensed under [CC BY 4.0](#)

## Standards and Conceptual Approach

The Open RN *Nursing Mental Health and Community Concepts*, 2e textbook is based on several external standards and uses a conceptual approach.

### External Standards

#### American Nurses Association (ANA)

The ANA establishes Standards for Professional Nursing Practice and the Code of Ethics for Nurses.<sup>12</sup>

- <https://www.nursingworld.org/ana/about-ana/standards/>

#### American Psychiatric Nurses Association (APNA)

The APNA advances the science and education of psychiatric-mental health nursing. APNA is committed to the practice of psychiatric-mental health nursing, health and wellness promotion through identification of mental health issues, prevention of mental health problems, and the care and treatment of persons with mental health disorders.<sup>3</sup>

- <https://www.apna.org/>

#### American Psychiatric Nurses Association Education Council, Undergraduate Branch

1. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.
2. American Nurses Association. (2025). *Code of ethics for nurses*. American Nurses Association. <https://codeofethics.ana.org/provisions>
3. American Psychiatric Nurses Association. <https://www.apna.org/>

The APNA created a toolkit to help define and integrate psychiatric-mental health nursing content into undergraduate nursing curricula.<sup>4</sup>

- <https://www.apna.org/resources/undergraduate-education-toolkit/>

## **Substance Abuse and Mental Health Services Administration (SAMHSA)**

The Substance Abuse and Mental Health Services Administration (SAMHSA) is the agency within the U.S. Department of Health and Human Services that leads public health efforts to advance the behavioral health of the nation and to improve the lives of individuals living with mental health and substance use disorders and their families. SAMHSA's mission is to reduce the impact of substance abuse and mental illness on America's communities and draws advice from public members and professionals in the field of substance abuse and mental health.<sup>5</sup>

- <https://www.samhsa.gov/>

## **The National Council Licensure Examination for Registered Nurses: NCLEX-RN Test Plans**

The NCLEX-RN test plans are updated every three years to reflect fair, comprehensive, current, and entry-level nursing competency.<sup>6</sup>

4. American Psychiatric Nurses Association Education Council, Undergraduate Branch. (2016). *Crosswalk toolkit: Defining and using psychiatric-mental health nursing skills in undergraduate nursing education*. <https://www.apna.org/resources/undergraduate-education-toolkit/>
5. Substance Abuse and Mental Health Services Administration. (n.d.). *Strategic plan FY2019-FY2023*. [https://www.samhsa.gov/sites/default/files/samhsa\\_strategic\\_plan\\_fy19-fy23\\_final-508.pdf](https://www.samhsa.gov/sites/default/files/samhsa_strategic_plan_fy19-fy23_final-508.pdf)
6. NCSBN. (n.d.). *Test plans*. <https://www.nclex.com/test-plans.page>



- <https://www.ncsbn.org/nclex.htm>

## **The National League of Nursing (NLN): Competencies for Graduates of Nursing Programs**

NLN competencies guide nursing curricula to position graduates in a dynamic health care arena with practice that is informed by a body of knowledge to help ensure the public receives safe, quality care.<sup>7</sup>

- <https://www.nln.org/education/nursing-education-competencies/competencies-for-graduates-of-nursing-programs>

## **American Association of Colleges of Nursing (AACN): The Essentials: Competencies for Professional Nursing Education**

The AACN provides a framework for preparing individuals as members of the discipline of nursing, reflecting expectations across the trajectory of nursing education and applied experience.

- <https://www.aacnnursing.org/Portals/42/AcademicNursing/pdf/Essentials-2021.pdf>

## **Quality and Safety Education for Nurses (QSEN) Institute: Prelicensure Competencies**

Quality and safety competencies include knowledge, skills, and attitudes to be developed in nursing prelicensure programs. QSEN competencies include patient-centered care, teamwork and collaboration, evidence-based practice, quality improvement, safety, and informatics.<sup>8</sup>

7. National League of Nursing. *Competencies for graduates of nursing programs*. <https://www.nln.org/education/nursing-education-competencies/competencies-for-graduates-of-nursing-programs>
8. QSEN. (n.d.). *About*. <https://qsen.org/about-qsen/>

- <https://qsen.org/competencies/>

## Wisconsin State Legislature, Administrative Code Chapter N6

The Wisconsin Administrative Code governs the Registered Nursing and Practical Nursing professions in Wisconsin.<sup>9</sup>

- [https://docs.legis.wisconsin.gov/code/admin\\_code/n/6](https://docs.legis.wisconsin.gov/code/admin_code/n/6)

## Healthy People 2030

Healthy People 2030 envisions a society in which all people can achieve their full potential for health and well-being across the life span. Healthy People provides objectives based on national data and includes social determinants of health.<sup>10</sup>

- <https://health.gov/healthypeople>

## Conceptual Approach

The Open RN *Nursing Mental Health and Community Concepts*, 2e textbook incorporates the following concepts:

- **Holism.** Florence Nightingale taught nurses to focus on the principles of holism, including wellness and the interrelationship of human beings and their environment. This textbook encourages holistic nursing care by addressing the impact of social determinants of health (SDOH) on mental

9. Wisconsin State Legislature. (2018). *Chapter 6: Standards of practice for registered nurses and licensed practical nurses*. Board of Nursing. <https://docs.legis.wisconsin.gov/statutes/statutes/441>

10. Healthy People 2030. (n.d.). *Social determinants of health*. U.S. Department of Health and Human Services. <https://health.gov/healthypeople/objectives-and-data/social-determinants-health>

health.

- **Evidence-Based Practice (EBP).** Evidence-based practices are referenced by footnotes throughout the textbook. To promote the development of digital literacy, hyperlinks are provided to credible, free online resources that supplement content. The Open RN textbooks will be updated as new EBP is established and after the release of updated NCLEX Test Plans every three years.
- **Clinical Judgment.** Associated unfolding case studies are written to reflect the NCSBN Clinical Judgment Measurement Model used on the NCLEX-RN. Formative assessments encourage students to recognize cues, analyze cues, prioritize hypotheses, generate solutions, take action, and evaluate outcomes.<sup>11</sup>
- **Cultural Competency.** Nurses have an ethical obligation to practice with cultural humility and provide culturally responsive care to the clients and communities they serve based on the ANA Code of Ethics<sup>12</sup> and the ANA Scope and Standards of Practice.<sup>13</sup>
- **Safe, Quality, Patient-Centered Care.** Content reflects the priorities of safe, quality, patient-centered care.
- **Clear and Inclusive Language.** Clear language is used based on preferences expressed by prelicensure nursing students to enhance understanding of complex concepts.<sup>14</sup> “They” is used as a singular

11. Dickison, P., Haerling, K. A., & Lasater, K. (2019). Integrating the national council of state boards of nursing clinical judgment model into nursing educational frameworks. *Journal of Nursing Education*. 58(2), 72-78. <https://doi.org/10.3928/01484834-20190122-03>

12. American Nurses Association. (2025). *Code of ethics for nurses*. American Nurses Association. <https://codeofethics.ana.org/provisions>

13. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

14. Verkuyl, M., Lapum, J., St-Amant, O., Bregstein, J., & Hughes, M. (2020).

pronoun to refer to a person whose gender is unknown or irrelevant to the context of the usage, as endorsed by APA style. It is inclusive of all people and helps writers avoid making assumptions about gender.<sup>15</sup>

- **Open-Source Images and Fair Use.** Images are included to promote visual learning. Students and faculty can reuse open-source images by following the terms of their associated [Creative Commons licensing](#). Some images are included based on Fair Use as described in the “[Code of Best Practices for Fair Use and Fair Dealing in Open Education](#)” presented at the OpenEd 2020 conference. Refer to the footnotes of images for source and licensing information throughout the text.
- **Open Pedagogy.** Students are encouraged to contribute to the Open RN project in meaningful ways by reviewing content for clarity and assisting in the creation of open-source images.<sup>16</sup>

## Supplementary Materials

Several supplementary resources are provided with this textbook:

- Links to supplementary videos promote student understanding of concepts and procedures.
- Online, interactive, and written learning activities in every chapter provide immediate formative feedback.
- Critical thinking questions encourage the development of clinical judgment as students apply content to authentic client care scenarios.

Healthcare students’ use of an e-textbook open educational resource on vital sign measurement: A qualitative study. *Open Learning: The Journal of Open, Distance and e-Learning*. <https://doi.org/10.1080/02680513.2020.1835623>

15. American Psychological Association. (2021). *Singular “They.”* <https://apastyle.apa.org/style-grammar-guidelines/grammar/singular-they>

16. [The Open Pedagogy Notebook](#) by Steel Wagstaff is licensed under [CC BY 4.0](#)

- Formative Next Generation-style case studies are provided in every chapter to help students prepare for the NCLEX-RN.
- Free downloadable textbook versions can be downloaded from the home page for offline use.
- Affordable soft cover print versions can be purchased from [Kendall-Hunt Publishing Company](#)









## 1.1 Introduction

### Learning Objectives

- Identify safety/protective interventions for the client and others
- Identify safety/protective concerns for the nurse
- Explain various mental health treatment approaches
- Support diversity across the lifespan in client-centered care

Mental health is an important part of everyone's overall health and well-being. Mental health includes our emotional, psychological, and social well-being. It affects how we think, feel, and act. It also helps determine how we handle stress, relate to others, and make healthy choices. Mental health is important at every stage of life, from childhood to adolescence and through adulthood.<sup>1</sup> This chapter will provide an overview of mental health, mental illness, and mental health nursing. As with all areas of nursing, when caring for a person with a mental health diagnosis, it is important to focus on client-centered care and evaluate the effectiveness of care in terms of the highest level of functioning that person is able to achieve.

### Reflective Questions

1. Centers for Disease Control and Prevention. (2025). *About mental health*. <https://www.cdc.gov/mental-health/about/index.html>  
[https://www.cdc.gov/mental-health/?CDC\\_AAref\\_Val=https://www.cdc.gov/mentalhealth/index.htm](https://www.cdc.gov/mental-health/?CDC_AAref_Val=https://www.cdc.gov/mentalhealth/index.htm)

As we begin this chapter, reflect on the following questions:

1. How do you define mental health?
2. How do you define mental illness?
3. How can you tell the difference between someone experiencing general mental health challenges and someone with a mental illness by looking at how they function day to day?
4. Consider how you communicate with clients. Taking into consideration individual differences, which therapeutic communication techniques have you found to be the most effective? What factors have you found interfere with effective communication, and how do you address them in practice?
5. How does ineffective communication impact client care? How can it affect your nursing license or create legal implications?

## 1.2 Mental Health and Mental Illness

Mental health is an essential component of health. The World Health Organization (WHO) defines **health** as a state of complete physical, mental, and social well-being and not merely the absence of disease or infirmity. **Mental health** is a state of well-being in which an individual realizes their own abilities, copes with the normal stresses of life, works productively, and contributes to their community. The promotion, protection, and restoration of mental health is a vital concern of individuals, nurses, communities, and societies throughout the world.<sup>1</sup>

According to the American Psychiatric Association, **mental illness** is a health condition involving changes in emotion, thinking, or behavior (or a combination of these) associated with emotional distress and problems functioning in social, work, or family activities.<sup>2</sup> The term mental illness is an older term that is often used interchangeably with mental health disorders. The term mental health disorders is a more modern clinical term used in diagnostic manuals like the *Diagnostic and Statistical Manual of Mental Disorders (DSM-5-TR)*.

Mental illness is common in the United States. Nearly one in five (19 percent) of adults experience some form of mental illness, one in twelve (8.5 percent) have a substance use disorder, and one in 24 (4 percent) have a serious mental illness.<sup>3</sup>

1. World Health Organization. (2018). *Mental health: Strengthening our response*. <https://www.who.int/news-room/fact-sheets/detail/mental-health-strengthening-our-response>
2. American Psychiatric Association. (n.d.). *What is mental illness?* <https://www.psychiatry.org/patients-families/what-is-mental-illness>
3. American Psychiatric Association. (n.d.). *What is mental illness?* <https://www.psychiatry.org/patients-families/what-is-mental-illness>

Mental health disorders increase the risk of chronic physical illnesses, such as heart disease, diabetes, respiratory diseases, cancer, and strokes, and can lead to thoughts and intentions of suicide.<sup>4</sup> Suicide is a common symptom associated with mental illness and is the second leading cause of death in Americans aged 15-34.<sup>5</sup>



View the following YouTube video on An Update About WHO's Special Initiative on Mental Health<sup>6</sup>: [WHO Special Initiative on Mental Health: An update on the latest figures, learnings, and challenges](https://youtu.be/ti7OIMq7V9I)

## Mental Health Continuum

Mental health fluctuates over the course of an individual's life span and can

4. Scott, K. M., Lim, C., Al-Hamzawi, A., Alonso, J., Bruffaerts, R., Caldas-de-Almeida, J. M., Florescu, S., de Girolamo, G., Hu, C., de Jonge, P., Kawakami, N., Medina-Mora, M. E., Moskalewicz, J., Navarro-Mateu, F., O'Neill, S., Piazza, M., Posada-Villa, J., Torres, Y., & Kessler, R. C. (2016). Association of mental disorders with subsequent chronic physical conditions: World mental health surveys from 17 countries. *JAMA Psychiatry*, 73(2), 150-8. <https://doi:10.1001/jamapsychiatry.2015.2688>.
5. Centers for Disease Control and Prevention. (2025). *About mental health*. <https://www.cdc.gov/mental-health/about/index.html>
6. World Health Organization. (2023, December 6). *WHO special initiative on mental health: An Update on the latest figures, learnings, and challenges* [Video]. YouTube. All rights reserved. <https://youtu.be/ti7OIMq7V9I>

range from well-being to emotional problems and/or mental illness as indicated on the **mental health continuum** illustrated in Figure 1.1.<sup>7,8,9</sup>




Figure 1.1 Mental Health Continuum (Used with permission.)

**Well-being** is in the “healthy” range of the mental health continuum in which individuals are experiencing a state of good mental and emotional health. They may experience stress and discomfort resulting from occasional

7. “continuum.jpg” by [University of Michigan](https://hr.umich.edu/benefits-wellness/health-well-being/mhealthy/faculty-staff-well-being/mental-emotional-health/mental-emotional-health-classes-training-events/online-tutorial-supervisors/section-1-what-you-need-know-about-mental-health-problems-substance-misuse) is used with permission. Access the original at <https://hr.umich.edu/benefits-wellness/health-well-being/mhealthy/faculty-staff-well-being/mental-emotional-health/mental-emotional-health-classes-training-events/online-tutorial-supervisors/section-1-what-you-need-know-about-mental-health-problems-substance-misuse>

8. William, S. (2021). The continuum between temperament and mental illness as dynamical phases and transitions. *Frontiers in Psychiatry*, 11, 1617. <https://doi.org/10.3389/fpsyt.2020.614982>

9. University of Michigan Human Resources. (n.d.). *Section 1: What you need to know about mental health problems and substance misuse*. <https://hr.umich.edu/benefits-wellness/health-well-being/mhealthy/faculty-staff-well-being/mental-emotional-health/mental-emotional-health-classes-training-events/online-tutorial-supervisors/section-1-what-you-need-know-about-mental-health-problems-substance-misuse>

problems of everyday life, but they are able to cope effectively with these stressors and experience no impairments to daily functioning.

On the other end of the mental health continuum are mental health problems where individuals have progressively more difficulty coping with problems and stressors. Within this range are two categories: emotional problems/concerns and mental illness. For individuals experiencing emotional problems, discomfort has risen to a level of mild to moderate distress, and they are experiencing mild or temporary impairments in functioning, such as insomnia, lack of concentration, or loss of appetite. As their level of distress increases, they may seek treatment and often start with visiting their primary health care provider.

Emotional problems are classified as a mental illness when a person experiences significant distress, along with moderate to severe impairment in daily functioning at work, school, or home. This diagnosis is made when the individual meets specific clinical criteria and their symptoms cannot be better explained by a medical condition or substance use. Mental illness includes relatively common disorders, such as depression and anxiety, as well as less common disorders such as schizophrenia. Mental illness is characterized by alterations in thinking, mood, or behavior. The term **serious mental illness** refers to mental illness that causes disabling functional impairment that substantially interferes with one or more major life activities. The Americans With Disabilities Act defines **major life activities** as, “caring for oneself, performing manual tasks, seeing, hearing, eating, sleeping, walking, standing, lifting, bending, speaking, breathing, learning, reading, concentrating, thinking, communicating, and working.”<sup>10</sup> Examples of serious mental illnesses that commonly interfere with major life activities include major

10. Office of Federal Contract Compliance Programs. (2009). *ADA Amendments Act of 2008 frequently asked questions*. U.S. Department of Labor. <https://www.dol.gov/general/topic/disability/ada>

depressive disorder, schizophrenia, and bipolar disorder.<sup>11</sup> Individuals with serious mental illnesses may experience long-term impairments ranging from moderate to disabling in nature, but many can lead productive lives with effective treatment. For example, roughly half of schizophrenia clients recovered or significantly improved over the long-term, suggesting that functional remission is possible.<sup>12,13</sup>

Mental health providers, such as psychiatrists, psychologists, therapists, social workers, or advanced practice mental health nurses, use the *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5-TR)* published by the American Psychiatric Association to assess a client's signs and symptoms and determine a mental health diagnosis. The manual lists diagnostic criteria including feelings, behaviors, and time frames to be officially classified as a mental health disorder.<sup>14</sup>

While the exact number of mental health disorders can vary slightly depending on the classification system used, the DSM-5-TR lists approximately 157 distinct disorders.<sup>15</sup> People can experience different types of

11. American Psychiatric Association. (n.d.). *What is mental illness?*  
<https://www.psychiatry.org/patients-families/what-is-mental-illness>
12. Vita, A., & Barlati, S. (2018). Recovery from schizophrenia: Is it possible? *Current Opinion in Psychiatry*, 31(3), 246–255. <https://doi.org/10.1097/YCO.0000000000000407>
13. Rakitzi, S., Georgila, P., & Becker-Woitag, A. P. (2021). The recovery process for individuals with schizophrenia in the context of evidence-based psychotherapy and rehabilitation: A systematic review. *European Psychologist*, 26(2), 96–111. <https://doi.org/10.1027/1016-9040/a000400>
14. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR*.
15. Borgogna, N. C., Owen, T., & Aita, S. L. (2024). The absurdity of the latent

mental health disorders, and different disorders can occur at the same time or vary in intensity over time. Mental illness can be ongoing, occur over a short period of time, or be episodic (i.e., it comes and goes with discrete beginnings and ends).<sup>16</sup>

► Read more information about specific mental health disorders at the [Medline Plus Mental Health and Behavior webpage](#).

## Assessing Dysfunction and Impairment

Mental health disorders have been defined as a type of dysfunction that causes distress or impaired functioning and deviates from typical or expected behavior according to societal or cultural standards. This definition includes three components referred to as dysfunction, distress, and deviance.<sup>17</sup>

**Dysfunction** includes disturbances in a person's thinking, emotional regulation, or behavior that reflects significant dysfunction in psychological, biological, or developmental processes underlying mental functioning. In other words, dysfunction refers to a breakdown in cognition, emotion, and/or behavior. For instance, an individual experiencing the delusion of being an all powerful deity demonstrates a breakdown in cognition, as their thought processes are disconnected from reality. An individual who is unable to

disease model in mental health: 10,130,814 ways to have a DSM-5-TR psychological disorder. *Journal of Mental Health*. 33(4):451-459. doi: 10.1080/09638237.2023.2278107

16. Centers for Disease Control and Prevention. (2025). *About mental health*. <https://www.cdc.gov/mental-health/about/index.html>

17. Worthy, L. D., Lavigne, T., & Romero, F. (2025). *Culture and psychology*. Maricopa Community College. <https://open.maricopa.edu/culturepsychology/>



experience pleasure has a breakdown in emotion, and an individual who is unable to leave home and attend work due to fear of having a panic attack is exhibiting a breakdown in behavior.<sup>18</sup>

**Distress** refers to psychological and/or physical pain. Simply put, distress refers to suffering. For example, the loss of a loved one causes anyone to experience emotional pain, distress, and a temporary impairment in functioning. **Impairment** refers to a limited ability to engage in activities of daily living (i.e., they cannot maintain personal hygiene, prepare meals, or pay bills) or participate in social events, work, or school. Impairment can also interfere with the ability to perform important life roles such as a caregiver, parent, or student.<sup>19</sup>

**Deviance** refers to behavior that violates social norms or cultural expectations because one's culture determines what is "normal." When a person is described as "deviant," it means they are not following the stated and unstated rules of their society (referred to as **social norms**).<sup>20</sup>

Nurses complete and document initial and ongoing assessments of dysfunction, distress, and behavior associated with an individual's diagnosed mental health disorder. The World Health Organization Disability Assessment Scale (WHODAS) is a recommended tool for assessing impairments resulting from mental illness.<sup>21</sup> The **WHODAS** is a generic assessment instrument that

18. Worthy, L. D., Lavigne, T., & Romero, F. (2025). *Culture and psychology*. Maricopa Community College. <https://open.maricopa.edu/culturepsychology/>
19. Worthy, L. D., Lavigne, T., & Romero, F. (2025). *Culture and psychology*. Maricopa Community College. <https://open.maricopa.edu/culturepsychology/>
20. Worthy, L. D., Lavigne, T., & Romero, F. (2025). *Culture and psychology*. Maricopa Community College. <https://open.maricopa.edu/culturepsychology/>
21. Office of Federal Contract Compliance Programs. (2009). *ADA Amendments Act of 2008 frequently asked questions*. U.S. Department of Labor.

provides a standardized method for measuring health and disability across cultures.<sup>22</sup> The WHODAS assesses functioning in six domains: cognition, mobility, self-care, getting along, life activities, and participation.<sup>23</sup>

► View the [WHODAS 2.0 webpage](#).

The Global Assessment of Functioning (GAF) was historically used to rate the seriousness of a mental illness and measure how symptoms affect an individual's day-to-day life on a scale of 0 to 100. It is an overall (global) measure of how clients are doing and rates psychological, social, and occupational functioning on the continuum from mental well-being to serious mental illness. The higher the score, the better the daily functioning. The GAF was omitted from the *DSM-5-TR* because it had questionable validity

<https://www.dol.gov/agencies/ofccp/faqs/americans-with-disabilities-act-amendments>

22. World Health Organization. (2012). *Measuring health and disability: Manual for WHO disability assessment schedule (WHODAS 2.0)*. [Manual]. [https://www.who.int/publications/i/item/measuring-health-and-disability-manual-for-who-disability-assessment-schedule-\(-whodas-2.0\)](https://www.who.int/publications/i/item/measuring-health-and-disability-manual-for-who-disability-assessment-schedule-(-whodas-2.0))
23. National Academies of Sciences, Engineering, and Medicine. (2016). *Measuring specific mental illness diagnoses with functional impairment: Work-shop summary* [PDF]. J. C. Rivard and K. Marton, Rapporteurs. Committee on National Statistics and Board on Behavioral, Cognitive, and Sensory Sciences, Division of Behavioral and Social Sciences and Education. Board on Health Sciences Policy, Institute of Medicine. The National Academies Press. <http://elibrary.pcu.edu.ph:9000/digi/NA02/2016/21920.pdf>

and reliability, but some government agencies and insurance companies continue to include it in paperwork to assess client functioning.<sup>24</sup>

## Recovery

Mental illness is treatable. Research shows that people with mental illness can get better, and many recover completely.<sup>25</sup> The majority of individuals with mental illness continue to function in their daily lives. **Recovery** refers to a process of change through which individuals improve their health and wellness, live a self-directed life, and strive to reach their full potential.<sup>26</sup> Dimensions that support a life in recovery include the following:

- **Health:** Overcoming or managing one's disease(s), as well as living in a physically and emotionally healthy way
- **Home:** Having a stable and safe place to live
- **Purpose:** Participating in meaningful daily activities, such as a job, school, volunteerism, family caretaking, or creative endeavors; and the independence, income, and resources to participate in society
- **Community:** Enjoying relationships and social networks that provide support, friendship, love, and hope

## Early Signs of Mental Health Problems

Mental health problems are common. We all experience problems and stressors from daily living at the mild end of the mental health continuum,

24. Smith, M. (2025). *What is the global assessment of functioning (GAF) scale?* WebMD. <https://www.webmd.com/mental-health/gaf-scale-facts>
25. Centers for Disease Control and Prevention. (2025). *About mental health.* <https://www.cdc.gov/mental-health/about/index.html>
26. Center for Substance Abuse Treatment. (2014). *Trauma-informed care in behavioral health services.* <https://www.ncbi.nlm.nih.gov/books/NBK207201/>

and each year one in five Americans experience mental illness.<sup>27</sup> Nurses in all care settings must recognize signs and symptoms of diagnosed and undiagnosed emotional and mental health problems in clients. Each mental health disorder has specific signs and symptoms, but common signs of mental health problems in adults and adolescents are as follows<sup>28</sup>:

- Excessive worrying or fear
- Excessive sad or low feelings
- Confused thinking or problems concentrating and learning
- Extreme mood changes, including uncontrollable “highs” or feelings of euphoria
- Prolonged or strong feelings of irritability or anger
- Avoidance of friends and social activities
- Difficulty understanding or relating to other people
- Changes in sleeping habits or feeling tired and low energy
- Changes in eating habits, such as increased hunger or lack of appetite
- Changes in sex drive
- Disturbances in perceiving reality, referred to as hallucinations (i.e., when a person senses things that don’t exist in reality)
- Inability to perceive changes in one’s own feelings, behavior, or personality (i.e., lack of insight)
- Misuse of substances like alcohol, drugs, or prescription medications
- Multiple physical ailments without obvious causes (such as headaches, stomachaches, or vague and ongoing “aches and pains”)
- Thoughts of suicide
- Inability to carry out daily activities or handle daily problems and stress
- Intense fear of weight gain or being overly concerned with appearance

27. National Alliance on Mental Illness. (2023). *Mental health by the numbers*.  
<https://www.nami.org/about-mental-illness/mental-health-by-the-numbers/>

28. National Alliance on Mental Illness. (n.d.). *Warning signs and symptoms*.  
<https://nami.org/About-Mental-Illness/Warning-Signs-and-Symptoms>

Mental health problems can also be present in young children. Because children are still learning how to identify and talk about thoughts and emotions, their most obvious symptoms are behavioral or complaints of physical symptoms. Behavioral symptoms in children can include the following<sup>29</sup> :

- Changes in school performance
- Excessive worry or anxiety, for example, fighting to avoid going to bed or school
- Hyperactive behavior
- Frequent nightmares or a marked change in sleeping habits
- Frequent disobedience or aggression
- Frequent temper tantrums

View the following YouTube video about warning signs of mental health problems<sup>30</sup> : [10 Common Warning Signs of a Mental Health Condition](https://youtu.be/zt4sOjWwV3M)

## Cultural Impact

**Culture** is a complex context of knowledge, beliefs, and behaviors that provide structure to specific ethnic, racial, religious, geographic, or social groups. Culture encompasses the personal identity, language, thoughts,

29. National Alliance on Mental Illness. (n.d.). *Warning signs and symptoms*. <https://nami.org/About-Mental-Illness/Warning-Signs-and-Symptoms>

30. NAMI. (2015, February 2). *10 common warning signs of a mental health condition* [Video]. YouTube. All rights reserved. <https://youtu.be/zt4sOjWwV3M>

communications, actions, customs, beliefs, values, and institutions of a variety of groups of people, as well as smaller groups of individuals within a larger community. Cultural values and beliefs impact how a person views certain ideas or behaviors. In the case of mental health, it can impact whether or not the individual seeks help, the type of help sought, and the support available. Every individual has different cultural beliefs and faces a unique journey to recovery. In general, historically marginalized communities in the United States are less likely to access mental health treatment, or they wait until symptoms are severe before seeking assistance.<sup>31</sup>

Four ways that culture can impact mental well-being are the following<sup>32</sup>:

- **Cultural stigma.** Every culture has a different perspective on mental health, and many cultures have a stigma surrounding mental health. Mental health challenges may be considered a weakness and something to hide, which can make it harder for those struggling to talk openly and ask for help. For example, in some Asian cultures, mental health challenges are often viewed as a sign of weakness or a failure to meet familial or societal expectations. For instance, a young woman in Japan might experience severe depression due to the pressure of excelling academically and meeting the high standards of her family. She may feel discouraged from openly discussing her condition, believing it reflects poorly on their upbringing or family reputation.
- **Describing symptoms.** Culture can influence how people describe or feel about their symptoms. It can affect whether someone chooses to recognize and talk openly about physical symptoms, emotional

31. Mental Health First Aid. (2019). *Four ways culture impacts mental health*. National Council for Mental Wellbeing. <https://www.mentalhealthfirstaid.org/2019/07/four-ways-culture-impacts-mental-health/>

32. Mental Health First Aid. (2019). *Four ways culture impacts mental health*. National Council for Mental Wellbeing. <https://www.mentalhealthfirstaid.org/2019/07/four-ways-culture-impacts-mental-health/>

symptoms, or both. For example, members of the Amish community are typically stoic and endure physical and emotional pain without complaining.

- **Community support.** Cultural factors can determine how much support someone gets from their family and community when it comes to mental health. Because of existing stigma, it can be challenging for individuals to find mental health treatment and support. For example, in a close-knit Latinx community, the value placed on family unity and support is a central cultural norm. A young man begins experiencing anxiety and panic attacks and decides to confide in his family. However, when he brings up the idea of seeking therapy, his parents are skeptical, and a trusted elder—a respected family friend—dismisses the idea, saying that mental health struggles are a sign of personal weakness and can be overcome with willpower.
- **Resources.** When looking for mental health treatment, it can be difficult to find resources and treatment options that take into account a specific culture's concerns and needs. For example, A Somali immigrant living in the United States begins experiencing symptoms of post-traumatic stress disorder (PTSD) due to their experiences during the civil war in their home country. They want to seek help but face several barriers related to their cultural concerns and needs. While they find a few local clinics offering mental health services, none have therapists who are familiar with Somali culture, language, or the unique traumas faced by refugees. The therapists use approaches that the individual feels are disconnected from their values, such as emphasizing individualistic coping mechanisms rather than community-based healing, which is more aligned with their culture.

Nurses can help clients by understanding the role culture plays in their mental health. They should also be aware of how their own culture might impact their perception of their client's mental health status. This represents cultural humility. **Cultural humility** is a lifelong process of self-reflection and self-critique whereby individuals actively engage in learning about different

cultural perspectives while acknowledging the limitations of their own knowledge.

- ▶ Read more about cultural diversity and providing culturally responsive care in the “[Diverse Patients](#)” chapter of *Open RN Nursing Fundamentals*.
- ▶ Read more about improving your cultural competency at “[Improving Cultural Competency for Behavioral Health Professionals](#)”

## Causes of Mental Illness

Mental health researchers have developed several theories to explain the causes of mental health disorders, but they have not reached consensus. One factor in which they all agree is that an individual is not at fault for the condition, and they cannot simply turn symptoms on or off at will. There are likely several factors that combine to trigger a mental health disorder, including environmental, biological, and genetic factors.<sup>33</sup> Another significant role in the development and exacerbation of mental illness is psychological factors, which include both interpersonal and coping mechanisms. Interpersonal conflicts include chronic interpersonal stress, such as disagreements, rejections, and feeling let down, can significantly contribute to mental health issues. These stressors are associated with increased negative affect, which in turn can lead to various health problems, including

33. University of Michigan Human Resources. (n.d.). *Section 1: What you need to know about mental health problems and substance misuse*.  
<https://hr.umich.edu/benefits-wellness/health-well-being/mhealthy/faculty-staff-well-being/mental-emotional-health/mental-emotional-health-classes-training-events/online-tutorial-supervisors/section-1-what-you-need-know-about-mental-health-problems-substance-misuse>



anxiety and depression. Interpersonal problems are particularly impactful in conditions like anxious depression, where dysfunctional attitudes mediate the relationship between interpersonal stress and anxiety symptoms. An individual's coping mechanisms reflect the way a person deals with stress and adversity. Coping mechanisms can either mitigate or exacerbate mental health problems. Mal-adaptive coping strategies, such as emotion-focused coping, can worsen psychological distress and increase the likelihood of mental disorders.<sup>34</sup>

## Environmental Factors

Individuals are affected by broad social and cultural factors, as well as by unique factors in their personal environments. Social factors such as racism, discrimination, poverty, and violence (often referred to as “social determinants of health”) can contribute to mental illness.

- ▶ Read more about addressing social determinants of health in the “[Advocacy](#)” chapter of *Open RN Nursing Management and Professional Concepts*.

Additionally, it is estimated that 61% of adults have experienced early **adverse childhood experiences (ACEs)** such as abuse, neglect, or growing up in a household with violence, mental illness, substance misuse, incarceration, or divorce. Chronic stress from ACEs can change brain development and affect how the body responds to stress. ACEs are linked to chronic health problems,

34. Woo, J., Whyne, E. Z., & Steinhardt, M. A. (2024). Psychological distress and self-reported mental disorders: The partially mediating role of coping strategies. *Anxiety Stress Coping*, 37(2), 180-191. [doi: 10.1080/10615806.2023.2258805](https://doi.org/10.1080/10615806.2023.2258805)

mental illness, and substance misuse in adulthood.<sup>35,36</sup> See Figure 1.2<sup>37</sup> for an image of adverse childhood experiences.

35. University of Michigan Human Resources. (n.d.). *Section 1: What you need to know about mental health problems and substance misuse*.  
<https://hr.umich.edu/benefits-wellness/health-well-being/mhealthy/faculty-staff-well-being/mental-emotional-health/mental-emotional-health-classes-training-events/online-tutorial-supervisors/section-1-what-you-need-know-about-mental-health-problems-substance-misuse>
36. National Human Trafficking Training and Technical Assistance Center. (n.d.). *The original ACE study*. [https://nhttac.acf.hhs.gov/soar/eguide/stop/adverse\\_childhood\\_experiences](https://nhttac.acf.hhs.gov/soar/eguide/stop/adverse_childhood_experiences)
37. “ACEs.png” by unknown author for [Centers for Disease Control and Prevention](#) is licensed in the [Public Domain](#). Access for free at [https://www.cdc.gov/injury/pdfs/priority/ACEs-Strategic-Plan\\_Final\\_508.pdf](https://www.cdc.gov/injury/pdfs/priority/ACEs-Strategic-Plan_Final_508.pdf)

## Figure 1. What are Adverse Childhood Experiences?




Figure 1.2 Adverse Childhood Experiences (ACEs)

Individual trauma resulting from an event, series of events, or set of circumstances that is experienced as physically or emotionally harmful can have lasting adverse effects on the individual's functioning and mental, physical, social, emotional, or spiritual well-being.<sup>38</sup> Read more about ACEs and addressing individual trauma in the "[Introduction to Trauma-Informed Care](#)" section of this chapter.

► Visit the following PDF and take the [Adverse Childhood](#)

38. U.S. Department of Health & Human Services. (2020). *Strategic goal 3: Strengthen the economic and social well-being of Americans across the lifespan*. <https://www.hhs.gov/about/strategic-plan/strategic-goal-3/index.html>

- ▶ [Experiences Questionnaire for Adults](#) to better understand how previous experiences can affect one's well-being.

Current stressors such as relationship difficulties, the loss of a job, the birth of a child, a move, or prolonged problems at work can also be important contributory environmental factors.<sup>39</sup>

- ▶ Read more about stress in the “[Stress, Coping, and Crisis Intervention](#)” chapter.

## Biological Factors

Scientists believe the brain can have an imbalance of neurotransmitters, such as dopamine, acetylcholine, gamma-aminobutyric acid (GABA), norepinephrine, glutamate, and serotonin, resulting in changes in behavior, mood, and thought. While causes of fluctuations in brain chemicals aren't fully understood, contributing factors can include physical illness, hormonal changes, reactions to medication, substance misuse, diet, and stress.<sup>40</sup>

39. University of Michigan Human Resources. (n.d.). *Section 1: What you need to know about mental health problems and substance misuse*.  
<https://hr.umich.edu/benefits-wellness/health-well-being/mhealthy/faculty-staff-well-being/mental-emotional-health/mental-emotional-health-classes-training-events/online-tutorial-supervisors/section-1-what-you-need-know-about-mental-health-problems-substance-misuse>

40. University of Michigan Human Resources. (n.d.). *Section 1: What you need to*

- ▶ Read more about neurotransmitters and the central nervous system in the “[Psychotropic Medications](#)” chapter.

Some studies also suggest that depressive and bipolar disorders are accompanied by immune system dysregulation and inflammation.<sup>41</sup>

## Genetics

There appears to be a hereditary pattern to some mental illnesses. For example, individuals with major depressive disorder often have parents or other close relatives with the same illness. Research continues to investigate genes involved in specific disorders so that treatment can be effectively targeted to the individual.<sup>42</sup>

*know about mental health problems and substance misuse.*

<https://hr.umich.edu/benefits-wellness/health-well-being/mhealthy/faculty-staff-well-being/mental-emotional-health/mental-emotional-health-classes-training-events/online-tutorial-supervisors/section-1-what-you-need-know-about-mental-health-problems-substance-misuse>

41. Kraybill, O. (2019). Inflammation and mental health symptoms. *Psychology Today*. <https://www.psychologytoday.com/us/blog/expressive-trauma-integration/201905/inflammation-and-mental-health-symptoms>
42. University of Michigan Human Resources. (n.d.). *Section 1: What you need to know about mental health problems and substance misuse.* <https://hr.umich.edu/benefits-wellness/health-well-being/mhealthy/faculty-staff-well-being/mental-emotional-health/mental-emotional-health-classes-training-events/online-tutorial-supervisors/section-1-what-you-need-know-about-mental-health-problems-substance-misuse>



View the following YouTube video on causes of mental illness<sup>43</sup> : [Understanding the Biology of Mental Illness](https://youtu.be/LLUoG9Se77w)

## WHO Guidelines for Mental Health Care

Nurses protect and promote the mental well-being of all individuals and address the needs of individuals diagnosed with mental disorders. The World Health Organization (WHO) published the *Mental Health Intervention Guide* for nurses and primary health care providers that provides evidence-based guidance and tools for assessing and managing priority mental health and substance use disorders using clinical decision-making protocols. Essential principles for providing mental health care include promoting respect and dignity for the individuals seeking care; using effective communication skills to ensure care is provided in a nonjudgmental, non-stigmatizing, and supportive manner; and conducting comprehensive assessments.<sup>44</sup>

### Promoting Respect and Dignity

Individuals with mental health and substance use conditions should be treated with respect and dignity in a culturally appropriate manner. Health care professionals should promote the preferences of people with mental health and substance use disorders and support them, their family members, and their loved ones in an inclusive and equitable manner. The following box contains tips discussed in the WHO *Mental Health Intervention Guide*.

43. Alabama Department of Health. (2011, July 29). *Understanding the biology of mental illness* [Video]. YouTube. All rights reserved. <https://youtu.be/LLUoG9Se77w> .

44. World Health Organization. (2019). mhGAP intervention guide – version 2.0. <https://www.who.int/publications/i/item/9789241549790>

## Tips for Supporting Individuals with Mental Illness<sup>45</sup>

### Do:

- Treat people with mental health and substance use conditions with respect and dignity.
- Protect confidentiality.
- Ensure privacy.
- Advocate for individual preferences.
- Provide access to information and explain the proposed treatment risks and benefits in writing when possible.
- Make sure the person provides consent to treatment.
- Promote autonomy and independent living in the community.
- Provide access to decision-making options.

### Don't:

- Discriminate against people with mental health and substance use conditions.
- Make decisions for or on behalf of individuals.
- Use overly technical language when explaining proposed treatment.

In addition to the tips discussed in the WHO *Mental Health Intervention Guide*, nurses should approach individuals with compassion and provide emotional support in order to foster dignity and respect in their care of

<sup>45</sup>. World Health Organization. (2019). mhGAP intervention guide – version 2.0. <https://www.who.int/publications/i/item/9789241549790>

individuals with mental illness and substance use disorders. It is also important to provide holistic care, focusing on treating the unique needs of the individual and acknowledging their unique experience. Nurses should also refrain from imposing their personal or cultural values on their clients.

## Using Effective Communication Skills

Using effective communication skills promotes quality mental health care for all individuals. Tips for effective communication from the WHO *Mental Health Intervention Guide* are described in the following box.

### **Effective Communication Tips for Quality Mental Health Care<sup>46</sup>**

- **Create an environment that facilitates open communication.**
  - Meet the person in a private space, if possible.
  - Be welcoming and conduct introductions in a culturally appropriate manner.
  - Use culturally appropriate eye contact, body language, and facial expressions that facilitate trust.
  - Explain to adults that information discussed during the visit will be kept confidential. (Special considerations regarding “conditional confidentiality” and mandatory reporting for minors are discussed in the “[Childhood and](#)

46. World Health Organization. (2019). mhGAP intervention guide – version 2.0. <https://www.who.int/publications/i/item/9789241549790>



[Adolescence Disorders](#)” chapter.)

- If caregivers are present, suggest speaking with the client alone (except for young children) and obtain consent from the client to share clinical information.
- When interviewing a young person, consider having another person present who identifies with the same gender to maintain feelings of a psychologically safe environment.

- **Involve the person.**

- Include the person (and with their consent, their caregivers and family members) in all aspects of assessment and management as much as possible. This includes children, adolescents, adults, and older adults.

- **Start by listening.**

- Actively listen. Be empathic and sensitive. (Read more about active listening in the “[Therapeutic Communication and the Nurse-Client Relationship](#)” chapter.)
- Allow the person to speak without interruption.
- Be patient and ask for clarification of unclear information.
- For children, use language that they can understand. For example, ask about their interests (toys, friends, school, etc.).
- Convey that you understand their feelings and situation.

- **Be friendly, respectful, and nonjudgmental.**
  - Always be respectful.
  - Be nonjudgmental about an individual's behaviors and appearances.
  - Remain calm and professional.
- **Use good verbal communication skills.**
  - Use simple language. Be clear and concise. Avoid medical terminology only understood by health care professionals.
  - Use open-ended questions and other therapeutic communication techniques. (Read more about specific techniques in the ["Therapeutic Communication and the Nurse-Client Relationship"](#) chapter.) For example:
    - Use open-ended questions: "Tell me more about what happened?"
    - Summarize: "So, your brother pushed you off your bike and then laughed when you fell and started crying?"
    - Clarify: "To clarify, were you at home or a neighbor's house when this happened?"
  - Summarize and repeat key points at the end of the conversation.
  - Allow the person to ask questions about the information provided. For example, "What questions do you have about what we have

discussed today?”

- **Respond with sensitivity when people disclose traumatic experiences (e.g., sexual assault, violence, or self-harm).**
  - Thank the person for sharing this sensitive information.
  - Show extra sensitivity when discussing difficult topics.
  - Remind the person that what they tell you will only be shared with the immediate treatment team to provide the best possible care.
  - Acknowledge that it may have been difficult for the person to disclose the information.

## THERAPEUTIC RELATIONSHIP

In all nursing care, the therapeutic relationship with the client is essential. This is especially so in psychiatric care, where the therapeutic relationship is considered to be the foundation of client care and healing.<sup>47</sup> Although nurse generalists are not expected to perform advanced psychiatric interventions, all nurses are expected to engage in compassionate, supportive relationships

47. Ross, C. A., & Goldner, E. M. (2009). Stigma, negative attitudes and discrimination towards mental illness within the nursing profession: A review of the literature. *Journal of Psychiatric and Mental Health Nursing*, 16(6), 558-567. <https://doi.org/10.1111/j.1365-2850.2009.01399.x>

with their clients and use therapeutic communication as part of the “art of nursing.”<sup>48</sup>

The nurse-client relationship establishes trust and rapport with a specific purpose. It facilitates therapeutic communication and engages the client in decision-making regarding their plan of care. Read more about therapeutic communication and the nurse-client relationship in the “[Therapeutic Communication and the Nurse-Client Relationship](#)” chapter.

## Conducting Comprehensive Assessments

Clients undergo comprehensive assessments related to their disorder, including mental status examination, psychosocial assessment, physical examination, and review of laboratory results. Specific nursing assessments are further discussed in the “[Application of the Nursing Process in Mental Health Care](#)” chapter as well in each “Disorder” chapter. Persons with severe mental health and substance use disorders are two to three times more likely to die of preventable disease like infections and cardiovascular disorders, so it is also important for nurses to advocate for the medical treatment of existing physical disorders.<sup>49</sup>

▶ View the [WHO’s Mental Health Gap Intervention Guide](#).

48. Centers for Disease Control & Prevention. (2025). *Managing stress*.  
<https://www.cdc.gov/mental-health/living-with/index.html>

49. World Health Organization. (2019). mhGAP intervention guide – version 2.0.  
<https://www.who.int/publications/i/item/9789241549790>

## 1.3 Introduction to Trauma-Informed Care

Individuals may experience trauma during their lifetimes that can have a lasting impact on their mental health. **Trauma** results from an event, series of events, or set of circumstances that are experienced by an individual as physically or emotionally harmful and can have lasting adverse effects on the individual's functioning and physical, social, emotional, or spiritual well-being. Events may be human-made, such as war, terrorism, sexual abuse, violence, or medical trauma; or they can be the products of nature (e.g., flooding, hurricanes, and tornadoes).

It's not just the event itself that determines if it is traumatic, but the individual's experience of the event. Two people may be exposed to the same event or series of events but experience and interpret these events in vastly different ways. Various biopsychosocial and cultural factors influence an individual's immediate response and long-term reactions to trauma. For example, one's personal history, support system, coping mechanisms, personality traits, age and developmental stage, etc. all can influence trauma response. For most individuals, regardless of the severity of the trauma, the effects of trauma are met with **resilience**, defined as the ability to rise above circumstances or meet challenges with fortitude. Resilience includes the process of using available resources to negotiate hardship and/or the consequences of adverse events.<sup>1</sup>

Trauma can affect people of any culture, age, gender, or sexual orientation. Individuals may also experience trauma even if the event didn't happen to them. A traumatic experience can be a single event, a series of events, or adverse childhood experiences (ACEs). Review information about ACEs in the "[Mental Health and Mental Illness](#)" section of this chapter. There has been an increased focus on the ways in which trauma, psychological distress, quality of life, health, mental illness, and substance misuse are linked. For example, the

1. Center for Substance Abuse Treatment. (2014). *Trauma-informed care in behavioral health services*. <https://www.ncbi.nlm.nih.gov/books/NBK207201/>

terrorist attacks of September 11, 2001, the wars in Iraq and Afghanistan, disastrous hurricanes, and the COVID pandemic have moved traumatic experiences to the forefront of national consciousness. Trauma can affect individuals, families, groups, communities, specific cultures, and generations. It can overwhelm an individual's ability to cope; stimulate the "fight, flight, or freeze" stress reaction; and produce a sense of fear, vulnerability, and helplessness.<sup>2</sup>

► Read more information about the stress reaction in the "[Stress, Coping, and Crisis Intervention](#)" chapter.

For some people, reactions to a traumatic event are temporary, whereas other people have prolonged reactions to trauma with enduring mental health consequences, such as post-traumatic stress disorder, anxiety disorder, substance use disorder, mood disorder, or psychotic disorder. Others may exhibit culturally mediated physical symptoms referred to as **somatization**, in which psychological stress is expressed through physical concerns such as chronic headaches, pain, and stomachaches. Traumatic experiences can significantly impact how an individual functions in daily life and how they seek medical care.<sup>3</sup> Nurses must keep in mind to not interject their own experiences or perspectives because something minor to them may be major to the client.

Individuals may not recognize the significant effects of trauma or may avoid the topic altogether. Likewise, nurses may not ask questions that elicit a

2. Center for Substance Abuse Treatment. (2014). *Trauma-informed care in behavioral health services*. <https://www.ncbi.nlm.nih.gov/books/NBK207201/>
3. Center for Substance Abuse Treatment. (2014). *Trauma-informed care in behavioral health services*. <https://www.ncbi.nlm.nih.gov/books/NBK207201/>

client's history of trauma. They may feel unprepared to address trauma-related issues proactively or struggle to effectively address traumatic experiences within the constraints of their agency's policies.<sup>4</sup>

By recognizing that traumatic experiences are closely tied to mental health, nurses can provide trauma-informed care and promote resilience. **Trauma-informed care (TIC)** is a strengths-based framework that acknowledges the prevalence and impact of traumatic experiences in clinical practice. TIC emphasizes physical, psychological, and emotional safety for both survivors and health professionals and creates opportunities for survivors to rebuild a sense of control and empowerment (i.e., resilience).<sup>5</sup> TIC acknowledges that clients can be retraumatized by unexamined agency policies and practices and stresses the importance of providing client-centered care rather than applying general treatment approaches.<sup>6</sup>

TIC enhances therapeutic communication between the client and the nurse. It decreases risks associated with misunderstanding clients' reactions or underestimating the need for referrals for trauma-specific treatment. TIC encourages client-centered care by involving the client in setting goals and planning care that optimizes therapeutic outcomes and minimizes adverse effects. Clients are more likely to feel empowered, invested, and satisfied when they receive TIC.<sup>7</sup>

4. Center for Substance Abuse Treatment. (2014). *Trauma-informed care in behavioral health services*. <https://www.ncbi.nlm.nih.gov/books/NBK207201/>
5. Center for Substance Abuse Treatment. (2014). *Trauma-informed care in behavioral health services*. <https://www.ncbi.nlm.nih.gov/books/NBK207201/>
6. Center for Substance Abuse Treatment. (2014). *Trauma-informed care in behavioral health services*. <https://www.ncbi.nlm.nih.gov/books/NBK207201/>
7. Center for Substance Abuse Treatment. (2014). *Trauma-informed care in behavioral health services*. <https://www.ncbi.nlm.nih.gov/books/NBK207201/>

Implementing TIC requires specific training, but it begins with the first contact a person has with an agency. It requires all staff members (e.g., receptionists, direct client-care staff, nurses, supervisors, and administrators) to recognize that an individual's traumatic experiences can greatly influence their receptivity and engagement with health services. It can affect their interactions with staff, as well as their responsiveness to care plans and interventions.<sup>8</sup>

View the following YouTube video on trauma-informed approach to health care<sup>9</sup>: [Dr. Pickens Explains Trauma-Informed Approach](#)

▶ Read more details about trauma-informed care (TIC) in the "[Trauma, Abuse, and Violence](#)" chapter.

8. Center for Substance Abuse Treatment. (2014). *Trauma-informed care in behavioral health services*. <https://www.ncbi.nlm.nih.gov/books/NBK207201/>

9. Washington State Health Care Authority. (2019, June 24). *Dr. Pickens explains trauma-informed approach* [Video]. YouTube. All rights reserved. <https://youtu.be/6syEFO4OSFU>



## 1.4 Stigma

Despite a recent focus on mental health in the United States, there are still many harmful attitudes and misunderstandings surrounding mental illnesses that can cause people to ignore their mental health and make it more difficult for them to reach out for help.<sup>1,2</sup> **Stigma** has been defined as a cluster of negative attitudes and beliefs that motivates the general public to fear, reject, avoid, and discriminate against people with mental health disorders.<sup>3</sup>

It estimated that nearly two-thirds of people with diagnosable mental health disorders do not seek treatment due to the stigma of mental illness. The *U.S. Surgeon General's Report* in 1999 was a milestone report that sought to dispel the stigma of mental illness and its impact on those seeking care.<sup>4</sup> The National Alliance on Mental Illness (NAMI) seeks to improve the lives of those

1. Centers for Disease Control and Prevention. (2025). *Mental health stigma*. <https://www.cdc.gov/mental-health/stigma/>
2. Corrigan, P. W., & Watson, A. C. (2002). Understanding the impact of stigma on people with mental illness. *World Psychiatry: Official Journal of the World Psychiatric Association (WPA)*, 1(1), 16-20. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1489832/>
3. Substance Abuse and Mental Health Services Administration, Center for Mental Health Services. (2004). *School materials for a mental health friendly classroom: Training package* [PDF]. SAMHSA Pub. No. P040478M. Substance Abuse and Mental Health Services Administration. <https://www.prevention.org/Resources/2e2ba6ce-1e47-40ec-b49d-f7952b1f95a1/School%20Materials%20for%20a%20Mental%20Health%20Friendly%20Classroom.pdf>
4. Hegner, R. E. (2000). Dispelling the myths and stigma of mental illness: The surgeon general's report on mental health. *Issue Brief*, 754. <https://www.ncbi.nlm.nih.gov/books/NBK559750/>

with mental illness and reduce stigma through education, support, and advocacy. NAMI encourages people to share their stories to discredit stereotypes, break the silence, and document discrimination.<sup>5</sup>

- ▶ View SAMHSA's [Mental Health: Get the Facts](#) to review myths and facts about mental health

However, stigma and negative attitudes toward mental illness can still be found among nurses. A review of nursing literature by Ross and Golder explored negative attitudes and discrimination towards mental illness in the nursing profession. Several studies from a variety of countries indicated that health care professionals can be classified in three categories in relation to stigma, including “stigmatizers,” “the stigmatized,” and “de-stigmatizers.” “Stigmatizers” refer to nurses in medical settings with stereotypical attitudes towards clients with mental illnesses, psychiatric-mental health nurses, and/or psychiatry. Nurses classified as “the stigmatized” have mental health disorders or perceive stigma regarding their roles as psychiatric-mental health nurses. “De-stigmatizers” actively work to reduce stigma surrounding mental health disorders. The authors found that many nurses share commonly held stereotypical beliefs portrayed in the media. For example, clients with mental health disorders have been portrayed in the media as dangerous, unpredictable, violent, or bizarre, and these portrayals can cause fearful attitudes. Nurses in the studies were also concerned about inadvertently saying or doing “the wrong thing” or “setting off” uncontrollable behavior. Many nurses in general medical settings felt they lacked the skills to confidently and competently manage behavioral symptoms of clients with mental health disorders. The authors of the review reported that nursing

5. Abderholden, S. (2019). *It's not stigma, it's discrimination* [Blog]. National Alliance on Mental Illness. <https://www.nami.org/Blogs/NAMI-Blog/March-2019/It-s-Not-Stigma-It-s-Discrimination>

literature supports additional mental health education for entry-level nurses and practicing nurses to enhance their knowledge base on mental health.<sup>6</sup>

Nurses can reduce stigma and advocate for a client's needs and dignity by establishing a therapeutic nurse-client relationship. Nursing actions that can be taken include participating in or facilitating training on mental health awareness as well providing education and awareness about mental health disorders to clients, families, and communities. A therapeutic nurse-client relationship is essential in all settings, but it is especially important in mental health care where the therapeutic relationship is considered the foundation of client care and healing. Although nurse generalists are not expected to perform advanced psychiatric-mental health nursing interventions, all nurses are expected to engage in compassionate, supportive relationships with their clients.<sup>7</sup> In fact, in *Nursing: Scope and Standards of Practice (2021)*, the American Nurses Association states, "The nursing profession, rooted in caring relationships, demands that nurses reflect unconditional positive regard for every patient."<sup>8</sup>

6. Ross, C. A., & Goldner, E. M. (2009). Stigma, negative attitudes and discrimination towards mental illness within the nursing profession: A review of the literature. *Journal of Psychiatric and Mental Health Nursing*, 16(6), 558-567. <https://doi.org/10.1111/j.1365-2850.2009.01399.x>
7. Ross, C. A., & Goldner, E. M. (2009). Stigma, negative attitudes and discrimination towards mental illness within the nursing profession: A review of the literature. *Journal of Psychiatric and Mental Health Nursing*, 16(6), 558-567. <https://doi.org/10.1111/j.1365-2850.2009.01399.x>
8. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

- ▶ Read more about establishing a therapeutic nurse-client relationship in the “[Therapeutic Communication and the Nurse-Client Relationship](#)” chapter.

The first step in resolving stigma is to become aware of one’s personal beliefs. Take the survey in the following box to become more aware of your own attitudes and biases toward mental health care.

- ▶ Visit the following PDF document and take the [Personal Attitudes Survey](#) (page 8) from the Mental Health and High School Curriculum Guide by the Canadian Mental Health Association.

## 1.5 Boundaries

**Boundaries** are limits we set as individuals that define our levels of comfort when interacting with others. Personal boundaries include limits in physical, sexual, intellectual, emotional, material, and financial areas of our lives. Boundaries promote psychological safety in relationships at work, home, and with partners by protecting one's well-being and limiting the stress response. For example, if you come away from a meeting or conversation with someone feeling depleted, anxious, or tense, consider if your boundaries were crossed. A lack of healthy personal boundaries can lead to emotional and physical fatigue.<sup>1</sup>

Five major types of personal boundaries include the following<sup>2</sup>:

- **Physical:** Physical boundaries refer to one's personal space, privacy, and body. For example, some people are comfortable with public displays of affection (hugs, kisses, and hand-holding), while others prefer not to be touched in public.
- **Sexual:** Sexual boundaries refer to one's comfort level with intimacy and attention of a sexual nature. This can include sexual comments and touch, not just sexual acts.
- **Intellectual:** Intellectual boundaries refer to one's thoughts and beliefs. Intellectual boundaries are not respected when someone dismisses another person's ideas and opinions.
- **Emotional:** Emotional boundaries refer to a person's feelings. For example, an individual might not feel comfortable sharing feelings with

1. Pattemore, C. (2021). *10 ways to build and preserve better boundaries*. PsychCentral. <https://psychcentral.com/lib/10-way-to-build-and-preserve-better-boundaries#types>

2. Pattemore, C. (2021). *10 ways to build and preserve better boundaries*. PsychCentral. <https://psychcentral.com/lib/10-way-to-build-and-preserve-better-boundaries#types>

another person and prefer to share information gradually over time.

- **Material:** Physical and financial limits that we set around our personal possessions, space, and resources.
- **Financial:** Financial boundaries refer to how one prefers to spend or save money.

When caring for clients with mental health disorders, it is common to notice problems with setting appropriate boundaries. For example, a client experiencing bipolar disorder may exhibit a lack of financial and sexual boundaries. When they are experiencing a manic episode, they may spend thousands of dollars on a credit card over a weekend or have sexual relations with someone they just met. Another example of boundary issues is an individual with a depressive disorder who is treated poorly by their partner but does not leave or assert boundaries because they don't feel that they deserve to be treated any better.

Nurses must establish professional boundaries with all clients while also maintaining a respectful and caring relationship. Due to their professional role, nurses have authority and access to sensitive information that can make clients feel vulnerable. *A Nurses Guide to Professional Boundaries* by the National Council of State Boards of Nursing (NCSBN) states that it is the nurse's responsibility to use clinical judgment to determine and maintain professional boundaries. Nurses should limit self-disclosure of personal information and avoid situations where they have a personal or business relationship with a client. The difference between a caring nurse-client relationship and an over-involved relationship can be difficult to discern, especially in small communities or in community health nursing where roles may overlap. In these circumstances, it is important for the nurse to openly acknowledge their dual relationship and emphasize when they are performing in a professional capacity. Signs of inappropriate boundaries include the following<sup>3</sup>:

3. National Council of State Boards of Nursing. (2025). *Professional boundaries*.

- Self-disclosing intimate or personal issues with a client
- Engaging in behaviors that could be interpreted as flirting
- Keeping secrets with a client
- Believing you are the only one who truly understands or can help the client
- Spending more time than is necessary with a particular client
- Speaking poorly about colleagues or your employment setting with the client and/or their family
- Showing favoritism to a particular client
- Meeting a client in settings outside of work
- Contacting a client and/or their family members using social media
- Developing a social or romantic relationship with a client
- Sharing personal contact information
- Displaying excessive emotional attachment to the client
- Sharing confidential information about the client with unauthorized individuals.
- Making treatment decisions based on personal feelings rather than client needs

Establishing professional boundaries with clients diagnosed with mental health disorders is essential due to the vulnerability of the client population, as well as the behavioral manifestations of some disorders. For safety purposes, all healthcare workers should keep their last name, home address, personal telephone number, and social media handles private.



View the NCSBN video: "[Professional Boundaries in Nursing.](https://www.ncsbn.org/nursing-regulation/practice/professional-boundaries.page)"

<https://www.ncsbn.org/nursing-regulation/practice/professional-boundaries.page>

- ▶ Read [\*A Nurse's Guide to Professional Boundaries\*](#) PDF from the National Council of State Boards of Nursing (NCSBN)



## 1.6 Establishing Safety

Suicidal thoughts are a common symptom of mental health disorders and typically resolve with effective treatment. However, despite increased national attention on mental health care and efforts to reduce stigma, there has been no significant decline in suicide rates in the United States. Suicide remains a leading cause of death nationwide, highlighting the persistent gap between awareness and measurable outcomes.<sup>1</sup> Given this reality, establishing a safe and supportive environment for individuals experiencing suicidal ideation is not only a clinical priority but also a moral imperative. This includes consistent risk assessments, staff training, safety planning, and reducing access to lethal means, all of which are essential strategies for suicide prevention in health care and community settings.

### Recognizing Risk Factors for Suicide

Everyone can help prevent suicide by recognizing risk factors for suicide and intervening appropriately. Warning signs of suicide include client statements or nurse observations of the following<sup>2</sup>:

- Previous suicide attempt(s)
- History of depression or other mental illness
- Serious illness or chronic pain
- Criminal/legal problems
- Job/financial problems or loss
- Impulsive or aggressive tendencies

1. The Joint Commission. (2025). *Behavioral health care and human services: National patient safety goals*. <https://www.jointcommission.org/standards/national-patient-safety-goals/behavioral-health-care-national-patient-safety-goals/>

2. Centers for Disease Control and Prevention. (2024). *Risk and protective factors for suicide*. <https://www.cdc.gov/suicide/risk-factors/>

- Substance use
- Current or prior history of adverse childhood experiences
- Sense of hopelessness
- Violence victimization or perpetration

See Figure 1.3<sup>3</sup> for five action steps for anyone to take to prevent suicide in someone experiencing suicidal thoughts or ideations. Nurses can educate others to take the following steps if they believe someone may be in danger of suicide<sup>4</sup>:

- Call 911 if danger for self-harm seems imminent.
- Ask the person if they are thinking about killing themselves. Although asking this question can feel invasive, it is common for individuals with mental health problems to share their thoughts and plans regarding suicide. Asking them about suicide will not “put the idea into their head” or make it more likely that they will attempt suicide. In fact, by responding appropriately, you can help save their life by asking this question.
- Listen without judging and show you care.
- Stay with the person or make sure the person is in a private, secure place with another caring person until you can get further help.
- Remove any objects that could be used in a suicide attempt.
- Call or text 988 to reach the new nationwide Suicide and Crisis Line for a direct connection with compassionate, accessible care and support for anyone experiencing mental health-related distress.

3. “5actionsteps\_t.jpg” by unknown author for [National Institute of Mental Health](https://www.nimh.nih.gov/health/topics/suicide-prevention) is licensed in the [Public Domain](https://www.nimh.nih.gov/health/topics/suicide-prevention). Access for free at <https://www.nimh.nih.gov/health/topics/suicide-prevention>

4. Substance Abuse and Mental Health Services Administration. (2021). *Help prevent suicide*. <https://www.samhsa.gov/suicide>



Figure 1.3 Preventing Suicide

## Establishing a Safe Care Environment for Clients

In addition to encouraging these general action steps to prevent suicide, nurses can further prevent suicide by establishing a safe care environment. Establishing a safe care environment is a priority nursing intervention.

Reducing the risk for suicide is one of the National Patient Safety Goals for Behavioral Health Care established by The Joint Commission. New requirements were established in 2020 that apply to clients in psychiatric hospitals, clients being evaluated or treated for behavioral health conditions as their primary reason for care in general hospital units or critical access hospitals, and all clients who express **suicidal ideation** during their course of care.<sup>5</sup> These requirements include performing an environmental risk assessment, screening for suicidal ideation, assessing suicide risk, documenting risk of suicide, developing a safety plan, following evidence-based written policies and procedures, providing information on follow-up

5. The Joint Commission. (2025). *Behavioral health care and human services: National patient safety goals*. <https://www.jointcommission.org/standards/national-patient-safety-goals/behavioral-health-care-national-patient-safety-goals/>

care on discharge, and monitoring effectiveness of these actions in preventing suicides. These requirements are discussed in further detail in the following subsections.<sup>6</sup>

► Read more about suicide prevention at [Joint Commission's Suicide Prevention webpage](#).

## Perform Environmental Risk Assessment

An **environmental risk assessment** identifies physical environment features that could be used by clients to attempt suicide. Nurses implement actions to safeguard individuals identified at a high risk of suicide from environmental risks, such as continuous monitoring, routinely removing objects from rooms that could be used for self-harm, assessing objects brought into a facility by clients and visitors, and using safe transportation procedures when moving clients to other parts of the hospital.

In psychiatric hospitals and on psychiatric units within general hospitals, additional measures are taken to prevent suicide by hanging by removing anchor points, door hinges, and hooks. The Veteran's Health Administration showed that the use of a Mental Health Environment of Care Checklist to facilitate a thorough, systematic environmental assessment reduced the rate of suicide from 4.2 per 100,000 admissions to 0.74 per 100,000 admissions.<sup>7</sup>

6. The Joint Commission. (2019). *R3 report / Requirement, rationale, reference* [PDF]. [https://www.jointcommission.org/-/media/tjc/documents/standards/r3-reports/r3\\_18\\_suicide\\_prevention\\_hap\\_bhc\\_cah\\_11\\_4\\_19\\_final.pdf](https://www.jointcommission.org/-/media/tjc/documents/standards/r3-reports/r3_18_suicide_prevention_hap_bhc_cah_11_4_19_final.pdf)
7. The Joint Commission. (2019). *R3 report / Requirement, rationale, reference* [PDF]. [https://www.jointcommission.org/-/media/tjc/documents/standards/r3-reports/r3\\_18\\_suicide\\_prevention\\_hap\\_bhc\\_cah\\_11\\_4\\_19\\_final.pdf](https://www.jointcommission.org/-/media/tjc/documents/standards/r3-reports/r3_18_suicide_prevention_hap_bhc_cah_11_4_19_final.pdf)

- ▶ Read more about the VA Mental Health at the [Mental Health Environment of Care Checklist \(MHEOCC\) webpage](#)<sup>8</sup>

## Screen for Suicidal Ideation with a Validated Tool

Clients being evaluated or treated for mental health conditions often have suicidal ideation (i.e., thoughts of killing themselves). Additionally, clients being treated for medical conditions often have coexisting mental health disorders or psychosocial issues that can cause suicidal ideation. Therefore, all clients aged 12 and older admitted for acute health care should be screened for suicidal ideation with a validated tool. An example of a validated screening tool is the Patient Safety Screener.<sup>9,10</sup> View more information about the Patient Safety Screener tool in the following boxes.

- ▶ Visit the Suicide Prevention Resource Center's webpage to read

8. U.S. Department of Veterans Affairs. (2021). *VHA national center for patient safety*. <https://www.patientsafety.va.gov/professionals/onthejob/mentalhealth.asp>
9. The Joint Commission. (2019). *R3 report / Requirement, rationale, reference* [PDF]. [https://www.jointcommission.org/-/media/tjc/documents/standards/r3-reports/r3\\_18\\_suicide\\_prevention\\_hap\\_bhc\\_cah\\_11\\_4\\_19\\_final1.pdf](https://www.jointcommission.org/-/media/tjc/documents/standards/r3-reports/r3_18_suicide_prevention_hap_bhc_cah_11_4_19_final1.pdf)
10. Suicide Prevention Resource Center. (n.d.). *The patient safety screener: A brief tool to detect suicide risk*. <https://sprc.org/micro-learning/patientsafetysscreeener>

▶ more about the [The Patient Safety Screener: A Brief Tool to Detect Suicide Risk](#)<sup>11</sup>



View the following YouTube video on administering the Patient Safety Screener:<sup>12</sup> [The Patient Safety Screener 3](#)

## Assess Suicide Risk

An evidence-based **suicide risk assessment** should be completed on clients who have screened positive for suicidal ideation. Clients with suicidal ideation vary widely in their risk for a suicide attempt depending upon whether they have a plan, intent, or past history of attempts. An in-depth assessment of clients who screen positive for suicide risk must be completed to determine how to appropriately keep them safe from harm. Assessment for suicide risk includes asking about their suicidal ideation (i.e., thoughts of suicide), if they have a plan for committing suicide, their intent on completing the plan, previous suicidal or self-harm behaviors, risk factors, and protective factors.<sup>13</sup>

11. Suicide Prevention Resource Center. (n.d.). *The patient safety screener: A brief tool to detect suicide risk*. <https://sprc.org/micro-learning/the-patient-safety-screener-a-brief-tool-to-detect-suicide-risk/>

12. SPRC. (2018, April 26). *The Patient Safety Screener 3* [Video]. YouTube. All rights reserved. <https://youtu.be/4GmGiRBMnYc>

13. The Joint Commission. (2019). *R3 report / Requirement, rationale, reference*

When assessing for a suicide plan, notice if the plan is specific and the method they plan to use. The risk of acting on suicide thoughts increases with a specific plan. The risk also increases if the plan includes use of a lethal method that is accessible to the client.

An example of an evidence-based suicide risk assessment tool that anyone can use with anyone, anywhere is the Columbia Protocol, also known as the Columbia-Suicide Severity Rating Scale (C-SSRS). Read more about the C-SSRS in the following box. The C-SSRS uses a series of simple, plain-language questions that anyone can ask. The answers help identify if a person is at risk for suicide, assess the severity and immediacy of that risk, and gauge the level of support that the person needs. Examples of questions include the following<sup>14</sup>:

- Have you had thoughts of killing yourself?
- Have you thought about how you might do this?
- Have you done anything, started to do anything, or prepared to do anything to end your life?

### **Columbia Suicide Severity Rating Scale (C-SSRS)<sup>15</sup>**

[PDF]. [https://www.jointcommission.org/-/media/tjc/documents/standards/r3-reports/r3\\_18\\_suicide\\_prevention\\_hap\\_bhc\\_cah\\_11\\_4\\_19\\_final1.pdf](https://www.jointcommission.org/-/media/tjc/documents/standards/r3-reports/r3_18_suicide_prevention_hap_bhc_cah_11_4_19_final1.pdf)

14. The Columbia Lighthouse Project. (n.d.). *Identify risk. Prevent suicide.* <https://cssrs.columbia.edu/>

15. The Columbia Lighthouse Project. (n.d.). *Identify risk. Prevent suicide.* <https://cssrs.columbia.edu/>

- ▶ Read more about using the C-SSRS at [Columbia Lighthouse Project web site](#).

View the following YouTube video on C-SSRS<sup>16</sup> at [Saving Lives Worldwide – A Call to Action – The Columbia Lighthouse Project](#)

## Develop a Safety Plan

If a client is assessed as high risk for suicide, a safety plan (or crisis response plan) should be created in collaboration with the client. A **safety plan** is a prioritized written list of warning signs, coping strategies, and sources of support that clients can use before or during a suicidal crisis. The plan should be brief, in the client's own words, and easy to read. After the plan is developed, the nurse should problem solve with the client to identify barriers or obstacles to using the plan. It should be discussed where the client will keep the safety plan and how it will be located during a crisis.<sup>17,18</sup>

16. The Columbia Lighthouse Project. (2016, October 19). *Saving lives worldwide – A call to action – The Columbia Lighthouse Project* [Video]. YouTube. All rights reserved. <https://youtu.be/csPPsstf2og>

17. Western Interstate Commission for Higher Education. (2008). *Safety planning guide* [PDF Handout]. <https://www.sprc.org/sites/default/files/SafetyPlanningGuide%20Quick%20Guide%20for%20Clinicians.pdf>
18. Schuster, H., Jones, N., & Qadri, S. F. (2021). Safety planning: Why it is essential on the day of discharge from in-patient psychiatric hospitalization in reducing future risks of suicide. *Cureus*, 13(12), e20648. <https://doi.org/10.7759/cureus.20648>



- ▶ Read the [Safety Planning Guide PDF](#) by the Western Interstate Commission for Higher Education.<sup>19</sup>

## Document Level of Risk for Suicide

After suicide screening and suicide risk are assessed, it should be documented and communicated with the treatment team, along with the plan to keep the client safe. It is vital for all health care team members caring for the client to be aware of their level of risk and plans to reduce that risk as they provide care.<sup>20</sup> Nurses complete documentation regarding the level of a client's suicide risk and associated interventions every shift or more frequently as needed, depending upon the client status.

## Follow Written Policies and Procedures

Nurses must strictly follow agency policies and procedures addressing the care of individuals who are identified at risk for suicide to keep them safe. The Joint Commission's National Patient Safety Goal 15.01.01 Reduce the Risk for Suicide requires screening of all clients being evaluated for behavioral health conditions ages 12 and older. For example, in some suicide cases reported to The Joint Commission, the root cause was a failure of staff to adhere to

19. Western Interstate Commission for Higher Education. (2008). *Safety planning guide* [PDF]. <https://www.sprc.org/sites/default/files/SafetyPlanningGuide%20Quick%20Guide%20for%20Clinicians.pdf>

20. The Joint Commission. (2019). *R3 report / Requirement, rationale, reference* [PDF]. [https://www.jointcommission.org/-/media/tjc/documents/standards/r3-reports/r3\\_18\\_suicide\\_prevention\\_hap\\_bhc\\_cah\\_11\\_4\\_19\\_final1.pdf](https://www.jointcommission.org/-/media/tjc/documents/standards/r3-reports/r3_18_suicide_prevention_hap_bhc_cah_11_4_19_final1.pdf)

agency policies, such as one-to-one monitoring for clients identified as high risk for suicide.<sup>21</sup>

## Provide Information for Follow-Up Care on Discharge

Nurses should provide written information at discharge, in addition to the safety plan, regarding follow-up care to clients identified at risk for suicide and share it with their family members and loved ones as appropriate. Studies have shown that a client's risk for suicide is high after discharge from psychiatric inpatient or emergency department settings. Developing a safety plan with the client and providing the number of crisis call centers can decrease suicidal behavior after the client leaves the care of the organization.<sup>22</sup>

## Monitor Effectiveness of Suicide Prevention Interventions

The effectiveness of policies and protocols regarding suicide prevention should be evaluated on a periodic basis as part of overall quality improvement initiatives of the agency.<sup>23</sup> Research demonstrates implementation of the Zero Suicide Model results in lower suicidal behaviors.

21. The Joint Commission. (2019). *R3 report | Requirement, rationale, reference* [PDF]. [https://www.jointcommission.org/-/media/tjc/documents/standards/r3-reports/r3\\_18\\_suicide\\_prevention\\_hap\\_bhc\\_cah\\_11\\_4\\_19\\_final1.pdf](https://www.jointcommission.org/-/media/tjc/documents/standards/r3-reports/r3_18_suicide_prevention_hap_bhc_cah_11_4_19_final1.pdf)
22. The Joint Commission. (2019,). *R3 report | Requirement, rationale, reference* [PDF]. [https://www.jointcommission.org/-/media/tjc/documents/standards/r3-reports/r3\\_18\\_suicide\\_prevention\\_hap\\_bhc\\_cah\\_11\\_4\\_19\\_final1.pdf](https://www.jointcommission.org/-/media/tjc/documents/standards/r3-reports/r3_18_suicide_prevention_hap_bhc_cah_11_4_19_final1.pdf)
23. The Joint Commission. (2019). *R3 report | Requirement, rationale, reference* [PDF]. [https://www.jointcommission.org/-/media/tjc/documents/standards/r3-reports/r3\\_18\\_suicide\\_prevention\\_hap\\_bhc\\_cah\\_11\\_4\\_19\\_final1.pdf](https://www.jointcommission.org/-/media/tjc/documents/standards/r3-reports/r3_18_suicide_prevention_hap_bhc_cah_11_4_19_final1.pdf)

## Zero Suicide Toolkit<sup>24</sup>

- ▶ Read the American Psychiatric Association *Psych News Alert*, “‘Zero Suicide’ Practices at Mental Health Clinics Reduce Suicide Among Patients”.
- ▶ Visit the [Zero Suicide Toolkit webpage](#).



View the following WHO YouTube video<sup>25</sup> on preventing suicide by health care workers:



One or more interactive elements has been excluded from this version of the text. You can view them online here: <https://wtcs.pressbooks.pub/nursingmhcc/?p=73#oembed-1>

## Establishing a Safe Care Environment for Nurses and Other Health Care Team Members

The American Nurses Association states, “No staff nurse should have to deal

24. Zero Suicide. (n.d.). *Zero suicide toolkit*. <https://zerosuicide.edc.org/toolkit/zero-suicide-toolkitsm>

25. World Health Organization (WHO). (2019, October 8). *Preventing suicide: Information for health workers* [Video]. YouTube. Licensed in the [Public Domain](#). [https://youtu.be/Ey7n8SfwS\\_A](https://youtu.be/Ey7n8SfwS_A)

with violence in the workplace, whether from staff, clients, or visitors.”<sup>26</sup>

**Workplace violence** is the act or threat of violence, ranging from verbal abuse to physical assaults directed toward persons at work or on duty. The impact of workplace violence can range from psychological issues to physical injury or even death. Violence can occur in any workplace and among any type of worker, but the risk for nonfatal violence resulting in days away from work is greatest for health care workers.<sup>27</sup> Research indicates the rate of physical assaults on nurses is 13.2 per 100 nurses per year, and 25% of psychiatric nurses experienced disabling injuries from client assault. Many experts believe these figures are under represented and that most incidents of violence go unreported.<sup>28</sup> See Figure 1.4<sup>29</sup> for an illustration of safety first.

26. American Nurses Association. (n.d.). *Safety on the job*.

<https://www.nursingworld.org/practice-policy/work-environment/health-safety/safety-on-the-job/>

27. Centers for Disease Control and Prevention. (2024). *About workplace violence*.

[https://www.cdc.gov/niosh/violence/about/?CDC\\_AAref\\_Val=https://www.cdc.gov/niosh/topics/violence/default.html](https://www.cdc.gov/niosh/violence/about/?CDC_AAref_Val=https://www.cdc.gov/niosh/topics/violence/default.html)

28. Centers for Disease Control and Prevention. (2024). *About workplace violence*.

[https://www.cdc.gov/niosh/topics/violence/training\\_nurses.html](https://www.cdc.gov/niosh/topics/violence/training_nurses.html)

29. “Safety-First–Arvin61r58.png” by unknown author at freesvg.org is licensed under [CC0 1.0](https://creativecommons.org/licenses/by/4.0/). Access for free at <https://freesvg.org/safety-first>



Figure 1.4 Safety First

Safety strategies for nurses providing client care include the following<sup>30</sup>:

- **Dress for Safety**
  - Tuck away long hair so that it can't be grabbed
  - Avoid piercings or necklaces that can be pulled
  - Avoid overly tight clothing that can restrict movement or overly loose clothing or scarves that can be caught
  - Use breakaway safety lanyards for glasses, keys, or name tags
  - Do not wear your stethoscope around your neck
- **Be Aware of Your Work Environment**
  - When in a room with a client or visitor who is demonstrating warning signs of escalation, position yourself between the door and the client so you can exit quickly if needed
  - Note exits and emergency phone numbers, especially if you float to

30. Centers for Disease Control and Prevention. (2024). About *workplace violence*. [https://www.cdc.gov/niosh/topics/violence/training\\_nurses.html](https://www.cdc.gov/niosh/topics/violence/training_nurses.html)

other areas

- Recognize that confusion, background noises, and crowding can increase clients' stress levels
- Be aware that mealtimes, shift changes, and transporting clients are times of increased disruptive behaviors

- **Be Attuned to Client Behaviors**

- Most violent behavior is preceded by warning signs, including verbal cues and nonverbal cues. The greater the number of cues, the greater the risk for violence. Be aware of these verbal and nonverbal cues indicating a client's potential escalation to violence:

- **Verbal Cues**

- Speaking loudly or yelling
    - Swearing
    - Using a threatening tone of voice

- **Nonverbal and Behavioral Cues**

- Evidence of confusion or disorientation
    - Irritability or easily angered
    - Boisterous behavior (i.e., overly loud, shouting, slamming doors)
    - Disheveled physical appearance (i.e., neglected hygiene)
    - Holding arms tightly across chest
    - Clenching fists
    - Heavy breathing
    - Pacing or agitated, restlessness
    - Looking terrified (signifying fear and high anxiety)
    - Staring with a fixed look
    - Holding oneself in an aggressive or threatening posture
    - Throwing objects
    - Exhibiting sudden changes in behavior or signs of being under the influence of a substance

- **Use Violence Risk Assessment Tools**

- Use risk assessment tools to evaluate individuals for potential violence, enabling all health care providers to share a common frame of reference and understanding. This minimizes the possibility that communications regarding a person's potential for violence will be misinterpreted. These tools can be used as an initial assessment upon admission to determine potential risk for violence and repeated daily to assist in predicting imminent violent behavior within the next 24 hours. See sample risk assessment tools in the box at the end of this section.

- **Be Attuned to Your Own Responses**

- Be aware of your own feelings, responses, and sensitivities and pay attention to your instincts. For example, your “fight or flight” response can be an early warning sign of impending danger to get help or get out.
- Be aware of how you express yourself and how others respond to you. Those who know you well may respond differently than strangers. Effective therapeutic communication skills are an essential tool in preventing violence.
- Use self-awareness and acknowledge if you have a personal history of abuse, trauma, or adverse childhood experiences (ACEs) that can affect how you respond to situations.
- If coworkers are engaging in abusive behaviors, consider if you are exhibiting similar behaviors.
- Be aware that fatigue can diminish your alertness and your ability to respond appropriately to a challenging situation.

- **Check Your Cultural Biases**


- A key aspect of self-awareness is recognizing how our own particular cultural heritage, values, and belief systems affect how we respond to our clients and coworkers and how they, in turn, respond to us.

### Sample Violence Risk Assessment Tools from the CDC:

- [▶ Triage Tool PDF](#)
- [▶ Indicator for Violent Behavior PDF](#)
- [▶ Assault and Homicidal Danger Assessment Tool PDF](#)

If travelling to a home setting as a home health nurse, additional safety strategies are as follows<sup>31</sup>:

- Review agency files to confirm that a background check was done on a client regarding any history of violence or crime, drug or alcohol abuse, and mental health diagnoses. Also, check to see if a client's family member has a record of violence or arrest.
- If entering a situation assessed as potentially dangerous, you should be accompanied by a team member who has training in de-escalation and crisis intervention.
- Always carry a charged cell phone.
- Make sure someone always knows where you are.
- Have a code word to use with your office or coworkers to let them know you're in trouble if you can't call the police.

The CDC offers a free, online course called Workplace  
 Violence Prevention for Nurses to better understand the scope and nature of violence in the workplace. Access the

31. Centers for Disease Control and Prevention. (2024). About *workplace violence*. [https://www.cdc.gov/niosh/topics/violence/training\\_nurses.html](https://www.cdc.gov/niosh/topics/violence/training_nurses.html)





free CDC course on workplace violence with nurse videos  
at the [Workplace Violence Prevention for Nurses webpage](#)

- ▶ Read more about crisis interventions in the [Stress, Coping, and Crisis Intervention](#) chapter.

## 1.7 Psychiatric-Mental Health Nursing

### What is Psychiatric-Mental Health Nursing?

Registered nurses (RNs) in a variety of settings provide care for clients with medical illnesses who may also be experiencing concurrent mental health disorders. Nurses who specialize in psychiatric-mental health nursing promote clients' well-being through prevention strategies and client education, while also using the nursing process to provide care for clients with mental health and substance use disorders.<sup>1</sup> According to the American Psychiatric Nurses Association, psychiatric-mental health nurse specialists perform the following activities<sup>2</sup>:

- Partner with individuals to achieve their recovery goals
- Provide health promotion and maintenance
- Conduct intake screening, evaluation, and triage
- Provide case management
- Teach self-care activities
- Administer and monitor mental functioning and behavior treatment regimens
- Practice crisis intervention and stabilization
- Engage in psychiatric rehabilitation and intervention
- Educate clients, families, and communities
- Coordinate care
- Work within interdisciplinary teams

Within the specialty of psychiatric-mental health nursing, there is an

1. American Psychiatric Nurses Association. (n.d.). *About psychiatric-mental health nursing*. <https://www.apna.org/about-psychiatric-nursing/>

2. American Psychiatric Nurses Association. (n.d.). *About psychiatric-mental health nursing*. <https://www.apna.org/about-psychiatric-nursing/>

opportunity to become board certified. Eligibility requirements include a bachelor's degree, two years of full-time work, 30 hours of continuing education, and passing a certification exam. The nurse earns the credential of PMH-BC (Psychiatric-Mental Health-Board Certified) or RN-BC.

Psychiatric-mental health advanced practice registered nurses (PMH-APRN) and nurse practitioners (PMHNP-BC) are registered nurses with a Master of Science in Nursing (MSN) or Doctor of Nursing Practice (DNP) degree in psychiatric nursing. PMH-APRNs perform the following activities:

- Provide individual, group, couples, and/or family psychotherapy
- Prescribe medication for acute and chronic illnesses
- Conduct comprehensive assessments
- Provide clinical supervision
- Diagnose, treat, and manage chronic or acute illness
- Provide integrative therapy interventions
- Order, perform, and interpret lab tests and other diagnostic studies
- Provide preventative care, including screening
- Develop policies for programs and systems
- Make referrals for health problems outside their scope of practice
- Perform procedures

## Standards of Psychiatric-Mental Health Nursing

The American Psychiatric Nurses Association establishes standards of practice in psychiatric-mental health nursing that are built on the ANA Scope and Standards of Practice (2021). These standards are published in the *Psychiatric-Mental Health Nursing: Scope and Standards of Practice* document.<sup>3</sup> The standards are very similar to the ANA Scope and Standards of

3. American Nurses Association, American Psychiatric Nurses Association, and International Society of Psychiatric-Mental Health Nurses. (2014). *Psychiatric-mental health nursing: Scope and standards of practice* (2nd ed.). Nursebooks.org

Practice, with additional activities included in the *Intervention* standard of care. These interventions will be further discussed in the “[Implementation](#)” section of the “Application of the Nursing Process in Mental Health Care” chapter.

- ▶ Read the [About Psychiatric-Mental Health Nursing](#) webpage to learn more about the American Psychiatric Nursing Association.

There are specific legal and ethical considerations that apply to caring for clients with mental illness. See the “[Legal and Ethical Considerations in Mental Health Care](#)” chapter for further information.

## Treatment Settings

Treatment settings for mental health illnesses vary based on the severity and complexity of the condition, as well as the resources available. These settings can be broadly categorized into several types:

### Primary Care Settings

Mild to moderate mental health conditions are often managed in primary care settings. This includes the integration of mental health services within primary care practices, where general practitioners may provide initial assessments, prescribe medications, and offer brief counseling.

### Outpatient Services

These settings are suitable for individuals who require regular, but not intensive, mental health care. Outpatient clinics provide services such as psychotherapy, medication management, and follow-up visits. Community mental health teams often operate in these settings to offer coordinated care.

Clients often initially visit their primary care provider when concerned about their mental health. If a client has a more severe disorder, they are typically referred to specialized psychiatric care providers such as psychiatrists, psychiatric-mental health advanced practice registered nurses/nurse practitioners, psychologists, social workers, counselors, or other licensed therapists.

## Day Treatment Programs

These programs provide a higher level of care than outpatient services but do not require overnight stays. They are designed for individuals who need intensive treatment during the day but can return home in the evening.

## Inpatient Care Settings

Clients with acute mental health symptoms, or those who are at-risk for hurting themselves or others, may be hospitalized. They are often initially seen in the emergency department for emergency psychiatric care. Clients may seek voluntary admission, or in some situations, may be involuntarily admitted after referral for emergency evaluation by law enforcement, schools, friends, or family members. This setting is for individuals with severe mental health conditions that require 24-hour care and supervision. Inpatient treatment focuses on crisis stabilization, safety, and rapid discharge planning. It is important to note that individuals with suicidal ideation who are not medically stable may be admitted to medical surgical units for care. Specialized units may exist for different populations, such as children, adolescents, and those with dual diagnoses. Read more about involuntary admissions in the “[Patient Rights](#)” section of the “Legal and Ethical Considerations in Mental Health Care” chapter.

Acute-care psychiatric units in general hospitals are typically locked units on a separate floor of the hospital with the purpose of maintaining environmental safety for its clients. State-operated psychiatric hospitals serve clients who have chronic serious mental illness. They also provide court-related care for

criminal cases where the client was found “not guilty by reason of insanity.” This judgment means the client was deemed to be so mentally ill when they committed a crime that they cannot be held responsible for the act, but instead require treatment.<sup>4</sup>

## Community-Based Services

Community-based services include mobile crisis response teams, intensive in-home therapies, and therapeutic foster care. These services aim to provide support in the least restrictive environment possible, helping individuals remain in their communities while receiving necessary care.

## Telepsychiatry

This modality extends mental health services to remote or underserved areas, providing access to psychiatric care through telecommunication technologies. It is particularly useful in community settings such as schools and correctional facilities.

4. Halter, M. J. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

## 1.8 Learning Activities



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=86#h5p-2>

1



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=86#h5p-1>

2



*An interactive H5P element has been excluded from this version of the text. You can view it*

1. “MH Therapeutic Foundations Glossary Cards” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
2. “MH Therapeutic Question Set 1” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

online here:

<https://wtcs.pressbooks.pub/nursingmhcc/?p=86#h5p-3>

3

- ▶ Test your clinical judgment with a NCLEX Next-Generation style case study: [Chapter 1, Case Study 1](#)<sup>4</sup>



3. “MH Therapeutic Question Set 3” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

4. “MH Therapeutic Foundations Case Study” by Kellea Ewen is licensed under [CC BY-NC 4.0](#)



## I Glossary

**Adverse childhood experiences (ACEs):** Traumatic circumstances experienced during childhood such as abuse, neglect, or growing up in a household with violence, mental illness, substance use, incarceration, or divorce.

**Boundaries:** Limits that we set as individuals that define our levels of comfort when interacting with others. Personal boundaries include limits in physical, sexual, intellectual, emotional, sexual, and financial areas of our lives.

**Culture:** A complex context of knowledge, beliefs, and behaviors that provide structure to specific ethnic, racial, religious, geographic, or social groups.

**Cultural humility:** A lifelong process of self-reflection and self-critique whereby individuals actively engage in learning about different cultural perspectives while acknowledging the limitations of their own knowledge.

**Deviance:** Behavior that violates social norms or cultural expectations because one's culture determines what is "normal."

**Distress:** Psychological and/or physical pain.

**Dysfunction:** Disturbances in a person's thinking, emotional regulation, or behavior that reflects significant dysfunction in psychological, biological, or developmental processes underlying mental functioning.

**Environmental risk assessment:** Identification of physical environment features that could be used to attempt suicide in clients identified as at a high risk for suicide.

**Health:** A state of complete physical, mental, and social well-being and not merely the absence of disease or infirmity.

**Impairment:** A limited ability to engage in activities of daily living (i.e., they cannot maintain personal hygiene, prepare meals, or pay bills) or participate in social events, work, or school.

**Major life activities:** Activities of daily living such as caring for oneself, performing manual tasks, seeing, hearing, eating, sleeping, walking, standing, lifting, bending, speaking, breathing, learning, reading, concentrating, thinking, communicating, and working.<sup>1</sup>

**Mental health:** A state of well-being in which an individual realizes their own abilities, copes with the normal stresses of life, works productively, and contributes to their community.<sup>2</sup>

**Mental health continuum:** A continuum of mental health, ranging from well-being to emotional problems to mental illness.

**Mental illness:** A health condition involving changes in emotion, thinking, or behavior (or a combination of these) associated with emotional distress and problems functioning in social, work, or family activities.<sup>3</sup>

**Recovery:** A process of change through which individuals improve their health and wellness, live a self-directed life, and strive to reach their full potential.<sup>4</sup>

1. Office of Federal Contract Compliance Programs. (2009). *ADA Amendments Act of 2008 frequently asked questions*. U.S. Department of Labor. <https://www.dol.gov/agencies/ofccp/faqs/americans-with-disabilities-act-amendments>
2. World Health Organization. (2018). *Mental health: Strengthening our response*. <https://www.who.int/news-room/fact-sheets/detail/mental-health-strengthening-our-response>
3. American Psychiatric Association. (n.d.). *What is mental illness?* <https://www.psychiatry.org/patients-families/what-is-mental-illness>
4. Center for Substance Abuse Treatment. (2014). *Trauma-informed care in behavioral health services*. <https://www.ncbi.nlm.nih.gov/books/NBK207201/>

**Resilience:** The ability to rise above circumstances or meet challenges with fortitude.<sup>5</sup>

**Safety plan:** A prioritized written list of coping strategies and sources of support that clients can use before or during a suicidal crisis. The plan should be brief, in the client's own words, and easy to read. After the plan is developed, the nurse should problem solve with the client to identify barriers or obstacles to using the plan. It should be discussed where the client will keep the safety plan and how it will be located during a crisis.

**Serious mental illness:** Mental illness that causes disabling functional impairment that substantially interferes with one or more major life activities. Examples of serious mental illnesses that commonly interfere with major life activities include major depressive disorder, schizophrenia, and bipolar disorder.<sup>6</sup>

**Social norms:** Stated and unstated rules of an individual's society.

**Stigma:** A cluster of negative attitudes and beliefs that motivates the general public to fear, reject, avoid, and discriminate against people with mental health disorders.

**Suicidal ideation:** Thoughts of killing oneself.

**Suicide risk assessment:** Identifying the risk of a client dying by suicide by assessing suicidal ideation, plan, intent, suicidal or self-harm behaviors, risk factors, and protective factors.

**Trauma:** An event, series of events, or set of circumstances that is experienced by an individual as physically or emotionally harmful and can have lasting

5. Center for Substance Abuse Treatment. (2014). *Trauma-informed care in behavioral health services*. <https://www.ncbi.nlm.nih.gov/books/NBK207201/>

6. American Psychiatric Association. (n.d.). *What is mental illness?* <https://www.psychiatry.org/patients-families/what-is-mental-illness>

adverse effects on the individual's functioning and physical, social, emotional, or spiritual well-being.

**Trauma-informed care (TIC):** A strengths-based framework that acknowledges the prevalence and impact of traumatic experiences in clinical practice. TIC emphasizes physical, psychological, and emotional safety for both survivors and health professionals and creates opportunities for survivors to rebuild a sense of control and empowerment referred to as resilience.<sup>7</sup>

**Well-being:** The “healthy” range of the mental health continuum where individuals are experiencing a state of good mental and emotional health.

**Workplace violence:** The act or threat of violence, ranging from verbal abuse to physical assaults, directed toward persons at work or on duty.

**World Health Organization Disability Assessment Scale (WHODAS):** A generic assessment instrument that provides a standardized method for measuring health and disability across cultures.

7. Center for Substance Abuse Treatment. (2014). *Trauma-informed care in behavioral health services*. <https://www.ncbi.nlm.nih.gov/books/NBK207201/>

PART II

CHAPTER 2 THERAPEUTIC COMMUNICATION AND THE  
NURSE-CLIENT RELATIONSHIP



## 2.1 Introduction

### Learning Objectives

- Identify advanced therapeutic communication techniques
- Describe barriers to effective therapeutic communication
- Explore guidelines for effective communication during telehealth

Nurses engage in compassionate, supportive, professional relationships with their clients as part of the “art of nursing.”<sup>1</sup> This chapter will review the nurse-client relationship, therapeutic communication, and motivational interviewing. It will also introduce guidelines for effective communication during telehealth.

1. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

## 2.2 Basic Concepts of Communication

### Communication Standard of Professional Performance

The Standard of Professional Performance for *Communication* established by the American Nurses Association (ANA) is defined as, “The registered nurse communicates effectively in all areas of professional practice.”<sup>1</sup> See the following box for the competencies associated with the *Communication* standard.

#### **ANA’s Communication Competencies**

The registered nurse:

- Assesses one’s own communication skills and effectiveness.
- Demonstrates cultural humility, professionalism, and respect when communicating.
- Assesses communication ability, health literacy, resources, and preferences of health care consumers to inform the interprofessional team and others.
- Uses language translation resources to ensure effective communication.
- Incorporates appropriate alternative strategies to communicate effectively with health care consumers who have visual, speech, language, or communication difficulties.
- Uses communication styles and methods that demonstrate caring, respect, active listening, authenticity, and trust.
- Conveys accurate information to health care consumers,

1. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.



families, community stakeholders, and members of the interprofessional team.

- Advocates for the health care consumer and their preferences and choices when care processes and decisions do not appear to be in the best interest of the health care consumer.
- Maintains communication with interprofessional team members and others to facilitate safe transitions and continuity in care delivery.
- Confirms with the recipient if the communication was heard and if the recipient understands the message.
- Contributes the nursing perspective in interactions and discussions with the interprofessional team and other stakeholders.
- Promotes safety in the care or practice environment by disclosing and reporting concerns related to potential or actual hazards or deviations from the standard of care.
- Demonstrates continuous improvement of communication skills.

- ▶ Review basic communication concepts for nurses in the “[Communication](#)” chapter in *Open RN Nursing Fundamentals*.

## Nurse-Client Relationship

Establishment of the therapeutic nurse-client relationship is vital in nursing care. Nurses engage in compassionate, supportive, professional relationships

with their clients as part of the “art of nursing.”<sup>2</sup> This is especially true in psychiatric care, where the therapeutic relationship is considered to be the foundation of client care and healing.<sup>3</sup> The **nurse-client relationship** establishes trust and rapport with a specific purpose; it facilitates therapeutic communication and engages the client in decision-making regarding their plan of care.

Therapeutic nurse-client relationships vary in depth, length, and focus, and are influenced based on the care setting. Brief therapeutic encounters might last only a few minutes and focus on the client’s immediate needs, safety, current feelings, or behaviors. For example, in the emergency department setting, a nurse may therapeutically communicate with a client in crisis who recently experienced a situational trauma. During longer periods of time, such as inpatient care, nurses work with clients in setting short-term goals and outcomes that are documented in the nursing care plan and evaluated regularly. In long-term care settings, such as residential facilities, the therapeutic nurse-client relationship may last several months and include frequent interactions focusing on behavior modification.

► Read more about crisis and crisis intervention in the “[Stress, Coping, and Crisis Intervention](#)” chapter.

2. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.
3. Ross, C. A., & Goldner, E. M. (2009). Stigma, negative attitudes and discrimination towards mental illness within the nursing profession: A review of the literature. *Journal of Psychiatric and Mental Health Nursing*, 16(6), 558-567. <https://doi.org/10.1111/j.1365-2850.2009.01399.x>

# Phases of Development of a Therapeutic Relationship

The nurse-client relationship goes through three phases. A well-known nurse theorist named Hildegard Peplau described these three phases as orientation, working, and termination.<sup>4</sup>

## ORIENTATION PHASE

During the brief orientation phase, clients may realize they need assistance as they adjust to their current status. Simultaneously, nurses introduce themselves and begin to obtain essential information about clients as individuals with unique needs, values, beliefs, and priorities. During this brief phase, trust is established, and rapport begins to develop between the client and the nurse. Nurses ensure privacy when talking with the client and providing care and respect the client's values, beliefs, and personal boundaries.

A common framework used for introductions during client care is AIDET, a mnemonic for Acknowledge, Introduce, Duration, Explanation, and Thank You.

- **Acknowledge:** Greet the client by the name documented in their medical record. Make eye contact, smile, and acknowledge any family or friends in the room. Ask the client their preferred way of being addressed. For example, knock gently and enter the room calmly. Approach with a non-threatening demeanor and open body language. "Hello, Alex. I see you've just arrived—welcome. I know being here might feel overwhelming at first, but I'm really glad you're here. May I ask how you'd like to be addressed and what pronouns you use?"

4. Hagerty, T. A., Samuels, W., Norcini-Pala, A., & Gigliotti, E. (2018). Peplau's theory of interpersonal relations: An alternate factor structure for patient experience data? *Nursing Science Quarterly*, 30(2), 160-167. <https://dx.doi.org/10.1177%2F0894318417693286>

- **Introduce:** Introduce yourself by your name and role. Build trust by being transparent and personable. For example, “My name is Jordan, and I’m the registered nurse assigned to support you today. My role is to help make sure you feel safe, listened to, and supported throughout your stay.”
- **Duration:** Estimate a timeline for how long it will take to complete the task you are doing. Provide structure without adding pressure, acknowledging the client’s emotional state. For example, “Right now, I’d like to spend about 15–20 minutes getting to know you a little better and going over a few admission questions. We’ll talk about how you’re feeling, your goals for being here, and how we can support you best.”
- **Explanation:** Explain step by step what to expect next and answer questions. Be clear and gentle, inviting collaboration. For example, “We’ll talk in a private space so you can share as much or as little as you’re comfortable with. I’ll ask about what brought you in, what’s been going on recently, and your support system. You’re welcome to take breaks or let me know if anything feels too much.”
- **Thank You:** At the end of the encounter, thank the client and ask if anything is needed before you leave. Affirm the client’s effort and courage in showing up. In an acute or long-term care setting, ensure the call light is within reach and the client knows how to use it. If family members are present, thank them for being there to support the client as appropriate. For example, ““Thank you for sharing your time with me today—it takes real strength to take this step. Before I go, is there anything I can get for you? I’ll make sure you know how to reach me if you need anything.”

## WORKING PHASE

The majority of a nurse’s time with a client is spent in the working phase of the therapeutic relationship. This is where trust deepens, and meaningful progress toward health and well-being begins. During this phase, nurses use active listening to understand the client’s needs, values, and goals, starting with open-ended questions to explore the reason for seeking care.

Assessment findings are used to develop or refine the nursing plan of care and to guide client education that is tailored to the individual. For example, after using the AIDET framework to greet and orient a new client like Alex, a

nurse might say: “Alex, thank you again for sharing some of your story with me earlier. I know being here is a big step. Based on what you’ve told me so far, I’d like to talk through a few ideas together that could help you feel supported during your stay.”

If a care plan has already been initiated, the nurse uses this phase to implement targeted interventions aimed at both short-term relief and long-term outcomes. As the relationship strengthens, clients begin to see nurses as educators, counselors, and trusted care providers—not just clinical staff. In Alex’s case, the nurse might continue: “Can you tell me more about what led you to seek care today and what you’re hoping will be different by the time you leave? Your voice really matters here—we’ll create a plan together that feels realistic and helpful for you.”

This collaborative approach is essential. Nurses use therapeutic communication techniques—such as reflection, clarification, silence, and motivational interviewing—to help clients become more aware of their thoughts, emotions, and options. Through nonjudgmental feedback, the nurse supports the client in exploring and clarifying their goals, strengths, and coping strategies.<sup>5</sup> For example, the nurse might gently reflect: “It sounds like you’ve been trying to manage everything on your own, and it’s been exhausting. It’s okay to need help—being here is a brave step.” In these moments, the nurse not only delivers clinical care but also fosters emotional healing and client empowerment. The working phase is where healing conversations and actions come together, providing a foundation for meaningful change and lasting outcomes.

5. Hagerty, T. A., Samuels, W., Norcini-Pala, A., & Gigliotti, E. (2018). Peplau’s theory of interpersonal relations: An alternate factor structure for patient experience data? *Nursing Science Quarterly*, 30(2), 160-167. <https://dx.doi.org/10.1177%2F0894318417693286>

## TERMINATION PHASE

The termination phase marks the conclusion of the therapeutic nurse-client relationship. It typically occurs at the end of a shift, during a transfer, or at discharge from care. When the working phase has been effective, the client's goals have been addressed collaboratively through the efforts of the client, nurse, and interprofessional team. This final phase provides an opportunity for reflection, reinforcement, and closure—essential for empowering the client to move forward independently with confidence.

In Alex's case, after several days of support and progress, the nurse prepares to end the therapeutic relationship at the end of the shift and as Alex nears discharge. The nurse approaches the conversation with sensitivity and structure. "Alex, I want to take a few minutes to talk with you before I end my shift. It's been a privilege working with you this week, and I want to thank you for the trust you've placed in me."

Because clients may sometimes attempt to return to the working phase to delay separation, the nurse remains aware and gently holds space for any emotions that arise. "I know transitions can bring up a lot of feelings. It's normal to feel a mix of relief, nervousness, or even sadness. Would it be helpful to talk through any of that together before I go?"

The nurse encourages Alex to reflect on progress and reinforces the client's strengths. "You've worked incredibly hard to identify what matters most to you, and you've been honest and brave every step of the way. Remember when we first talked and you weren't sure if you even belonged here? Now look at how far you've come—recognizing your needs, setting goals, and planning next steps."

Together, they review discharge goals and available community supports. "Before you leave, we've set up a follow-up appointment with the outpatient team for next Tuesday. I've also included a few community support groups you might want to explore. These resources are here to walk with you as you continue building on the work you started here."

The nurse concludes the relationship with warmth, professionalism, and

closure. “Alex, thank you again for allowing me to be part of your care. I truly believe in your ability to keep moving forward. If you ever need support again, know that it’s okay to reach out.”

This approach to the termination phase helps ensure that the client leaves with a clear sense of accomplishment, continuity of care, and confidence in their ability to self-advocate and engage support systems. It also models a healthy ending to a professional relationship—an important skill for both clients and nurses.

## 2.3 Therapeutic Communication

Therapeutic communication has roots going back to Florence Nightingale, who insisted on the importance of building trusting relationships with individuals. She taught that therapeutic healing resulted from nurses' presence with clients.<sup>1</sup> Since then, several professional nursing associations have highlighted therapeutic communication as one of the most vital elements in nursing. In psychiatric settings, effective communication has been shown to reduce violence against healthcare staff, highlighting its importance in maintaining a safe and supportive environment.<sup>2</sup> **Therapeutic communication** is a type of professional communication defined as the purposeful, interpersonal, information-transmitting process that leads to client understanding and participation.<sup>3</sup> Read an example of a nurse using therapeutic communication in the following box.

### Example of Nurse Using Therapeutic Listening

1. Karimi, H., & Masoudi Alavi, N. (2015). Florence Nightingale: The mother of nursing. *Nursing and Midwifery Studies*, 4(2), e29475. <https://doi.org/10.17795/nmsjournal29475>
2. Amara, S. S, Hansen, B., & Torres, J. (2024). Revisiting therapeutic communication as an evidence-based intervention to decrease violence by patients against staff on psychiatric wards – A quality improvement project. *Issues in Mental Health Nursing* 45(12), 1340-1352. [doi: 10.1080/01612840.2024.2414744](https://doi.org/10.1080/01612840.2024.2414744).
3. Abdolrahimi, M., Ghiyasvandian, S., Zakerimoghadam, M., & Ebadi, A. (2017). Therapeutic communication in nursing students: A Walker & Avant concept analysis. *Electronic Physician*, 9(8), 4968–4977. <https://doi.org/10.19082/4968>



Ms. Z. is a nurse (as simulated in Figure 2.1)<sup>4</sup> who enjoys interacting with clients. When she goes to clients' rooms, she greets them and introduces herself and her role in a calm tone. She kindly asks clients about their concerns and problems and notices their reactions. She provides information and answers their questions. Clients perceive that she wants to help them. She treats clients professionally by respecting boundaries and listening to them in a nonjudgmental manner. She addresses communication barriers and respects clients' cultural beliefs. She notices clients' health literacy and ensures they understand her messages and client education. As a result, clients trust her and feel as if she cares about them, so they feel comfortable sharing their health care needs with her.<sup>5</sup>

4. "[beautiful african nurse taking care of senior patient in wheelchair](#)" by [agilemktg1](#) is in the [Public Domain](#).

5. Abdolrahimi, M., Ghiyasvandian, S., Zakerimoghadam, M., & Ebadi, A. (2017). Therapeutic communication in nursing students: A Walker & Avant concept analysis. *Electronic Physician*, 9(8), 4968–4977. <https://doi.org/10.19082/4968>



*Figure 2.1 Nursing Using Therapeutic Communication*

Therapeutic communication is different from social interaction. Social interaction does not have a goal or purpose and includes casual sharing of information, whereas therapeutic communication has a goal or purpose for the conversation. An example of a nursing goal using therapeutic communication is, “The client will share feelings or concerns about their treatment plan by the end of the conversation.”

Therapeutic communication includes active listening, professional touch, and a variety of therapeutic communication techniques.

## Active Listening

Listening is an important part of communication. There are three main types of listening, including competitive, passive, and active listening. Competitive listening occurs when we are mostly focused on sharing our own point of view instead of listening to someone else. Passive listening occurs when we

are not interested in listening to the other person, and we assume we understand what the person is communicating correctly without verifying their message. During **active listening**, we communicate both verbally and nonverbally that we are interested in what the other person is saying while also actively verifying our understanding with them. For example, an active listening technique is to restate what the person said and then verify our understanding is correct. This feedback process is the major difference between passive listening and active listening.<sup>6</sup>

Nonverbal communication is an important component of active listening. **SOLER** is a mnemonic for establishing good nonverbal communication with clients. SOLER stands for the following<sup>7</sup>:

- **S:** Sitting and squarely facing the client
- **O:** Using open posture (i.e., avoid crossing arms)
- **L:** Leaning towards the client to indicate interest in listening
- **E:** Maintaining good eye contact
- **R:** Maintaining a relaxed posture

## Touch

Professional touch is a powerful way to communicate caring and empathy if done respectfully while also being aware of the client's preferences, cultural beliefs, and personal boundaries. Nurses use professional touch when assessing, expressing concern, or comforting clients. For example, simply holding a client's hand during a painful procedure can effectively provide comfort.

6. Libretext. (2022). *Human Resources (Dias)*. [https://socialsci.libretexts.org/Bookshelves/Communication/Book:\\_Human\\_Relations\\_\(Dias\)](https://socialsci.libretexts.org/Bookshelves/Communication/Book:_Human_Relations_(Dias))

7. Stickley, T. (2011). From SOLER to SURETY for effective non-verbal communication. *Nurse Education in Practice*, 11(6), 395–398. <https://doi.org/10.1016/j.nepr.2011.03.021>

For individuals with a history of trauma, touch can be negatively perceived, so it is important to always ask permission before touching. Inform the person before engaging in medical procedures requiring touch such as, “May I hold your arm still during the blood draw?”

Nurses should avoid using touch with individuals who are becoming agitated or experiencing a manic or psychotic episode because it can cause escalation. It is also helpful to maintain a larger interpersonal distance when interacting with an individual who is experiencing paranoia or psychosis.

## Therapeutic Communication Techniques

There are a variety of therapeutic techniques that nurses use to engage clients in verbalizing emotions, establishing goals, and discussing coping strategies. See Table 2.3a for definitions of various therapeutic communication techniques discussed in the *American Nurse*, the official journal of the American Nurses Association.

Table 2.3a Therapeutic Communication Techniques<sup>8</sup>

8. American Nurse. (n.d.). *Therapeutic communication techniques*.  
<https://www.myamericannurse.com/therapeutic-communication-techniques/>

Therapeutic Techniques	Definition	Examples
<b>Acceptance</b>	Acceptance acknowledges a client's emotions or message and affirms they have been heard. Acceptance isn't necessarily the same thing as agreement; it can be enough to simply make eye contact and say, "I hear what you are saying." Clients who feel their nurses are listening to them and taking them seriously are more likely to be receptive to care.	<p>Client: "I hate taking all this medicine. It makes me feel numb."</p> <p>Nurse (making eye contact): "Yes, I understand."</p>
<b>Clarification</b>	Clarification asks the client to further define what they are communicating. Similar to active listening, asking for clarification when a client says something confusing or ambiguous is important. It helps nurses ensure they understand what is actually being said and can help clients process their ideas more thoroughly.	<p>Client: "I feel useless to everyone and everything."</p> <p>Nurse: "I'm not sure I understand what you mean by useless. Can you give an example of a time you felt useless?"</p>
<b>Focusing</b>	Focusing on a specific statement made by a client that seems particularly important prompts them to discuss it further. Clients don't always have an objective perspective on their situation or past experiences, but as impartial observers, nurses can more easily pick out important topics on which to focus.	<p>Client: "I grew up with five brothers and sisters. We didn't have much money, so my mom was always working and never home. We had to fend for ourselves, and there was never any food in the house."</p> <p>Nurse: "It sounds as if you experienced some stressful conditions growing up."</p>

<b>Exploring</b>	Exploring gathers more information about what the client is communicating.	<p>Client: "I had to lie when I found out a dark secret about my sister."</p> <p>Nurse: "If you feel comfortable doing so, tell me more about the situation and your sister's dark secret."</p>
<b>Giving Recognition</b>	Giving recognition acknowledges and validates the client's positive health behaviors. Recognition acknowledges a client's behavior and highlights it without giving an overt compliment. A compliment can sometimes be taken as condescending, especially when it concerns a routine task like making the bed.	Nurse: "I noticed you took all of your medications."
<b>Open-Ended Questions/ Offering General Leads</b>	Using open questions or offering general leads provides keywords to "open" the discussion while also seeking more information. Therapeutic communication is most effective when clients direct the flow of conversation and decide what to talk about. Giving clients a broad opening such as "What's on your mind today?" or "What would you like to talk about?" is a good way to encourage clients to discuss what's on their mind.	<p>Client: "I'm unsure of what to do next."</p> <p>Nurse: "Tell me more about your concerns."</p>
<b>Paraphrasing</b>	Paraphrasing rephrases the client's words and key ideas to clarify their message and encourage additional communication.	<p>Client: "I've been way too busy today."</p> <p>Nurse: "Participating in the support groups today has kept you busy."</p>

<b>Presenting Reality</b>	Presenting reality restructures the client's distorted thoughts with valid information.	<p>Client: "I can't go in that room; there are spiders on the walls."</p> <p>Nurse: "I see no evidence of spiders on the walls."</p>
<b>Restating</b>	Restating uses different word choices for the same content stated by the client to encourage elaboration.	<p>Client: "The nurses hate me here."</p> <p>Nurse: "You feel as though the nurses dislike you?"</p>
<b>Reflecting</b>	Reflecting asks clients what they think they should do, encourages them to be accountable for their own actions, and helps them come up with solutions.	<p>Client: "Do you think I should do this new treatment or not?"</p> <p>Nurse: "What do you think the pros and cons are for the new treatment plan?"</p>
<b>Providing Silence</b>	Providing silence allows quiet time for self-reflection by the client.	The nurse does not verbally respond after a client makes a statement, although they may nod or use other nonverbal communication to demonstrate active listening and validation of the client's message.
<b>Making Observations</b>	Observations about the appearance, demeanor, or behavior of clients can help draw attention to areas that might pose a problem for them.	<p>Nurse: "You look tired today."</p> <p>Client: "I haven't been getting much sleep lately because of so many racing thoughts in my head at night."</p>

<b>Offering Self/Providing Presence</b>	Offering self provides support by being present. Inpatient care can be lonely and stressful at times. When nurses provide presence and spend time with their clients, it shows clients they value them and are willing to give them time and attention.	Offering to simply sit with clients for a few minutes is a powerful way to create a caring connection.
<b>Encouraging Descriptions of Perceptions</b>	Asking about perceptions in an encouraging, nonjudgmental way is important for clients experiencing sensory issues or hallucinations. It gives clients a prompt to explain what they're perceiving without casting their perceptions in a negative light. It is also important to establish safety by ensuring the hallucinations are not encouraging the client to harm themselves or others.	<p>The client looks distracted and frightened as if they see or hear something.</p> <p>Nurse: "It looks as though you might be hearing something. What do you hear now?" or "It looks as if you might be seeing something. What does it look like to you?"</p>
<b>Encouraging Comparisons</b>	Encouraging comparisons helps clients reflect on previous situations in which they have coped effectively. In this manner, nurses can help clients discover solutions to their problems.	<p>Nurse: "It must have been difficult when you went through a divorce. How did you cope with that?"</p> <p>Client: "I walked my dog outside a lot."</p> <p>Nurse: "It sounds as though walking your dog outside helps you cope with stress and feel better?"</p>
<b>Offering Hope</b>	Offering hope encourages a client to persevere and be resilient.	Nurse: "I remember you shared with me how well you coped with difficult situations in the past."



<b>Offering Humor</b>	<p>Humor can lighten the mood and contribute to feelings of togetherness, closeness, and friendliness. However, it is vital for the nurse to tailor humor to the client's sense of humor.</p>	<p>Nurse: "Knock, knock."</p> <p>Client: "Who's there?"</p> <p>Nurse: "Orange."</p> <p>Client: "Orange who?"</p> <p>Nurse: "Orange you glad to see me?" (Laughs with the client)</p> <p>Or</p> <p>Nurse (smiling after finishing teaching and inserting the IV):</p> <p>"Well, not too bad for my first time, right?"</p> <p>(Pauses for dramatic effect, then adds)</p> <p>"Kidding! I've done this a hundred times—just wanted to keep things exciting!"</p>
<b>Confronting</b>	<p>Confronting presents reality or challenges a client's assumptions. Nurses should only apply this technique during the working phase after they have established trust. Confrontation, when used correctly, can help clients break destructive routines or understand the state of their current situation.</p>	<p>Client: "I haven't drunk much this year."</p> <p>Nurse: "Yesterday you told me that every weekend you go out and drink so much you don't know where you are when you wake up."</p>

<b>Summarizing</b>	Summarizing demonstrates active listening to clients and allows the nurse to verify information. Ending a discussion with a phrase such as “Does that sound correct?” gives clients explicit permission to make corrections if they’re necessary.	Client: “I don’t like to take my medications because they make me tired, and I gain a lot of weight.”  Nurse: “You haven’t been taking your medications this month because of the side effects of fatigue and weight gain. Is that correct?”
--------------------	---	--

## Nontherapeutic Responses

Nurses must be aware of potential barriers to communication and avoid nontherapeutic responses. Nonverbal communication such as looking at one’s watch, crossing arms across one’s chest, or not actively listening may be perceived as barriers to communication. Nontherapeutic verbal responses often block the client’s communication of feelings or ideas. See Table 2.3b for a description of nontherapeutic responses to avoid.

Table 2.3b Nontherapeutic Responses<sup>9,10</sup>

9. Abdolrahimi, M., Ghiyasvandian, S., Zakerimoghadam, M., & Ebadi, A. (2017). Therapeutic communication in nursing students: A Walker & Avant concept analysis. *Electronic Physician*, 9(8), 4968–4977. <https://doi.org/10.19082/4968>

10. Sharma, N. P., & Gupta, V. (2023). Therapeutic communication. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK567775/>

Nontherapeutic Response	Description	Examples
<b>Asking Personal Questions</b>	Asking personal questions that are not relevant to the situation is not professional or appropriate. Don't ask questions just to satisfy your curiosity.	<p>Nontherapeutic: "Why have you and Mary never gotten married?"</p> <p>Therapeutic: "How would you describe your relationship with Mary?"</p>
<b>Giving Personal Opinions</b>	Giving personal opinions takes away the decision-making from the client. Effective problem-solving must be accomplished by the client and not provided by the nurse.	<p>Nontherapeutic: "If I were you, I would put your father in a nursing home to reduce your stress."</p> <p>Therapeutic: "Let's explore options for your father's care."</p>
<b>Changing the Subject</b>	Changing the subject when someone is trying to communicate with you demonstrates lack of empathy and blocks further communication. It communicates that you don't care about what they are sharing.	<p>Nontherapeutic: "Let's not talk about your insurance problems; it's time for your walk now."</p> <p>Therapeutic: "After your walk, let's look into what is going on with your insurance company."</p>

<b>Stating Generalizations and Stereotypes</b>	Generalizations and stereotypes can threaten nurse-client relationships.	<p>Nontherapeutic: "Older adults are always confused."</p> <p>Therapeutic: "Tell me more about your concerns about your father's confusion."</p>
<b>Providing False Reassurances</b>	When a client is seriously ill or distressed, the nurse may be tempted to offer false hope with statements that everything will be alright. These comments can discourage further expressions of a client's feelings.	<p>Nontherapeutic: "You'll be fine; don't worry."</p> <p>Therapeutic: "It must be difficult not to know what will happen next. What can I do to help?"</p>
<b>Showing Sympathy</b>	Sympathy focuses on the nurse's feelings rather than the client. It demonstrates pity rather than trying to help the client cope with the situation.	<p>Nontherapeutic: "I'm so sorry about your amputation; I can't imagine losing my leg due to a car crash."</p> <p>Therapeutic: "The loss of your leg is a major change. How do you think this will affect your life?"</p>

<b>Asking “Why” Questions</b>	<p>A nurse may be tempted to ask the client to explain “why” they believe, feel, or act in a certain way. However, clients and family members can interpret “why” questions as accusations and become defensive. It is best to rephrase a question to avoid using the word “why.”</p>	<p>Nontherapeutic: “Why are you so upset?”</p> <p>Therapeutic: “You seem upset. Tell me more about that.”</p>
<b>Approving or Disapproving</b>	<p>Nurses should not impose their own attitudes, values, beliefs, and moral standards on others while in the professional nursing role. Judgmental messages contain terms such as “should,” “shouldn’t,” “ought to,” “good,” “bad,” “right,” or “wrong.” Agreeing or disagreeing sends the subtle message that a nurse has the right to make value judgments about the client’s decisions. Approving implies that the behavior being praised is the only acceptable one, and disapproving implies that the client must meet the nurse’s expectations or standards. Instead, the nurse should assist the client to explore their own values, beliefs, goals, and decisions.</p>	<p>Nontherapeutic: “You shouldn’t consider elective surgery; there are too many risks involved.”</p> <p>Therapeutic: “You are considering having elective surgery. Tell me more about the pros and cons of surgery.”</p>

<b>Giving Defensive Responses</b>	<p>When clients or family members express criticism, nurses should listen to the message. Listening does not imply agreement. To discover reasons for the client's anger or dissatisfaction, the nurse should listen without criticizing, avoid being defensive or accusatory, and attempt to defuse anger.</p>	<p>Client: "Everyone is lying to me!"</p> <p>Nontherapeutic: "No one here would intentionally lie to you."</p> <p>Therapeutic: "You believe people have been dishonest with you. Tell me more about what happened." (After obtaining additional information, the nurse may elect to follow the chain of command at the agency and report the client's concerns for follow-up.)</p>
<b>Providing Passive or Aggressive Responses</b>	<p>Passive responses serve to avoid conflict or sidestep issues, whereas aggressive responses provoke confrontation. Nurses should use assertive communication.</p>	<p>Nontherapeutic: "It's your fault you are feeling ill because you didn't take your medicine."</p> <p>Therapeutic: "Taking your medicine every day can prevent these symptoms from returning."</p>

<b>Arguing</b>	<p>Arguing against client perceptions denies that they are real and valid. They imply that the other person is lying, misinformed, or uneducated. The skillful nurse can provide information or present reality in a way that avoids argument.</p>	<p>Nontherapeutic: “How can you say you didn’t sleep last night when I heard you snoring!”</p> <p>Therapeutic: “You don’t feel rested this morning? Let’s talk about ways to improve the quality of your rest.”</p>
----------------	--	---

See the following box for a summary of tips for using therapeutic communication and avoiding common barriers to therapeutic communication.

### **Tips for Effective Therapeutic Communication**

- Establish a goal for the conversation.
- Be self-aware of one’s nonverbal behaviors and communication
- Observe the client’s nonverbal behaviors and actions as ‘cues’ for assessments and planning interventions.
- Avoid self-disclosure of personal information and use professional boundaries. (Review boundary setting in the “[Boundaries](#)” section of Chapter 1.)
- Be client-centered and actively listen to what the client is expressing (e.g., provide empathy, not sympathy; show respect; gain the client’s trust; and accept the person as who they are as an individual).
- Be sensitive to the values, cultural beliefs, attitudes, practices, and problem-solving strategies of the client.

- Use open-ended questions and active listening to encourage client expression and build trust.
- Recognize themes in a conversation (e.g., Is there a theme emerging of poor self-esteem, guilt, shame, loneliness, helplessness, hopelessness, or suicidal thoughts?).

### **Common Barriers to Therapeutic Communication**

- Using a tone of voice that is disengaged, condescending, or disapproving.
- Using medical jargon or too many technical terms.
- Asking yes/no questions instead of open-ended questions.
- Continually asking “why,” causing the client to become defensive or feel challenged by your questions.
- Using too many probing questions, causing the client to feel you are interrogating them, resulting in defensiveness or refusal to talk with the nurse.
- Lacking awareness of one’s biases, fears, feelings, or insecurities.
- Causing sensory overload in the client with a high emotional level of the content.
- Giving advice.
- Blurring the nurse-client relationship boundaries (e.g., assuming control of the conversation, disclosing personal information, practicing outside one’s scope of practice).

## **Recognizing and Addressing Escalation**

When communicating therapeutically with a client, it is important to



recognize if the client is escalating with increased agitation and becoming a danger to themselves, staff, or other clients. When escalation occurs, providing safety becomes the nurse's top priority. Read more information in the "[Crisis and Crisis Intervention](#)" section of the "Stress, Coping, and Crisis Intervention" chapter.

## Cultural Considerations

Recall the discussion from Chapter 1 on how cultural values and beliefs can impact a client's mental health in many ways. Every culture has a different perspective on mental health. For many cultures, there is stigma surrounding mental health. Mental health challenges may be considered a weakness and something to hide, which can make it harder for those struggling to talk openly and ask for help. Culture can also influence how people describe and feel about their symptoms. It can affect whether someone chooses to recognize and talk openly about physical symptoms, emotional symptoms, or both. Cultural factors can determine how much support someone gets from their family and community when it comes to mental health.<sup>11</sup>

Nurses can help clients understand the role culture plays in their mental health by encouraging therapeutic communication about their symptoms and treatment. For example, a nurse should ask, "Can you tell me more about how your culture influences your values and beliefs?" or "Can you tell me more about how your culture influences your options for treatment?"

► Read more about providing culturally responsive care in the

11. Mental Health First Aid. (2019). *Four ways culture impacts mental health*. National Council for Mental Wellbeing. <https://www.mentalhealthfirstaid.org/2019/07/four-ways-culture-impacts-mental-health/>

- ▶ [“Diverse Clients”](#) chapter of Open RN *Nursing Fundamentals*, 2e.

## 2.4 Motivational Interviewing

Client education and health promotion are core nursing interventions.

**Motivational interviewing (MI)** is a communication skill used to elicit and emphasize a client's personal motivation for modifying behavior to promote health. MI has been effectively used for several health issues such as smoking cessation, diabetes, substance use disorders, and adherence to a treatment plan.<sup>1</sup>

The spirit of motivational interviewing is a collaborative partnership between nurses and clients, focused on client-centered care, autonomy, and personal responsibility. It is a technique that explores a client's motivation, confidence, and roadblocks to change. During motivational interviewing, nurses pose questions, actively listen to client responses, and focus on where the client is now with current health behavior and where they want to be in the future.<sup>2</sup>

Motivational interviewing uses these principles<sup>3</sup>:

- **Express empathy.** Use reflective listening to convey acceptance and a nonjudgmental attitude. Rephrase client comments to convey active listening and let clients know they are being heard.
- **Highlight discrepancies.** Help clients become aware of the gap between

1. Rubak, S., Sandbaek, A., Lauritzen, T., & Christensen, B. (2005). Motivational interviewing: A systematic review and meta-analysis. *The British Journal of General Practice: The Journal of the Royal College of General Practitioners*, 55(513), 305–312. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1463134/>
2. Droppa, M., & Lee, H. (2014). Motivational interviewing: A journey to improve health. *Nursing*, 44(3), 40–46. <https://doi.org/10.1097/01.NURSE.0000443312.58360.82>
3. Droppa, M., & Lee, H. (2014). Motivational interviewing: A journey to improve health. *Nursing*, 44(3), 40–46. <https://doi.org/10.1097/01.NURSE.0000443312.58360.82>

their current behaviors and their values and goals. Present objective information that highlights the consequences of continuing their current behaviors to motivate them to change their behavior.

- **Adjust to resistance.** Adjust to a client's resistance and do not argue. The client may demonstrate resistance by avoiding eye contact, becoming defensive, interrupting you, or seeming distracted by looking at their watch or cell phone. Arguing can place the client on the defensive and in a position of arguing against the change. Focus on validating the client's feelings.
- **Understand motivations.** Uncover a client's personal reasons for making behavioral changes and build on them.
- **Support self-efficacy.** Encourage the client's optimistic belief in the prospect of change and encourage them to commit to positive behavioral changes. Ask clients to elaborate on past successes to build self-confidence and support self-efficacy.
- **Resist the reflex to provide advice.** Avoid imposing your own perspective and advice.

When implementing motivational interviewing, it is important to assess the client's readiness for change. Motivational interviewing is especially useful for clients in the contemplation stage who are feeling ambivalent about making change. Recall these five stages of behavioral change<sup>4</sup>:

- **Precontemplation:** Not considering change.
- **Contemplation:** Ambivalent about making change.
- **Preparation:** Taking steps toward implementing change.
- **Action:** Actively involved in the change process.
- **Maintenance:** Sustaining the target behavior.

4. Droppa, M., & Lee, H. (2014). Motivational interviewing: A journey to improve health. *Nursing*, 44(3), 40–46. <https://doi.org/10.1097/01.NURSE.0000443312.58360.82>

Identify clients who are ambivalent about making a behavioral change or following a treatment plan by listening for the phrase, “Yes, but.” The “but” holds the key for opening the conversation about ambivalence. For example, a client may state, “I want to take my medication, but I hate gaining weight.” The content in the sentence after the “but” reveals the client’s personal roadblock to making a change and should be taken into consideration when planning outcomes and interventions.<sup>5</sup>

See the following box for an example of a nurse using motivational interviewing with a client.

### **Example of Motivational Interviewing<sup>6</sup>**

Mr. L. had been in treatment for bipolar I disorder with medication management and supportive therapy for many years. He had a history of alcohol dependence but was in full recovery. Mr. L. was admitted to the intensive care unit with a toxic lithium level. He had been seen in the emergency room the preceding evening and was noted to have a very high blood alcohol level. The next day the nurse asked the client about his alcohol use using motivational interviewing.

**Client:** I am so sick of everyone always blaming everything on my drinking!

5. Droppa, M., & Lee, H. (2014). Motivational interviewing: A journey to improve health. *Nursing*, 44(3), 40–46. <https://doi.org/10.1097/01.NURSE.0000443312.58360.82>

6. Griffith, L. J. (2008). The psychiatrist’s guide to motivational interviewing. *Psychiatry*, 5(4), 42–47. <https://www.ncbi.nlm.nih.gov/pubmed/19727309>

**Nurse (Using reflective listening):** You seem pretty angry about the perception that you were hospitalized because you had been drinking.

**Client:** You better believe it! I am a man! I can have a few drinks if I want to!

**Nurse: (Expressing empathy and acceptance):** You want to be respected even when you are drinking.

**Client:** I have had some trouble in the past with drinking, but that is not now. I can quit if I want to! Compared to what I used to drink, this is nothing.

**Nurse (Rolling with resistance):** So, you see yourself as having had drinking problems in the past, but the drinking you've done recently is not harmful for you.

**Client:** Well, I guess I did end up in the hospital.

**Nurse (Using open-ended questioning):** Tell me more about what happened.

**Client:** I was pretty angry after an argument with my girlfriend, and I decided to buy a bottle of whiskey.

**Nurse (Exploring):** And then?

**Client:** Well, I meant to have a couple of shots, but I ended up drinking the whole fifth. I really don't remember what happened next. They said I nearly died.

**Nurse (Summarizing):** So, after many years of not drinking, you decided to have a couple of drinks after the argument with your girlfriend, but unintentionally drank enough to have a blackout and nearly die.

**Client:** I guess that does sound like a problem...but I don't want anyone else telling me whether or not I can drink!

**Nurse (Emphasizing autonomy):** If you would be able to go back in time, would you like to approach the situation the same or differently?



View the following supplementary YouTube videos about motivational interviewing:

- [Introduction to Motivational Interviewing](#)<sup>7</sup>
- [Motivational Interviewing – Good Example – Alan Lyme](#)<sup>8</sup>

- ▶ Complete Western Region Public Health Training Center's [Motivational Interviewing](#) course and receive a certificate of completion.

7. Matulich, B. (2013, May 30). *Introduction to motivational interviewing* [Video]. YouTube. All rights reserved. <https://youtu.be/s3MCJZ7OGRk>

8. TheRETACHannel. (2013, July 18). *Motivational interviewing – Good example – Alan Lyme* [Video]. YouTube. All rights reserved. <https://youtu.be/67l6g1l7Zao>

## 2.5 Teletherapy and Telehealth

**Telehealth** is the use of digital technologies to deliver medical care, health education, and public health services by remotely connecting multiple users in separate locations. Nurses must be aware of potential barriers affecting client use of telehealth (such as lack of Internet access or lack of support for individuals learning new technologies), as well as state and federal policies regarding telehealth and using their nursing license across state lines.

- ▶ Read more about telehealth licensing requirements and interstate compacts at the [Telehealth.hhs.gov](https://telehealth.hhs.gov) webpage.

**Teletherapy** is mental health counseling over the phone or online with videoconferencing. COVID-19 has led to reduced access to medical and mental health care, so delivering behavioral health care via telehealth is one way to address this issue. When using teletherapy, nurses should treat clients as if they are sitting across from them and focus on eye contact and empathetic expressions to build a connection, just as one would do during a face-to-face encounter.<sup>1</sup> Effective teletherapy involves adherence to professional guidelines, appropriate training, managing technology, ensuring privacy, and engaging caregivers when necessary.

Group therapy can also be accomplished via telehealth. Connecting clients through telehealth creates a group dynamic that can build community, reduce feelings of isolation, and offer new perspectives. Group therapy via telehealth can create a sense of belonging and build a trusted support system.

1. Telehealth.HHS.gov. (2024). *Telehealth for behavioral health care*. Health Resources & Services Administration. <https://telehealth.hhs.gov/providers/telehealth-for-behavioral-health/individual-teletherapy/>



Here are a few guidelines for group therapy telehealth sessions<sup>2</sup>:

- **Prescreen group members:** Group members may have various needs, experiences, or personalities. It is helpful to screen each potential client to ensure every member can benefit from group therapy and that their needs match the goals of the group.
- **Require completion of online consent forms:** Group telehealth sessions involve multiple people and are conducted outside of a controlled setting like an office. Client consent forms should be required and available online. The consent forms should outline any associated risks, benefits, and limits to confidentiality.
- **Develop group guidelines:** Make clear ground rules covering what is acceptable and what is not acceptable. Some common ground rules include requiring all participants to have their camera on, attend from a room where they can be alone during the session, and use the digital “raise hand” feature (or raise their hand) when they want to speak. Prohibiting recording of the session is a common ground rule to protect confidentiality. Address logistical topics like how many missed sessions are allowed and how to contact the group leader(s).
- **Select your settings and technology:** Choose the telehealth video platform that best suits your needs for encryption and privacy, user controls, and more. Go through all of the settings ahead of time to select the options that provide the highest level of privacy. Think about what will help you and the group communicate effectively such as screen sharing options or a virtual whiteboard.
- **Be engaging:** When you are on screen instead of in person, it is even more important to be conscious of the group dynamic and take steps to keep group members interested, energized, and engaged. Start with introductions and greetings using first names only for privacy. Make eye

2. Telehealth.HHS.gov. (2024). *Telehealth for behavioral health care*. Health Resources & Services Administration. <https://telehealth.hhs.gov/providers/best-practice-guides/telehealth-for-behavioral-health/group-teletherapy>

contact with group members by looking into the camera and use body language and hand gestures to help express your ideas. Build in moments for clients to interact and contribute to the conversation, such as breakout rooms or paired discussions.

- **Group participation:** Some group members might be more willing to talk than others. Make sure to pause so that everyone has a chance to contribute to the discussion. Let participants know that they can always decide to pass if they do not want to talk when they are called upon.

## 2.6 Learning Activities



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=194#h5p-4>

1



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=194#h5p-6>

2



*An interactive H5P element has been excluded from this version of the text. You can view it*

1. “MH Therapeutic Communications Glossary Cards” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
2. “MH Therapeutic Communications Drag and Drop” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

online here:

<https://wtcs.pressbooks.pub/nursingmhcc/?p=194#h5p-9>

3



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=194#h5p-10>

4

▶ Test your clinical judgment with a NCLEX  
Next-Generation style question: [Chapter 2,](#)  
[Assignment 1](#)<sup>5</sup>



3. “MH Therapeutic Communication Question Set 1” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

4. “MH Therapeutic Communications Question Set 2” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

5. “MH Next-Generation Style Question ” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

- ▶ Test your clinical judgment with a NCLEX Next-Generation style question: [Chapter 2, Assignment 2](#)<sup>6</sup>



- ▶ Test your clinical judgment with a NCLEX Next-Generation style case study: [Chapter 2, Case Study](#)<sup>7</sup>



- ▶ **Telehealth Virtual Simulations**

These virtual, screen-based simulations demonstrate telehealth phone calls made by nurses to clients who were recently hospitalized and live in rural areas. It is assumed the clients may not have internet access and may experience barriers to accessing care due to their geographical location. During each virtual simulation, the student has the opportunity to make decisions on how to best use therapeutic communication to establish rapport and gain trust with the client, while also

6. "MH Next-Generation Style Question " by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

7. "MH Next-Generation Therapeutic Communication Case Study " by Kellea Ewen is licensed under [CC BY-NC 4.0](#)

- ▶ considering the potential impact of the client's cultural beliefs and social determinants of health. Immediate feedback and rationale are provided for correct and incorrect answers as the student proceeds through the simulation. The first telehealth visit is a post-hospitalization welcome call where the nurse follows up on client concerns post-hospitalization. During the second telehealth call, the nurse identifies the client's specific learning needs. During the third telehealth visit, the nurse evaluates the effectiveness of the health teaching provided. The virtual simulations can be accessed by clicking on the following links.<sup>8</sup>

**Client 1 :** Alejandro Hernandez, Hispanic Male with Heart Failure: [Visit 1](#) – [Visit 2](#) – [Visit 3](#)

**Client 2:** Chen Xiulan, Asian Female with COPD: [Visit 1](#) – [Visit 2](#) – [Visit 3](#)

**Client 3:** Carolyn Smith, Caucasian Female with Mild Dementia and Chronic Kidney Disease: [Visit 1](#) – [Visit 2](#) – [Visit 3](#)

**Client 4:** Dakotah Thunderhawk, Native American Male Post-Cerebrovascular Accident: [Visit 1](#) – [Visit 2](#) – [Visit 3](#)

**Client 5:** Marian Johnson, African American Female with Diabetes and Amputation: [Visit 1](#) – [Visit 2](#) – [Visit 3](#)

9

8. “Telehealth Virtual Simulations” by OpenRN is licensed under [CC BY-NC 4.0](#)

9. This workforce product was funded by a grant awarded by the U.S. Department of Labor’s Employment and Training Administration. The product was created by the grantee and does not necessarily reflect the official position of the U.S. Department of Labor. The U.S. Department of Labor makes no guarantees, warranties, or assurances of any kind, express or implied, with respect to such information, including any

information on linked sites and including, but not limited to, accuracy of information or its completeness, timeliness, usefulness, adequacy, continued availability, or ownership.

## II Glossary

**Active listening:** Communicating both verbally and nonverbally that we are interested in what the other person is saying while also actively verifying our understanding with them.

**Motivational interviewing (MI):** A communication skill used to elicit and emphasize a client's personal motivation for modifying behavior to promote health.

**Nurse-client relationship:** A relationship that establishes trust and rapport with a specific purpose of facilitating therapeutic communication and engaging the client in decision-making regarding their plan of care.

**SOLER:** A mnemonic for effective nonverbal communication that stands for the following<sup>1</sup>:

- S: Sit and squarely face the client
- O: Open posture
- L: Lean towards the client to indicate interest in listening
- E: Eye contact
- R: Relax

**Telehealth:** The use of digital technologies to deliver medical care, health education, and public health services by remotely connecting multiple users in separate locations.

**Teletherapy:** Mental health counseling over the phone or online with videoconferencing tools.

**Therapeutic communication:** A type of professional communication defined

1. Stickley, T. (2011). From SOLER to SURETY for effective non-verbal communication. *Nurse Education in Practice*, 11(6), 395–398. <https://doi.org/10.1016/j.nepr.2011.03.021>



as the purposeful, interpersonal, information-transmitting process that leads to client understanding and participation.<sup>2</sup>

2. Abdolrahimi, M., Ghiyasvandian, S., Zakerimoghadam, M., & Ebadi, A. (2017). Therapeutic communication in nursing students: A Walker & Avant concept analysis. *Electronic Physician*, 9(8), 4968–4977. <https://doi.org/10.19082/4968>







## 3.1 Introduction

### Learning Objectives

- Describe the role of stress in exacerbating symptoms of mental health disorders
- Identify safety/protective interventions for the client and others
- Provide health teaching on stress management techniques and adaptive coping strategies
- Recognize the use of defense mechanisms
- Recognize a client in crisis
- Describe crisis intervention

Nurses support the emotional, mental, and social well-being of all clients experiencing stressful events and those with acute and chronic mental illnesses.<sup>1</sup> This chapter will review stressors, stress management, coping strategies, defense mechanisms, and crisis intervention.

1. NCLEX. (n.d.). *Test plans*. <https://www.nclex.com/test-plans.page>

## 3.2 Stress

Everyone experiences stress during their lives. High levels of stress can cause symptoms like headaches, back pain, and gastrointestinal symptoms. Chronic stress can contribute to the development of chronic illnesses, as well as acute physical illnesses, due to decreased effectiveness of the immune system and impact on the hypothalamic-pituitary-adrenal (HPA) axis. The HPA impact is discussed in greater detail below. It is important for nurses to recognize signs and symptoms of stress in themselves and others, as well as encourage effective stress management strategies. This section begins with an overview of the stress response and its signs and symptoms, followed by a discussion of stress management techniques.

We will begin this section by reviewing the stress response and signs and symptoms of stress and then discuss stress management techniques.

### Stress Response

**Stressors** are any internal or external event, force, or condition that results in physical or emotional stress.<sup>1</sup> The body's sympathetic nervous system (SNS) responds to actual or perceived stressors with the “fight, flight, freeze, or fawn” stress response. Several reactions occur during the **stress response** that help the individual to achieve the purpose of either fighting or running.

Several physiological changes occur:

- The respiratory, cardiovascular, and musculoskeletal systems activate to increase breathing, heart rate, and blood pressure, ensuring oxygenated blood reaches the muscles.
- The liver releases glucose to provide energy for immediate action.
- Pupils dilate to enhance vision.
- Sweating prevents overheating due to increased muscle activity.

1. American Psychological Association. (n.d.). Stressor. *APA Dictionary of Psychology*. <https://dictionary.apa.org>

- The digestive system slows, redirecting blood flow to essential muscles.

These responses are coordinated by the hypothalamic-pituitary-adrenal (HPA) axis, which releases hormones like epinephrine, norepinephrine, and glucocorticoids (including cortisol, the “stress hormone”). These hormones target SNS receptors throughout the body, ensuring a rapid, systemic reaction.<sup>2</sup>

In the “fawn” response, which is particularly common in individuals exposed to relational trauma, the body may still activate stress-related pathways, but the behavioral focus shifts toward pleasing or placating others to prevent harm. Though less visibly physical, fawning is still driven by the same biological systems seeking safety in a perceived threat.

Once the threat subsides, the parasympathetic nervous system (PNS) counteracts the SNS, restoring the body to its pre-stress state. See Figure 3.1 for a comparison of SNS and PNS effects.

2. Betts, J. G., Young, K. A., Wise, J. A., Johnson, E., Poe, B., Kruse, D. H., Korol, O., Johnson, J. E., Womble, M., & DeSaix, P. (2022). *Anatomy and physiology 2e*. OpenStax. <https://openstax.org/books/anatomy-and-physiology-2e/pages/1-introduction>




Figure 3.1 SNS and PNS Simulation

## Signs and Symptoms of Chronic Stress

Nurses are often the first to notice signs and symptoms of stress and can help make their clients aware of these symptoms. Common signs and symptoms of chronic stress are as follows<sup>34</sup>:

- Shaw, W., Labott-Smith, S., Burg, M. M., Hostinar, C., Alen, N., van Tilburg, M. A. L., Berntson, G. G., Tavian, S. M., & Spirito, M. (2018). *Stress effects on the body*. American Psychological Association. <https://www.apa.org/topics/stress/body>
- Kelly, J. F., & Coons, H. L. (2019). *Stress won't go away? Maybe you are suffering*



- Irritability
- Fatigue
- Headaches
- Difficulty concentrating
- Rapid, disorganized thoughts
- Difficulty sleeping
- Digestive problems
- Changes in appetite
- Feeling helpless
- A perceived loss of control
- Low self-esteem
- Loss of sexual desire
- Nervousness
- Frequent infections or illnesses
- Vocalized suicidal thoughts

## Effects of Chronic Stress

The “fight or flight or freeze” stress response prepares the body to react quickly to immediate threats. However, exposure to long-term stress can cause serious effects on the cardiovascular, musculoskeletal, endocrine, gastrointestinal, and reproductive systems.<sup>5</sup>

**Cardiovascular System:** Sustained increases in heart rate, blood pressure, and stress hormones contribute to arterial inflammation, raising the risk of hypertension, heart attack, and stroke.<sup>6</sup>

*from chronic stress.* American Psychological Association. <https://www.apa.org/topics/stress/chronic>

5. Shaw, W., Labott-Smith, S., Burg, M. M., Hostinar, C., Alen, N., van Tilburg, M. A. L., Berntson, G. G., Tavian, S. M., & Spirito, M. (2018). *Stress effects on the body.* American Psychological Association. <https://www.apa.org/topics/stress/body>
6. Shaw, W., Labott-Smith, S., Burg, M. M., Hostinar, C., Alen, N., van Tilburg, M. A.

**Musculoskeletal System:** During acute stress, muscles tense and relax once the threat passes. However, chronic stress keeps muscles in a constant state of tension, which may contribute to stress-related disorders. For example tension and migraine headaches are linked to persistent muscle tightness in the shoulders, neck, and head. Musculoskeletal pain in the lower back and upper extremities has been associated with job-related stress.<sup>7</sup>

**Endocrine & Immune Systems:** In acute stress, cortisol provides energy to cope with immediate challenges. However, chronic stress weakens the immune system, increasing susceptibility to conditions such as chronic fatigue, metabolic disorders (e.g., diabetes, obesity), depression, and immune dysfunction.<sup>8</sup>

**Gastrointestinal System:** Chronic stress can disrupt eating patterns, leading to eating much more or much less than usual. It may also cause:

- Acid reflux, often due to increased consumption of food, alcohol, or tobacco.
- Bowel irregularities—stress can induce muscle spasms in the bowel and affect how quickly food moves through the gastrointestinal system, leading to diarrhea or constipation.
- Aggravation of chronic gastrointestinal disorders (e.g., inflammatory bowel disease, irritable bowel syndrome) due to factors such as increased

L., Berntson, G. G., Tavian, S. M., & Spirito, M. (2018). *Stress effects on the body*. American Psychological Association. <https://www.apa.org/topics/stress/body>

7. Shaw, W., Labott-Smith, S., Burg, M. M., Hostinar, C., Alen, N., van Tilburg, M. A. L., Berntson, G. G., Tavian, S. M., & Spirito, M. (2018). *Stress effects on the body*. American Psychological Association. <https://www.apa.org/topics/stress/body>

8. Shaw, W., Labott-Smith, S., Burg, M. M., Hostinar, C., Alen, N., van Tilburg, M. A. L., Berntson, G. G., Tavian, S. M., & Spirito, M. (2018). *Stress effects on the body*. American Psychological Association. <https://www.apa.org/topics/stress/body>

nerve sensitivity, disturbed microbiota, altered gastrointestinal motility, and/or disrupted immune function.<sup>9</sup>

**Reproductive System:** Excess cortisol can disrupt normal reproductive function in both men and women.

**Men:** Chronic stress can affect testosterone production, resulting in:

- Reduced libido, erectile dysfunction, or impotence.
- Impaired sperm production and maturation.
- Decreased sperm quality—studies indicate that, compared to men who did not experience stressful events in the past year, those who experienced two or more stressful events within the same period had:
  - Lower sperm motility.
  - A reduced percentage of sperm with normal morphology (size and shape).<sup>10</sup>

**Women:** Chronic stress can affect reproductive health, leading to:

- Menstrual irregularities, including absent or irregular cycles, more painful periods, and changes in cycle length.
- Worsened premenstrual symptoms, such as cramping, fluid retention, bloating, negative mood, and mood swings.
- Reduced sexual desire.
- Impaired fertility and pregnancy health—stress can negatively impact:

9. Shaw, W., Labott-Smith, S., Burg, M. M., Hostinar, C., Alen, N., van Tilburg, M. A. L., Berntson, G. G., Tavian, S. M., & Spirito, M. (2018). *Stress effects on the body*. American Psychological Association. <https://www.apa.org/topics/stress/body>

10. Shaw, W., Labott-Smith, S., Burg, M. M., Hostinar, C., Alen, N., van Tilburg, M. A. L., Berntson, G. G., Tavian, S. M., & Spirito, M. (2018). *Stress effects on the body*. American Psychological Association. <https://www.apa.org/topics/stress/body>

- A woman's ability to conceive.
- Pregnancy health and postpartum recovery.
- Fetal development.
- Maternal-infant bonding following delivery.
- The risk of postpartum depression.<sup>11</sup>

## Adverse Childhood Experiences

Adults with adverse childhood experiences or exposure to adverse life events often experience ongoing chronic stress with an array of physical, mental, and social health problems throughout adulthood. Some of the most common health risks include physical and mental illness, substance use disorder, and a high level of engagement in risky sexual behavior.<sup>12</sup>

As previously discussed in [Chapter 1](#), **adverse childhood experiences (ACEs)** include sexual abuse, physical abuse, emotional abuse, physical neglect, emotional neglect, parental loss, or parental separation before the child is 18 years old. Individuals who have experienced four or more ACEs are at a significantly higher risk of developing mental, physical, and social problems in adulthood. Research has established that early life stress is a predictor of smoking, alcohol consumption, and drug dependence. Adults who experienced ACEs related to maladaptive family functioning (parental mental illness, substance use disorder, criminality, family violence, physical and sexual abuse, and neglect) are at higher risk for developing mood, substance abuse, and anxiety disorders. ACEs are also associated with an increased risk of the

11. Shaw, W., Labott-Smith, S., Burg, M. M., Hostinar, C., Alen, N., van Tilburg, M. A. L., Berntson, G. G., Tavian, S. M., & Spirito, M. (2018). *Stress effects on the body*. American Psychological Association. <https://www.apa.org/topics/stress/body>
12. Amnie, A. G. (2018). Emerging themes in coping with lifetime stress and implication for stress management education. *SAGE Open Medicine*, 6. <https://doi.org/10.1177%2F2050312118782545>

development of malignancy, cardiovascular disease, metabolic syndrome, and other chronic debilitating conditions.<sup>13</sup>

## Stress Management

Recognizing signs and symptoms of stress allows individuals to implement stress management strategies. Nurses can educate clients about effective strategies for reducing the stress response. Relaxation techniques and other stress-relieving activities have been shown to effectively reduce muscle tension, decrease the incidence of stress-related disorders, and increase a sense of well-being. For individuals with chronic pain conditions, stress-relieving activities have been shown to improve mood and daily function.<sup>14</sup> Effective strategies include the following<sup>15 16</sup>:

- Set personal and professional boundaries
- Maintain a healthy social support network
- Select healthy food choices
- Engage in regular physical exercise

13. Amnie, A. G. (2018). Emerging themes in coping with lifetime stress and implication for stress management education. *SAGE Open Medicine*, 6. <https://doi.org/10.1177%2F2050312118782545>
14. Shaw, W., Labott-Smith, S., Burg, M. M., Hostinar, C., Alen, N., van Tilburg, M. A. L., Berntson, G. G., Tavian, S. M., & Spirito, M. (2018). *Stress effects on the body*. American Psychological Association. <https://www.apa.org/topics/stress/body>
15. Shaw, W., Labott-Smith, S., Burg, M. M., Hostinar, C., Alen, N., van Tilburg, M. A. L., Berntson, G. G., Tavian, S. M., & Spirito, M. (2018). *Stress effects on the body*. American Psychological Association. <https://www.apa.org/topics/stress/body>
16. Kelly, J. F., & Coons, H. L. (2019). *Stress won't go away? Maybe you are suffering from chronic stress*. American Psychological Association. <https://www.apa.org/topics/stress/chronic>

- Get an adequate amount of sleep each night
- Set realistic and fair expectations
- Mindfulness activities

Setting limits is essential for effectively managing stress. Individuals should list all of the projects and commitments making them feel overwhelmed, identify essential tasks, and cut back on nonessential tasks. For work-related projects, responsibilities can be discussed with supervisors to obtain input on priorities. Encourage individuals to refrain from accepting any more commitments until they feel their stress is under control.<sup>17</sup>

Maintaining a healthy social support network with friends and family can provide emotional support.<sup>18</sup> Caring relationships and healthy social connections are essential for achieving resilience.

Eating nutritiously is a valuable tool in stress management. Clients should be taught to eat at regular intervals to ensure steady blood glucose levels to ensure optimal performance. Foods high in fatty acids, such as nuts and fish oil, can help combat anxiety and depression. Foods that are high in vitamins and minerals, such as leafy greens can also work to combat stress. A diet that is high in fiber has also been shown to result in lower perceived levels of stress. Additionally, caffeine should be avoided as it can heighten anxiety. Lastly, prepare for stress inducing or hectic times by ensuring that high

17. Kelly, J. F., & Coons, H. L. (2019). *Stress won't go away? Maybe you are suffering from chronic stress*. American Psychological Association. <https://www.apa.org/topics/stress/chronic>

18. Kelly, J. F., & Coons, H. L. (2019). *Stress won't go away? Maybe you are suffering from chronic stress*. American Psychological Association. <https://www.apa.org/topics/stress/chronic>

nutrient food is readily available. Good options are hummus with carrots, yogurt with granola or fruit and cheese.<sup>19</sup>

Physical activity increases the body's production of endorphins that boost the mood and reduce stress. Nurses can educate clients that a brisk walk or other aerobic activity can increase energy and concentration levels and lessen feelings of anxiety.<sup>20</sup>

People who are chronically stressed often suffer from lack of adequate sleep and, in some cases, stress-induced insomnia. Nurses can educate individuals how to take steps to increase the quality of sleep. Experts recommend going to bed at a regular time each night, striving for at least 7-8 hours of sleep, and, if possible, eliminating distractions, such as television, cell phones, and computers from the bedroom. Begin winding down an hour or two before bedtime and engage in calming activities such as listening to relaxing music, reading an enjoyable book, taking a soothing bath, or practicing relaxation techniques like meditation. Avoid eating a heavy meal or engaging in intense exercise immediately before bedtime. If a person tends to lie in bed worrying, encourage them to write down their concerns and work on quieting their thoughts.<sup>21</sup> For many individuals, sleeping in a cool room facilitates sleep.

Nurses can encourage clients to set realistic expectations, look at situations more positively, see problems as opportunities, and refute negative thoughts to stay positive and minimize stress. Setting realistic expectations and

19. The University of North Carolina at Chapel Hill. (2025). *Nutrition and stress*. [https://campushealth.unc.edu/health\\_topic/nutrition-and-stress/](https://campushealth.unc.edu/health_topic/nutrition-and-stress/)

20. Kelly, J. F., & Coons, H. L. (2019). *Stress won't go away? Maybe you are suffering from chronic stress*. American Psychological Association. <https://www.apa.org/topics/stress/chronic>

21. Kelly, J. F., & Coons, H. L. (2019). *Stress won't go away? Maybe you are suffering from chronic stress*. American Psychological Association. <https://www.apa.org/topics/stress/chronic>

positively reframing the way one looks at stressful situations can make life seem more manageable. Clients should be encouraged to keep challenges in perspective and do what they can reasonably do to move forward.<sup>22</sup>

Mindfulness is a form of meditation that uses breathing and thought techniques to create an awareness of one's body and surroundings. Research suggests that mindfulness can have a positive impact on stress, anxiety, and depression.<sup>23</sup> Additionally, guided imagery may be helpful for enhancing relaxation. The use of guided imagery provides a narration that the mind can focus on during the activity. For example, as the nurse encourages a client to use mindfulness and relaxation breathing, they may say, "As you breathe in, imagine waves rolling gently in. As you breathe out, imagine the waves rolling gently back out to sea." Read more about mindfulness techniques in the [Coping](#) section of this chapter.

## WHO Stress Management Guide

In addition to the stress management techniques discussed in the previous section, the World Health Organization (WHO) shares additional techniques in a guide titled *Doing What Matters in Times of Stress*. This guide consists of five categories. Each category includes techniques and skills that, based on evidence and field testing, can reduce overall stress levels even if only used for

22. Kelly, J. F., & Coons, H. L. (2019). *Stress won't go away? Maybe you are suffering from chronic stress*. American Psychological Association. <https://www.apa.org/topics/stress/chronic>
23. Kandola, A. (2018). *What are the health effects of chronic stress?* MedicalNewsToday. <https://www.medicalnewstoday.com/articles/323324#treatment>



a few minutes each day. These categories include 1) Grounding, 2) Unhooking, 3) Acting on our values, 4) Being kind, and 5) Making room.<sup>24</sup>

Nurses can educate clients that powerful thoughts and feelings are a natural part of stress, but problems can occur if we get “hooked” by them. For example, one minute you might be enjoying a meal with family, and the next moment you get “hooked” by angry thoughts and feelings. Stress can make someone feel as if they are being pulled away from the values of the person they want to be, such as being calm, caring, attentive, committed, persistent, and courageous.<sup>25</sup>

There are many kinds of difficult thoughts and feelings that can “hook us,” such as, “This is too hard,” “I give up,” “I am never going to get this,” “They shouldn’t have done that,” or memories about difficult events that have occurred in our lives. When we get “hooked,” our behavior changes. We may do things that make our lives worse, like getting into more disagreements, withdrawing from others, or spending too much time lying in bed. These are called “away moves” because they move us away from our values. Sometimes emotions become so strong they feel like emotional storms. However, we can “unhook” ourselves by focusing and engaging in what we are doing, referred to as “grounding.”<sup>26</sup>

## GROUNDING

“Grounding” is a helpful tool when feeling distracted or having trouble focusing on a task and/or the present moment. The first step of grounding is

24. World Health Organization. (2020). *Doing what matters in times of stress: An illustrated guide*. <https://www.who.int/publications/i/item/9789240003927>
25. World Health Organization. (2020). *Doing what matters in times of stress: An illustrated guide*. <https://www.who.int/publications/i/item/9789240003927>
26. World Health Organization. (2020). *Doing what matters in times of stress: An illustrated guide*. <https://www.who.int/publications/i/item/9789240003927>

to notice how you are feeling and what you are thinking. Next, slow down and connect with your body by focusing on your breathing. Exhale completely and wait three seconds, and then inhale as slowly as possible. Slowly stretch your arms and legs and push your feet against the floor. The next step is to focus on the world around you. Notice where you are and what you are doing. Use your five senses. What are five things you can see? What are four things you can hear? What can you smell? Tap your leg or squeeze your thumb and count to ten. Touch your knees or another object within reach. What does it feel like? Grounding helps us engage in life, refocus on the present moment, and realign with our values.<sup>27</sup>

## UNHOOKING

At times we may have unwanted, intrusive, negative thoughts that negatively affect us. “Unhooking” is a tool to manage and decrease the impact of these unwanted thoughts. First, NOTICE that a thought or feeling has hooked you, and then NAME it. Naming it begins by silently saying, “Here is a thought,” or “Here is a feeling.” By adding “I notice,” it unhooks us even more. For example, “I notice there is a knot in my stomach.” The next step is to REFOCUS on what you are doing, fully engage in that activity, and pay full attention to whoever is with you and whatever you are doing. For example, if you are having dinner with family and notice feelings of anger, note “I am having feelings of anger,” but choose to refocus and engage with family.<sup>28</sup>

## ACTING ON OUR VALUES

The third category of skills is called “Acting on Our Values.” This means, despite challenges and struggles we are experiencing, we will act in line with

27. World Health Organization. (2020). *Doing what matters in times of stress: An illustrated guide*. <https://www.who.int/publications/i/item/9789240003927>

28. World Health Organization. (2020). *Doing what matters in times of stress: An illustrated guide*. <https://www.who.int/publications/i/item/9789240003927>

what is important to us and our beliefs. Even when facing difficult situations, we can still make the conscious choice to act in line with our values. The more we focus on our own actions, the more we can influence our immediate world and the people and situations we encounter every day. We must continually ask ourselves, “Are my actions moving me toward or away from my values?” Remember that even the smallest actions have impact, just as a giant tree grows from a small seed. Even in the most stressful of times, we can take small actions to live by our values and maintain or create a more satisfying and fulfilling life. These values should also include self-compassion and care. By caring for oneself, we ultimately have more energy and motivation to then help others.<sup>29</sup>

## BEING KIND

“Being Kind” is a fourth tool for reducing stress. Kindness can make a significant difference to our mental health by being kind to others, as well as to ourselves. Being kind to others helps build strong relationships, which makes people happier and more resilient. Studies show that people who are kind tend to live longer and have better health. Acts of kindness benefit both the person giving and receiving them. Even just witnessing kindness can make people feel good and encourage them to be more kind themselves. It creates a positive cycle of connection and caring within communities.<sup>30</sup>

## MAKING ROOM

“Making Room” is a fifth tool for reducing stress. Sometimes trying to push away painful thoughts and feelings does not work very well. In these

29. World Health Organization. (2020). *Doing what matters in times of stress: An illustrated guide*. <https://www.who.int/publications/i/item/9789240003927>

30. Fryburg, D. A. (2021). Kindness as a stress reduction-health promotion intervention: A review of the psychobiology of caring. *American Journal of Lifestyle Medicine*, 16(1):89-100. [doi: 10.1177/1559827620988268](https://doi.org/10.1177/1559827620988268).

situations, it is helpful to notice and name the feeling, and then “make room” for it. “Making room” means allowing the painful feeling or thought to come and go like the weather. Nurses can educate clients that as they breathe, they should imagine their breath flowing into and around their pain and making room for it. Instead of fighting with the thought or feeling, they should allow it to move through them, just like the weather moves through the sky. If clients are not fighting with the painful thought or feeling, they will have more time and energy to engage with the world around them and do things that are important to them.<sup>31</sup>

▶ Read *Doing What Matters in Times of Stress* by the World Health Organization (WHO).<sup>32</sup>

▶ View the following YouTube video on the WHO Stress Management Guide<sup>33</sup>:



One or more interactive elements has been excluded from this version of the text. You can view them online here: <https://wtcs.pressbooks.pub/nursingmhcc/?p=135#oembed-1>

31. World Health Organization. (2020). *Doing what matters in times of stress: An illustrated guide*. <https://www.who.int/publications/i/item/9789240003927>
32. World Health Organization. (2020). *Doing what matters in times of stress: An illustrated guide*. <https://www.who.int/publications/i/item/9789240003927>
33. World Health Organization (WHO). (2020, November 4). *Doing what matters in times of stress: An illustrated guide* [Video]. YouTube. Licensed in the Public Domain. <https://youtu.be/E3Cts45FNrk>

# Stress Related to the COVID-19 Pandemic and World Events

The COVID-19 pandemic had a major effect on many people's lives. Many health care professionals faced challenges that were stressful, overwhelming, and caused strong emotions.<sup>34</sup> See Figure 3.2<sup>35</sup> for a message from the World Health Organization regarding stress and health care workers.



Figure 3.2 Stress and Healthcare Workers

Learning to cope with stress in a healthy way can increase feelings of

34. Centers for Disease Control and Prevention. (2021). *Healthcare personnel and first responders: How to cope with stress and build resilience during the COVID-19 pandemic*. <https://stacks.cdc.gov/view/cdc/95184>
35. "89118597-2933268333403014-548082632068431872-n.jpg" by unknown author for [World Health Organization \(WHO\)](https://www.who.int/campaigns/connecting-the-world-to-combat-coronavirus/healthyathome/healthyathome—mental-health?gclid=Cj0KCQiA0MD_BRCTARIsADXoopa7YZIdaIqCtKIGrxDV8YcUBtpVSD2HaOtT9NsdT8ajyCXbnPot-bsaAvlQEALw_wcB) is licensed in the [Public Domain](https://www.who.int/campaigns/connecting-the-world-to-combat-coronavirus/healthyathome/healthyathome—mental-health?gclid=Cj0KCQiA0MD_BRCTARIsADXoopa7YZIdaIqCtKIGrxDV8YcUBtpVSD2HaOtT9NsdT8ajyCXbnPot-bsaAvlQEALw_wcB). Access for free at [https://www.who.int/campaigns/connecting-the-world-to-combat-coronavirus/healthyathome/healthyathome—mental-health?gclid=Cj0KCQiA0MD\\_BRCTARIsADXoopa7YZIdaIqCtKIGrxDV8YcUBtpVSD2HaOtT9NsdT8ajyCXbnPot-bsaAvlQEALw\\_wcB](https://www.who.int/campaigns/connecting-the-world-to-combat-coronavirus/healthyathome/healthyathome—mental-health?gclid=Cj0KCQiA0MD_BRCTARIsADXoopa7YZIdaIqCtKIGrxDV8YcUBtpVSD2HaOtT9NsdT8ajyCXbnPot-bsaAvlQEALw_wcB).

resiliency for health care professionals. Here are ways to help manage stress resulting from world events<sup>36</sup>:

- Take breaks from watching, reading, or listening to news stories and social media. It's good to be informed but consider limiting news to just a couple times a day and disconnecting from phones, TVs, and computer screens for a while.
  - It can be important to do a self check-in before reading any news. "Do I have the emotional energy to handle a difficult headline if I see one?"
- Take care of your body.
  - Take deep breaths, stretch, or meditate
  - Try to eat healthy, well-balanced meals
  - Exercise regularly
  - Get plenty of sleep
  - Avoid excessive alcohol, tobacco, and substance use
  - Continue routine preventive measures (such as vaccinations, cancer screenings, etc.) as recommended by your health care provider
- Make time to unwind. Plan activities you enjoy.
- Purposefully connect with others. It is especially important to stay connected with your friends and family. Helping others cope through phone calls or video chats can help you and your loved ones feel less lonely or isolated. Connect with your community or faith-based organizations.
- Use the techniques described in the WHO stress management guide.<sup>37</sup>

36. Centers for Disease Control and Prevention. (2021). *Healthcare personnel and first responders: How to cope with stress and build resilience during the COVID-19 pandemic*. <https://stacks.cdc.gov/view/cdc/95184>

37. World Health Organization. (2020). *Doing what matters in times of stress: An illustrated guide*. <https://www.who.int/publications/i/item/9789240003927>

## Strategies for Self-Care

By becoming self-aware regarding signs of stress, you can implement self-care strategies to prevent compassion fatigue and burnout. Use the following “A’s” to assist in building resilience, connection, and compassion<sup>38</sup>:

- **Attention:** Become aware of your physical, psychological, social, and spiritual health. What are you grateful for? What are your areas of improvement? This protects you from drifting through life on autopilot.
- **Acknowledgement:** Honestly look at all you have witnessed as a health care professional. What insight have you experienced? Acknowledging the pain of loss you have witnessed protects you from invalidating the experiences.
- **Affection:** Choose to look at yourself with kindness and warmth. Affection and self-compassion prevent you from becoming bitter and “being too hard” on yourself.
- **Acceptance:** Choose to be at peace and welcome all aspects of yourself. By accepting both your talents and imperfections, you can protect yourself from impatience, victim mentality, and blame.

38. Lowey, S. E. (2015). *Nursing care at the end of life*. Open SUNY Textbooks. <https://ecampusontario.pressbooks.pub/nursingcare/>

### 3.3 Coping

The health consequences of chronic stress depend on an individual's coping styles and their resilience to real or perceived stress. **Coping** refers to cognitive and behavioral efforts made to master, tolerate, or reduce external and internal demands and conflicts.<sup>1</sup>

**Coping strategies** are actions, a series of actions, or thought processes used in meeting a stressful or unpleasant situation or in modifying one's reaction to such a situation. Coping strategies are classified as adaptive or maladaptive. **Adaptive coping strategies** include problem-focused coping and emotion-focused coping. **Problem-focused coping** typically focuses on seeking treatment such as counseling or cognitive behavioral therapy.

**Emotion-focused coping** includes strategies such as mindfulness, meditation, and yoga; using humor and jokes; seeking spiritual or religious pursuits; engaging in physical activity or breathing exercises; and seeking social support.

**Maladaptive coping responses** include avoidance of the stressful condition, withdrawal from a stressful environment, disengagement from stressful relationships, and misuse of drugs and/or alcohol.<sup>2</sup> Nurses can educate individuals and their family members about adaptive and emotion-focused coping strategies. Additionally, nurses can make referrals to interprofessional team members for problem-focused coping and treatment options for individuals experiencing maladaptive coping responses to stress.

1. Amnie, A. G. (2018). Emerging themes in coping with lifetime stress and implication for stress management education. *SAGE Open Medicine*, 6. <https://doi.org/10.1177%2F2050312118782545>
2. Amnie, A. G. (2018). Emerging themes in coping with lifetime stress and implication for stress management education. *SAGE Open Medicine*, 6. <https://doi.org/10.1177%2F2050312118782545>



# Emotion-Focused Coping Strategies

Nurses can educate clients about many emotion-focused coping strategies, such as meditating, practicing yoga, journaling, praying, spending time in nature, nurturing supportive relationships, and practicing mindfulness.

## Meditation

Meditation can induce feelings of calm and clearheadedness and improve concentration and attention. Research has shown that meditation increases the brain's gray matter density, which can reduce sensitivity to pain, enhance the immune system, help regulate difficult emotions, and relieve stress. Meditation has been proven helpful for people with depression and anxiety, cancer, fibromyalgia, chronic pain, rheumatoid arthritis, type 2 diabetes, chronic fatigue syndrome, and cardiovascular disease.<sup>3</sup> See Figure 3.3<sup>4</sup> for an image of an individual participating in meditation.

3. Delagran, L. (n.d.). *What is spirituality?* University of Minnesota.  
<https://www.takingcharge.csh.umn.edu/what-spirituality>

4. “[yoga-class-a-cross-legged-palms-up-meditation-position-850×831.jpg](#)” by Amanda Mills, USCDCCP on Pixnio is licensed under CC0



Figure 3.3 Meditation

## Yoga

Yoga is a centuries-old spiritual practice that creates a sense of union within the practitioner through physical postures, ethical behaviors, and breath expansion. The systematic practice of yoga has been found to reduce inflammation and stress, decrease depression and anxiety, lower blood pressure, and increase feelings of well-being.<sup>5</sup> See Figure 3.4<sup>6</sup> for an image of an individual participating in yoga.

5. Delagran, L. (n.d.). *What is spirituality?* University of Minnesota.  
<https://www.takingcharge.csh.umn.edu/what-spirituality>

6. “9707554768.jpg” by [Dave Rosenblum](#) is licensed under [CC BY 2.0](#)



Figure 3.4 Yoga

## Journaling

Journaling can help a person become more aware of their inner life and feel more connected to experiences. Studies show that writing during difficult times may help a person find meaning in life's challenges and become more resilient in the face of obstacles. When journaling, it can be helpful to focus on three basic questions: What experiences give me energy? What experiences drain my energy? Were there any experiences today where I felt alive and experienced “flow”? Allow yourself to write freely, without stopping to edit or worry about spelling and grammar.<sup>7</sup>

## Prayer

Prayer can elicit the relaxation response, along with feelings of hope, gratitude, and compassion, all of which have a positive effect on overall well-being. There are several types of prayer rooted in the belief that there is a higher power. This belief can provide a sense of comfort and support in

<sup>7</sup>. Delagran, L. (n.d.). *What is spirituality?* University of Minnesota.  
<https://www.takingcharge.csh.umn.edu/what-spirituality>

difficult times. A recent study found that adults who were clinically depressed who believed their prayers were heard by a concerned presence responded much better to treatment than those who did not believe.<sup>8</sup>

Individuals can be encouraged to find a spiritual community, such as a church, synagogue, temple, mosque, meditation center, or other local group that meets to discuss spiritual issues. The benefits of social support are well-documented, and having a spiritual community to turn to for fellowship can provide a sense of belonging and support.<sup>9</sup>

## Spending Time in Nature & Spirituality

Spending time in nature is cited by many individuals as a spiritual practice that contributes to their mental health.<sup>10</sup> Spirituality is defined as a dynamic and intrinsic aspect of humanity through which persons seek ultimate meaning, purpose, and transcendence and experience relationship to self, family, others, community, society, nature, and the significant or sacred.<sup>11</sup> Spiritual needs and spirituality are often mistakenly equated with religion, but

8. Delagran, L. (n.d.). *What is spirituality?* University of Minnesota. <https://www.takingcharge.csh.umn.edu/what-spirituality>
9. Delagran, L. (n.d.). *What is spirituality?* University of Minnesota. <https://www.takingcharge.csh.umn.edu/what-spirituality>
10. Yamada, A., Lukoff, D., Lim, C. S. F., & Mancuso, L. L. (2020). Integrating spirituality and mental health: Perspectives of adults receiving public mental health services in California. *Psychology of Religion and Spirituality*, 12(3), 276–287. <https://doi.org/10.1037/rel0000260>
11. Puchalski, C. M., Vitillo, R., Hull, S. K., & Reller, N. (2014). Improving the spiritual dimension of whole person care: Reaching national and international consensus. *Journal of Palliative Medicine*, 17(6), 642–656. <https://doi.org/10.1089/jpm.2014.9427>

spirituality is a broader concept. Other elements of spirituality include meaning, love, belonging, forgiveness, and connectedness.<sup>12</sup>

Spending time in nature is a powerful expression of spirituality for many individuals, offering a space for reflection, connection, and renewal. In the natural world, people often find a sense of awe and wonder that transcends the ordinary, fostering a deep awareness of something greater than themselves. This experience can be profoundly grounding and healing, particularly for those navigating mental health challenges or seeking greater emotional balance. Nature provides a unique setting where individuals can engage in contemplative practices such as walking, mindfulness, journaling, or simply being present. These moments often lead to a heightened sense of purpose, inner peace, and clarity. The rhythmic patterns of the natural environment—such as waves, birdsong, rustling leaves, or the rising and setting of the sun—can evoke feelings of harmony and connectedness that are essential to many people’s spiritual well-being.

## Supportive Relationships

Individuals should be encouraged to nurture supportive relationships with family, significant others, and friends. Relationships aren’t static – they are living, dynamic aspects of our lives that require attention and care. To benefit from strong connections with others, individuals should take charge of their relationships and devote time and energy to support them. It can be helpful to create rituals together. With busy schedules and the presence of online social media that offer the façade of real contact, it’s very easy to drift from friends. Research has found that people who deliberately make time for gatherings enjoy stronger relationships and more positive energy. An easy

12. Rudolfsson, G., Berggren, I., & da Silva, A. B. (2014). Experiences of spirituality and spiritual values in the context of nursing – An integrative review. *The Open Nursing Journal*, 8, 64–70. <https://dx.doi.org/10.2174%2F1874434601408010064>

way to do this is to create a standing ritual that you can share and that doesn't create more stress, such as talking on the telephone on Fridays or sharing a walk during lunch breaks.<sup>13</sup>

## Mindfulness

Mindfulness has been defined as, "Awareness that arises through paying attention, on purpose, in the present moment, and nonjudgmentally."

Mindfulness has also been described as, "Non-elaborative, nonjudgmental, present-centered awareness in which each thought, feeling, or sensation that arises is acknowledged and accepted as it is." Mindfulness helps us be present in our lives and gives us some control over our reactions and repetitive thought patterns. It helps us pause, get a clearer picture of a situation, and respond more skillfully. Compare your default state to mindfulness when studying for an exam in a difficult course or preparing for a clinical experience. What do you do? Do you tell yourself, "I am not good at this" or "I am going to look stupid"? Does this distract you from paying attention to studying or preparing? How might it be different if you had an open attitude with no concern or judgment about your performance? What if you directly experienced the process as it unfolded, including the challenges, anxieties, insights, and accomplishments, while acknowledging each thought or feeling and accepting it without needing to figure it out or explore it further? If practiced regularly, mindfulness helps a person start to see the habitual patterns that lead to automatic negative reactions that create stress. By observing these thoughts and emotions instead of reacting to them, a person can develop a broader perspective and can choose a more effective response.<sup>14</sup>

13. Delagran, L. (n.d.). *What is spirituality?* University of Minnesota.  
<https://www.takingcharge.csh.umn.edu/what-spirituality>

14. Delagran, L. (n.d.). *What is spirituality?* University of Minnesota.  
<https://www.takingcharge.csh.umn.edu/what-spirituality>

- ▶ Try free mindfulness activities at the [Free Mindfulness Project](#).

## Coping with Loss and Grief

In addition to assisting individuals recognize and cope with their stress and anxiety, nurses can also use this knowledge regarding coping strategies to support clients and their family members as they cope with life changes, grief, and loss that can cause emotional problems and feelings of distress.

- ▶ Review concepts and nursing care related to coping with grief and loss in the “[Grief and Loss](#)” chapter of *Open RN Nursing Fundamentals*.

### 3.4 Defense Mechanisms

When providing clients with stress management techniques and effective coping strategies, nurses must be aware of common defense mechanisms.

**Defense mechanisms** are reaction patterns used by individuals to protect themselves from anxiety that arises from stress and conflict.<sup>1</sup> Excessive use of defense mechanisms is associated with specific mental health disorders. With the exception of suppression, all other defense mechanisms are unconscious and out of the awareness of the individual. See Table 3.4 for a description of common defense mechanisms.

Table 3.4 Common Defense Mechanisms

1. American Psychological Association. (n.d.). Stressor. *APA Dictionary of Psychology*. <https://dictionary.apa.org>



Defense Mechanisms	Definitions	Examples
<b>Conversion</b>	Anxiety caused by repressed impulses and feelings are converted into a physical symptoms. <sup>2</sup>	An individual scheduled to see their therapist to discuss a past sexual assault experiences a severe headache and cancels the appointment.
<b>Denial</b>	Unpleasant thoughts, feelings, wishes, or events are ignored or excluded from conscious awareness to protect themselves from overwhelming worry or anxiety. <sup>3,4</sup>	A client recently diagnosed with cancer states there was an error in diagnosis and they don't have cancer.  Other examples include denial of a financial problem, an addiction, or a partner's infidelity.
<b>Dissociation</b>	A feeling of being disconnected from a stressful or traumatic event – or feeling that the event is not really happening – to block out mental trauma and protect the mind from too much stress. <sup>5</sup>	A person experiencing physical abuse may feel as if they are floating above their bodies observing the situation.
<b>Displacement</b>	Unconscious transfer of one's emotions or reaction from an original object to a less-threatening target to discharge tension. <sup>6</sup>	An individual who is angry with their partner kicks the family dog. An angry child breaks a toy or yells at a sibling instead of attacking their father. A frustrated employee criticizes their spouse instead of their boss. <sup>7</sup>
<b>Introjection</b>	Unconsciously incorporating the attitudes, values, and qualities of another person's personality. <sup>8</sup>	A client talks and acts like one of the nurses they admire.

2. American Psychological Association. (n.d.). Stressor. *APA Dictionary of Psychology*.  
<https://dictionary.apa.org>

3. American Psychological Association. (n.d.). Stressor. *APA Dictionary of Psychology*.  
<https://dictionary.apa.org>
4. Sissons, C. (2020). *Defense mechanisms in psychology: What are they?*  
MedicalNewsToday. <https://www.medicalnewstoday.com/articles/defense-mechanisms>
5. Sissons, C. (2020). *Defense mechanisms in psychology: What are they?*  
MedicalNewsToday. <https://www.medicalnewstoday.com/articles/defense-mechanisms>
6. American Psychological Association. (n.d.). Stressor. *APA Dictionary of Psychology*.  
<https://dictionary.apa.org>
7. American Psychological Association. (n.d.). Stressor. *APA Dictionary of Psychology*.  
<https://dictionary.apa.org>
8. American Psychological Association. (n.d.). Stressor. *APA Dictionary of Psychology*.  
<https://dictionary.apa.org>

<b>Projection</b>	A process when one attributes their individual positive or negative characteristics, affects, and impulses to another person or group. <sup>9</sup>	A person conflicted over expressing anger changes “I hate him” to “He hates me.” <sup>10</sup>
<b>Rationalization</b>	Logical reasons are given to justify unacceptable behavior to defend against feelings of guilt, maintain self-respect, and protect oneself from criticism. <sup>11</sup>	A client who is overextended on several credit cards rationalizes it is okay to buy more clothes to be in style when spending money that was set aside to pay for the monthly rent and utilities. A student caught cheating on a test rationalizes, “Everybody cheats.”
<b>Reaction Formation</b>	Unacceptable or threatening impulses are denied and consciously replaced with an opposite, acceptable impulse. <sup>12</sup>	A client who hates their mother writes in their journal that their mom is a wonderful mother.
<b>Regression</b>	A return to a prior, lower state of cognitive, emotional, or behavioral functioning when threatened with overwhelming external problems or internal conflicts. <sup>13</sup>	A child who was toilet trained reverts to wetting their pants after their parents’ divorce.
<b>Repression</b>	Painful experiences and unacceptable impulses are unconsciously excluded from consciousness as a protection against anxiety. <sup>14</sup>	A victim of incest indicates they have always hated their brother (the molester) but cannot remember why.
<b>Splitting</b>	Objects provoking anxiety and ambivalence are viewed as either all good or all bad. <sup>15</sup>	A client tells the nurse they are the most wonderful person in the world, but after the nurse enforces the unit rules with them, the client tells the nurse they are the worst person they have ever met.

9. American Psychological Association. (n.d.). Stressor. *APA Dictionary of Psychology*.  
<https://dictionary.apa.org>

<b>Suppression</b>	A conscious effort to keep disturbing thoughts and experiences out of mind or to control and inhibit the expression of unacceptable impulses and feelings. Suppression is similar to repression, <sup>16,17</sup> but it is a conscious process.	An individual has an impulse to tell their boss what they think about them and their unacceptable behavior, but the impulse is suppressed because of the need to keep the job.
<b>Sublimation</b>	Unacceptable sexual or aggressive drives are unconsciously channeled into socially acceptable modes of expression that indirectly provide some satisfaction for the original drives and protect individuals from anxiety induced by the original drive. <sup>18</sup>	An individual with an exhibitionistic impulse channels this impulse into creating dance choreography. A person with a voyeuristic urge completes scientific research and observes research subjects. An individual with an aggressive drive joins the football team. <sup>19</sup>
<b>Symbolization</b>	The substitution of a symbol for a repressed impulse, affect, or idea. <sup>20</sup>	A client unconsciously wears red clothing due a repressed impulse to physically harm someone.

10. American Psychological Association. (n.d.). Stressor. *APA Dictionary of Psychology*.  
<https://dictionary.apa.org>

11. American Psychological Association. (n.d.). Stressor. *APA Dictionary of Psychology*.  
<https://dictionary.apa.org>

12. American Psychological Association. (n.d.). Stressor. *APA Dictionary of Psychology*.  
<https://dictionary.apa.org>

13. American Psychological Association. (n.d.). Stressor. *APA Dictionary of Psychology*.  
<https://dictionary.apa.org>

14. American Psychological Association. (n.d.). Stressor. *APA Dictionary of Psychology*.  
<https://dictionary.apa.org>

15. American Psychological Association. (n.d.). Stressor. *APA Dictionary of Psychology*.  
<https://dictionary.apa.org>

16. American Psychological Association. (n.d.). Stressor. *APA Dictionary of Psychology*.  
<https://dictionary.apa.org>
17. Sissons, C. (2020). *Defense mechanisms in psychology: What are they?*  
MedicalNewsToday. <https://www.medicalnewstoday.com/articles/defense-mechanisms>
18. American Psychological Association. (n.d.). Stressor. *APA Dictionary of Psychology*.  
<https://dictionary.apa.org>
19. American Psychological Association. (n.d.). Stressor. *APA Dictionary of Psychology*.  
<https://dictionary.apa.org>
20. American Psychological Association. (n.d.). Stressor. *APA Dictionary of Psychology*.  
<https://dictionary.apa.org>

## 3.5 Crisis and Crisis Intervention

If you were asked to describe someone in crisis, what would come to your mind? Many of us might draw on traditional images of someone anxiously wringing their hands, pacing the halls, having a verbal outburst, or acting erratically. Health care professionals should be aware that crisis can be reflected in these types of behaviors, but it can also be demonstrated in various verbal and nonverbal signs. There are many potential causes of crisis, and there are four phases an individual progresses through to crisis. Nurses and other health care professionals are often the frontline care providers when an individual faces a crisis, so it is important to recognize signs of crisis, know what to assess, intervene appropriately, and evaluate crisis resolution.

### Definition of Crisis

A **crisis** can be broadly defined as the inability to cope or adapt to a stressor. Historically, the first examination of crisis and development of formal crisis intervention models occurred among psychologists in the 1960s and 1970s. Although definitions of crisis have evolved, there are central tenets related to an individual's stress management.

Consider the historical context of crisis as first formally defined in the literature by Gerald Caplan. Crisis was defined as a situation that produces psychological disequilibrium in an individual and constitutes an important problem in which they can't escape or solve with their customary problem-solving resources.<sup>1</sup> This definition emphasized the imbalance created by situation stressors.

Albert Roberts updated the concept of crisis management in more recent years to include a reflection on the level of an individual's dysfunction. He defined crisis as an acute disruption of psychological homeostasis in which one's usual coping mechanisms fail with evidence of distress and functional

1. Caplan, G. (1964). *Principles of preventive psychiatry*. Basic Books.

impairment.<sup>2</sup> A person's subjective reaction to a stressful life experience compromises their ability (or inability) to cope or function.

## Causes of Crisis

A crisis can emerge for individuals due to a variety of events. It is also important to note that events may be managed differently by different individuals. For example, a stressful stimulus occurring for Client A may not induce the same crisis response as it does for Client B. Therefore, nurses must remain vigilant and carefully monitor each client for signs of emerging crisis.

A crisis commonly occurs when individuals experience some sort of significant life event. These events may be unanticipated, but that is not always the case. An example of anticipated life events that may cause a crisis include the birth of a baby. For example, the birth (although expected) can result in a crisis for some individuals as they struggle to cope with and adapt to this major life change. Predictable, routine schedules from before the child was born are often completely upended. Priorities shift to an unyielding focus on the needs of the new baby. Although many individuals welcome this change and cope effectively with the associated life changes, it can induce crises in those who are unprepared for such a change.

Crisis situations are more commonly associated with unexpected life events. Individuals who experience a newly diagnosed critical or life-altering illness are at risk for experiencing a crisis. For example, a client experiencing a life-threatening myocardial infarction or receiving a new diagnosis of cancer may experience a crisis. Additionally, the crisis may be experienced by family and loved ones of the client as well. Nurses should be aware that crisis

2. Roberts, A. R. (2005). Bridging the past and present to the future of crisis intervention and crisis management. In A. R. Roberts (Ed.), *Crisis intervention handbook: Assessment, treatment, and research* (3rd ed.). Oxford University Press. pp. 3-34.

intervention and the need for additional support may occur in these types of situations and often extend beyond the needs of the individual client.

Other events that may result in crisis development include stressors such as the loss of a job, loss of one's home, divorce, or death of a loved one. It is important to be aware that clustering of multiple events can also cause stress to build sequentially so that individuals can no longer successfully manage and adapt, resulting in crisis.

## Categories of Crises

Due to a variety of stimuli that can cause the emergence of a crisis, crises can be categorized to help nurses and health care providers understand the crisis experience and the resources that may be most beneficial for assisting the client and their family members. Crises can be characterized into one of three categories: maturational, situational, or social crisis. Table 3.5a explains characteristics of the different categories of crises and provides examples of stressors associated with that category.

Table 3.5a Categories of Crises



Category	Characteristics	Examples
<b>Maturation Crisis (also known as Developmental crisis)</b>	<ul style="list-style-type: none"> <li>• The result of normal processes of growth and development.</li> <li>• Commonly occurs at specific developmental periods of life.</li> <li>• It is predictable in nature and normally occurs as a part of life.</li> <li>• An individual is vulnerable based on their equilibrium.</li> </ul>	<ul style="list-style-type: none"> <li>• Birth</li> <li>• Adolescence</li> <li>• Marriage</li> <li>• Death</li> </ul>
<b>Situational Crisis</b>	<ul style="list-style-type: none"> <li>• An unexpected personal stressful event occurs with little advance warning.</li> <li>• It is less predictable in nature.</li> <li>• The event threatens an individual's equilibrium.</li> </ul>	<ul style="list-style-type: none"> <li>• Accident</li> <li>• Illness or serious injury of self or family member</li> <li>• Loss of a job</li> <li>• Bankruptcy</li> <li>• Relocation/geographical move</li> <li>• Divorce</li> </ul>
<b>Social Crisis (also known as Adventitious crisis)</b>	<ul style="list-style-type: none"> <li>• An event that is uncommon or unanticipated.</li> <li>• The event often involves multiple losses or extensive losses.</li> <li>• It can occur due to a major natural or man-made event.</li> <li>• It is unpredictable in nature.</li> <li>• The event poses a severe threat to an individual's equilibrium.</li> </ul>	<ul style="list-style-type: none"> <li>• Flood</li> <li>• Fire</li> <li>• Tornado</li> <li>• Hurricane</li> <li>• Earthquake</li> <li>• War</li> <li>• Riot</li> <li>• Violent crime</li> </ul>

## Phases of Crisis

The process of crisis development can be described as four distinct phases. The phases progress from initial exposure to the stressor, to tension escalation, to an eventual breaking point. These phases reflect a sequential progression in which resource utilization and intervention are critical for assisting a client in crisis. Table 3.5b describes the various phases of crisis, their defining characteristics, and associated signs and symptoms that individuals may experience as they progress through each phase.

Table 3.5b Crisis Phases<sup>3,4</sup>

3. Caplan, G. (1964). *Principles of preventive psychiatry*. Basic Books.
4. Centers for Disease Control and Prevention. (2018). *The National Institute for Occupational Health and Safety*. <https://www.cdc.gov/niosh/>

Crisis Phase	Defining Characteristics	Signs and Symptoms
<b>Phase 1:</b>  <b>Normal Stress &amp; Anxiety</b>	<p>Exposure to a precipitating stressor.</p> <p>Stressors may be considered minor annoyances and inconveniences of everyday life.</p>	<p>Anxiety levels or the stress response begin to elevate.</p> <p>Individuals try using previously successful problem-solving techniques to attempt resolution of the stressor.</p> <p>Individuals are rational and in control of their behavior and emotions.</p>
<b>Phase 2:</b>  <b>Rising Anxiety Level</b>	<p>Problem-solving techniques do not relieve the stressor.</p> <p>Use of past coping strategies are not successful.</p>	<p>Anxiety levels increase and individuals experience increased discomfort.</p> <p>Feelings of helplessness, confusion, and disorganized thinking may occur. Individuals may complain of “feeling lost” in how to proceed.</p> <p>Individuals may experience elevated heart rate and respiration rate. Their voice pitch may be higher with a more rapid speech pattern.</p> <p>Nervous habits such as finger or foot tapping may occur.</p>
<b>Phase 3:</b>  <b>Severe Level of Stress and Anxiety</b>	<p>Individuals use all possible internal and external resources.</p> <p>Problems are explored from different perspectives, and new problem-solving techniques are attempted.</p>	<p>Equilibrium may be restored if new problem-solving approaches are successful. Individuals experience decreased anxiety if resolution occurs.</p> <p>If new problem-solving techniques are not successful, the level of anxiety worsens, and functioning is impaired as the stressor continues to impact the individual.</p> <p>Capacity to reason becomes significantly diminished, and behaviors become more disruptive.</p> <p>Communication processes may include yelling and swearing. Individuals may become very argumentative or use threats.</p> <p>Individuals may pace; clench their fists; perspire heavily; or demonstrate rapid, shallow, panting breaths.</p>

<b>Phase 4: Crisis</b>	If resolution is not achieved, tension escalates to a critical breaking point.	<p>Individuals experience unbearable anxiety, increased feelings of panic, and disordered thinking processes. There is an urgent need to end emotional discomfort. Many cognitive functions are impaired as the crisis event becomes thought consuming. Emotions are labile, and some clients may experience psychotic thinking.</p> <p>*It is important to note that some individuals at this level of crisis may be a danger to themselves and others.</p>
----------------------------	--	--

## Crisis Assessment

Nurses must be aware of the potential impact of stressors for their clients and the ways in which they may manifest in a crisis. The first step in assessing for a crisis occurs with the basic establishment of a therapeutic nurse-client relationship. Understanding who your client is, what is occurring in their life, what resources are available to them, and their individual beliefs, supports, and general demeanor can help a nurse determine if a client is at risk for ineffective coping and possible progression to crisis.

Crisis symptoms can manifest in various ways and can vary significantly depending on the underlying condition and context. Nurses should carefully monitor for signs of the progression through the phases of crisis such as the following:

- Escalating anxiety
- Denial
- Confusion or disordered thinking
- Anger and hostility
- Helplessness and withdrawal
- Inefficiency
- Hopelessness and depression
- Steps toward resolution and reorganization
- Physiological symptoms such as headache or shortness of breath

When a nurse identifies these signs in a client or their family members, it is important to carefully explore the symptoms exhibited and the potential

stressors. Collecting information regarding the severity of the stress response, the individual's or family's resources, and the crisis phase can help guide the nurse and health care team toward appropriate intervention.

## Crisis Interventions

Crisis intervention is an important role for the nurse and health care team to assist clients and families toward crisis resolution. Resources are employed, and interventions are implemented to therapeutically assist the individual in whatever phase of crisis they are experiencing. A crisis state is time-limited, usually lasting several days but no longer than four to six weeks. Depending on the stage of the crisis, various strategies and resources are used.

The goals of crisis intervention are the following:

- Ensure safety: Identify, assess, and intervene
- Provide support: Return the individual to a prior level of functioning as quickly as possible
- Lessen negative impact on future mental health, manage symptoms

During the crisis intervention process, new skills and coping strategies are acquired, resulting in change. Various factors can influence an individual's ability to resolve a crisis and return to equilibrium, such as realistic perception of an event, adequate situational support, and adequate coping strategies to respond to a problem. Nurses can implement strategies to reinforce these factors.

## Strategies for Crisis Phase 1 and 2

Table 3.5c describes strategies and techniques for early phases of a crisis that can help guide the individual toward crisis resolution.

## Table 3.5c Phase 1 & 2 Early Crisis Intervention Strategies<sup>5</sup>

5. Centers for Disease Control and Prevention. (2018). *The National Institute for Occupational Health and Safety*. <https://www.cdc.gov/niosh/>

Verbal	Strategies	Examples
Therapeutic use of words holds significant power to defuse the stress response.	Encourage the person to express their thoughts and concerns.	<p>"I understand how hard this must be for you."</p> <p>"Can you tell me more about how you are feeling?"</p>
Be attuned to the individual's tone of voice and body language.	Use a shared problem-solving approach. Avoid being defensive.	"I understand your feelings of frustration. How can we correct this problem?"
Be attuned to word choice.	Use empathetic inquiry.	"You seem to be upset. Tell me more about what is bothering you."
Nonverbal	Strategies	Example
Be aware of your nonverbal messages and be in control of your body positions.	Be calm and act calm. Invite the client to sit to help them calm down and demonstrate you are calm.	Maintain nonthreatening eye contact, smile, and keep hands open and visible.
	Listen.	Nod your head to demonstrate that you are engaged with the individual.
	Respect personal space.	Maintain distance and avoid touching an individual who is upset.
	Approach the client from an angle or from the side.	Avoid directly approaching an individual, as it can feel confrontational.
	Avoid threatening gestures.	Avoid finger pointing or crossing arms.

	Demonstrate respect.	Mirror the individual's nonverbal messaging. Avoid laughing or joking.
--	----------------------	---

## Strategies for Crisis Phase 3

If an individual continues to progress in severity to higher levels of crisis, the previously identified verbal and nonverbal interventions for Phase 1 and Phase 2 may be received with a variability of success. For example, for a receptive individual who is still in relative control of their emotions, the verbal and nonverbal interventions may still be well-received. However, if an individual has progressed to Phase 3 with emotional lability, the nurse must recognize this escalation and take additional measures to protect oneself. If an individual demonstrates loss of problem-solving ability or the loss of control, the nurse must take measures to ensure safety for themselves and others in all interactions with the client. This can be accomplished by calling security or other staff to assist when engaging with the client. It is important to always note the location of exits in the client's room and ensure the client is never between the nurse and the exit. Rapid response devices may be worn, and nurses should feel comfortable using them if a situation begins to escalate.

Verbal cues can still hold significant power even in a late phase of crisis. The



nurse should provide direct cues to an escalating client such as, “Mr. Andrews, please sit down and take a few deep breaths. I understand you are angry. You need to gain control of your emotions, or I will have to call security for assistance.” This strategy is an example of limit-setting that can be helpful for de-escalating the situation and defusing tension. Setting limits is important for providing behavioral guidance to a client who is escalating, but it is very different from making threats. Limit-setting describes the desired behavior whereas making threats is nontherapeutic. See additional examples contrasting limit-setting and making threats in the following box.<sup>6</sup>

### Examples of Limit-Setting Versus Making Threats<sup>7</sup>

- **Threat:** “If you don’t stop, I’m going to call security!”
- **Limit-Setting:** “Please sit down. I will have to call for assistance if you can’t control your emotions.”
  
- **Threat:** “If you keep pushing the call button over and over like that, I won’t help you.”
- **Limit-Setting:** “Ms. Ferris, I will come as soon as I am able when you need assistance, but please give me a chance to get to your room.”

6. Centers for Disease Control and Prevention. (2018). *The National Institute for Occupational Health and Safety*. <https://www.cdc.gov/niosh/>

7. Centers for Disease Control and Prevention. (2018). *The National Institute for Occupational Health and Safety*. <https://www.cdc.gov/niosh/>

- **Threat:** “That type of behavior won’t be tolerated!”
- **Limit-Setting:** “Mr. Barron, please stop yelling and screaming at me. I am here to help you.”

## Strategies for Crisis Phase 4

A person who is experiencing an elevated phase of crisis is not likely to be in control of their emotions, cognitive processes, or behavior. It is important to give them space so they don’t feel trapped. Many times these individuals are not responsive to verbal intervention and are solely focused on their own fear, anger, frustration, or despair. Don’t try to argue or reason with them. Individuals in Phase 4 of crisis often experience physical manifestations such as rapid heart rate, rapid breathing, and pacing.

If you can’t successfully de-escalate an individual who is becoming increasingly more agitated, seek assistance. If you don’t believe there is an immediate danger, call a psychiatrist, psychiatric-mental health nurse specialist, therapist, case manager, social worker, or family physician who is familiar with the person’s history. The professional can assess the situation and provide guidance, such as scheduling an appointment or admitting the person to the hospital. If you can’t reach someone and the situation continues to escalate, consider calling your county mental health crisis unit, crisis response team, or other similar contacts. If the situation is life-threatening or if serious property damage is occurring, call 911 and ask for immediate assistance. When you call 911, tell them someone is experiencing a mental health crisis and explain the nature of the emergency, your relationship to the person in crisis, and whether there are weapons involved. Ask the 911 operator

to send someone trained to work with people with mental illnesses such as a **Crisis Intervention Training (CIT) officer.**<sup>8</sup>

A nurse who assesses a client in this phase should observe the client's behaviors and take measures to ensure the client and others remain safe. A person who is out of control may require physical or chemical restraints to be safe. Nurses must be aware of organizational policies and procedures, as well as documentation required for implementing restraints, if the client's or others' safety is in jeopardy. Read more about ANA guidelines on using restraints in the "[Client Rights](#)" section of the "Legal and Ethical Considerations in Mental Health Care" chapter and information on safely implementing restraints in the "[Workplace Violence](#)" section of the "Trauma, Abuse, and Violence" chapter.

- ▶ Review guidelines for safe implementation of restraints in the "[Restraints](#)" section of *Open RN Nursing Fundamentals*, 2e.

## Crisis Resources

Depending on the type of stressors and the severity of the crisis experienced, there are a variety of resources that can be offered to clients and their loved ones. Nurses should be aware of community and organizational resources that are available in their practice settings. Support groups, hotlines, shelters, counseling services, and other community resources like the Red Cross may

8. Brister, T. (2018). *Navigating a mental health crisis: A NAMI resource guide for those experiencing a mental health emergency*. National Alliance on Mental Illness. [https://www.nami.org/Support-Education/Publications-Reports/Guides/Navigating-a-Mental-Health-Crisis/Navigating-A-Mental-Health-Crisis?utm\\_source=website&utm\\_medium=cta&utm\\_campaign=crisisguide](https://www.nami.org/Support-Education/Publications-Reports/Guides/Navigating-a-Mental-Health-Crisis/Navigating-A-Mental-Health-Crisis?utm_source=website&utm_medium=cta&utm_campaign=crisisguide)

be helpful. Read more about potential national and local resources in the following box.

### **Mental Health Crisis Resources**

- ▶ [NAMI: National Alliance on Mental Health](#)
- ▶ [ADRC of Central Wisconsin](#)
- ▶ [Wisconsin County Crisis Lines](#)
- ▶ [Wisconsin Suicide & Crisis Hotlines](#)

## **Mental Health Crisis**

When an individual is diagnosed with a mental health disorder, the potential for crisis is always present. Risk of suicide is always a priority concern for people with mental health disorders in crisis. Any talk of suicide should always be taken seriously. Most people who attempt suicide have given some warning. If someone has attempted suicide before, the risk is even greater. Read more about assessing suicide risk in the “[Establishing Safety](#)” section of Chapter 1. Encouraging someone who is having suicidal thoughts to get help is a safety priority.

Common signs that a mental health crisis is developing are as follows:

- Inability to perform daily tasks like bathing, brushing teeth, brushing hair, or changing clothes
- Rapid mood swings, increased energy level, inability to stay still, pacing, suddenly depressed or withdrawn, or suddenly happy or calm after period of depression
- Increased agitation with verbal threats; violent, out-of-control behavior or destruction of property
- Abusive behavior to self and others, including substance misuse or self-harm (cutting)
- Isolation from school, work, family, or friends

- Loss of touch with reality (psychosis) – unable to recognize family or friends, confused, doesn't understand what people are saying, hearing voices, or seeing things that aren't there
- Paranoia

Clients with mental illness and their loved ones need information for what to do if they are experiencing a crisis. *Navigating a Mental Health Crisis: A NAMI Resource Guide for Those Experiencing a Mental Health Emergency* provides important, potentially life-saving information for people experiencing mental health crises and their loved ones. It outlines what can contribute to a crisis, warning signs that a crisis is emerging, strategies to help de-escalate a crisis, and available resources.

- ▶ Read NAMI's [\*Navigating a Mental Health Crisis: A NAMI Resource Guide for Those Experiencing a Mental Health Emergency\*](#).

## 3.6 Applying the Nursing Process to Stress and Coping

This section will review the nursing process as it applies to stress and coping.

### Assessments (Recognize Cues)

Here are several nursing assessments used to determine an individual's response to stress and their strategies for stress management and coping:

- Recognize nonverbal cues of physical or psychological stress
- Assess for environmental stressors affecting client care
- Assess for signs of abuse or neglect
- Assess client's ability to cope with life changes
- Assess family dynamics
- Assess the potential for violence
- Assess client's support systems and available resources
- Assess client's ability to adapt to temporary/permanent role changes
- Assess client's reaction to a diagnosis of acute or chronic mental illness (e.g., rationalization, hopefulness, anger)
- Assess constructive use of defense mechanisms by a client
- Assess if the client has successfully adapted to situational role changes (e.g., accept dependency on others)
- Assess client's ability to cope with end-of-life interventions
- Recognize the need for psychosocial support to the family/caregiver
- Assess clients for maladaptive coping such as substance abuse
- Identify a client in crisis

### Diagnoses (Analyze Cues)

Nursing diagnoses related to stress and coping are *Stress Overload* and *Ineffective Coping*. See Table 3.6 to compare the definitions and defining characteristics for these nursing diagnoses.

Table 3.6 Stress and Coping Nursing Diagnoses

Nursing Diagnosis	Definition	Selected Defining Characteristics
<b>Stress Overload</b>	Excessive amounts and types of demands that require action.	<ul style="list-style-type: none"> <li>• Excessive stress</li> <li>• Impaired decision-making</li> <li>• Impaired functioning</li> <li>• Increase in anger</li> <li>• Increased impatience</li> </ul>
<b>Ineffective Coping</b>	A pattern of invalid appraisal of stressors, with cognitive and/or behavioral efforts, that fails to manage demands related to well-being.	<ul style="list-style-type: none"> <li>• Alteration in concentration</li> <li>• Alteration in sleep pattern</li> <li>• Change in communication pattern</li> <li>• Fatigue</li> <li>• Inability to ask for help</li> <li>• Inability to deal with a situation</li> <li>• Ineffective coping strategies</li> <li>• Insufficient social support</li> <li>• Substance misuse</li> </ul>

## Outcomes Identification (Generate Solutions)

An outcome is a measurable behavior demonstrated by the client in response to nursing interventions. Outcomes should be identified before nursing interventions are planned. Outcomes identification includes setting short- and long-term goals and then creating specific expected outcome statements for each nursing diagnosis. Goals are broad, general statements, and outcomes are specific and measurable. Expected outcomes are statements of measurable action for the client within a specific time frame that are responsive to nursing interventions.

Expected outcome statements should contain five components easily remembered using the “SMART” mnemonic:

- Specific
- Measurable
- Attainable/Action oriented
- Relevant/Realistic
- Time frame

An example of a SMART outcome related to *Stress Overload* is, “The client will identify two stressors that can be modified or eliminated by the end of the week.”

An example of a SMART outcome related to *Ineffective Coping* is, “The client will identify three preferred coping strategies to implement by the end of the week.”

Read more information about establishing SMART outcome statements in the [“Outcome Identification”](#) section of Chapter 4.



## Planning Interventions Related to Stress and Coping (Generate Solutions)

Common nursing interventions that are implemented to facilitate effective coping in their clients include the following:

- Implement measures to reduce environmental stressors
- Teach clients about stress management techniques and coping strategies
- Provide caring interventions for a client experiencing grief or loss, as well as resources to adjust to loss/bereavement
- Identify the client in crisis and tailor crisis intervention strategies to assist them to cope
- Guide the client to resources for recovery from crisis (i.e., social supports)

## Implementation (Take Action)

When implementing nursing interventions to enhance client coping, it is important to recognize signs of a crisis and maintain safety for the client, oneself, and others. Review signs of a client in crisis and crisis intervention strategies in the “[Crisis and Crisis Intervention](#)” section of this chapter.

## Evaluation (Evaluate Outcome)

After implementing individualized interventions for a client, it is vital to evaluate their effectiveness. Review the specific SMART outcomes and deadlines that have been established for a client and determine if interventions were effective in meeting these outcomes or if the care plan requires modification.

## 3.7 Learning Activities



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=165#h5p-26>

1



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=165#h5p-12>

2



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=165#h5p-13>

1. “MH Stress, Coping, and Crisis Intervention Glossary Cards ” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

2. “MH Stress, Coping, and Crisis Drag and Drop ” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

- ▶ Test your clinical judgment with an NCLEX Next Generation-style question: [Chapter 3, Assignment 1](#)<sup>4</sup>



- ▶ Test your clinical judgment with an NCLEX Next Generation-style case study: [Chapter 3, Case Study 1](#)<sup>5</sup>



3. “MH Stress, Coping, and Crisis Question Set 1 ” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

4. “MH Next-Generation Style Question Stress, Coping, Crisis ” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

5. “MH Next-Generation Style Stress, Coping and Crisis Intervention ” by Kellea Ewen is licensed under [CC BY-NC 4.0](#)

### III Glossary

**Adaptive coping strategies:** Coping strategies, including problem-focused coping and emotion-focused coping.

**Adverse childhood experiences:** Potentially traumatic events that occur in childhood such as sexual abuse, physical abuse, emotional abuse, physical neglect, emotional neglect, parental loss, or parental separation before the child is 18 years old.

**Coping:** Cognitive and behavioral efforts made to master, tolerate, or reduce external and internal demands and conflicts.<sup>1</sup>

**Coping strategies:** An action, series of actions, or a thought process used in meeting a stressful or unpleasant situation or in modifying one's reaction to such a situation.<sup>2</sup>

**Crisis:** The inability to cope or adapt to a stressor.

**Crisis Intervention Training Officer:** A law enforcement or public safety professional who is specially trained to respond to individuals experiencing a mental health crisis or other behavioral emergencies.

**Defense mechanisms:** Unconscious reaction patterns used by individuals to protect themselves from anxiety that arises from stress and conflict.<sup>3</sup>

1. Amnie, A. G. (2018). Emerging themes in coping with lifetime stress and implication for stress management education. *SAGE Open Medicine*, 6. <https://doi.org/10.1177%2F2050312118782545>
2. American Psychological Association. (n.d.). Stressor. *APA Dictionary of Psychology*. <https://dictionary.apa.org>
3. American Psychological Association. (n.d.). Stressor. *APA Dictionary of Psychology*. <https://dictionary.apa.org>

**Emotion-focused coping:** Adaptive coping strategies such as practicing mindfulness, meditation, and yoga; using humor and jokes; seeking spiritual or religious pursuits; engaging in physical activity or breathing exercises; and seeking social support.

**Maladaptive coping responses:** Ineffective responses to stressors such as avoidance of the stressful condition, withdrawal from a stressful environment, disengagement from stressful relationships, and misuse of drugs and/or alcohol.

**Maturational crisis:** A type of developmental crisis that occurs during normal life transitions or stages of growth.

**Problem-focused coping:** Adaptive coping strategies that typically focus on seeking treatment such as counseling or cognitive behavioral therapy.

**Stress response:** The body's physiological response to a real or perceived stressor. For example, the respiratory, cardiovascular, and musculoskeletal systems are activated to breathe rapidly, stimulate the heart to pump more blood, dilate the blood vessels, and increase blood pressure to deliver more oxygenated blood to the muscles.

**Stressors:** Any internal or external event, force, or condition that results in physical or emotional stress.



PART IV

CHAPTER 4 APPLICATION OF THE NURSING PROCESS TO MENTAL  
HEALTH CARE





### Learning Objectives

- Apply the nursing process to mental health care
- Apply the subcategories of the *Implementation* standard of care to mental health care
- Explain various mental health treatment approaches
- Promote a therapeutic environment
- Compare NCLEX Next Generation terminology to the nursing process

**Psychiatric-mental health nursing** is, “The nursing practice specialty committed to promoting mental health through the assessment, diagnosis, and treatment of behavioral problems, mental disorders, and comorbid conditions across the life span. Psychiatric-mental health nursing intervention is an art and a science, employing a purposeful use of self and a wide range of nursing, psychosocial, and neurobiological evidence to produce effective outcomes.”<sup>1</sup>

In 2014, the American Psychiatric Nurses Association (APNA) and the International Society of Psychiatric-Mental Health Nurses (ISPN) published the *Psychiatric-Mental Health Nursing: Scope and Standards of Practice* resource in alignment with the second edition of the ANA's *Nursing: Scope*

1. American Nurses Association, American Psychiatric Nurses Association, and International Society of Psychiatric-Mental Health Nurses. (2014). *Psychiatric-mental health nursing: Scope and standards of practice* (2nd ed.). Nursebooks.org

and *Standard of Practice Nursing*. The *Psychiatric-Mental Health Nursing: Scope and Standards of Practice* resource guides psychiatric-mental health nurses in the application of their professional skills and responsibilities and should be reviewed in conjunction with state Board of Nursing policies and practices that govern the practice of nursing.<sup>2</sup> The Standards of Practice for Psychiatric-Mental Health Nursing mirror the ANA Standards of Professional Nursing Practice of *Assessment, Diagnosis, Outcome Identification, Planning, Implementation, and Evaluation*, but also have additional competencies for Psychiatric-Mental Health Registered Nurse Specialists (PMH-RNs) and Advanced Practice Registered Nurse Specialists (PMH-APRNs) and additional components for the *Implementation* standard of care.

This chapter will review how nurses apply the nursing process and the ANA Standards of Professional Nursing Practice to clients experiencing a mental health condition. Assessments, nursing diagnoses, expected outcomes, and interventions pertaining to mental health will be reviewed while incorporating life span and cultural considerations. For assessments, nursing diagnoses, expected outcomes, and interventions related to specific mental health conditions, see each corresponding “disorder” chapter.

2. American Nurses Association, American Psychiatric Nurses Association, and International Society of Psychiatric-Mental Health Nurses. (2014). *Psychiatric-mental health nursing: Scope and standards of practice* (2nd ed.). Nursebooks.org

## 4.2 Applying the Nursing Process

The nursing process is a critical thinking model based on a systematic approach to client-centered care. Nurses use the **nursing process** to perform clinical reasoning and make clinical judgments when providing client care. The nursing process is based on the Standards of Professional Nursing Practice established by the American Nurses Association (ANA). These standards are authoritative statements of the actions and behaviors that all registered nurses, regardless of role, population, specialty, and setting, are expected to perform competently.<sup>1</sup>

The mnemonic **ADOPIE** is an easy way to remember the six ANA standards regarding the nursing process. Each letter refers to one of the six components of the nursing process: *Assessment, Diagnosis, Outcomes Identification, Planning, Implementation, and Evaluation*. The nursing process is a continuous, cyclic process that is constantly adapting to the client's current health status. See Figure 4.1<sup>2</sup> for an illustration of the nursing process.

1. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

2. "The Nursing Process" by Kim Ernstmeier at [Chippewa Valley Technical College](#) is licensed under [CC BY 4.0](#)




Figure 4.1 The Nursing Process

- ▶ Review using the nursing process in the “[Nursing Process](#)” chapter in *Open RN Nursing Fundamentals, 2e*.

## 4.3 Assessment

The *Assessment* Standard of Practice established by the American Nurses Association (ANA) states, “The registered nurse collects pertinent data to the health care and information relative to the health care consumer’s health or the situation.”<sup>1</sup> Review the competencies for the *Assessment* Standard of Practice for registered nurses in the following box.

### **ANA’s Assessment Competencies<sup>2</sup>**

The registered nurse:

- Creates the safest environment possible for conducting assessments.
- Collects pertinent data related to health and quality of life in a systematic, ongoing manner, with compassion and respect for the wholeness, the inherent dignity, worth, and unique attributes of every person, including, but not limited to, demographics, environmental and occupational exposures, social determinants of health, health disparities, physical, functional, psychosocial, emotional, cognitive, spiritual/transpersonal, sexual, sociocultural, age-related, environmental, and lifestyle/economic assessments.
- Utilizes a health and wellness model of assessment that incorporates integrative approaches to data collection and

1. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

2. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

honors the whole person.

- Recognizes the health care consumer or designated person as the decision-maker regarding their own health.
- Explores the health care consumer's culture, values, preferences, expressed and unexpressed needs, and knowledge of the health care situation.
- Assesses the impact of family dynamics on the health care consumer's health and wellness.
- Identifies enhancements and barriers to effective communication based on personal, cognitive, physiological, psychosocial, literacy, financial, and cultural considerations.
- Engages the health care consumer, family, significant others, and interprofessional team members in holistic, culturally sensitive data collection.
- Integrates knowledge from current local, regional, national, and global health initiatives and environmental factors into the assessment process.
  - State and local departments of health
  - ▶ [World Health Organization](#)
  - ▶ [World Health Organization health topics](#)
  - ▶ [Healthy People](#)
  - ▶ [Centers for Disease Control and Prevention](#)
- Prioritizes data collection based on the health care consumer's immediate condition, the anticipated needs of the health care consumer or situation, or both.
- Uses evidence-based assessment techniques and available data and information to identify patterns and variances in the consumer's health.
- Remains knowledgeable about constantly changing technologies that impact the assessment process (e.g.,

telehealth, artificial intelligence).

- Analyzes assessment data to identify patterns, trends, and situations that impact the person's health and wellness.
- Validates the analysis with the health care consumer.
- Documents data accurately and makes accessible to the interprofessional team in a timely manner.
- Communicates changes in a person's condition to the interprofessional team.
- Applies the provisions of the ANA Code of Ethics, legal guidelines, and policies to the collection, maintenance, use, and dissemination of data and information.
- Recognizes the impact of one's own personal attitudes, values, beliefs, and biases on the assessment process.

Nursing assessments related to mental health disorders differ from physiological assessments with a greater focus on collecting subjective data. For example, prior to administering a cardiac medication to a client with a heart condition, a nurse will assess objective data such as blood pressure and an apical heart rate to determine the effectiveness of the medication treatment. However, prior to administering an antidepressant, a nurse uses therapeutic communication to ask questions and gather subjective data about how the client is feeling to determine the effectiveness of the medication. The nurse will also observe client behaviors, speech, mood, and thought processes as part of the assessment.

As a nurse, you cannot directly measure a neurotransmitter to determine the effects of the medication, but you can ask questions to determine how your client is feeling emotionally and perceiving the world, which are influenced by neurotransmitter levels. An example of a nurse using therapeutic communication to perform subjective assessment is, "Tell me more about how you are feeling today." The nurse may also use general survey techniques such as simply observing the client to assess for cues of behavior. Examples of

objective data collected by a general survey could be assessing the client's mood, hygiene, appearance, or movement.

Recall the mental health continuum introduced in the “[Foundational Mental Health Concepts](#)” chapter (see Figure 4.2<sup>3</sup>). Nurses in any setting holistically assess their clients' physical, emotional, and mental health, as well as any impairments impacting their functioning. They must recognize subtle cues of undiagnosed or poorly managed physical and mental disorders and follow up appropriately with other members of the interprofessional health care team.




Figure 4.2 Mental Health Continuum (Used with permission.)

When assessing a client's mental health, the nurse incorporates a variety of assessments, in addition to the traditional physical examination. Assessments may include the following:

- Performing a mental status examination

3. “continuum.jpg” by [University of Michigan](#) is used with permission. Access the original at <https://hr.umich.edu/benefits-wellness/health-well-being/mhealthy/faculty-staff-well-being/mental-emotional-health/mental-emotional-health-classes-training-events/online-tutorial-supervisors/section-1-what-you-need-know-about-mental-health-problems-substance-misuse>.



- Completing a psychosocial assessment including:
  - Reviewing the client's reason for seeking healthcare
  - Screening the client for suicidal ideation, exposure to trauma or violence, signs of self injury, and substance misuse
  - Complete a cultural assessment, spiritual assessment, and assess client family dynamics
  - Use of psychotropic medications (drugs that treat psychiatric symptoms) and/or other medications that can cause psychiatric symptoms as side effects
  - History of mental disorders, family history of mental illness, and previous hospitalizations
  - Educational background, occupational background
  - Coping mechanisms and functional ability
- Reviewing specific laboratory results related to the client's use of psychotropic and other medications
- Incorporating life span and developmental considerations

## Mental Status Examination

A **mental status examination** assesses a client's level of consciousness and orientation, appearance and general behavior, speech, motor activity, affect and mood, thought and perception, attitude and insight, and cognitive abilities. The examiner should also monitor their personal reaction to a client when performing a mental status examination. For example, the examiner should be aware of their own facial expressions in order to not influence a client's reaction or willingness to respond to questions. The structured components of a mental status examination are outlined in Table 4.3 and further described in the following subsections.

Registered nurses must use effective clinical interviewing skills while performing a mental status assessment and developing a therapeutic nurse-client relationship.<sup>4</sup> Read more about establishing a therapeutic nurse-client

4. American Nurses Association, American Psychiatric Nurses Association, and

relationship in the “[Therapeutic Communication and the Nurse-Client Relationship](#)” chapter. Assessing a client with a suspected or previously diagnosed mental health disorder focuses on both verbal and nonverbal assessments. New assessment findings are compared to the baseline admission findings to determine if the client’s condition is improving, worsening, or remaining the same.

When conducting a focused assessment on a client’s mental health, the mental status examination is a priority component of the overall assessment. A successful nurse develops a style in which the bulk of the mental status examination is performed through unstructured observations made during the routine physical examination, also referred to as the “general survey.” When the nurse recognizes cues of possible mental health disorders, such as aberrant behavior or difficulties in day-to-day functioning, then a focused mental status examination should be completed.

► Read about the components of a general survey in the “[General Survey](#)” chapter of *Open RN Nursing Skills*.

Table 4.3 Mental Status Examination<sup>5</sup>

International Society of Psychiatric-Mental Health Nurses. (2014). *Psychiatric-Mental Health Nursing: Scope and Standards of Practice* (2nd ed.). Nursebooks.org

5. Martin, D. C. (1990). The mental status examination. In Walker, H. K., Hall, W. D., Hurst, J. W., (Eds.), *Clinical methods: The history, physical, and laboratory examinations*. (3rd ed.). Butterworths. <https://www.ncbi.nlm.nih.gov/books/NBK320/>

Assessment	Expected Findings/Optimal Level of Functioning	Unexpected Findings/Impaired Functioning
<b>Signs of Distress</b>	<ul style="list-style-type: none"> <li>• Calm and comfortable with no signs of distress</li> </ul>	<ul style="list-style-type: none"> <li>• Unresponsive</li> <li>• Difficulty breathing</li> <li>• Chest pain</li> <li>• New onset of confusion</li> <li>• Moaning</li> <li>• Grimacing</li> </ul>
<b>Level of Consciousness and Orientation</b>	<ul style="list-style-type: none"> <li>• Alert</li> <li>• Oriented to person, place, and time</li> <li>• Aware of the situation</li> </ul>	<ul style="list-style-type: none"> <li>• Unable to provide name, location, or day</li> <li>• Clouded consciousness</li> <li>• Delirium</li> <li>• Obtundation</li> <li>• Stupor</li> <li>• Coma</li> </ul>
<b>Appearance and General Behavior</b>	<ul style="list-style-type: none"> <li>• Appears stated age</li> <li>• Well-groomed</li> <li>• Dressed appropriately for the weather and situation</li> <li>• Erect posture</li> <li>• Good oral hygiene</li> <li>• Culturally appropriate eye contact</li> <li>• Socializes with others</li> <li>• No threatening behaviors</li> </ul>	<ul style="list-style-type: none"> <li>• Appears older than stated age</li> <li>• Unkempt or poor hygiene</li> <li>• Not dressed appropriately for the weather and/or situation</li> <li>• Slumped posture</li> <li>• Poor eye contact</li> <li>• Does not socialize with others</li> <li>• Demonstrates threatening behavior</li> </ul>

<b>Speech</b>	<ul style="list-style-type: none"> <li>• Exhibiting spontaneous speech</li> <li>• Even speech rate, rhythm, and tone</li> <li>• Responds to verbal questions</li> <li>• Speech is clear and understandable</li> <li>• Follows instructions appropriately for developmental level</li> </ul>	<ul style="list-style-type: none"> <li>• Does not respond to verbal questions</li> <li>• Does not follow instructions appropriately for development level</li> <li>• Speech is unclear</li> <li>• Rapid or pressured speech</li> <li>• Halting speech</li> <li>• Brief or reduced replies</li> </ul>
<b>Motor Activity</b>	<ul style="list-style-type: none"> <li>• Good balance</li> <li>• Moves extremities equally bilaterally</li> <li>• Smooth gait</li> </ul>	<ul style="list-style-type: none"> <li>• Poor balance</li> <li>• Uneven gait</li> <li>• Slow movements</li> <li>• Lack of spontaneous movement</li> <li>• Motor restlessness</li> <li>• Repetitive movements</li> <li>• Tremors</li> <li>• Pacing</li> <li>• Uncontrolled, involuntary movement</li> </ul>

<b>Affect and Mood</b>	<ul style="list-style-type: none"> <li>• Displays wide range of emotions that are appropriate to situation</li> <li>• Congruent with mood</li> <li>• Bright</li> <li>• Hopeful with goals</li> <li>• Positive self-worth</li> </ul>	<ul style="list-style-type: none"> <li>• Inappropriate or incongruent with the situation</li> <li>• Subdued</li> <li>• Tearful</li> <li>• Labile</li> <li>• Blunted</li> <li>• Flat</li> <li>• Feeling uneasy or unhappy</li> <li>• Intense excitement or happiness</li> <li>• Lack of motivation</li> <li>• Lack of interest to engage in social activities</li> <li>• Diminished feelings of pleasure in everyday life</li> <li>• Decreased interest in activities that would be interesting</li> </ul>
<b>Thought and Perception</b>	<ul style="list-style-type: none"> <li>• Realistic</li> <li>• Logical</li> <li>• Goal-directed</li> <li>• Organized</li> <li>• Ability to focus or concentrate</li> <li>• Absence of suicidal ideation</li> <li>• Absence of homicidal ideation</li> <li>• Absence of violence ideation</li> </ul>	<ul style="list-style-type: none"> <li>• Inability to focus or concentrate</li> <li>• Irrational fear</li> <li>• Exaggerated response</li> <li>• Delusions</li> <li>• Hallucinations</li> <li>• Illusions</li> <li>• Obsessions</li> <li>• Racing thoughts</li> <li>• Flight of ideas</li> <li>• Loose associations</li> <li>• Clang associations</li> <li>• Suicidal ideation</li> <li>• Homicidal ideation</li> </ul>

<b>Attitude and Insight</b>	<ul style="list-style-type: none"> <li>• Looks toward improvement and/or recovery</li> <li>• Demonstrates understanding of the situation</li> </ul>	<ul style="list-style-type: none"> <li>• Exhibits hostility, anger, helplessness, pessimism, overdramatization, self-centeredness, or passivity</li> <li>• Demonstrates little or no understanding of the situation</li> <li>• Inability to recognize one is ill</li> </ul>
<b>Cognitive Abilities</b>	<ul style="list-style-type: none"> <li>• Focused attention</li> <li>• Good immediate recall, short-term memory, and long-term storage</li> </ul>	<ul style="list-style-type: none"> <li>• Distractibility</li> <li>• Inability to think abstractly</li> <li>• Impaired reasoning</li> <li>• Poor immediate recall</li> <li>• Poor short-term memory</li> <li>• Poor long-term memory</li> </ul>
<b>Examiner's Reaction to Client</b>	<ul style="list-style-type: none"> <li>• Noticing and managing examiner's internal responses to the client such as frustration, boredom, sadness, anxiousness, or countertransference</li> <li>• Non-judgmental, neutral approach</li> <li>• Professional demeanor with control of subjective emotional and cognitive responses to ensure objectivity.</li> </ul>	<ul style="list-style-type: none"> <li>• Lack of awareness of examiner's internal responses to the client such as frustration, boredom, sadness, anxiousness, countertransference</li> </ul>

## Signs of Distress

If a client is exhibiting signs of distress during an examination, the nurse must quickly obtain focused assessment data and obtain additional assistance based on the level of emergency care required and agency policy. For example, if a client is found unresponsive, a “code” is typically called during inpatient care, or 911 is called in an outpatient setting as the nurse begins cardiopulmonary resuscitation (CPR). If a client is demonstrating difficulty breathing, new onset confusion, or other signs of a deteriorating condition, the rapid response team may be called, or other emergency assistance may be obtained per agency policy. Keep in mind that the emergency administration of naloxone may be required in cases of a suspected opioid overdose.

## Level of Consciousness and Orientation

A normal level of consciousness is when the client is alert (i.e., the ability to respond to stimuli at the same level as most people) and oriented to person, place, and time while remaining aware of their situation. **Clouded consciousness** refers to a state of reduced awareness to stimuli. **Delirium** is an acute onset of an abnormal mental state, often with fluctuating levels of consciousness, disorientation, irritability, and hallucinations. Delirium is often associated with infection, metabolic disorders, or toxins in the central nervous system. **Obtundation** refers to a moderate reduction in the client’s level of awareness so that mild to moderate stimuli do not awaken the client. When arousal does occur, the client is slow to respond. **Stupor** refers to unresponsiveness unless a vigorous stimulus is applied, such as a sternal rub. The client quickly drifts back into a deep sleep-like state on cessation of the stimulation. **Coma** refers to unarousable unresponsiveness, where vigorous noxious stimuli may not elicit reflex motor responses. For example, a client in a coma may not pull their foot away from a painful prick of their toe with a needle. When documenting reduced levels of consciousness, note the type of

stimulus required to arouse the client and the degree to which the client can respond when aroused.<sup>6</sup>

## Appearance and General Behavior

This component refers to an overall impression of the client, including their physical appearance regarding their age, grooming, dressing, posture, eye contact, ability to socialize with others, and general behaviors. There are several terms used to describe a client's appearance and behavior. For example, the appearance of one's age can be altered due to chronic illness and pain. Providers may document that a client "appears their stated age" or "appears older than their stated age." Clients may be described as well-groomed (i.e., exhibit good hygiene) or **disheveled** (i.e., their hair, clothes, or hygiene appears untidy, disordered, unkempt, or messy). Their dress may be described as "appropriate" or "inappropriate" according to the weather and situation. A client's posture may be described as "erect" or "slumped." Clients may be described as having "good eye contact" (i.e., they maintain a direct gaze into the examiner's eyes) or "poor eye contact" (i.e., they avoid direct eye contact).<sup>7</sup> Life span and cultural considerations must always be kept in mind when assessing a client's appearance and general behavior. For example, some cultures consider direct eye contact disrespectful.

6. Martin, D. C. (1990). The mental status examination. In Walker, H. K., Hall, W. D., Hurst, J. W., (Eds.), *Clinical methods: The history, physical, and laboratory examinations*. (3rd ed.). Butterworths. <https://www.ncbi.nlm.nih.gov/books/NBK320/>
7. Martin, D. C. (1990). The mental status examination. In Walker, H. K., Hall, W. D., Hurst, J. W., (Eds.), *Clinical methods: The history, physical, and laboratory examinations*. (3rd ed.). Butterworths. <https://www.ncbi.nlm.nih.gov/books/NBK320/>



# Speech

Evaluating speech as the client answers open-ended questions provides useful information. A client demonstrates normal speech when responding to verbal questions appropriately with an even rate, rhythm, and tone. Their speech is clear and understandable, and the client follows instructions appropriately.

Characteristics of speech can be described as normal, rapid, slow (i.e., delayed rhythm of conversation), loud, or soft. Stuttering and aphasia may occur. Examples of speech difficulties include lack of appropriate responses to verbal questions, rapid and/or pressured speech of a client experiencing mania or amphetamine intoxication, or halting speech of a client experiencing word-finding difficulties due to a previous stroke.<sup>8</sup>

Other terms used to describe speech include **circumstantial** (i.e., speaking with many unnecessary or tedious details without getting to the point of the conversation), **poverty of content** (i.e., a conversation in which the client talks without stating anything related to the question, or their speech in general is vague and meaningless), and **alogia** (i.e. brief or reduced replies).

## Motor Activity

Overall motor activity should be noted, including any tics or unusual mannerisms. Normal motor activity refers to the client having good balance, moving all extremities equally bilaterally, and walking with a smooth gait. Slow movements or lack of spontaneity in movement can occur due to depression or dementia. **Dyskinesia** (uncontrolled, involuntary movement)

8. Martin, D. C. (1990). The mental status examination. In Walker, H. K., Hall, W. D., Hurst, J. W., (Eds.), *Clinical methods: The history, physical, and laboratory examinations*. (3rd ed.). Butterworths. <https://www.ncbi.nlm.nih.gov/books/NBK320/>

and **akathisia** (i.e., motor restlessness) may occur if the client is experiencing extrapyramidal syndrome related to psychotropic medication use.<sup>9</sup>

Terminology used to describe motor activity includes **psychomotor agitation** (i.e., a condition of purposeless, non goal-directed activity) and **psychomotor retardation** (i.e., a condition of extremely slow physical movements, slumped posture, or slow speech patterns).

## Affect and Mood

**Affect** refers to the client's expression of emotion, and mood refers to the predominant emotion expressed by an individual.<sup>10</sup> Sustained emotions influence a person's behavior, personality, and perceptions. **Mood** can be described using various terms such as neutral, elevated, or **labile** (i.e., a rapid change in emotional responses, mood, or affect that are inappropriate for the moment or the situation). It can also be described as anxious, angry, sad, irritable, **dysphoric** (i.e., exhibiting depression), or **euphoric** (i.e., a pathologically elevated sense of well-being). People may express feelings of emptiness, impaired self-esteem, indecisiveness, or crying spells.<sup>11</sup>

9. Martin, D. C. (1990). The mental status examination. In Walker, H. K., Hall, W. D., Hurst, J. W., (Eds.), *Clinical methods: The history, physical, and laboratory examinations*. (3rd ed.). Butterworths. <https://www.ncbi.nlm.nih.gov/books/NBK320/>
10. Martin, D. C. (1990). The mental status examination. In Walker, H. K., Hall, W. D., Hurst, J. W., (Eds.), *Clinical methods: The history, physical, and laboratory examinations*. (3rd ed.). Butterworths. <https://www.ncbi.nlm.nih.gov/books/NBK320/>
11. Martin, D. C. (1990). The mental status examination. In Walker, H. K., Hall, W. D., Hurst, J. W., (Eds.), *Clinical methods: The history, physical, and laboratory examinations*. (3rd ed.). Butterworths. <https://www.ncbi.nlm.nih.gov/books/NBK320/>

Normal affect and mood are described as **euthymic** (i.e., displays a wide range of emotion that is appropriate for the situation). Abnormal findings related to affect include inappropriateness for the situation (e.g., laughing at the recent death of a loved one) or incongruent. **Congruence** refers to the consistency of verbal and nonverbal communication. Affect may also be described as subdued, tearful, labile, **blunted** (i.e., diminished range and intensity), or **flat** (no emotional expression).

Other terminology related to documenting a client's mood includes **alexithymia** (i.e., the inability to describe emotions with how one is feeling), **anhedonia** (i.e., the lack of experiencing pleasure in activities normally found enjoyable), **apathy** (i.e., a lack of feelings, emotions, interests, or concerns), **avolition** (i.e. total lack of motivation) and **asociality** (i.e. a lack of motivation to engage in social activities).

## Thoughts and Perceptions

The manner in which a client perceives and responds to stimuli is a critical psychiatric assessment. The inability to process information accurately is a component of the definition of psychotic thinking. For example, does the client harbor realistic concerns or are their concerns elevated to the level of irrational fear? Is the client responding in an exaggerated fashion to actual events? Is there no discernible basis in reality for the client's beliefs or behavior?<sup>12</sup>

Clients with mental health disorders may experience intrusive thoughts, delusions, and/or obsessions. **Delusions** are a fixed, false belief not held by cultural peers and persisting in the face of objective contradictory evidence. For example, a client may have the delusion that the CIA is listening to their

12. Martin, D. C. (1990). The mental status examination. In Walker, H. K., Hall, W. D., Hurst, J. W., (Eds.), *Clinical methods: The history, physical, and laboratory examinations*. (3rd ed.). Butterworths. <https://www.ncbi.nlm.nih.gov/books/NBK320/>

conversations via satellites. **Grandiose delusions** refer to a state of false attribution to the self of great ability, knowledge, importance or worth, identity, prestige, power, or accomplishment.<sup>13</sup> Clients may withdraw into an inner fantasy world that's not equivalent to reality, where they have inflated importance, powers, or a specialness that is opposite of what their actual life is like.<sup>14</sup> **Paranoia** is a condition characterized by delusions of persecution.<sup>15</sup> Clients often experience extreme suspiciousness or mistrust or express fear. For example, a resident of a long-term care facility may have delusions that the staff is trying to poison them.

**Obsessions** are persistent thoughts, ideas, images, or impulses that are experienced as intrusive or inappropriate and result in anxiety, distress, or discomfort. Common obsessions include repeated thoughts about contamination, a need to have things in a particular order or sequence, repeated doubts, aggressive impulses, and sexual imagery. Obsessions are distinguished from excessive worries about everyday occurrences because they are not concerned with real-life problems.<sup>16</sup> **Rumination** is obsessional thinking involving excessive, repetitive thoughts that interfere with other forms of mental activity.<sup>17</sup>

13. American Psychological Association. (n.d.). *APA Dictionary of Psychology*.  
<https://dictionary.apa.org/>

14. Raypole, C. (2019, November 20). *10 signs of covert narcissism*. Healthline.  
<https://www.healthline.com/health/covert-narcissist#fantasies>

15. American Psychological Association. (n.d.). *APA Dictionary of Psychology*.  
<https://dictionary.apa.org/>

16. American Psychological Association. (n.d.). *APA Dictionary of Psychology*.  
<https://dictionary.apa.org/>

17. American Psychological Association. (n.d.). *APA Dictionary of Psychology*.  
<https://dictionary.apa.org/>

Clients may also experience altered perceptions such as hallucinations and illusions. **Hallucinations** are false sensory perceptions not associated with real external stimuli and can include any of the five senses (auditory, visual, gustatory, olfactory, and tactile). For example, a client may see spiders climbing on the wall or hear voices telling them to do things. These are referred to as “visual hallucinations” or “auditory hallucinations.”

**Illusions** are misperceptions of real stimuli. For example, a client may misperceive tree branches blowing in the wind at night to be the arms of monsters trying to grab them.

It is important for nurses to remember that delusions, hallucinations, and illusions feel very real to clients and cause internal emotional reactions, even when a caregiver reassures them they are not based in reality. Because clients often conceal these experiences, it is helpful to ask leading questions, such as, “Have you ever seen or heard things that other people could not see or hear? Have you ever seen or heard things that later turned out not to be there?”<sup>18</sup>

Other terms used to document clients’ thought processes include racing thoughts, flight of ideas, loose associations, and clang associations. **Racing thoughts** are fast moving and often repetitive thought patterns that can be overwhelming. They may focus on a single topic, or they may represent multiple different lines of thought. For example, a client may have racing thoughts about a financial issue or an embarrassing moment.

**Flight of ideas** indicates the client frequently shifts from one topic to another with rapid speech, making it seem fragmented. The examiner may feel the client is rambling and changing topics faster than they can keep track, and

18. Martin, D. C. (1990). The mental status examination. In Walker, H. K., Hall, W. D., Hurst, J. W., (Eds.), *Clinical methods: The history, physical, and laboratory examinations*. (3rd ed.). Butterworths. <https://www.ncbi.nlm.nih.gov/books/NBK320/>

they probably can't get a word in edgewise.<sup>19</sup> An example of a client exhibiting a flight of ideas is, "My father sent me here. He drove me in a car. The car is yellow in color. Yellow looks good on me."<sup>20</sup>

**Loose associations** refers to jumping from one idea to an unrelated idea in the same sentence. For example, the client might state, "I like to dance, and my feet are wet."<sup>21</sup> The term **word salad** refers to severely disorganized and virtually incomprehensible speech or writing, marked by severe loosening of associations.<sup>22</sup>

**Clang associations** refers to stringing words together that rhyme without logical association and do not convey rational meaning. For example, a client exhibiting clang associations may state, "Here she comes with a cat catch a rat match."

Other disturbances in thought and perception may include **echolalia** (i.e. repetition of words or phrases spoken to others) and **magical thinking** (i.e. false belief that reality can be changed by one's thoughts).

Clients with altered perceptions, especially when experiencing hallucinations and delusions, may have violent thoughts regarding themselves or others. If a

19. Martin, D. C. (1990). The mental status examination. In Walker, H. K., Hall, W. D., Hurst, J. W., (Eds.), *Clinical methods: The history, physical, and laboratory examinations*. (3rd ed.). Butterworths. <https://www.ncbi.nlm.nih.gov/books/NBK320/>
20. PsychologGenie. (n.d.). *The true meaning of flight of ideas explained with examples*. <https://psychologenie.com/flight-of-ideas-meaning-examples>
21. American Psychological Association. (n.d.). *APA Dictionary of Psychology*. <https://dictionary.apa.org/>
22. American Psychological Association. (n.d.). *APA Dictionary of Psychology*. <https://dictionary.apa.org/>

client is having auditory hallucinations, it is vital for the nurse to determine if the voices are encouraging the client to hurt themselves or others. **Homicidal ideation** refers to threats or acts of life-threatening harm toward another person. **Suicidal ideation** is used to describe an individual who has been thinking about suicide but does not necessarily have an intention to act on that idea. **Suicide attempt** is a term used to describe an individual who harms themselves with intent to end their life but does not die as a result of their actions. **Suicide plan** refers to an individual who has a plan for suicide, has the means to injure oneself, and has the intent to die.

Of all portions of the mental status examination, the evaluation of thought disorders is the most difficult and requires a thorough assessment.<sup>23</sup> Psychiatric-mental health nurse specialists receive additional training in assessing thought disorders. These types of thought disorders are associated with mental illnesses like bipolar disorder and schizophrenia and may precede an episode of psychosis, so it is important to obtain further assistance if you notice a client is newly exhibiting these types of behaviors.<sup>24</sup>

- ▶ Read more information about how to help individuals experiencing hallucinations and delusions in the “[Applying the Nursing Process to Schizophrenia](#)” section of the “Psychosis and Schizophrenia” chapter.

23. Martin, D. C. (1990). The mental status examination. In Walker, H. K., Hall, W. D., Hurst, J. W., (Eds.), *Clinical methods: The history, physical, and laboratory examinations*. (3rd ed.). Butterworths. <https://www.ncbi.nlm.nih.gov/books/NBK320/>
24. Joy, R. (2020, January 20). *Clang association: When a mental health condition disrupts speech*. Healthline. <https://www.healthline.com/health/clang-association#whats-it-sound-like>

## Attitude and Insight

The client's attitude is the emotional tone displayed toward the examiner, other individuals, or their illness. It may convey a sense of hostility, anger, helplessness, pessimism, overdramatization, self-centeredness, or passivity. It is important to determine the client's attitude toward emotional problems or diagnosed mental health disorders. Does the client look forward to improvement and recovery or are they resigned to suffer?<sup>25</sup>

**Insight** is the client's ability to identify the existence of a problem and to have an understanding of its nature.

Nurses must also be aware of transference. **Transference** occurs when the client projects (i.e., transfers) their feelings onto the nurse. For example, a client is feeling angry at a family member related to a previous disagreement and displaces the anger onto the nurse during the interview.

## Cognitive Abilities

**Cognition** is the mental action or process of acquiring knowledge and understanding through thought, experience, and the senses. It includes thinking, knowing, remembering, judging, and problem-solving. When performing focused assessments on cognition, the examiner assesses attention and memory.<sup>26</sup> A term related to assessing attention is

25. Martin, D. C. (1990). The mental status examination. In Walker, H. K., Hall, W. D., Hurst, J. W., (Eds.), *Clinical methods: The history, physical, and laboratory examinations*. (3rd ed.). Butterworths. <https://www.ncbi.nlm.nih.gov/books/NBK320/>

26. Martin, D. C. (1990). The mental status examination. In Walker, H. K., Hall, W. D., Hurst, J. W., (Eds.), *Clinical methods: The history, physical, and laboratory examinations*. (3rd ed.). Butterworths. <https://www.ncbi.nlm.nih.gov/books/NBK320/>



**distractibility**, referring to the client's attention being easily drawn to unimportant or irrelevant external stimuli.

Memory disturbance is a common complaint and is often a presenting symptom in the elderly. Memory can be grouped into three categories: immediate recall, short-term memory, and long-term storage. Short-term memory is the most clinically pertinent, and the most important to be tested. Short-term retention requires that the client process and store information so that they can move on to a second intellectual task and then call up the remembrance after completion of the second task. For example, short-term memory may be tested by having the client repeat the names of four unrelated objects and then asking the client to recall the information in 3 to 5 minutes after performing a second, unrelated mental task.<sup>27</sup>

## Examiner's Reaction to the Client

Assessing a client sometimes results in the nurse developing subtle and easily overlooked feelings toward the client. For example, it can be difficult to repeatedly address a client's negative state. Examiners may experience feelings of frustration, which can be taken by clients to mean there's something wrong with them. In such cases, nurses should examine their reactions to the client and be alert to feelings of distraction, boredom, or frustration. They should also be aware that clients perceive a nurse's feelings through their nonverbal communication, such as facial expressions, posture, tone of voice, and lack of eye contact.<sup>28</sup>

27. Martin, D. C. (1990). The mental status examination. In Walker, H. K., Hall, W. D., Hurst, J. W., (Eds.), *Clinical methods: The history, physical, and laboratory examinations*. (3rd ed.). Butterworths. <https://www.ncbi.nlm.nih.gov/books/NBK320/>

28. DeAngelis, T. (2019). Better relationships with patients lead to better outcomes. *Monitor on Psychology*, 50(10), 38. <https://www.apa.org/monitor/2019/11/ce-corner-relationships>

Nurses should also be aware of countertransference. **Countertransference** refers to a tendency for the examiner to displace (transfer) their own feelings onto the client and then these feelings may influence the client. For example, a nurse finds themselves providing advice about how to raise children to a client. Upon self-reflection, they realize it is a countertransference reaction related to their previous parenting experience.<sup>29</sup>

- ▶ Review a brief [mental status examination PDF form](#) from TherapistAid.

## Psychosocial Assessment

A **psychosocial assessment** (also referred to as a health history) is a component of the nursing assessment process that obtains additional subjective data to detect risks and identify treatment opportunities and resources. Agencies have specific forms used for psychosocial assessments/health histories that typically consist of several components<sup>30, 31</sup>:

29. DeAngelis, T. (2019). Better relationships with patients lead to better outcomes. *Monitor on Psychology*, 50(10), 38. <https://www.apa.org/monitor/2019/11/ce-corner-relationships>
30. GW School of Medicine & Health Sciences. (n.d.). *Clinical FICA tool*. <https://smhs.gwu.edu/spirituality-health/program/transforming-practice-health-settings/clinical-fica-tool>
31. Glasner, J., Baltag, V., & Ambresin, A. E. (2021). Previsit multidomain psychosocial screening tools for adolescents and young adults: A system review. *Journal of Adolescent Health*, 68, 449-459. [https://www.jahonline.org/article/S1054-139X\(20\)30600-5/pdf](https://www.jahonline.org/article/S1054-139X(20)30600-5/pdf)

- Reason for seeking health care (i.e., “chief complaint”)
- Cultural assessment
- Spiritual assessment
- Family dynamics
- Thoughts of self-injury or suicide
- Current and past medical history
- Current medications
- History of previously diagnosed mental health disorders
- Previous hospitalizations
- Educational background
- Occupational background
- History of exposure to psychological trauma, violence, and domestic abuse
- Substance use (tobacco, alcohol, recreational drugs, misused prescription drugs)
- Family history of mental illness
- Coping mechanisms
- Functional ability/Activities of daily living

▶ Review specific questions used during a psychosocial assessment/health history in the “[Health History](#)” chapter in *Open RN Nursing Skills, 2e*.

## Reason for Seeking Health Care

It is helpful to begin the psychosocial assessment by obtaining the reason why the client is seeking health care in their own words. During a visit to a clinic or emergency department or on admission to a health care agency, the client’s primary reasons for seeking care are referred to as the **chief complaint**. Assessing a client’s chief complaint recognizes that clients are complex beings, with potentially multiple coexisting health needs, but there

is often a pressing issue that requires most immediate care. Questions used to evaluate a client's chief complaint are as follows:

- What brought you in today?
- How long has this been going on?
- How is this affecting you?

After identifying the reason the client is seeking health care, additional focused questions are used to obtain detailed information about priority concerns, such as persistent nausea or other symptoms causing discomfort. The mnemonic PQRSTU is used to ask the client questions in an organized fashion. See Table 4.3b for examples of questions used to assess a client's report of nausea.

Table 4.3b Sample PQRSTU Questions

PQRSTU	Sample Questions
<b>Provocation/ Palliation</b>	What were you doing when the nausea started?  Does anything make it better or worse (e.g., food, position, movement, m
<b>Quality</b>	Describe the feeling? (e.g., queasy, churning, lightheaded, urge to vomit?
<b>Region</b>	Do you feel the nausea in a specific part of your body (e.g., upper stomach
<b>Severity</b>	On a scale of 0 to 10, how bad is the nausea right now? Has it gotten worse
<b>Timing/Treatment</b>	When did the nausea start?  Is it constant or does it come and go?  Does it occur at specific times (e.g., after eating, in the morning)?
<b>Understanding</b>	What do you think is causing the nausea?

## Cultural Formulation Interview Questions

While performing a psychosocial assessment, it is important to performing a cultural assessment. We all bring our own cultural beliefs, values, and expectations to the clinical encounter, which influences how we approach specific aspects of care. The American Psychiatric Association developed evidence-based Cultural Formulation Interview (CFI) questions as a way to incorporate cultural assessment into the care of all clients that enhances

clinical understanding and decision-making.<sup>32</sup> The CFI questions are used to clarify key aspects of the presenting clinical problem from the point of view of the individual and other members of the individual's social network (e.g., family, friends, or others involved in the current problem). This includes the problem's meaning, potential sources of help, and expectations for health care services.

CFI questions used with all clients include the following<sup>33</sup>:

- What brings you here today?
- What troubles you most about this problem?
- What do you think is the cause of this problem?
- Are there any kinds of support that make this problem better, such as support from family, friends, or others?
- Are there any kinds of stresses that make this problem worse, such as difficulties with money or family problems?
- Sometimes aspects of people's background or identity can make their problem better or worse, such as the communities they belong to, the languages they speak, where they or their family are from, their race or ethnic background, their gender or sexual orientation, or their faith or religion. Are there any aspects of your background or identity that make a difference to this problem?
- Sometimes people have various ways of dealing with problems. What have you done on your own to cope with this problem?

32. DeSilva, R., Aggarwall, N. K., & Lewis-Fernandez, R. (2015). The DSM-5 cultural formulation interview and the evolution of cultural assessment in psychiatry. *Psychiatric Times*, 32(6). <https://www.psychiatristimes.com/view/dsm-5-cultural-formulation-interview-and-evolution-cultural-assessment-psychiatry>

33. American Psychiatric Association. (2013). Cultural formulation. In *Diagnostic and Statistical Manual of Mental Disorders* (5th ed.). American Psychiatric Association. pp. 749-759.

- Often, people look for help from many different sources, including different kinds of doctors, helpers, or healers. In the past, what kinds of treatment, help, advice, or healing have you sought for this problem?
- Has anything prevented you from getting the help you need?
- What kinds of help do you think would be most useful to you at this time for this problem?
- Are there other kinds of help that your family, friends, or other people have suggested that would be helpful for you now?
- Sometimes health care professionals and clients misunderstand each other because they come from different backgrounds or have different expectations. Have you been concerned about misaligned care expectations and is there anything that we can do to provide you with the care you need?

Findings from the cultural formulation interview are used to individualize a client's plan of care to their preferences, values, beliefs, and goals.

## Spiritual Assessment

Spiritual assessment is included in a psychosocial assessment. It is common for people in the process of recovery from mental health disorders and substance use to search for spiritual support.<sup>34</sup> **Spirituality** includes a sense of connection to something larger than oneself and typically involves a search for meaning and purpose in life. Basic questions used to assess spirituality include the following:

- Who or what provides you with strength or hope?
- How do you express your spirituality?

34. Neto, G. L., Rodrigues, L., Rozendo da Silva, D. A., Turato, E. R., & Campos, C. J. G. (2018). Spirituality review on mental health and psychiatric nursing. *Revista Brasileira de Enfermagem*, 71 (Suppl 5), 2323-2333. <https://doi.org/10.1590/0034-7167-2016-0429>

- What spiritual needs can we advocate for you during this health care experience?

Over the past decade, research has demonstrated the importance of spirituality in health care.<sup>35, 36</sup> Spiritual distress is very common for clients experiencing serious illness, injury, or the dying process, and nurses are on the front lines as they assist these individuals to cope with these life events. Addressing a client's spirituality and advocating spiritual care have been shown to improve clients' health and quality of life, including how they experience pain, cope with stress and suffering associated with serious illness, and approach end of life.<sup>37, 38</sup>

35. Pilger, C., Molzahn, A. E., de Oliveira, M. P., & Kusumota, L. (2016). The relationship of the spiritual and religious dimensions with quality of life and health of patients with chronic kidney disease: An integrative literature review. *Nephrology Nursing Journal: Journal of the American Nephrology Nurses' Association*, 43(5), 411–426. <https://pubmed.ncbi.nlm.nih.gov/30550069/>
36. Puchalski, C., Jafari, N., Buller, H., Haythorn, T., Jacobs, C., & Ferrell, B. (2020). Interprofessional spiritual care education curriculum: A milestone toward the provision of spiritual care. *Journal of Palliative Medicine*, 23(6), 777–784. <https://doi.org/10.1089/jpm.2019.0375>
37. Pilger, C., Molzahn, A. E., de Oliveira, M. P., & Kusumota, L. (2016). The relationship of the spiritual and religious dimensions with quality of life and health of patients with chronic kidney disease: An integrative literature review. *Nephrology Nursing Journal: Journal of the American Nephrology Nurses' Association*, 43(5), 411–426. <https://pubmed.ncbi.nlm.nih.gov/30550069/>
38. Puchalski, C., Jafari, N., Buller, H., Haythorn, T., Jacobs, C., & Ferrell, B. (2020). Interprofessional spiritual care education curriculum: A milestone toward the provision of spiritual care. *Journal of Palliative Medicine*, 23(6), 777–784. <https://doi.org/10.1089/jpm.2019.0375>



The FICA Spiritual History Tool<sup>39</sup> is a common tool used to gather information about a client's spiritual history and preferences. FICA<sup>40</sup> is a mnemonic for Faith, Importance, Community, and Address in Care. Read more about the FICA<sup>40</sup> tool in the following box.

### **The FICA<sup>40</sup> Spiritual History Tool**

**F – Faith and Belief:** Determine if the client identifies with a particular belief system or spirituality.

**I – Importance:** Ask, “Is this belief important to you? Does it influence how you think about health and illness? Does it influence your health care decisions?”

**C – Community:** Determine if the client belongs to a spiritual community (e.g., a church, temple, mosque, or other group). If not, ask, “Would it be helpful to you to find a spiritual community?”

**A – Address in Care:** Evaluate what should be addressed during the client's care. Ask, “What should be included in your treatment plan? Are there spiritual practices you want to develop? Would you like to see a chaplain, spiritual director, or pastoral counselor while you are here?”

39. GW School of Medicine & Health Sciences. (n.d.). *Clinical FICA tool*. <https://smhs.gwu.edu/spirituality-health/program/transforming-practice-health-settings/clinical-fica-tool>

40. GW School of Medicine & Health Sciences. (n.d.). *Clinical FICA tool*. <https://smhs.gwu.edu/spirituality-health/program/transforming-practice-health-settings/clinical-fica-tool>

Based on the assessment findings, nurses may refer clients to agency chaplains or to the client's religious leaders for spiritual support to enhance coping.

- ▶ Read more about spiritual assessment and providing spiritual care in the "[Spirituality](#)" chapter of *Open RN Nursing Fundamentals, 2e*.

## Family Dynamics

Family dynamics are included in a psychosocial assessment, especially for children, adolescents, and older adults. **Family dynamics** refers to the patterns of interactions among relatives, their roles and relationships, and the various factors that shape their interactions. Because family members rely on each other for emotional, physical, and economic support, they are primary sources of relationship security or stress. Family dynamics and the quality of family relationships can have either a positive or negative impact on an individual's health. For example, secure and supportive family relationships can provide love, advice, and care, whereas stressful family relationships can be burdened with arguments, unhealthy relationships, and a lack of support.<sup>41</sup>

Unhealthy family dynamics can cause children to experience trauma and stress as they grow up. This type of exposure, known as adverse childhood experiences (ACEs), is linked to an increased risk of developing physical and mental health problems such as heart, lung, and liver disease; depression; and

41. This work is a derivative of [StatPearls](#) by Jabbari & Rouster and is licensed under [CC BY 4.0](#)

anxiety. Unhealthy family dynamics also correlate with an increased risk of substance use and addiction among adolescents.<sup>42</sup>

- ▶ Review information about adverse childhood experiences (ACEs) in the “[Mental Health and Mental Illness](#)” section of Chapter 1.

Effectively assessing and addressing a client’s family dynamic and its role in health and disease require an interprofessional team of health professionals, including nurses, physicians, social workers, and therapists. Nurses are in a unique position to observe and document interaction patterns, assess family relationships, and tend to family concerns in the clinical setting because they are in frequent contact with family members.<sup>43</sup>

## Suicide Screening

As discussed in Chapter 1, all clients aged 12 and older presenting for acute care should be screened for suicidal ideation. Clients being evaluated or treated for mental health conditions often have suicidal ideation, and up to 10 percent of emergency department clients presenting with medical issues have a hidden risk for suicide, such as recent suicidal ideation or previous

42. This work is a derivative of [StatPearls](#) by Jabbari & Rouster and is licensed under [CC BY 4.0](#)

43. This work is a derivative of [StatPearls](#) by Jabbari & Rouster and is licensed under [CC BY 4.0](#)

suicide attempts.<sup>44</sup> Universal screening allows for the detection of suicide risk and implementation of early interventions before a person attempts suicide.

It is important to introduce suicide screening in a way that helps the client understand its purpose and normalize questions that might otherwise seem intrusive. A nurse might introduce the topic in the following way: “Now I’m going to ask you some questions that we ask everyone treated here, no matter what problem they are here for. It is part of the hospital’s policy, and it helps us to make sure we are not missing anything important.”<sup>45</sup>

The Patient Safety Screener (PSS-3) is an example of a brief screening tool to detect suicide risk in all client presenting to acute care settings.<sup>46</sup> See Figure 4.3<sup>47</sup> for an image of the PSS-3.

The PSS-3 consists of assessing for three items: depression, active suicidal ideation, and lifetime suicide attempt. Each of these items explores a different aspect of suicide risk<sup>48</sup>:

- 44. Suicide Prevention Resource Center. (n.d.). *The patient safety screener: A brief tool to detect suicide risk*. <https://sprc.org/micro-learning/patientsafetyscreener>
- 45. Suicide Prevention Resource Center. (n.d.). *The patient safety screener: A brief tool to detect suicide risk*. <https://sprc.org/micro-learning/patientsafetyscreener>
- 46. Suicide Prevention Resource Center. (n.d.). *The patient safety screener: A brief tool to detect suicide risk*. <https://sprc.org/micro-learning/patientsafetyscreener>
- 47. “Patient Safety Screener (PSS-3) Pocket Card” by University of Massachusetts Medical School (UMass Medical) is used on the basis of Fair Use.
- 48. Suicide Prevention Resource Center. (n.d.). *The patient safety screener: A brief*

- Depression is a common precipitant of suicidal ideation and behavior and is the most common diagnosis among those who die by suicide.
- Suicidal ideation (i.e., thoughts about killing oneself) is a precondition for suicidal behavior.
- A previous suicide attempt is one of the most consistent risk factors for suicide.

Use this pocket card as a job aid or training tool when implementing universal suicide screening in acute care settings.

The Patient Safety Screener can be used during the Triage or Primary Nursing Assessment in acute care settings.  
Ask all three screening questions. Do not skip items.

**Introduction** "Now I'm going to ask you some questions that we ask everyone treated here, no matter what problem they are here for. It is part of the hospital's policy, and it helps us to make sure we are not missing anything important."

**Depression** ❶ Over the past 2 weeks, have you felt down, depressed, or hopeless?  
☐ Yes ☐ No ☐ Refused ☐ Patient unable to complete

**Suicidal ideation** ❷ Over the past 2 weeks, have you had thoughts of killing yourself?  
☐ Yes ☐ No ☐ Refused ☐ Patient unable to complete

**Suicide attempt** ❸ Have you ever attempted to kill yourself?  
☐ Yes ☐ No ☐ Refused ☐ Patient unable to complete  
 ...3a. If Yes to item 3, ask: when did this last happen?  
☐ Within the past 24 hours (including today) ☐ More than 6 months ago  
☐ Within the last month (but not today) ☐ Refused  
☐ Between 1 and 6 months ago ☐ Patient unable to complete

**TIPS**

- ✓ Ask all questions exactly as worded
- ✓ Do not bundle or re-word questions
- ✓ Treat the patient with empathy

**"Yes" to Item 1 = positive screen for Depression.**

**"Yes" to Item 2 OR "last 6 months" to Item 3 = positive screen for Suicide Risk.**

**Apply site protocol for further evaluation and management.**

**Patient Safety Screener (PSS-3) Pocket Card**

*The Patient Safety Screener 3 (PSS-3) has been validated in prospective studies and is detailed in Boudreaux et al. (2015)*

Figure 4.3 Patient Safety Screener for Suicide Risk. Used under Fair Use.

## Self-Injury

**Non-suicidal self-injury (NSSI)** refers to intentional self-inflicted destruction of body tissue without suicidal intention and for purposes not socially

tool to detect suicide risk. <https://sprc.org/micro-learning/patientsafetyscreener>

sanctioned. Common forms of NSSI include behaviors such as cutting, burning, scratching, and self-hitting. It is considered a maladaptive coping strategy without the desire to die. NSSI is a common finding among adolescents and young adults in psychiatric inpatient settings.<sup>49</sup>

## Current and Past Medical History

A thorough review of a client's current and past medical conditions provides critical insight into their overall health and potential influences on mental well-being. Chronic illnesses such as diabetes, cardiovascular disease, or neurological disorders can contribute to or exacerbate psychiatric symptoms. Understanding this history helps nurses identify connections between physical and mental health.

## Medication Review

When evaluating a client's mental health status, it is essential to review their current use of psychotropic medications, as well as any other prescribed or over-the-counter drugs that may contribute to psychiatric symptoms. Psychotropic medications, which are used to manage conditions such as depression, anxiety, bipolar disorder, and schizophrenia, can have complex effects and may interact with other substances in ways that influence mood, cognition, or behavior. Additionally, certain non-psychiatric medications—such as corticosteroids, antihypertensives, or stimulants—can produce side effects that mimic or exacerbate psychiatric conditions. A thorough medication review helps ensure accurate assessment, prevents misdiagnosis, and guides appropriate treatment planning.

49. Cipriano, A., Cella, S., & Cotrufo, P. (2017). Nonsuicidal self-injury: A systematic review. *Frontiers in Psychology*, 8. <https://www.frontiersin.org/articles/10.3389/fpsyg.2017.01946/full>

# History of Previously Diagnosed Mental Health Disorders

Documenting any formally diagnosed mental health conditions, such as depression, anxiety, schizophrenia, or bipolar disorder, offers a foundation for understanding the client's mental health journey. This history supports continuity of care and helps identify patterns or triggers related to past episodes. Documenting any formally diagnosed mental health conditions, such as depression, anxiety, schizophrenia, or bipolar disorder, offers a foundation for understanding the client's mental health journey. This history supports continuity of care and helps identify patterns or triggers related to past episodes.

## Previous Hospitalizations

Exploring prior psychiatric or medical hospitalizations provides context for the severity and frequency of past health episodes. This information can help the nurse assess the progression of mental health conditions and the effectiveness of previous interventions.

## Educational Background

Understanding a client's level of education assists in planning appropriate communication strategies and assessing cognitive development and health literacy. Educational attainment may also impact employment, income, and social opportunities, influencing mental wellness.

## Occupational Background

A client's work history can reflect their ability to maintain structure, social roles, and economic stability. Job satisfaction or stress, unemployment, or disability due to mental health issues may reveal important psychosocial stressors or strengths.

# History of Exposure to Psychological Trauma, Violence, and Domestic Abuse

Exposure to trauma—such as childhood abuse, domestic violence, or combat experiences—can profoundly impact an individual's emotional regulation, cognitive development, and overall mental health. Traumatic experiences often disrupt the brain's ability to process stress and emotions effectively, leading to heightened anxiety, depression, difficulties in forming and maintaining relationships, and in some cases, the development of post-traumatic stress disorder (PTSD). Individuals who have experienced trauma may exhibit hypervigilance, emotional numbness, mood swings, or difficulty trusting others. These long-lasting effects can persist well into adulthood, influencing personal well-being, social functioning, and physical health outcomes. Understanding the deep and complex consequences of trauma is essential for recognizing behavioral responses and addressing mental health needs with empathy and evidence-based support.

## Substance Use

A comprehensive assessment of tobacco use, alcohol consumption, recreational drug use, and the misuse of prescription medications is essential in evaluating an individual's overall health, identifying risk factors, and detecting co-occurring substance use disorders. Substance use often does not occur in isolation and may be intertwined with underlying mental health conditions such as depression, anxiety, or trauma-related disorders. These substances can alter neurochemical pathways in the brain, leading to changes in mood, behavior, and cognitive function. For example, stimulant use may increase irritability or anxiety, while depressants such as alcohol or opioids can contribute to emotional blunting or memory impairment.

Accurate assessment enables healthcare providers to recognize patterns of use, understand the potential impact on physical and mental health, and tailor treatment approaches accordingly. Furthermore, substance use can interfere with the effectiveness of medical and psychiatric treatments, reduce



adherence to care plans, and increase the risk of adverse outcomes. Early identification and integrated treatment planning are critical to improving prognosis, enhancing client safety, and supporting long-term recovery and wellness.

## Family History of Mental Illness

Exploring family mental health history helps identify potential genetic or environmental influences on the client's condition. It also provides insight into the support systems or stigma the client may face in their home environment.

## Coping Mechanisms

Identifying the client's typical coping strategies—both adaptive and maladaptive—offers a window into their resilience and vulnerability. Understanding how individuals manage stress or adversity informs the development of effective care plans and interventions.

## Functional Ability/Activities of Daily Living (ADLs)

Assessing a client's ability to perform basic self-care tasks and daily routines—such as bathing, dressing, preparing meals, maintaining personal hygiene, managing medications, or handling finances—is a critical component of evaluating overall functioning. These activities, often referred to as Activities of Daily Living (ADLs) and Instrumental Activities of Daily Living (IADLs), provide important insight into a person's level of independence and the extent to which mental, emotional, or cognitive challenges may be affecting day-to-day life.

Functional impairments in these areas can serve as early warning signs of a worsening mental health condition, such as major depressive disorder, schizophrenia, or dementia, and may reflect cognitive decline, lack of motivation, or impaired judgment. Identifying deficits in daily functioning

helps clinicians determine the need for supportive services, such as occupational therapy, case management, home health care, or placement in a more structured living environment. In many cases, a multidisciplinary intervention that includes mental health professionals, medical providers, and social services is necessary to promote safety, improve quality of life, and support recovery.

In order to complete a robust mental health assessment, screening tools may be utilized to identify symptoms and standardize evaluation. Screening tools are evidence-based methods to assess specific information related to mental health disorders. These tools may be used on admission to the hospital or treatment facility, as well as at different times throughout the client's stay. Common examples include PHQ-9 for depression, GAD-7 for anxiety, and the CAGE questionnaire for alcohol use. Findings from the tool's results may be used to compare client progress during the hospital stay or from a previous admission. The registered nurse often conducts these screening tools as part of the interprofessional health care treatment team. Read more about specific screening tools in each "disorder" chapter.

## Laboratory and Diagnostic Testing

Nurses review laboratory and diagnostic testing results as part of the assessment process. Laboratory and diagnostic testing are essential components of a comprehensive client assessment to exclude underlying medical causes, identify co-morbid conditions, and guide appropriate treatment. For example, nurses routinely monitor electrolytes to assess fluid and electrolyte balance, which is crucial in many clinical conditions. An example of this would be serum sodium levels, as abnormal sodium can indicate dehydration, kidney dysfunction, or other systemic issues that may impact client care and treatment decisions.

Specific laboratory and diagnostic tests will be discussed in each “disorder” chapter, as well as in the “[Psychotropic Medications](#)” chapter.

## Life Span Considerations

Life span considerations influence nursing assessments, care planning, and interventions. Mental health disorders occur across the life span, from the very young to the very old, and developmental stages must be considered when identifying impairments. Assessments and interventions must be individualized to the age and developmental level of the client. **Development** encompasses physical, social, and cognitive changes that occur continuously throughout one’s life. See Figure 4.4<sup>50</sup> for an image of the human life cycle.



Figure 4.4 Human Life Cycle

There are multiple factors that affect human development with expected milestones along the way. Cognitive development encompasses several different skills that develop at different rates. Each human has their own individual experience that influences development of intelligence and

50. “[shutterstock\\_149010437.jpg](#)” by [Robert Adrian Hillman](#) is used under license from [Shutterstock.com](#)

reasoning as they interact with one another. With these unique experiences, everyone has a memory of feelings and events that is exclusive to them.<sup>51</sup>

There are many theories regarding how infants and children grow and develop into happy, healthy adults. Three major theories that have historically impacted nursing care are Freud's Psychosexual Theory of Development, Erikson's Psychosocial Stages of Development, and Piaget's Cognitive Theory of Development.

## Freud's Psychosexual Theory of Development

Sigmund Freud (1856–1939) believed that personality develops during early childhood, and childhood experiences shape our personalities and behavior as adults. Freud believed that each individual must pass through a series of stages during childhood, and if we lack proper nurturance and parenting during a stage, we may become stuck, or fixated, in that stage. According to Freud, children's pleasure-seeking urges are focused on different areas of the body, called erogenous zones, at each of the five stages of development: oral, anal, phallic, latency, and genital.<sup>52</sup>

While most of Freud's ideas are not supported by research and modern psychologists dispute Freud's psychosexual stages as a legitimate explanation

51. Vallotton, C. D., & Fischer, K. W. (2008). Cognitive development. *Encyclopedia of Infant and Early Childhood Development*. Academic Press. pp. 286-298.  
<https://doi.org/10.1016/B978-012370877-9.00038-4>

52. This work is a derivative of Psychology 2e by [OpenStax](https://openstax.org/books/psychology-2e) and is licensed under CC BY 4.0. Access for free at <https://openstax.org/books/psychology-2e/pages/1-introduction>

for how one's personality develops, Freud's original theory supported that one's personality is shaped, in some part, by childhood experiences.<sup>53</sup>

## Erikson's Psychosocial Stages of Development

Erik Erikson (1902–1994) took Freud's theory and modified it as psychosocial theory. Erikson's psychosocial development theory emphasizes the social nature of our development rather than its sexual nature. It describes eight sequential stages of individual human development influenced by biological, psychological, and social factors throughout the life span that contribute to an individual's personality. Erikson's stages of development are trust versus mistrust, autonomy versus shame, initiative versus guilt, industry versus inferiority, identity versus identity confusion, intimacy versus isolation, generativity versus stagnation, and integrity versus despair.<sup>54, 55</sup>

- **Trust vs. Mistrust**

The first stage that develops is trust (or mistrust) that basic needs, such as nourishment and affection, will be met. Trust is the basis of our development during infancy (birth to 12 months). Infants are dependent upon their caregivers for their needs. Caregivers who are responsive and sensitive to their infant's needs help their baby to develop a sense of trust, and the infant will perceive the world as a safe, predictable place. Unresponsive caregivers who do not meet their baby's needs can

53. Psychology 2e by [OpenStax](#) is licensed under [CC BY 4.0](#). Access for free at <https://openstax.org/books/psychology-2e/pages/1-introduction>

54. This work is a derivative of [StatPearls](#) by Orenstein & Lewis and is licensed under [CC BY 4.0](#)

55. Psychology 2e by [OpenStax](#) is licensed under [CC BY 4.0](#). Access for free at <https://openstax.org/books/psychology-2e/pages/1-introduction>

engender feelings of anxiety, fear, and mistrust, and the infant will perceive the world as unpredictable.<sup>56</sup>

- **Autonomy vs. Shame**

Toddlers begin to explore their world and learn that they can control their actions and act on the environment to get results. They begin to show clear preferences for certain elements of the environment, such as food, toys, and clothing. A toddler's main task is to resolve the issue of autonomy versus shame and doubt by working to establish independence. For example, we might observe a budding sense of autonomy in a two-year-old child who wishes to choose their own clothes and dress themselves. Although the outfits might not be appropriate for the situation, the input in basic decisions has an effect on the toddler's sense of independence. If denied the opportunity to act on their environment, they may begin to doubt their abilities, which could lead to low self-esteem and feelings of shame.<sup>57</sup>

- **Initiative vs. Guilt**

After children reach the preschool stage (ages 3–6 years), they are capable of initiating activities and asserting control over their world through social interactions and play. By learning to plan and achieve goals while interacting with others, preschool children can master a

56. Psychology 2e by [OpenStax](https://openstax.org/books/psychology-2e) is licensed under [CC BY 4.0](https://creativecommons.org/licenses/by/4.0/). Access for free at <https://openstax.org/books/psychology-2e/pages/1-introduction>

57. Psychology 2e by [OpenStax](https://openstax.org/books/psychology-2e) and is licensed under [CC BY 4.0](https://creativecommons.org/licenses/by/4.0/). Access for free at <https://openstax.org/books/psychology-2e/pages/1-introduction>

feeling of initiative and develop self-confidence and a sense of purpose. Those who are unsuccessful at this stage may develop feelings of guilt.<sup>58</sup>

- **Industry vs. Inferiority**

During the elementary school stage (ages 7–11), children begin to compare themselves to their peers to see how they measure up. They either develop a sense of pride and accomplishment in their schoolwork, sports, social activities, and family life, or they may feel inferior and inadequate if they feel they don't measure up to their peers.<sup>59</sup>

- **Identity vs. Identity Confusion**

In adolescence (ages 12–18), children develop a sense of self. Adolescents struggle with questions such as, “Who am I?” and “What do I want to do with my life?” Along the way, adolescents try on many different selves to see which ones fit. Adolescents who are successful at this stage have a strong sense of identity and are able to remain true to their beliefs and values in the face of problems and other people's perspectives. Teens who do not make a conscious search for identity, or those who are pressured to conform to their parents' ideas for the future, may have a weak sense of self and experience role confusion as they are unsure of their identity and confused about the future.<sup>60</sup>

- **Intimacy vs. Isolation**

58. Psychology 2e by [OpenStax](#) is licensed under [CC BY 4.0](#). Access for free at <https://openstax.org/books/psychology-2e/pages/1-introduction>

59. Psychology 2e by [OpenStax](#) is licensed under [CC BY 4.0](#). Access for free at <https://openstax.org/books/psychology-2e/pages/1-introduction>

60. Psychology 2e by [OpenStax](#) is licensed under [CC BY 4.0](#). Access for free at <https://openstax.org/books/psychology-2e/pages/1-introduction>

People in early adulthood (i.e., 20s through early 40s) are ready to share their lives and become intimate with others after they have developed a sense of self. Adults who do not develop a positive self-concept in adolescence may experience feelings of loneliness and emotional isolation.<sup>61</sup>

- **Generativity vs. Stagnation**

When people reach their 40s, they enter a time period known as middle adulthood that extends to the mid-60s. The developmental task of middle adulthood is generativity versus stagnation. Generativity involves finding your life's work and contributing to the development of others through activities such as volunteering, mentoring, and raising children. Adults who do not master this developmental task may experience stagnation with little connection to others and little interest in productivity and self-improvement.<sup>62</sup>

- **Integrity vs. Despair**

The mid-60s to the end of life is a period of development known as late adulthood. People in late adulthood reflect on their lives and feel either a sense of satisfaction or a sense of failure. People who feel proud of their accomplishments feel a sense of integrity and often look back on their lives with few regrets. However, people who are not successful at this stage may feel as if their life has been wasted. They focus on what “would

61. Psychology 2e by [OpenStax](https://openstax.org/books/psychology-2e/pages/1-introduction) is licensed under [CC BY 4.0](https://creativecommons.org/licenses/by/4.0/). Access for free at <https://openstax.org/books/psychology-2e/pages/1-introduction>

62. Psychology 2e by [OpenStax](https://openstax.org/books/psychology-2e/pages/1-introduction) is licensed under [CC BY 4.0](https://creativecommons.org/licenses/by/4.0/). Access for free at <https://openstax.org/books/psychology-2e/pages/1-introduction>



have,” “should have,” or “could have” been. They face the end of their lives with feelings of bitterness, depression, and despair.<sup>63</sup>

## Piaget’s Cognitive Theory of Development

Jean Piaget (1896–1980) studied childhood development by focusing on children’s cognitive growth. He believed that thinking is a central aspect of development and that children are naturally inquisitive but do not think and reason like adults. Children explore the world as they attempt to make sense of their experiences. His theory explains that humans move from one stage to another as they seek cognitive equilibrium and mental balance. There are four stages in Piaget’s theory of development that occur in children from all cultures<sup>64, 65</sup>:

- **Sensorimotor period.** The first stage extends from birth to approximately two years and is a period of rapid cognitive growth. During this period, infants develop an understanding of the world by coordinating sensory experiences (seeing, hearing) with motor actions (reaching, touching). The main development during the sensorimotor stage is the understanding that objects exist, and events occur in the world independently of one’s own actions. Infants develop an understanding of what they want and what they must do to have their needs met. They begin to understand language used by those around them to make needs met.

63. Psychology 2e by [OpenStax](#) is licensed under [CC BY 4.0](#). Access for free at <https://openstax.org/books/psychology-2e/pages/1-introduction>

64. Vallotton, C. D., & Fischer, K. W. (2008). Cognitive development. *Encyclopedia of Infant and Early Childhood Development*. Academic Press. pp. 286-298. <https://doi.org/10.1016/B978-012370877-9.00038-4>

65. McLeod, S. (2025). *Piaget’s theory and stages of development*. SimplyPsychology. <https://www.simplypsychology.org/piaget.html>

- **Preoperational period.** The second stage begins in the toddler years. This continues through early school-age years. This is the time frame when children learn to think in images and symbols. Play is an important part of cognitive development during this period.
- **Concrete Operations period.** Older school-age children (age 7 years to 11 years) learn to think in terms of processes and can understand that there is more than one perspective when discussing a concept. This stage is considered a major turning point in the child's cognitive development because it marks the beginning of logical or operational thought.
- **Formal Operations period.** Children enter this stage around age 12 as they become self-conscious and egocentric. Adolescents gain the ability to think in an abstract manner by manipulating ideas in their head. Moving toward adulthood, this further develops into the ability to critically reason.

## Cognitive Impairment

**Cognitive impairment** is a term used to describe impairment in mental processes that drive how an individual understands and acts in the world, affecting the acquisition of information and knowledge. Cognitive impairments can range from mild impairments, such as impairments in cognitive operations, to profound intellectual impairments causing minimal independent functioning. Components of cognitive functioning include attention, decision-making, general knowledge, judgment, language, memory, perception, planning, and reasoning.<sup>66</sup>

▶ Review information about cognitive impairments associated

66. Schofield, D. W. (2018, December 26). *Cognitive deficits*. Medscape.  
<https://emedicine.medscape.com/article/917629-overview>

- ▶ with dementia and Alzheimer’s disease in the “[Cognitive Impairments](#)” chapter of *Open RN Nursing Fundamentals, 2e*.

**Intellectual disability** (formerly referred to as mental retardation) is a diagnostic term that describes intellectual and adaptive functioning deficits identified during the developmental period. In the United States, the developmental period refers to the span of time prior to the age of 18 years. Children with intellectual disabilities may demonstrate a delay in developmental milestones (e.g., sitting, speaking, walking) or demonstrate mild cognitive impairments that may not be identified until school-age. Intellectual disability is typically nonprogressive and lifelong. It is diagnosed by multidisciplinary clinical assessments and standardized testing and is treated with a multidisciplinary treatment plan that maximizes quality of life.<sup>67</sup>

## Resilience

When assessing an individual’s developmental level, it is important to consider possible effects of adverse childhood events (ACEs) on their development. Science tells us that some children develop **resilience**, the ability to overcome serious hardship or traumatic experiences, while others do not. One way to understand the development of resilience is to visualize a seesaw. Protective experiences and coping skills on one side counterbalance significant adversity on the other. Resilience is evident when a child’s health

67. Schofield, D. W. (2018, December 26). *Cognitive deficits*. Medscape. <https://emedicine.medscape.com/article/917629-overview>

and development tip toward positive outcomes – even when a heavy load of factors is stacked on the negative outcome side.<sup>68</sup>

The most common factor for children who develop resilience is at least one stable and committed relationship with a supportive parent, caregiver, or other adult. These relationships provide the personalized responsiveness and protection that buffer children from developmental disruption. They also build their ability to plan, monitor, and regulate behavior that enables children to respond adaptively to adversity and thrive. This combination of supportive relationships, adaptive skill-building, and positive experiences is the foundation of resilience.<sup>69</sup>

The capabilities that underlie resilience can be strengthened at any age. It is never too late to build resilience. Age-appropriate, health-promoting activities can significantly improve the chances that an individual will recover from stress-inducing experiences. For example, regular physical exercise, stress management activities, and programs that actively promote self-regulation skills can improve the abilities of children and adults to cope with adversity in their lives.<sup>70</sup>

► Read more about promoting resilience across the life span at [Harvard's Center on the Developing Child \*Resilience\* webpage.](https://developingchild.harvard.edu/science/key-concepts/resilience/)

68. Center on the Developing Child. (n.d.). *Resilience*. Harvard University. <https://developingchild.harvard.edu/science/key-concepts/resilience/>

69. Center on the Developing Child. (n.d.). *Resilience*. Harvard University. <https://developingchild.harvard.edu/science/key-concepts/resilience/>

70. Center on the Developing Child. (n.d.). *Resilience*. Harvard University. <https://developingchild.harvard.edu/science/key-concepts/resilience/>

## Cultural Considerations

Cultures and communities exhibit and explain symptoms of mental illness and manifest stress in various ways. Nurses should be aware of relevant contextual information stemming from a client's culture, race, ethnicity, religion, or geographical origin. For example, uncontrollable crying and headaches are symptoms of panic attacks in some cultures, whereas difficulty breathing may be the primary symptom in another culture. Understanding such distinctions will help nurses effectively treat them.<sup>71</sup>

At the center of client-centered care is practicing with cultural humility and inclusiveness. In the 2021 edition of *Nursing: Scope and Standards of Practice*, the American Nurses Association (ANA) established a Standard of Professional Performance called *Respectful and Equitable Practice*. This standard is defined as, "The registered nurse practices with cultural humility and inclusiveness." **Cultural humility** is "a humble and respectful attitude toward individuals of other cultures that pushes one to challenge their own cultural biases, realize they cannot know everything about other cultures, and approach learning about other cultures as a life-long goal and process."<sup>72</sup>

**Inclusiveness** is defined as "the practice of providing equal access to opportunities and resources for people who might otherwise be excluded or marginalized, such as those having physical or mental disabilities or belonging to other minority groups."<sup>73</sup> Read the ANA competencies for the *Respectful and Equitable Practice* standard in the following box.

71. American Psychiatric Association. (n.d.). *DSM-5 fact sheets*.

<https://www.psychiatry.org/psychiatrists/practice/dsm/educational-resources/dsm-5-fact-sheets>

72. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association

73. Oxford Learner's Dictionaries. (n.d.). *Inclusion*. Oxford University Press.

<https://www.oxfordlearnersdictionaries.com/us/definition/english/inclusion>

## ANA's Respectful and Equitable Practice Competencies<sup>74</sup>

The registered nurse:

- Demonstrates respect, equity, and empathy in actions and interactions with all health care consumers.
- Respects consumer decisions without bias.
- Participates in life-long learning to understand cultural preferences, worldviews, choices, and decision-making processes of diverse consumers.
- Reflects upon personal and cultural values, beliefs, biases, and heritage.
- Applies knowledge of differences in health beliefs, practices, and communication patterns without assigning values to the differences.
- Addresses the effects and impact of discrimination and oppression on practice within and among diverse groups.
- Uses appropriate skills and tools for the culture, literacy, and language of the individuals and population served.
- Communicates with appropriate language and behaviors, including the use of qualified health care interpreters and translators in accordance with consumer needs and preferences.
- Serves as a role model and educator for cultural humility and the recognition and appreciation of diversity and inclusivity.
- Identifies the cultural-specific meaning of interactions,

<sup>74</sup>. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association

terms, and content.

- Advocates for policies that promote health and prevent harm among diverse health care consumers and groups.
- Promotes equity in all aspects of health and health care.
- Advances organizational policies, programs, services, and practices that reflect respect, equity, and values for diversity and inclusion.

- ▶ Read more about cultural humility and advocating for the values, beliefs, and preferences of diverse clients in the “[Diverse Clients](#)” chapter of *Open RN Nursing Fundamentals, 2e*.

## 4.4 Diagnosis

The *Diagnosis* Standard of Practice by the American Nurses Association states, “The registered nurse analyzes assessment data to determine actual or potential diagnoses, problems, and issues.”<sup>1</sup> Review the competencies for the *Diagnosis* Standard of Practice for registered nurses in the following box.

### **ANA’s Diagnosis Competencies<sup>2</sup>**

The registered nurse:

- Identifies actual or potential risks to the health care consumer’s health and safety or barriers to health, which may include, but are not limited to, interpersonal, systematic, cultural, socioeconomic, or environmental circumstances.
- Uses assessment data, standardized classification systems, technology, and clinical decision support tools to articulate actual or potential diagnoses, problems, and issues.
- Identifies the health care consumer’s strengths and abilities, including, but not limited to, support systems, health literacy, and engagement in self-care.
- Verifies the diagnoses, problems, and issues with the health care consumer and interprofessional colleagues.
- Prioritizes diagnoses, problems, and issues based on

1. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

2. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.



mutually established goals to meet the needs of the health care consumer across the health-illness continuum and the care continuum.

- Documents diagnoses, problems, strengths, and issues in a manner that facilitates the development of the expected outcomes and collaborative plan.

- ▶ Review how to analyze assessment data, make hypotheses, and create nursing diagnoses statements in the “[Diagnosis](#)” section of the “Nursing Process” chapter in *Open RN Nursing Fundamentals*.

## Nursing Diagnoses

A nursing diagnosis is “a clinical judgment concerning a human response to health conditions/life processes, or a vulnerability to that response, by an individual, family, group, or community.”<sup>3</sup> Nursing diagnoses are customized to each client and drive the development of the nursing care plan. The nurse should refer to an evidence-based care planning resource and review the definitions and defining characteristics of the hypothesized nursing diagnoses.

Recall that nursing diagnoses are different from medical diagnoses and

3. Herdman, T. H., & Kamitsuru, S. (Eds.). (2021). *Nursing diagnoses: Definitions and classification, 2021-2023*. Thieme Publishers.

mental health diagnoses. Medical diagnoses focus on medical problems that have been identified by the physician, physician's assistant, or advanced nurse practitioner. Mental health diagnoses are established by mental health experts, such as psychiatrists, psychologists, and advanced practice psychiatric-mental health nurses, using the *Diagnostic and Statistical Manual of Mental Disorders (DSM-5-TR)*. See Figure 4.5<sup>4</sup> for an illustration of several mental health diagnoses.




Figure 4.5 Mental Health Diagnoses

Nursing diagnoses focus on the human response to health conditions and life processes and are established by registered nurses. Clients with the same mental health diagnosis will often respond differently and thus have different nursing diagnoses. For example, two clients may have the same diagnosis of Major Depressive Disorder. However, one client may demonstrate a high risk

4. “[Mental Disorder Silhouette.png](#)” by [Paget Michael Creelman](#) is licensed under [CC BY-SA 4.0](#)

for suicide whereas another client may experience impaired nutrition due to lack of appetite. The nurse must consider these different responses when creating an individualized nursing care plan.

## Prioritization

After identifying nursing diagnoses, the next step is prioritizing specific needs of the client. Prioritization is the process of identifying the most significant problems and the most important interventions to implement based on a client's current status.

It is essential that life-threatening concerns and crises are quickly identified and addressed immediately. Depending on the severity of a problem, the steps of the nursing process may be performed in a matter of seconds for life-threatening concerns. Nurses must recognize cues signaling a change in client condition, apply evidence-based practices in a crisis, and communicate effectively with interprofessional team members. Most client care situations fall somewhere between a crisis and routine care.

**Maslow's Hierarchy of Needs** is commonly used to prioritize the most urgent client needs. It is based on the theory that people are motivated by five levels of needs: physiological, safety, love, esteem, and self-actualization. The bottom levels of the pyramid represent the priority physiological needs intertwined with safety, whereas the upper levels focus on belonging, esteem, and self-actualization. Physiological needs must be met before focusing on higher level needs.<sup>5</sup> For example, priorities for a client experiencing mania are the need for food, fluid, and sleep, as well as controlling the agitation and impulsivity to ensure safety. These needs would need to be met before focusing on strategies to improve relationships with family and friends, build respect and acceptance for self and others, and engage in activities promoting personal growth. It is important to note that although safety is not

5. Maslow, A. H. (1943). A theory of human motivation. *Psychological Review*, 50(4), 370–396. <https://doi.org/10.1037/h0054346>

described as a top priority in this theory, nurses must always prioritize safety needs in addition to physiological needs. See Figure 4.6<sup>6</sup> for an image of Maslow’s Hierarchy of Needs.




Figure 4.6 Maslow's Hierarchy of Needs

## Common Nursing Diagnoses for Mental Health Conditions

Commonly used nursing diagnoses related to caring for clients with mental health conditions are included in Table 4.4. As always, when providing client care, refer to a current, evidence-based nursing care planning resource.

Table 4.4 Common Nursing Diagnoses Related to Mental Health<sup>7</sup>

6. “Maslow’s hierarchy of needs.svg” by J. Finkelstein is licensed under [CC BY-SA 3.0](#)

7. Herdman, T. H., & Kamitsuru, S. (Eds.). (2021). *Nursing diagnoses: Definitions and classification, 2021-2023*. Thieme Publishers.

Nursing Diagnosis	Definition	Selected Defining Characteristics
<b>Risk for Suicide</b>	Susceptible to self-inflicted, life-threatening injury.	<ul style="list-style-type: none"> <li>• Reports desire to die</li> <li>• Threats of killing self</li> <li>• Hopelessness</li> <li>• Social isolation</li> <li>• Giving away possessions</li> <li>• Substance misuse</li> <li>• Chronic pain</li> <li>• Sudden change in mood</li> </ul>
<b>Ineffective Coping</b>	A pattern of impaired appraisal of stressors with cognitive and/or behavioral efforts that fails to manage demands related to well-being.	<ul style="list-style-type: none"> <li>• Alteration in concentration</li> <li>• Alteration in sleep pattern</li> <li>• Change in communication pattern</li> <li>• Inability to meet basic needs</li> <li>• Ineffective coping strategies</li> <li>• Insufficient goal-directed behavior</li> <li>• Risk-taking behavior</li> <li>• Substance misuse</li> </ul>

<b>Readiness for Enhanced Coping</b>	<p>A pattern of effective appraisal of stressors with cognitive and behavioral efforts to manage demands related to well-being, which can be strengthened.</p>	<ul style="list-style-type: none"> <li>• Expresses desire to enhance: <ul style="list-style-type: none"> <li>◦ Knowledge of stress management strategies</li> <li>◦ Social support</li> <li>◦ Use of problem and emotion-oriented strategies</li> <li>◦ Use of spiritual resources</li> </ul> </li> </ul>
<b>Self-Neglect</b>	<p>A collection of culturally framed behaviors involving one or more self-care activities in which there is a failure to maintain a socially accepted standard of health and well-being.</p>	<ul style="list-style-type: none"> <li>• Insufficient personal hygiene</li> <li>• Insufficient environmental hygiene</li> <li>• Nonadherence to health activity</li> </ul>

<b>Fatigue</b>	An overwhelming sustained sense of exhaustion and decreased capacity for physical and mental work at the usual level.	<ul style="list-style-type: none"> <li>• Alteration in concentration</li> <li>• Alteration in libido</li> <li>• Apathy</li> <li>• Impaired ability to maintain usual routines</li> <li>• Ineffective role performance</li> <li>• Nonrestorative sleep patterns</li> </ul>
<b>Imbalanced Nutrition: Less than Body Requirements</b>	Intake of nutrients insufficient to meet metabolic needs.	<ul style="list-style-type: none"> <li>• Food intake less than recommended daily allowance</li> <li>• Insufficient interest in food</li> <li>• Body weight 20% or more below ideal weight range</li> <li>• Read more in the <a href="#">“Nutrition”</a> chapter in <i>Open RN Nursing Fundamentals</i></li> </ul>

<b>Constipation</b>	Decrease in normal frequency of defecation accompanied by difficult or incomplete passage of stool and/or passage of excessively dry, hard stool.	<ul style="list-style-type: none"> <li>• Read more in the “<a href="#">Constipation</a>” section of the “Elimination” chapter in <i>Open RN Nursing Fundamentals</i></li> <li>• Related factors: depression, emotional disturbance, medication side effects, imbalanced nutrition, and poor intake</li> </ul>
<b>Sleep Deprivation</b>	Prolonged periods of time without sustained natural, periodic suspension of relative consciousness that provides rest.	<ul style="list-style-type: none"> <li>• Anxiety</li> <li>• Agitation</li> <li>• Irritability</li> <li>• Restlessness</li> <li>• Alteration in concentration</li> <li>• Fatigue</li> <li>• Transient paranoia</li> <li>• Read more in the “<a href="#">Sleep and Rest</a>” chapter of <i>Open RN Nursing Fundamentals</i></li> </ul>



<b>Social Isolation</b>	Aloneness experienced by the individual and perceived as imposed by others and as a negative or threatening state.	<ul style="list-style-type: none"> <li>• Absence of support system</li> <li>• Flat or sad affect</li> <li>• Feeling different from others</li> <li>• Values incongruent with social norms</li> </ul>
<b>Chronic Low Self-Esteem</b>	Negative evaluation and/or feelings about one's own capabilities, lasting at least three months.	<ul style="list-style-type: none"> <li>• Repeatedly unsuccessful in life events</li> <li>• Underestimates ability to deal with situation</li> <li>• Exaggerates negative feedback about self</li> <li>• Excessive seeking of reassurance</li> <li>• Nonassertive behavior</li> </ul>
<b>Hopelessness</b>	Subjective state in which an individual sees limited or no alternatives or personal choices available and is unable to mobilize energy on own behalf.	<ul style="list-style-type: none"> <li>• Decrease in: <ul style="list-style-type: none"> <li>◦ Appetite</li> <li>◦ Affect</li> <li>◦ Verbalization</li> <li>◦ Initiative</li> </ul> </li> <li>• Shrugging in response to speaker</li> </ul>

<b>Spiritual Distress</b>	A state of suffering related to the impaired ability to experience meaning in life through connections with self, others, the world, or a superior being.	<ul style="list-style-type: none"> <li>• Ineffective coping strategies</li> <li>• Perceived insufficient meaning in life</li> <li>• Separation from support system</li> <li>• Anger</li> <li>• Hopelessness</li> </ul>
<b>Readiness for Enhanced Knowledge</b>	A pattern of cognitive information related to a specific topic or its acquisition, which can be strengthened.	<ul style="list-style-type: none"> <li>• Expresses desire to enhance learning</li> </ul>

### Sample Case A

During an interview with a 32-year-old client diagnosed with Major Depressive Disorder, Mr. J. exhibited signs of a sad affect and hopelessness. He expressed desire to die and reported difficulty sleeping and a lack of appetite with weight loss. He reports he has not showered in over a week, and his clothes have a strong body odor. The nurse analyzed this data and created four nursing diagnoses using a nurse care plan reference<sup>8</sup>:

- *Hopelessness* related to social isolation
- *Risk for Suicide* as manifested by reported desire to die

8. Ackley, B., Ladwig, G., Makic, M. B., Martinez-Kratz, M., & Zanotti, M. (2020). *Nursing diagnosis handbook: An evidence-based guide to planning care* (12th ed.). Elsevier.

- *Imbalanced Nutrition: Less than Body Requirements* related to insufficient dietary intake
  - *Self-Neglect* related to insufficient personal hygiene
- ▶ The nurse established the top priority nursing diagnosis of *Risk for Suicide* and immediately screened for suicidal ideation and a plan using the [Patient Safety Screener](#).

## 4.5 Outcome Identification

The *Outcomes Identification* Standard of Practice by the American Nurses Association states, “The registered nurse identifies expected outcomes for a plan individualized to the health care consumer or the situation.”<sup>1</sup> Review the competencies for the *Outcomes Identification* Standard of Practice for registered nurses in the following box.

### **ANA’s Outcomes Identification Competencies<sup>2</sup>**

The registered nurse:

- Engages with the health care consumer, interprofessional team, and others to identify expected outcomes.
- Collaborates with the health care consumer to define expected outcomes integrating the health care consumer’s culture, values, and ethical considerations.
- Formulates expected outcomes derived from assessments and diagnoses.
- Integrates evidence and best practices to identify expected outcomes.
- Develops expected outcomes that facilitate coordination of care.
- Identifies a time frame for the attainment of expected outcomes.

1. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

2. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

- Documents expected outcomes as measurable goals.
- Identifies the actual outcomes in relation to expected outcomes, safety, and quality standards.
- Modifies expected outcomes based on the evaluation of the status of the health care consumer and situation.

An **outcome** is a “measurable behavior demonstrated by the client who is responsive to nursing interventions.”<sup>3</sup> After nursing interventions are implemented, the nurse evaluates if the outcomes were met, partially met, or not met in the time frame indicated for that client.

Outcome identification includes setting short-term and long-term goals and then creating specific expected outcome statements for each nursing diagnosis. Outcome statements are always client-centered. They should be developed collaboratively with the client and individualized to meet the client’s unique needs, values, and cultural beliefs. They should start with the phrase “*The client will...*” Outcome statements should be directed at resolving the defining characteristics for that nursing diagnosis. Additionally, the outcome must be something the client is willing to cooperate in achieving.

- ▶ Read more about how to use the “[Motivational Interviewing](#)” technique in setting individualized goals and expected outcomes with a client in the “Therapeutic Communication and Nurse-Client Relationship” chapter.

3. Herdman, T. H., & Kamitsuru, S. (Eds.). (2021). *Nursing diagnoses: Definitions and classification, 2021-2023*. Thieme Publishers New York.

Outcome statements should contain five components easily remembered using the “SMART” mnemonic:

- **S**pecific
- **M**easurable
- **A**ttainable/Action-oriented
- **R**elevant/Realistic
- **T**ime frame

See Figure 4.7<sup>4</sup> for an image of the SMART components of outcome statements.



Figure 4.7 SMART Components

- ▶ Review how to create “SMART” expected outcomes in the “[Nursing Process](#)” chapter of *Open RN Nursing Fundamentals*.

4. “SMART-goals.png” by [Dungdm93](#) is licensed under [CC BY-SA 4.0](#)

## Unfolding Case A

Recall Sample Case A in the “[Diagnosis](#)” section regarding the 32-year-old male diagnosed with Major Depressive Disorder. The nurse created these four nursing diagnoses:

- *Hopelessness* related to social isolation
- *Risk for Suicide* as manifested by reported desire to die
- *Imbalanced Nutrition: Less than Body Requirements* related to insufficient dietary intake
- *Self-Neglect* related to insufficient personal hygiene

The nurse established the top priority nursing diagnosis of *Risk for Suicide* and immediately screened for suicidal ideation and a plan using the Patient Safety Screener.

The nurse then identified the following SMART expected outcome for the nursing diagnosis *Risk for Suicide* related to reported desire to die: *The client will remain free from self-harm self during the hospitalization stay.*

## 4.6 Planning

The *Planning* Standard of Practice by the American Nurses Association states, “The registered nurse develops a collaborative plan encompassing strategies to achieve expected outcomes.”<sup>1</sup>

Review the competencies for the *Planning* Standard of Practice for registered nurses in the following box.

### **ANA’s Planning Competencies<sup>2</sup>**

The registered nurse:

- Develops an individualized, holistic, evidence-based plan in partnership with the health care consumer, family, significant others, and interprofessional team.
- Designs innovative nursing practices that can be incorporated into the plan.
- Prioritizes elements of the plan based on the assessment of the health care consumer’s level of safety needs to include risks, benefits, and alternatives.
- Establishes the plan priorities with the health care consumer, family, significant others, and interprofessional team.
- Advocates for compassionate, responsible, and appropriate use of interventions to minimize unwarranted or unwanted

1. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

2. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.



treatment, health care consumer suffering, or both.

- Includes strategies designed to address each of the identified diagnoses, health challenges, issues, or opportunities. These strategies may include, but are not limited to, maintaining health and wellness; promotion of comfort; promotion of wholeness, growth, and development; promotion and restoration of health and wellness; prevention of illness, injury, disease, complications, and trauma; facilitation of healing; alleviation of suffering; supportive care; and mitigation of environmental or occupational risks.
- Incorporates an implementation pathway that describes an overall timeline, steps, and milestones.
- Provides for the coordination and continuity of care.
- Identifies cost and economic implications of the plan.
- Develops a plan that reflects compliance with current statutes, rules, regulations, and standards.
- Modifies the plan according to the ongoing assessment of the health care consumer's response and other outcome indicators.
- Documents the plan using standardized language or recognized terminology.
- Actively contributes at all levels in the development and continuous improvement of systems that support the planning process.

As always, consult a current, evidence-based nursing care planning resource when planning nursing interventions individualized to each client's needs. You might be asking yourself, "How do I know what evidence-based nursing interventions to include in the nursing care plan regarding mental health care?" There are several sources that can be used to select nursing

interventions. Many agencies have care planning tools and references included in the electronic health record that are easily documented in the client chart. Additionally, the Substance Abuse and Mental Health Services Administration (SAMHSA) maintains an evidence-based resource center.<sup>3</sup>

- ▶ Access the [\*Evidence-Based Resource Center\*](#) maintained by the Substance Abuse and Mental Services Administration (SAMHSA).

See sample planned nursing interventions for a client who has been diagnosed with *Risk for Suicide* in Table 4.6.

Table 4.6 Sample Nursing Interventions for Risk for Suicide<sup>4,5</sup>

3. Substance Abuse and Mental Health Services Administration. (n.d.). *Evidence-based practices resource center*. <https://www.samhsa.gov/resource-search/ebp>
4. DeAngelis, T. (2019). Better relationships with patients lead to better outcomes. *Monitor on Psychology*, 50(10), 38. <https://www.apa.org/monitor/2019/11/ce-corner-relationships>
5. Ackley, B., Ladwig, G., Makic, M. B., Martinez-Kratz, M., & Zanotti, M. (2020). *Nursing diagnosis handbook: An evidence-based guide to planning care* (12th ed.). Elsevier.

Nursing Intervention	Rationale
The nurse will...	
Use an evidence-based process to conduct a suicide risk assessment.	Clients with suicidal ideation vary widely in their risk for a suicide attempt depending on whether they have a plan, intent, and past history of attempts. In-depth assessment of clients who screen positive for suicide risk must be completed to determine how to appropriately treat them.
Document and communicate the client's overall level of risk for suicide with the treatment team and the plan for mitigating their risk for suicide.	All interprofessional health care team members who might come in contact with a client at risk for suicide must be aware of the level of risk and the mitigation plans to reduce that risk. This information should be explicitly documented in the client's record.
Perform an environmental risk assessment and remove features that could be used to attempt suicide.	The Veteran's Health Administration showed that use of a Mental Health Environment of Care Checklist to facilitate a thorough, systematic environmental assessment reduced the rate of suicide from 4.2 per 100,000 admissions to 0.74 per 100,000 admissions.
Administer prescribed treatment and collaboratively manage psychiatric symptoms that may be contributing to the client's suicidal ideation or behavior.	Symptoms of the disorder may require treatment with antidepressant, antipsychotic, or antianxiety medications. A systematic review has shown a significant effect for cognitive behavioral therapy in reducing suicidal behavior.
Express desire to help the client and validate the client's experience of psychological pain while maintaining a safe environment for the client.	The nurse must reconcile their goal of preventing suicide with recognition of the client's goal to alleviate their psychological pain.

Develop a positive therapeutic relationship with the client; do not make promises that may not be kept.	Nurses connect suicidal clients with humanity by guiding the client, encouraging effective coping strategies, and helping them connect appropriately with others.
Determine the client's need for supervision and assign a room near the nursing station as necessary.	Close assignment increases ease of observation and availability for a rapid response in the event of a suicide attempt.
Search the newly hospitalized client and the client's personal belongings for weapons or potential weapons and hoarded medications during the admission process and remove dangerous items.	Clients with suicidal ideation may bring the means with them. This action is necessary to maintain a hazard-free environment and client safety.
Limit access to windows and exits unless locked and shatterproof, as appropriate. Ensure exits are secure.	Suicidal behavior may include attempts to jump out of windows or escape to find other means of suicide.
Place the client in the least restrictive, safe, and monitored environment that allows for the necessary level of observation. Assess suicidal risk at least daily and more frequently as warranted.	Close observation of the client is necessary for safety as long as the intent remains high. Suicide risk should be assessed at frequent intervals to adjust suicide precautions and ensure restrictions continue to be appropriate.
Consider strategies to decrease isolation and opportunity to act on harmful thoughts (e.g., use of a sitter).	Clients have reported feeling safe and having their hope restored in response to close observation.
Create a safety plan that includes a no-suicide contract. Contract verbally or in writing with the client for no self-harm and recontract at appropriate intervals.	Discussing thoughts of suicide and self-harm with a trusted person can provide relief for the client. A safety plan gets the subject out in the open and places some of the responsibility for safety with the client. However, research has suggested that self-harm is not prevented by contracts, and ongoing assessment of suicide risk is necessary.

Explain suicide precautions and relevant safety issues to the client and family (purpose, duration, behavioral expectations, and behavioral consequences).	Suicide precautions may be viewed as restrictive. Clients have reported the loss of privacy as distressing. Explaining the reasoning for safety precautions helps the client understand why they are being used even though they may feel restrictive and distressing. When clients and family members understand the reasoning for the precautions, they are more likely to comply.
Verify the client has taken medications as ordered (e.g., conduct mouth checks after medication administration).	The client may attempt to hoard medications for a later suicide attempt.
Maintain increased surveillance of the client whenever the use of an antidepressant has been initiated or the dose increased.	Antidepressant medications take anywhere from 2 to 6 weeks to achieve full efficacy. During that period, the client's energy level may increase although the depression has not yet lifted, which increases the potential for suicide.
Involve the client in treatment planning and self-care management of psychiatric disorders.	Self-care management promotes feelings of self-efficacy. The more clients participate in their own care, the less powerless and hopeless they feel.
Assist the client in identifying a network of supportive persons and resources (e.g., family, clergy, care providers).	Social support and positive events were found to have a protective effect against suicidal ideation.
Document client behavior in detail to support involuntary admission if actively suicidal.	Read more about involuntary admissions in the " <a href="#">Patient Rights</a> " section of the "Legal and Ethical Considerations" chapter. Involuntary inpatient admissions serve to keep the client safe from harm. Involuntary outpatient commitment is also available in many states and can improve treatment, reduce the likelihood of hospital readmission, and reduce episodes of violent behavior in persons with severe psychiatric illnesses.

Involve the family in discharge planning (e.g., illness/medication teaching, recognition of increasing suicidal risk, client's plan for dealing with recurring suicidal thoughts, and community resources).	Family members must learn how to respond to cues early, support the treatment regimen, and encourage the client to initiate an emergency plan. When family members are aware of cues, treatments, and emergency plans, clients are less likely to act on thoughts of suicide or self-harm.
Before discharge from the hospital, ensure the client has a safety plan to use after discharge, including a supply of prescribed medications and a plan for outpatient follow-up. Ensure they understand the plan or have a caregiver able and willing to follow the plan, as well as the ability to access outpatient treatment.	Clients may have difficulty concentrating on the plan for follow-up. They may need assistance from others to ensure prescriptions are filled, appointments are attended, and transportation is available to appointments.
In the event of a client's suicide, refer the family to a support group for survivors of suicide.	Psychoeducational support group participants found relief in sharing their bereavement with others.

Review the "[Establishing Safety](#)" section for clients at risk for suicide in Chapter 1.

## 4.7 Implementation

The *Implementation* Standard of Practice by the American Nurses Association (ANA) states, “The registered nurse implements the identified plan.”<sup>1</sup> Review the competencies for the *Implementation* Standard of Practice for registered nurses in the following box.

### **ANA’s Implementation Competencies<sup>2</sup>**

The registered nurse:

- Demonstrates caring behaviors to develop therapeutic relationships.
- Provides care that focuses on the health care consumer.
- Advocates for the needs of diverse populations across the life span.
- Uses critical thinking and technology solutions to implement the nursing process to collect, measure, record, retrieve, trend, and analyze data and information to enhance health care consumer outcomes and nursing practice.
- Partners with the health care consumer to implement the plan in a safe, effective, efficient, timely, and equitable manner.
- Engages interprofessional team partners with

1. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

2. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

implementation of the plan through collaboration and communication across the continuum of care.

- Uses evidence-based interventions and strategies to achieve mutually identified goals and outcomes specific to the problem or needs.
- Delegates according to the health, safety, and welfare of the health care consumer.
- Delegates after considering the circumstance, person, task, direction or communication, supervision, and evaluation.
- Considers the state's Nurse Practice Act regulations, institution, and regulatory entities while maintaining accountability for the care.
- Documents implementation and any modifications, including changes or omissions, of the identified nursing care plan.

In addition to these competencies, the American Nurses Association established two additional subcategories for the *Implementation* standard: *Coordination of Care* and *Health Teaching and Health Promotion*. In addition to these basic subcategories, the American Psychiatric Association established additional subcategories for registered nurses working in psychiatric/mental health settings: *Pharmacological, Biological, and Integrative Therapies*; *Milieu Therapy*; and *Therapeutic Relationship and Counseling*. Each of these additional subcategories of *Implementation* is discussed in the following subsections.

## Coordination of Care

Review the competencies for the *Coordination of Care* Standard of Care in the following box.



## ANA's Coordination of Care Competencies<sup>3</sup>

The registered nurse:

- Collaborates with the health care consumer and the interprofessional team to help manage health care based on mutually agreed-upon outcomes.
- Organizes the components of the plan with input from the health care consumer and other stakeholders.
- Manages the health care consumer's care to reach mutually agreed-upon outcomes.
- Engages health care consumers in self-care to achieve preferred goals for quality of life.
- Assists the health care consumer to identify options for care and navigate the health care system and its services.
- Communicates with the health care consumer, interprofessional team, and community-based resources to effect safe transitions in continuity of care.
- Advocates for the delivery of dignified and person-centered care by the interprofessional team.
- Documents the coordination of care.

## Health Teaching and Health Promotion

Review the competencies for *Health Teaching and Health Promotion* in the following box.

3. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

## ANA's Health Teaching and Health Promotion Competencies<sup>4</sup>

The registered nurse:

- Provides opportunities for the health care consumer to identify needed health promotion, disease prevention, and self-management topics such as:
  - Healthy lifestyles
  - Self-care and risk management
  - Coping, adaptability, and resiliency
- Uses health promotion and health teaching methods in collaboration with the health care consumer's values, beliefs, health practices, developmental level, learning needs, readiness and ability to learn, language preference, spirituality, culture, and socioeconomic status.
- Uses feedback from the health care consumer and other assessments to determine the effectiveness of the employed strategies.
- Uses technologies to communicate health promotion and disease prevention information to the health care consumer.
- Provides health care consumers with information and education about intended effects and potential adverse effects of the plan of care.
- Engages consumer alliance and advocacy groups in health teaching and health promotion activities for health care

4. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

consumers.

- Provides anticipatory guidance to health care consumers to promote health and prevent or reduce risk.

## Pharmacological, Biological, and Integrative Therapies

**Biological therapies** are “any form of treatment for mental disorders that attempts to alter physiological functioning, including drug therapies, electroconvulsive therapy, and psychosurgery.”<sup>5</sup> **Integrative therapies** are defined by the American Psychiatric Association (APA) as “psychotherapy that selects theoretical models or techniques from various therapeutic schools to suit the client’s particular problems.”<sup>6</sup> **Psychotherapy interventions** include “all generally accepted and evidence-based methods of brief or long-term therapy, including individual therapy, group therapy, marital or couple therapy, and family therapy. These interventions use a range of therapy models, including, but not limited to, psychodynamic, cognitive, behavioral, and supportive interpersonal therapies to promote insight, produce behavioral change, maintain function, and promote recovery.”<sup>7</sup> Review the competencies for this Standard of Care in the following box.

5. American Psychological Association. (n.d.). *APA Dictionary of Psychology*.  
<https://dictionary.apa.org/>

6. American Psychological Association. (n.d.). *APA Dictionary of Psychology*.  
<https://dictionary.apa.org/>

7. American Nurses Association, American Psychiatric Nurses Association, and International Society of Psychiatric-Mental Health Nurses. (2014). *Psychiatric-mental health nursing: Scope and standards of practice* (2nd ed.).  
Nursebooks.org

## APNA's Pharmacological, Biological, and Integrative Therapies Competencies<sup>8</sup>

"The psychiatric-mental health registered nurse (PMH-RN) incorporates knowledge of pharmacological, biological, and complementary interventions with applied clinical skills to restore the health care consumer's health and prevent future disability."<sup>9</sup>

The PMH-RN:

- Applies current research findings to guide nursing actions related to pharmacology, other biological therapies, and integrative therapies.
- Assesses the health care consumer's response to biological interventions based on current knowledge of pharmacological agent's intended actions, interactive effects, potential untoward effects, and therapeutic doses.
- Includes health teaching for medication management to support health care consumers in managing their own medications and adhering to a prescribed regimen.

8. American Nurses Association, American Psychiatric Nurses Association, and International Society of Psychiatric-Mental Health Nurses. (2014). *Psychiatric-mental health nursing: Scope and standards of practice* (2nd ed.). Nursebooks.org

9. American Nurses Association, American Psychiatric Nurses Association, and International Society of Psychiatric-Mental Health Nurses. (2014). *Psychiatric-mental health nursing: Scope and standards of practice* (2nd ed.). Nursebooks.org

- Provides health teaching about mechanism of action, intended effects, potential adverse effects of the proposed prescription, ways to cope with transitional side effects, and other treatment options, including the selection of a no-treatment option.
- Directs interventions toward alleviating untoward effects of biological interventions.
- Communicates observations about the health care consumer's response to biological interventions to other health clinicians.

- ▶ Read more details about different types of psychotherapy treatments in [Psychology2e](#) by OpenStax and on the National Alliance of Mental Illness (NAMI) [website](#).
- ▶ Read more about medications in the “[Psychotropic Medications](#)” chapter.

## Milieu Therapy

A **therapeutic milieu** is defined by the American Psychiatric Nursing Association as, “A safe, welcoming, supportive, and functional physical treatment environment.” **Milieu therapy** includes nursing interventions used to assist health care consumers to make positive change and promote recovery in a therapeutic milieu. Nursing interventions include providing empathy, assisting in problem-solving, acting as a role model, demonstrating leadership, confronting discrepancies, encouraging self-efficacy, decreasing stimuli when necessary, and manipulating the environment so that the above

interventions can be effective.”<sup>10</sup> Review the APNA competencies for this Standard of Care in the following box.

### **APNA’s Milieu Therapy Competencies<sup>11</sup>**

“The psychiatric-mental health registered nurse (PMH-RN) provides, structures, and maintains a safe, therapeutic, recovery-oriented environment in collaboration with health care consumers, families, and other health care clinicians.”<sup>12</sup>

The PMH-RN:

- Orients the health care consumer and family to the care environment, including the physical environment, the roles of the different health care providers, self-involvement in the treatment and care delivery processes, schedules of events pertinent to their care and treatment, and

10. American Nurses Association, American Psychiatric Nurses Association, and International Society of Psychiatric-Mental Health Nurses. (2014). *Psychiatric-mental health nursing: Scope and standards of practice* (2nd ed.). Nursebooks.org

11. American Nurses Association, American Psychiatric Nurses Association, and International Society of Psychiatric-Mental Health Nurses. (2014). *Psychiatric-mental health nursing: Scope and standards of practice* (2nd ed.). Nursebooks.org

12. American Nurses Association, American Psychiatric Nurses Association, and International Society of Psychiatric-Mental Health Nurses. (2014). *Psychiatric-mental health nursing: Scope and standards of practice* (2nd ed.). Nursebooks.org

expectations regarding safe and therapeutic behaviors.

- Orients health care consumers to their rights and responsibilities particular to the treatment or care environment.
- Establishes a welcome, trauma-sensitive environment using therapeutic interventions, including, but not limited to, sensory or relaxation rooms.
- Conducts ongoing assessments of the health care consumer in relation to the environment to guide nursing interventions in maintaining a safe environment.
- Selects specific activities (both individual and group) that meet the health care consumer's physical and mental health needs for meaningful participation in the milieu and promotion of personal growth.
- Advocates that the health care consumer is treated in the least restrictive environment necessary to maintain the safety of the individual and others.
- Informs the health care consumer in a culturally sensitive manner about the need for limits related to safety and the conditions necessary to remove the restrictions.
- Provides support and validation to health care consumers when discussing their illness experience and seeks to prevent complications of illness.

## Therapeutic Relationship and Counseling

The American Nurses Association states, "Nursing integrates the art and science of caring... It facilitates healing and alleviates suffering through compassionate presence... The act of caring is the first step in the power to

heal.”<sup>13</sup> Jean Watson’s Human Caring Science Theory emphasizes the therapeutic relationship between the client and nurse and highlights the role of the nurse in defining the client as a unique human being to be valued, respected, nurtured, understood, and assisted. In a caring, therapeutic relationship, the nurse implements interventions to promote interpersonal connection, such as listening attentively, making eye contact, using verbal reassurances, and using professional touch with permission.<sup>14</sup> Nurses use several therapeutic techniques during a nurse-client relationship. Read more information in the “[Therapeutic Communication and the Nurse-Client Relationship](#)” chapter.

Read the APNA competencies regarding therapeutic relationship and counseling in the following box.

### **APNA’s Therapeutic Relationship and Counseling Competencies<sup>15</sup>**

“The psychiatric-mental health registered nurse (PMH-RN) uses the therapeutic relationship and counseling interventions to assist health care consumers in their individual recovery journeys by improving and regaining their previous coping abilities,

13. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

14. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

15. American Nurses Association, American Psychiatric Nurses Association, and International Society of Psychiatric-Mental Health Nurses. (2014). *Psychiatric-mental health nursing: Scope and standards of practice* (2nd ed.). Nursebooks.org



fostering mental health, and preventing mental disorder and disability.”<sup>16</sup>

The PMH-RN:

- Uses therapeutic relationship and counseling techniques to promote the health care consumer’s stabilization of symptoms and personal recovery goals.
- Uses the therapeutic relationship and counseling techniques, both in the individual and group setting, to reinforce healthy behaviors and interaction patterns and helps the health care consumer discover individualized health care behaviors to replace unhealthy ones.
- Documents counseling interventions, including, but not limited to, communication and interviewing techniques, problem-solving activities, crisis intervention, stress management, supportive skill building and educational groups, relaxation techniques, assertiveness training, and conflict resolution.

► Read more information about stress management, relaxation techniques, and crisis intervention in the “[Stress, Coping, and Crisis Management](#)” chapter.

16. American Nurses Association, American Psychiatric Nurses Association, and International Society of Psychiatric-Mental Health Nurses. (2014). *Psychiatric-mental health nursing: Scope and standards of practice* (2nd ed.). Nursebooks.org

## Categories of Interventions

Nurses implement several interventions related to each subcategory of the *Implementation* Standard of Care for clients in mental health settings. See Table 4.7 for common nursing interventions in mental health settings for each subcategory.

Table 4.7 Categories of Nursing Mental Health Interventions

Subcategories: Implementation Standard of Care	Sample Nursing Interventions
<b>Coordination of Care</b>	<ul style="list-style-type: none"> <li>• Refer to community support groups for optimal recovery.</li> <li>• Advocate for dignified care with the interprofessional team.</li> <li>• Communicate client trends with interprofessional team members such as cheeking (i.e., not swallowing medications), increased agitation, or propensity toward violence.</li> </ul>
<b>Health Teaching and Health Promotion</b>	<ul style="list-style-type: none"> <li>• Deliver health teaching to clients about self-care and stress management techniques.</li> <li>• Promote health by teaching about adaptive coping strategies such as journaling and daily exercise.</li> </ul>
<b>Pharmacological, Biological, and Integrative Therapies</b>	<ul style="list-style-type: none"> <li>• Provide health teaching about medications' mechanisms of action, intended effects, potential adverse effects, and ways to cope with transitional side effects.</li> </ul>

<b>Milieu Therapy</b>	<ul style="list-style-type: none"> <li>• Encourage client participation within the therapeutic milieu by attending support groups and exercise groups.</li> <li>• Perform intentional rounding at varying times between every 15-60 minutes and document.</li> <li>• Advocate for the least restrictive environment necessary to maintain the safety of the individual and others.</li> <li>• Perform environmental safety scans and eliminate any devices or objects that can cause injury. Remove strings, cords, and drawstrings.</li> </ul>
<b>Therapeutic Relationship and Counseling</b>	<ul style="list-style-type: none"> <li>• Observe for, document, and communicate changes in behavior.</li> <li>• Demonstrate caring behaviors.</li> <li>• Utilize therapeutic communication techniques.</li> </ul>

## Implementing Interventions

Implementation of interventions requires the RN to use critical thinking and clinical judgment. After the initial plan of care is developed, continual reassessment of the client is necessary to detect any changes in the client's condition requiring modification of the plan. The need for continual client reassessment underscores the dynamic nature of the nursing process and is crucial to providing safe care.

During the *Implementation* phase of the nursing process, the nurse prioritizes planned interventions, assesses client safety while implementing interventions, delegates interventions as appropriate, and documents interventions performed.

Prioritizing implementation of interventions follows a similar method as to prioritizing nursing diagnoses. Maslow's Hierarchy of Needs and the ABCs of

airway, breathing, and circulation are used to establish top priority interventions. When possible, least restrictive interventions are preferred. Read more about methods for prioritization under the “[Diagnosis](#)” section of this chapter.

It is essential to consider client safety when implementing interventions. At times, clients may experience a change in condition that makes a planned nursing intervention or provider prescription no longer safe to implement. For example, an established nursing care plan for a client states, “The nurse will ambulate the client 100 feet three times daily.” However, during assessment this morning, the client reports feeling dizzy today, and their blood pressure is 90/60. Using critical thinking and clinical judgment, the nurse decides to not implement the planned intervention of ambulating the client and notifies the provider of suspected side effects of the client’s antidepressant medication. This decision, supporting assessment findings, and notification of the provider should be documented in the client’s chart and also communicated during the shift handoff report.

- ▶ Read more about delegating interventions in the “[Delegation and Supervision](#)” chapter of *Open RN Nursing Management and Professional Concepts, 2e*.

## 4.8 Evaluation

The *Evaluation* Standard of Practice by the American Nurses Association states, “The registered nurse evaluates progress toward attainment of goals and outcomes.”<sup>1</sup> Review the competencies for the *Evaluation* Standard of Practice for registered nurses in the following box.

### **ANA’s Evaluation Competencies<sup>2</sup>**

The registered nurse:

- Uses applicable standards and defined criteria (e.g., Quality and Safety Education for Nurses [QSEN], Quadruple Aim, Institute for Healthcare Improvement [IHI]).
- Conducts a systematic, ongoing, and criterion-based evaluation of the goals and outcomes in relation to the structure, processes, and timelines prescribed in the plan.
- Collaborates with the health care consumer, stakeholders, interprofessional team, and others involved in the care or situation in the evaluation process.
- Determines, in partnership with the health care consumer and other stakeholders, the person-centeredness, effectiveness, efficiency, safety, timeliness, and equitability of the strategies in relation to the responses to the plan and attainment of outcomes.

1. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

2. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

- Uses ongoing assessment data, other data and information resources and benchmarks, research, and meta-analyses for the analytic activities to revise the diagnoses, outcomes, plan, implementation, and evaluation strategies as needed.
- Documents the results of the evaluation.
- Reports evaluation data in a timely fashion.
- Shares evaluation data and conclusions with the health care consumer and other stakeholders to promote clarity and transparency in accordance with state, federal, organizational, and professional requirements.

Evaluation focuses on the effectiveness of the nursing interventions by reviewing the expected outcomes to determine if they were met, partially met, or not met by the time frames indicated. Evaluation includes analysis of data from assessments, screening tools, laboratory results, and pharmacologic interventions, as well as the effectiveness of nursing interventions related to thought process and content. During the *Evaluation* phase, nurses use critical thinking to analyze reassessment data and determine if a client's expected outcomes have been met, partially met, or not met by the time frames established. If outcomes are not met or only partially met by the time frame indicated, the care plan should be revised. If revision is necessary, the nurse should consider which step of the nursing process requires modification. Have additional assessment data been obtained, or have assessment data changed? Has a different nursing diagnosis become a priority? Were the identified goals or expected outcomes unrealistic? Were any interventions not effective?

Reassessment should occur every time the nurse interacts with a client, discusses the care plan with others on the interprofessional team, or reviews updated laboratory or diagnostic test results. Nursing care plans should be updated as higher priority goals emerge. The results of the evaluation must be documented in the client medical record.

Ideally, when the planned interventions are implemented, the client will respond positively, and the expected outcomes are achieved. However, when interventions do not assist in progressing the client toward the expected outcomes, the nursing care plan must be revised to more effectively address the needs of the client. These questions can be used as a guide when revising the nursing care plan:

- Did anything unanticipated occur?
- Has the client's condition changed?
- Have the client's goals and priorities shifted?
- Were the expected outcomes and their time frames realistic?
- Are the nursing diagnoses accurate for this client at this time?
- Are the planned interventions appropriately focused on supporting outcome attainment?
- What barriers were experienced as interventions were implemented?
- Do ongoing assessment data indicate the need to revise diagnoses, outcome criteria, planned interventions, or implementation strategies?
- Are different interventions required?



## 4.9 NCLEX Next Generation

The National Council Licensure Examination for Registered Nurses (NCLEX-RN) is the exam that all nursing graduates must successfully pass to obtain their nursing license and become a registered nurse. The purpose of the NCLEX is to evaluate if a nursing graduate (i.e., candidate) is competent to provide safe, competent, entry-level nursing care. The NCLEX-RN is developed by the National Council of State Board of Nursing (NCSBN), an independent, nonprofit organization composed of the 50 state boards of nursing and other regulatory agencies.<sup>1</sup>

A new edition of the NCLEX called the “Next Generation NCLEX” was released in April 2023 that uses the Clinical Judgment Measurement Model (CJMM) to assess how well the candidate can think critically and use clinical judgment when providing safe nursing care.<sup>2</sup>

The CJMM complements the nursing process but uses different terminology in exam questions to assess the candidate’s clinical judgment. This terminology includes recognize cues, analyze cues, prioritize hypotheses,

1. NCLEX. (n.d.). NCSBN. <https://www.nclex.com/>

2. NCLEX. (n.d.). NCSBN. <https://www.nclex.com/>

generate solutions, take actions, and evaluate outcomes. See Table 4.9 for a comparison of these terms and actions to the nursing process.<sup>3, 4, 5</sup>

Table 4.9 Comparison of the NCSBN Clinical Judgment Measurement Model to the Nursing Process

3. NCLEX. (n.d.). *Clinical judgment measurement model*. NCSBN.  
<https://www.nclex.com/clinical-judgment-measurement-model.page>
4. Ignativicius, V. & Silvestri, L. (2022). *Preparing for the Next-Generation NCLEX (NGN): A “how-to” step-by-step faculty resource manual* [PDF]. Elsevier.  
[https://evolve.elsevier.com/education/wp-content/uploads/sites/2/NGN\\_FacultyGuide\\_Final.pdf](https://evolve.elsevier.com/education/wp-content/uploads/sites/2/NGN_FacultyGuide_Final.pdf)
5. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

NCSBN Clinical Judgment Skill	Description	Corresponding Step of the Nursing Process
<b>Recognize Cues</b>	<p><i>What data is clinically significant?</i></p> <p>Determining what client findings are significant, most important, and of immediate concern to the nurse (i.e., identifying “relevant cues”).</p>	Assessment
<b>Analyze Cues</b>	<p><i>What does the data mean?</i></p> <p>Analyzing data to determine if it is “expected” or “unexpected” or “normal” or “abnormal” for this client at this time according to their age, development, and clinical status.</p> <p>Making a clinical judgment concerning the clients’ “human response to health conditions/life processes, or a vulnerability for that response,” also referred to as “forming a hypothesis.”</p>	<p>Diagnosis</p> <p>(Analysis of Data)</p>
<b>Prioritize Hypotheses</b>	<p><i>What hypotheses should receive priority attention?</i></p> <p>Ranking client conditions and problems according to urgency, complexity, and time.</p>	Planning
<b>Generate Solutions</b>	<p><i>What should be done?</i></p> <p>Planning individualized interventions that meet the desired outcomes for the client; may include gathering additional assessment data.</p>	Planning
<b>Take Action</b>	<p><i>What will I do now?</i></p> <p>Implementing interventions that are safe and most appropriate for the client’s current priority conditions and problems.</p>	Implementation

<b>Evaluate Outcomes</b>	<p><i>Did the interventions work?</i></p> <p>Comparing actual client outcomes with desired client outcomes to determine effectiveness of care and making appropriate revisions to the nursing care plan.</p>	Evaluation
--------------------------	--	------------

It is important to note that NANDA Nursing Diagnoses are not specifically assessed on the NCLEX. However, the ability of a candidate to cluster client data, make hypotheses, prioritize hypotheses, and plan nursing interventions is based on a nursing knowledge base of potential human responses to health problems and life processes (otherwise known as nursing diagnoses).


There are also new “Next Generation” test questions called extended multiple response, extended drag and drop, cloze (drop-down), extended hot spot (highlighting), and matrix-grid<sup>6</sup>:

- **Extended Multiple Response:** Extended Multiple Response items allow candidates to select one or more answer options at a time and uses partial credit scoring.
- **Extended Drag and Drop:** Extended Drag and Drop items allow candidates to move or place response options into answer spaces.
- **Cloze (Drop – Down):** Cloze (Drop – Down) items allow candidates to select one option from a drop-down list. There can be more than one drop-down list in a cloze item. These drop-down lists can be used as words or phrases within a sentence or within tables and charts.
- **Enhanced Hot Spot (Highlighting):** Enhanced Hot Spot items allow candidates to select their answer by highlighting predefined words or phrases. Candidates can select and deselect the highlighted parts by clicking on the words or phrases. These types of items allow an individual to read a portion of a client medical record (e.g., a nursing note, medical

6. NCLEX. (n.d.). NCSBN. <https://www.nclex.com/>

history, lab values, medication record, etc.), and then select the words or phrases that answer the item.

- **Matrix/Grid:** Matrix/Grid items allow the candidate to select one or more answer options for each row and/or column. This item type can be useful in measuring multiple aspects of the clinical scenario with a single item.

View the following YouTube video<sup>7</sup> from NCSBN on Next  
 Generation test items: *The Right Decisions Come from the Right Questions*

Learning activities are incorporated throughout this book to assist students in learning how to use clinical judgment to client-based scenarios and respond to NCLEX Next Generation-style test questions.

<sup>7</sup>. NCSBN. (2019, December 17). *The Right Decisions Come from the Right Questions*. [Video]. YouTube. All rights reserved. <https://youtu.be/ZBXfkINIRF0>

## 4.10 Spotlight Application

Let's review how the nursing process can be applied to Sample Case A introduced in the "[Diagnosis](#)" section of this chapter regarding caring for a suicidal client:

### Assessment

During an interview with a 32-year-old male client diagnosed with Major Depressive Disorder, the client exhibited signs of a sad affect and hopelessness. He expressed desire to die and reported difficulty sleeping and a lack of appetite. He reports he has not showered in over a week and his clothes have a strong body odor.

### Diagnosis

The nurse analyzed this data and created four nursing diagnoses:

- *Hopelessness* related to social isolation
- *Risk for Suicide* as manifested by the reported desire to die
- *Imbalanced Nutrition: Less than Body Requirements* related to insufficient dietary intake
- *Self-Neglect* related to insufficient personal hygiene

The nurse established the top priority nursing diagnosis of *Risk for Suicide* and immediately screened for suicidal ideation and a plan using the Columbia Suicide Severity Rating Scale (C-SSRS).

### Outcome Identification

The nurse identified the following SMART expected outcomes:

- The client will verbalize feelings by the end of the shift.
- The client will remain free from injury during the hospitalization stay.

- The client will progressively gain at least one pound per week toward his ideal body weight (180 pounds).
- The client will participate in daily bathing.

## Planning and Implementation

The nurse implemented planned nursing interventions for *Risk for Suicide* as previously discussed in Table 4.6.

## Evaluation

Day 1: Outcomes partially met. By the end of the shift, the client verbalized feelings related to hopelessness and did not harm himself. He did not agree to participate in taking a bath and only ate 25% of his meal tray. Interventions will be re-attempted on Day 2 and reassessed for effectiveness.

## Sample Documentation

0900: 32-year-old male client diagnosed with Major Depressive Disorder admitted for active suicidal ideation with a plan to do so with a gun. He has the means to accomplish this plan at home. He has expressed the desire to die and reports difficulty sleeping and a lack of appetite for the past two weeks. He reports he has not showered in over a week, and his clothes have a strong body odor. Client was placed in a room near the nursing station and assigned a 1:1 sitter. His personal belongings were removed and placed in a secure area. An environmental scan was completed, and all hazards were removed from the room. He agreed to complete a no-harm contract. Dr. Delgado was notified at 0930. She assessed the client at 0945, and new orders for medications were received and administered. — Zerimiah Alimi, Nursing Student

## 4.11 Learning Activities



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=296#h5p-16>

1

Complete this learning activity to learn about typical clinical decisions a mental health nurse makes every day.



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=296#h5p-61>

2



Test your clinical judgment with a NCLEX Next Generation-style question:  
[Chapter 4, Assignment 1](#)

3

1. “MH Nursing Process Glossary Cards ” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

2. “MH A Day in the Life ” by Sue Dzubay s licensed under [CC BY-NC 4.0](#)

3. “MH Next Gen Nursing Process Question ” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)



Test your clinical judgment with a NCLEX Next Generation-style case study: [Chapter 4, Case Study 1](#)<sup>4</sup>



<sup>4</sup> “MH Next Gen Nursing Process Case Study ” by Kellea Ewen is licensed under [CC BY-NC](#)  
[4.0](#)

## IV Glossary

**ADOPIE:** A mnemonic for the components of the nursing process: Assessment, Diagnosis, Outcomes Identification, Planning, Implementation, and Evaluation.

**Affect:** A client's expression of emotion.

**Akathisia:** Motor restlessness.

**Alexithymia:** The inability to describe emotions with how one is feeling.

**Alogia:** Brief or reduced replies.

**Anhedonia:** The lack of experiencing pleasure in activities normally found enjoyable.

**Apathy:** A lack of feelings, emotions, interests, or concerns.

**Asociality:** A lack of interest in or withdrawal from social interactions and relationships.

**Avolition:** A lack of motivation or inability to initiate and persist in goal-directed activities.

**Biological therapies:** Any form of treatment for mental health disorders that attempts to alter physiological functioning, including drug therapies, electroconvulsive therapy, and psychosurgery.<sup>1</sup>

**Blunted:** A diminished range and intensity of affect or mood.

**Chief complaint:** The client's primary reasons for seeking care.

1. American Psychological Association. (n.d.). *APA Dictionary of Psychology*.  
<https://dictionary.apa.org/>

**Circumstantial:** Speaking with many unnecessary or tedious details without getting to the point of the conversation.

**Clang associations:** Stringing words together that rhyme without logical association and do not convey rational meaning. For example, a client exhibiting clang associations may state, “Here she comes with a cat catch a rat match.”

**Clouded consciousness:** A state of reduced awareness to stimuli.

**Cognition:** The mental action or process of acquiring knowledge and understanding through thought, experience, and the senses. It includes thinking, knowing, remembering, judging, and problem-solving.

**Cognitive impairment:** Impaired mental processes that drive how an individual understands and acts in the world, affecting the acquisition of information and knowledge. Components of cognitive functioning include attention, decision-making, general knowledge, judgment, language, memory, perception, planning, and reasoning.<sup>2</sup>

**Coma:** A state of unarousable unresponsiveness, where vigorous noxious stimuli may not elicit reflex motor responses.

**Congruence:** Consistency of verbal and nonverbal communication.

**Countertransference:** A tendency for the examiner to displace (transfer) their own feelings onto the client, and these feelings may influence the client.

**Cultural humility:** A humble and respectful attitude toward individuals of other cultures that pushes one to challenge their own cultural biases, realize

2. Schofield, D. W. (2018). *Cognitive deficits*. Medscape.  
<https://emedicine.medscape.com/article/917629-overview>

they cannot know everything about other cultures, and approach learning about other cultures as a life-long goal and process.<sup>3</sup>

**Delirium:** An onset of an abnormal mental state, often with fluctuating levels of consciousness, disorientation, irritability, and hallucinations. Delirium is often associated with infection, metabolic disorders, or toxins in the central nervous system.

**Delusions:** A fixed, false belief not held by cultural peers and persisting in the face of objective contradictory evidence. For example, a client may have the delusion that the CIA is listening to their conversations via satellites.

**Development:** Physical, social, and cognitive changes that occur continuously throughout one's life.

**Diagnostic and Statistical Manual of Mental Disorders (DSM-5):** The manual used to make mental health diagnoses established by mental health experts.

**Disheveled:** A client's hair, clothes, or hygiene appears untidy, disordered, unkempt, or messy.

**Distractibility:** A state when the client's attention is easily drawn to unimportant or irrelevant external stimuli.

**Dyskinesia:** Uncontrolled, involuntary movements.

**Dysphoric:** A client's mood or affect exhibiting persistent sadness or depression.

**Echolalia:** Repetitive imitation or echoing of words or phrases spoken by others.

**Euphoric:** A pathologically elevated sense of well-being.

3. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

**Euthymic:** Normal affect and mood with a wide range of emotion appropriate for the situation.

**Family dynamics:** Patterns of interactions among relatives, their roles and relationships, and the various factors that shape their interactions. Because family members rely on each other for emotional, physical, and economic support, they are primary sources of relationship security or stress. Family dynamics and the quality of family relationships can have either a positive or negative impact on an individual's health.

**Flat:** No emotional expression.

**Flight of ideas:** A state where the client frequently shifts from one topic to another with rapid speech, making it seem fragmented. The examiner may feel the client is rambling and changing topics faster than they can keep track, and they probably can't get a word in edgewise.<sup>4</sup> An example of a client exhibiting a flight of ideas is, "My father sent me here. He drove me in a car. The car is yellow in color. Yellow looks good on me."<sup>5</sup>

**Grandiose delusions:** A state of false attribution to the self of great ability, knowledge, importance or worth, identity, prestige, power, accomplishment.<sup>6</sup> Clients may withdraw into an inner fantasy world that's not equivalent to reality, where they have inflated importance, powers, or a specialness that is opposite of what their actual life is like.

4. Martin, D. C. (1990). The mental status examination. In Walker, H. K., Hall, W. D., Hurst, J. W., (Eds.), *Clinical methods: The history, physical, and laboratory examinations*. (3rd ed.). Butterworths. <https://www.ncbi.nlm.nih.gov/books/NBK320/>
5. PsycholoGenie. (n.d.). *The true meaning of flight of ideas explained with examples*. <https://psychologenie.com/flight-of-ideas-meaning-examples>
6. American Psychological Association. (n.d.). *APA Dictionary of Psychology*. <https://dictionary.apa.org/>

**Hallucinations:** False sensory perceptions not associated with real external stimuli that can include any of the five senses (auditory, visual, gustatory, olfactory and tactile). For example, a client may see spiders climbing on the wall or hear voices telling them to do things. These are referred to as “visual hallucinations” or “auditory hallucinations.”

**Homicidal ideation:** Threats or acts of life-threatening harm towards another person.

**Illusions:** Misperceptions of real stimuli. For example, a client may misperceive tree branches blowing in the wind at night to be the arms of monsters trying to grab them.

**Inclusiveness:** The practice of providing equal access to opportunities and resources for people who might otherwise be excluded or marginalized, such as those having physical or mental disabilities or belonging to other minority groups.<sup>7</sup>

**Insight:** The client demonstrates awareness of their situation.

**Integrative therapies:** Psychotherapy that selects theoretical models or techniques from various therapeutic schools to suit the client’s particular problems.<sup>8</sup>

**Intellectual disability:** A diagnostic term that describes intellectual and adaptive functioning deficits identified during the developmental period prior to the age 18.

**Labile:** Rapid changes in emotional responses, mood, or affect that are inappropriate for the moment or the situation.

7. Oxford Learner’s Dictionaries. (n.d.). *Inclusion*. Oxford University Press.  
<https://www.oxfordlearnersdictionaries.com/us/definition/english/inclusion>

8. American Psychological Association. (n.d.). *APA Dictionary of Psychology*.  
<https://dictionary.apa.org/>

**Loose associations:** Jumping from one idea to an unrelated idea in the same sentence. For example, the client might state, “I like to dance; my feet are wet.”<sup>9</sup> The term “word salad” refers to severely disorganized and virtually incomprehensible speech or writing, marked by severe loosening of associations.<sup>10</sup>

**Magical thinking:** A cognitive process in which an individual believes that their thoughts, words, or actions can directly influence events in the physical world in a way that defies the laws of cause and effect.

**Maslow’s Hierarchy of Needs:** A theory commonly used to prioritize the most urgent client needs.

**Mental status examination:** An assessment of a client’s level of consciousness and orientation, appearance and general behavior, speech, motor activity, affect and mood, thought and perception, attitude and insight, and cognitive abilities.

**Milieu therapy:** Nursing interventions used to assist health care consumers to make positive change and promote recovery by creating a therapeutic milieu. Milieu therapy includes interventions such as providing empathy, assisting in problem-solving, acting as a role model, demonstrating leadership, confronting discrepancies, encouraging self-efficacy, decreasing stimuli when necessary, and manipulating the environment so that the above interventions can be effective.<sup>11</sup>

9. American Psychological Association. (n.d.). *APA Dictionary of Psychology*.  
<https://dictionary.apa.org/>

10. American Psychological Association. (n.d.). *APA Dictionary of Psychology*.  
<https://dictionary.apa.org/>

11. American Nurses Association, American Psychiatric Nurses Association, and International Society of Psychiatric-Mental Health Nurses. (2014). *Psychiatric-*

**Mood:** The predominant emotion expressed by an individual.<sup>12</sup>

**Non-suicidal self-injury (NSSI):** Intentional self-inflicted destruction of body tissue without suicidal intention and for purposes not socially sanctioned. Common forms of NSSI include behaviors such as cutting, burning, scratching, and self-hitting.

**Nursing diagnosis:** A clinical judgment concerning a human response to health conditions/life processes or a vulnerability for that response, by an individual, family, group, or community.

**Nursing process:** A critical thinking model based on a systematic approach to client-centered care. Nurses use the nursing process to perform clinical reasoning and make clinical judgments when providing patient care.

**Obsessions:** Persistent thoughts, ideas, images, or impulses that are experienced as intrusive or inappropriate and result in anxiety, distress, or discomfort. Common obsessions include repeated thoughts about contamination, a need to have things in a particular order or sequence, repeated doubts, aggressive impulses, and sexual imagery. Obsessions are distinguished from excessive worries about everyday occurrences because they are not concerned with real-life problems.<sup>13</sup>

**Obtundation:** A moderate reduction in the client's level of awareness so that

*mental health nursing: Scope and standards of practice* (2nd ed.)  
Nursebooks.org.

12. Martin, D. C. (1990). The mental status examination. In Walker, H. K., Hall, W. D., Hurst, J. W., (Eds.), *Clinical methods: The history, physical, and laboratory examinations*. (3rd ed.). Butterworths. <https://www.ncbi.nlm.nih.gov/books/NBK320/>

13. American Psychological Association. (n.d.). *APA Dictionary of Psychology*. <https://dictionary.apa.org/>



mild to moderate stimuli do not awaken the client. When arousal does occur, the client is slow to respond.

**Outcome:** A measurable behavior demonstrated by the client who is responsive to nursing interventions.

**Paranoia:** A condition characterized by delusions of persecution.<sup>14</sup> Clients often experience extreme suspiciousness, mistrust, or expression of fear. For example, a resident of a long-term care facility may have delusions that the staff is trying to poison him.

**Poverty of content:** A conversation in which the client talks without stating anything related to the question, or their speech in general is vague and meaningless.

**Prioritization:** The process of identifying the most significant problems and the most important interventions to implement based on a client's current status.

**Psychiatric-mental health nursing:** The nursing practice specialty committed to promoting mental health through the assessment, diagnosis, and treatment of behavioral problems, mental disorders, and comorbid conditions across the life span. Psychiatric-mental health nursing intervention is an art and a science, employing a purposeful use of self and a wide range of nursing, psychosocial, and neurobiological evidence to produce effective outcomes.<sup>15</sup>

14. American Psychological Association. (n.d.). *APA Dictionary of Psychology*.  
<https://dictionary.apa.org/>

15. American Nurses Association, American Psychiatric Nurses Association, and International Society of Psychiatric-Mental Health Nurses. (2014). *Psychiatric-mental health nursing: Scope and standards of practice* (2nd ed.).  
Nursebooks.org

**Psychomotor agitation:** A condition of purposeless, non-goal-directed activity.

**Psychomotor retardation:** A condition of extremely slow physical movements, slumped posture, or slow speech patterns.

**Psychosocial assessment:** A component of the nursing assessment process that obtains additional subjective data to detect risks and identify treatment opportunities and resources.

**Psychotherapy interventions:** Generally accepted and evidence-based methods of brief or long-term therapy, including individual therapy, group therapy, marital or couple therapy, and family therapy. These interventions use a range of therapy models including, but not limited to, psychodynamic, cognitive, behavioral, and supportive interpersonal therapies to promote insight, produce behavioral change, maintain function, and promote recovery.<sup>16</sup>

**Racing thoughts:** Fast-moving and often repetitive thought patterns that can be overwhelming. They may focus on a single topic, or they may represent multiple different lines of thought. For example, a client may have racing thoughts about a financial issue or an embarrassing moment.

**Resilience:** The ability to overcome serious hardship or traumatic experiences.

**Rumination:** Obsessional thinking involving excessive, repetitive thoughts that interfere with other forms of mental activity.<sup>17</sup>

16. American Nurses Association, American Psychiatric Nurses Association, and International Society of Psychiatric-Mental Health Nurses. (2014). *Psychiatric-mental health nursing: Scope and standards of practice* (2nd ed.). Nursebooks.org

17. American Psychological Association. (n.d.). *APA Dictionary of Psychology*. <https://dictionary.apa.org/>

**Safety plan:** A prioritized written list of coping strategies and sources of support that clients can use before or during a suicidal crisis. The plan should be brief, in the client's own words, and easy to read. After the plan is developed, the nurse should problem solve with the client to identify barriers or obstacles to using the plan. It should be discussed where the client will keep the safety plan and how it will be located during a crisis.

**SMART outcomes:** Outcome statements should contain five components easily remembered using the "SMART" mnemonic: Specific, Measurable, Attainable/Action-oriented, Relevant/Realistic, with a Time frame.

**Spirituality:** A sense of connection to something larger than oneself that typically involves a search for meaning and purpose in life.

**Stupor:** A state of unresponsiveness unless a vigorous stimulus is applied, such as a sternal rub. The client quickly drifts back into a deep sleep-like state on cessation of the stimulation.

**Suicide attempt:** An action in which there is intent to end one's life but the individual does not die as a result of their actions.

**Suicidal ideation:** When an individual has been thinking about suicide but does not necessarily have an intention to act on that idea.

**Suicide plan:** An individual who has a plan for suicide, has the means to injure oneself, and has the intent to die.

**Therapeutic milieu:** A safe, welcoming, supportive, and functional physical treatment environment.

**Transference:** When the client projects (i.e., transfers) their feelings to the nurse. For example, a client is feeling angry at a family member related to a previous disagreement and displaces the anger to the nurse during the interview.



PART V

CHAPTER 5 LEGAL AND ETHICAL CONSIDERATIONS IN MENTAL  
HEALTH CARE



## 5.1 Introduction

### Learning Objectives

- Identify safety/protective interventions for the client and others
- Identify safety/protective concerns for the nurse
- Explain the nurse role as a collaborative advocate for the health needs of the community
- Support diversity across the lifespan in client-centered care
- Explore the nurse's role and legal and ethical responsibilities when providing care to clients with mental health disorders

Mental health nursing requires a deep understanding of the legal and ethical principles that guide safe, compassionate, and equitable care. This chapter explores the foundational concepts that shape ethical decision-making based on the American Nurses Association publication, *Code of Ethics for Nurses*.<sup>1</sup> It also reviews legal considerations specific to psychiatric settings, such as involuntary commitment, competency, mandatory reporting, duty to warn, and the use of seclusion and restraints during mental health treatment. By understanding these guidelines, nurses are prepared to advocate for their clients, uphold professional standards, and navigate the delicate balance between respecting individual freedoms and ensuring safety.

1. American Nurses Association. (2025). Code of ethics for nurses. American Nurses Association. <https://codeofethics.ana.org/provisions>

## 5.2 Ethical Principles

The American Nurses Association (ANA) designates “*Ethics*” as the first Standard of Professional Performance in the publication *Nursing: Scope and Standards of Practice*, stating, “The registered nurse integrates ethics in all aspects of practice.”<sup>1</sup> See the following box for nursing actions associated with the *Ethics* Standard of Professional Performance.

### **Ethics Standard of Professional Performance<sup>2</sup>**

The registered nurse:

- Uses the *Code of Ethics for Nurses* as a moral foundation to guide nursing practice and decision-making.
- Demonstrates that every person is worthy of nursing care through the provision of respectful, person-centered, compassionate care, regardless of personal history or characteristics (Beneficence).
- Advocates for health care consumer perspectives, preferences, and rights to informed decision-making and self-determination (Respect for autonomy).
- Demonstrates a primary commitment to the recipients of nursing and health care services in all settings and situations (Fidelity).
- Maintains therapeutic relationships and professional

1. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

2. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.



boundaries.

- Safeguards sensitive information within ethical, legal, and regulatory parameters (Nonmaleficence).
- Identifies ethics resources within the practice setting to assist and collaborate in addressing ethical issues.
- Integrates principles of social justice in all aspects of nursing practice (Justice).
- Refines ethical competence through continued professional education and personal self-development activities.
- Depicts one's professional nursing identity through demonstrated values and ethics, knowledge, leadership, and professional comportment.
- Engages in self-care and self-reflection practices to support and preserve personal health, well-being, and integrity.
- Contributes to the establishment and maintenance of an ethical environment that is conducive to safe, quality health care.
- Collaborates with other health professionals and the public to protect human rights, promote health diplomacy, enhance cultural sensitivity and congruence, and reduce health disparities.
- Represents the nursing perspective in clinic, institutional, community, or professional association ethics discussions.

## American Nurse Association Code of Ethics

The ANA *Code of Ethics* establishes the ethical standard for the nursing profession. It contains ten provisions that provide a guide for nurses to use in ethical practice and decision-making in all practice settings. The *Code of Ethics for Nurses* states that it stands as both a normative framework and an

aspirational guide, with the 2025 edition containing updated provisions based on the current health care environment and a new tenth provision that envisions the role of nursing in creating a healthier society and a healthier world. Read a summary of the ten provisions in the following box.<sup>3</sup>

### **Ten Provisions of the ANA's Code of Ethics for Nurses**<sup>4</sup>

**Provision 1:** The nurse practices with compassion and respect for the inherent dignity, worth, and unique attributes of every person.

**Provision 2:** The nurse's primary commitment is to the recipient(s) of nursing care, whether an individual, family, group, community, or population.

**Provision 3:** The nurse establishes a trusting relationship and advocates for the rights, health, and safety of the recipient(s) of nursing care.

**Provision 4:** Nurses have authority over nursing practice and are responsible and accountable for their practice consistent with their obligations to promote health, prevent illness, and to provide optimal care.

**Provision 5:** The nurse has moral duties to self as person of inherent dignity and worth including an expectation of a safe place to work that fosters flourishing, authenticity of self at work, and self-respect through integrity and professional competency.

3. American Nurses Association. (2025). *Code of ethics for nurses*. American Nurses Association. <https://codeofethics.ana.org/provisions>

4. American Nurses Association. (2025). *Code of ethics for nurses*. American Nurses Association. <https://codeofethics.ana.org/provisions>

**Provision 6:** Nurses through individual and collective effort, establish, maintain, and improve the ethical environment of the work setting that affects nursing care and the well-being of nurses.

**Provision 7:** Nurses advance the profession through multiple approaches to knowledge development, professional standards, and the generation of policies for nursing, health, and social concerns.

**Provision 8:** Nurses build collaborative relationships and networks with nurses, other healthcare and non-health care disciplines, and the public to achieve greater ends.

**Provision 9:** Nurses and their professional organizations work to enact and resource practices, policies, and legislation to promote social justice, eliminate health inequities, and facilitate human flourishing.

**Provision 10:** Nursing, through organizations and associations, participates in the global nursing and health community to promote human and environmental health, well-being, and flourishing.

► Read the full descriptions of the ten provisions in the online 2025 ANA [Code of Ethics for Nurses](#).

## Ethical Principles

Ethical principles are used to define right from wrong actions. They are used to help define nurses' moral duties and aid in ethical analysis and decision

making. Although there are many ethical principles that guide nursing practice, foundational ethical principles include respect for autonomy and self-determination, beneficence (do good), nonmaleficence (do no harm), justice (fairness), fidelity (keep promises), and veracity (tell the truth).

## Autonomy and Self Determination

The ANA *Ethics Standard of Professional Performance* states that nurses respect a client's **autonomy** by advocating for their perspectives, preferences, and rights to informed decision-making and self determination.<sup>5</sup>

Provision 1, Section 1.4 of the ANA *Code of Ethics for Nurses* states, "Respect for human dignity requires the recognition of specific patient rights, in particular, the right to self-determination. Recipients of care have the moral and legal right to determine what will be done with and to their own person; to be given accurate, complete, and understandable information in a manner that facilitates an informed decision; and to be assisted with weighing the benefits, burdens, and available options in their treatment, including the choice of no treatment. They also have the right to accept, refuse, or terminate treatment without undue influence, duress, deception, manipulation, coercion, or prejudice, and to be given necessary support throughout the decision-making and treatment process. Such support includes the opportunity to make decisions with family and persons of their choosing, and to partner with nurses and other healthcare professionals."<sup>6</sup> See examples of nurses implementing interventions based on the ethical principles of autonomy and self-determination in the following box.

5. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

6. American Nurses Association. (2025). *Code of ethics for nurses*. American Nurses Association. <https://codeofethics.ana.org/provisions>

## Case Applications of Autonomy and Self Determination

**Medical Example:** Sarah, an experienced registered nurse, is caring for Mr. Thompson, a 68-year-old man recently diagnosed with early-stage lung cancer. His oncologist has recommended surgery followed by chemotherapy, explaining that this approach offers the best chance of remission. However, after discussing the options with Sarah and his family, Mr. Thompson decides to decline aggressive treatment in favor of palliative care, prioritizing his quality of life over prolonged treatment. Although Sarah personally believes that pursuing treatment might extend his life, she respects Mr. Thompson's decision. She ensures that he fully understands the risks and benefits of his choice, answers his questions, and advocates for his wishes with the healthcare team. She arranges for a palliative care consultation and works to provide comfort-focused care, demonstrating her commitment to the ethical principle of autonomy.

**Mental Health Care Example:** A nurse is providing care for a client diagnosed with severe, treatment-resistant depression whose health care provider has recommended electroconvulsive therapy (ECT). The client tells the nurse they feel unsure about proceeding with ECT. The nurse provides health teaching about the ECT procedure and the potential side effects in a clear, compassionate manner and provides time for the client to ask questions and voice concerns. This action upholds the client's rights to self determination and autonomy and by respecting their right to make informed choices about their mental health treatment.

# Beneficence and Respect for Human Dignity

The ethical principle of **beneficence** is described in the ANA *Ethics Standard of Professional Performance* as the nurse “demonstrates that every person is worthy of nursing care through the provision of respectful, person-centered, compassionate care, regardless of personal history or characteristics.”<sup>7</sup>

Provision 1, Section 1.1 of the ANA *Code of Ethics for Nurses* states, “A fundamental principle that underlies all nursing practice is respect for the inherent dignity, worth, unique attributes, and human rights of all individuals; therefore, ethical nursing practice requires compassion for all humans as deserving of dignity and respect. Nurses maintain caring relationships and are committed to fair treatment, transparency, integrity-preserving compromise, building trust, and the best resolution of conflicts. The nurse is additionally committed to creating and sustaining an ethical environment where the nurse-patient relationship can flourish.”<sup>8</sup>

Section 1.3 of Provision 1 of the ANA *Code of Ethics for Nurses* further described beneficence as, “Optimal nursing care enables recipients to live with as much physical, emotional, social, religious, and/or spiritual well-being as possible, aligning with their preferences, values, and determination of quality of life. Nurses lead the implementation of responsible and appropriate evidence-based interventions across the lifespan to optimize the health and well-being of those in their care. When a recipient of care no longer sees a proportional benefit from the burdens of interventions, nurses are attentive and practice shared decision-making to arrive at medically achievable goals that reflect patient values. All human beings should have access to what they recognize as a good quality of life, which is subjective. Nurses appreciate that

7. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

8. American Nurses Association. (2025). *Code of ethics for nurses*. American Nurses Association. <https://codeofethics.ana.org/provisions>

what is right for one person may not be right for another. The nurse balances respect for values with harm mitigation and recognizes that every decision for each person is unique and situational.”<sup>9</sup>

When caring for clients with mental health disorders, nurses implement beneficence by actively advocating for evidence-based treatments that promote the best possible outcomes. For instance, if a client with schizophrenia is not responding well to their current medication regimen and is experiencing distressing side effects, the nurse collaborate with the healthcare team to recommend a more effective and better-tolerated alternative. By staying informed about current research and clinical guidelines, the nurse ensures the client receives the most appropriate care, demonstrating a commitment to the client’s well-being and overall mental health recovery. See an example of nurse implementing interventions based on the ethical principles of beneficence respect for human dignity in the following box.

### **Beneficence Case Application**

A nurse is caring for Jake, a 25-year-old client admitted for severe depression and suicidal ideation. After assessing Jake’s condition, the nurse notices that he is hesitant to engage in psychotherapy and has refused to take his prescribed antidepressant medication due to side effects he experienced in the past. The nurse provides health teaching to Jake about the prescribed antidepressants, psychotherapy, and lifestyle modifications to help treat depression, and also contacts his psychiatrist to share Jake’s concerns and sets up an appointment to discuss adjusting

9. American Nurses Association. (2025). *Code of ethics for nurses*. American Nurses Association. <https://codeofethics.ana.org/provisions>

his treatment plan to better suit his preferences. The nurse also continues to encourage Jake to participate in his scheduled cognitive behavioral therapy sessions, recognizing its proven effectiveness in treating depression. By proactively seeking the best care for Jake and ensuring he receives appropriate, individualized treatment, the nurse exemplifies the ethical principle of beneficence.

## Nonmaleficence

The ethical principle of **nonmaleficence** is commonly defined as a duty to do no harm and balancing avoidable harm with benefits of good achieved. Non-maleficence is further described in the ANA *Ethics Standard of Professional Performance* as the nurse “safeguards sensitive information within ethical, legal, and regulatory parameters.”<sup>10</sup>

Provision 1, Section 1.3 of the ANA *Code of Ethics for Nurses* further describes nonmaleficence related to health as, “Nurses promote health and wellness, address problems, and respect patient decisions. Respect for a patient’s decisions does not require that the nurse agrees with or supports all choices made by a recipient of care. When patient choices are assessed to be dangerous, risky, or self-destructive, nurses have a moral obligation to take appropriate actions to address the behavior, and provide accurate, evidence-based education and resources. In immediately dangerous situations, the

10. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.



nurse focuses on modifying the harmful behavior to either mitigate or eliminate the risk.”<sup>11</sup>

A classic example of doing no harm in nursing practice is reflected by nurses checking medication rights three times before administering medications. In this manner, medication errors can be avoided, and the duty to do no harm is met. Furthermore, nurses continually assess for side effects of medications and promptly report them to the health care provider. See an additional example of nursing implementing interventions based on the nonmaleficence ethical principle in the following box.

### **Nonmaleficence Case Application**

A nurse is caring for Lisa, a client diagnosed with bipolar disorder who has been taking lithium for mood stabilization. During the routine nursing assessment, the nurse notices that Lisa is experiencing potential signs of lithium toxicity, including symptoms of nausea, hand tremors, and mild confusion. Recognizing these cues, the nurse immediately reviews Lisa's most recent lab results and finds that her lithium level is approaching a toxic range. Acting quickly, the nurse notifies the prescribing provider and advocates for dosage adjustment to prevent further harm. The nurse educates Lisa about the importance of staying hydrated and recognizing and reporting early signs of lithium toxicity. By proactively identifying and addressing a potential medication-related harm, the nurse demonstrates the ethical principle of nonmaleficence, ensuring that Lisa's treatment remains both safe and effective.

11. American Nurses Association. (2025). *Code of ethics for nurses*. American Nurses Association. <https://codeofethics.ana.org/provisions>

# Justice

**Justice** is described in the ANA *Ethics* Standard of Professional Performance as the nurse “integrates principles of social justice in all aspects of nursing practice.”<sup>12</sup>

Provision 1, Section 1.2 of the ANA Code of Ethics for Nurses further describes justice as, “Nurses establish relationships of trust and provide nursing services according to need. Nurses engage in self-reflection to identify and mitigate bias or prejudice that interferes with or harms the nurse-patient relationship. The nurse recognizes that biases can exist both explicitly and unconsciously. Attributes such as the patient’s culture, value systems, religious and/or spiritual beliefs, lifestyle, social support system, preferred language, and sexual identity are to be considered when planning individual, family, and population-centered care. Nurses promote health and wellness, address problems, and respect patient decisions. Respect for a patient’s decisions does not require that the nurse agrees with or supports all choices made by a recipient of care.”<sup>13</sup>

Nurses have a social contract to “provide compassionate care that addresses the individual’s needs for protection, advocacy, empowerment, optimization of health, prevention of illness and injury, alleviation of suffering, comfort, and well-being.”<sup>14</sup> An example of a nurse upholding the principle of justice in all settings is ensuring that quality care is provided to all clients, even for those who do not have the cognitive ability to communicate their needs. See an

12. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

13. American Nurses Association. (2025). *Code of ethics for nurses*. American Nurses Association. <https://codeofethics.ana.org/provisions>

14. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

additional example of nursing implementing interventions based on the justice ethical principle in the following box.

### **Justice Case Application**

A nurse working in a busy emergency department is caring for two clients who arrived at the same time. One client is a well-dressed business professional with private insurance who is experiencing a panic attack, and the other is an unhoused individual with no health insurance who is experiencing a mental health crisis. Despite their differing backgrounds, the nurse treats both clients with the same level of care, professionalism, and respect.

The business professional can clearly articulate his concerns and symptoms. The unhoused client is disoriented and unable to clearly communicate his needs. The nurse takes time to thoroughly assess both clients, advocating for psychiatric evaluations and appropriate mental health care. For the unhoused individual, the nurse also implements interventions based on the client's physiological needs by making a referral to social work support services to advocate for food and housing resources. The nurse provides individualized health teaching to both clients and ensures they have follow-up care in place before discharge.

By providing equitable and compassionate care to both clients—regardless of their socioeconomic status—the nurse upholds the ethical principle of justice, ensuring that healthcare is delivered fairly and without bias.

# Fidelity

**Fidelity** is described in the ANA *Ethics* Standard of Professional Performance as the nurse “demonstrates a primary commitment to the recipients of nursing and healthcare services in all settings and services.”<sup>15</sup>

Provision 2, Section 2.1 of the ANA Code of Ethics for Nurses further describes fidelity as, “Within the context of nursing practice, the nurse prioritizes recipients of nursing care, placing them over institutions. Every clinical encounter and plan of care reflects the fundamental commitment of nursing to the inherent dignity, worth, unique attributes, and human rights of the patient. Nurses provide patients with opportunities to participate in ...planning and implementing their plan of care, and deciding what supportive services are acceptable to them.”<sup>16</sup>

Fidelity also means nurses must remain up-to-date with evidence-based practice in order to plan and implement effective nursing interventions. For example, a nurse is upholding the ethical standard of fidelity by staying informed about current cognitive-behavioral therapy (CBT) techniques and appropriately integrating them into nursing care plans for clients with mental health disorders.

## Fidelity Case Application

A nurse on a medical-surgical unit is caring for Mr. Thompson, a 70 year-old postoperative client recovering from orthopedic

15. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

16. American Nurses Association. (2025). *Code of ethics for nurses*. American Nurses Association. <https://codeofethics.ana.org/provisions>

surgery. During the assessment, the nurse notices non-verbal cues of pain, including wincing and guarding of the affected area, even though Mr. Thompson reports his pain level as a “1.” The nurse states this observation and incongruity in verbal and nonverbal pain indicators. Mr. Thompson confides in the nurse that he is in more pain than he has been reporting to the health care provider and nurses, but asks the nurse to not document it because he’s afraid it might delay his planned discharge to go home today.

The nurse listens carefully and empathetically to Mr. Thompson’s concerns. The nurse asks Mr. Thompson about his desire for discharge today and discovers he is mostly concerned about his cat, who is alone at home and needs to be fed. The nurse explains the importance of appropriate pain management for his recovery process and offers to contact the health care provider regarding the pain management plan offers. The nurse also offers to assist Mr. Thompson to use the phone in the room to call a trusted neighbor to feed the cat. The nurse documents the client’s verbal and nonverbal indicators of pain as well as his preferences for discharge today in the electronic medical record.

The nurse upholds the ethical principle of fidelity by maintaining honest, empathetic communication, honoring professional responsibilities, and maintaining trust in the nurse-client relationship.

## Veracity

**Veracity** means telling the truth. An example of veracity in health care is informed consent. Nurses ensure that clients have a good understanding of the benefits and risks of a prescribed procedure or psychotropic medication. By providing honest, clear, and complete information, nurses support clients

in making informed decisions about their mental health treatment, fostering trust and ethical care.

### **Veracity Case Application**

A nurse is providing preoperative care while preparing Mr. Jackson for an elective colonoscopy. The nurse reviews the client's chart and notices that he has not yet signed the informed consent form. When asking Mr. Jackson if he has any questions about the procedure or post-operative care, he admits he doesn't understand the risks and benefits of the procedure. The nurse waits to administer preoperative sedative medications and notifies the health care provider about Mr. Jackson's concerns. After the health care provider visits the client, answers his questions, and addresses his concerns, Mr. Jackson signs the informed consent form. The nurse witnesses the signature then administers the prescribed sedative medication. By ensuring that the client receives complete information before signing the informed consent form and ensuring the form is signed before preoperative medications are administered, the nurse upholds the ethical principle of veracity.

## **Role of Caring**

Nurses use a client-centered, care-based, ethical approach to nursing care that focuses on the specific circumstances of each situation. This approach aligns with the foundational nursing concepts of holism and caring in a nurse-client relationship rooted in dignity, respect, kindness, and

compassion.<sup>17</sup> It is grounded in dignity, respect, kindness, and compassion, ensuring that care is both ethically sound and deeply humanistic.

### **Role of Caring Case Application**

Elena, a nurse on an oncology unit, is caring for Mrs. Rivera, a client undergoing chemotherapy who is experiencing significant fatigue, hair loss, and emotional distress. During morning rounds, Elena notices that Mrs. Rivera is unusually quiet and tearful.

Instead of simply focusing on clinical tasks, Elena pulls up a chair, holds Mrs. Rivera's hand, and asks how she's truly feeling. Mrs. Rivera opens up about her fears, body image concerns, and feelings of isolation. Elena listens with empathy, validates her emotions, and offers support by involving the hospital counselor and arranging a visit from a volunteer who also underwent cancer treatment.

Later that day, Elena brings in a soft headscarf gifted by a local support group and helps Mrs. Rivera feel more comfortable and cared for. She checks in regularly—not just to assess physical symptoms, but to ensure emotional and spiritual needs are met.

By going beyond physical care to foster connection, compassion, and comfort, Elena exemplifies the role of caring in nursing, nurturing a healing environment that supports the whole person, not just the illness.

17. Legal Information Institute. (n.d.). *Welcome to LII*. Cornell Law School. <https://www.law.cornell.edu>

## 5.3 Standards of Care

Standards of care in nursing are guidelines that provide a foundation as to how a nurse should act and what they should and should not do in their professional capacity. These guidelines establish a baseline of quality client care and provide an objective standard of accountability within the profession. Standards of care are enforced by courts of law and state Boards of Nursing, who evaluate a nurse's practice against these standards. If a nurse's actions (or lack of actions) do not meet the accepted standard of care, their conduct may be found to be negligent.<sup>1</sup>

Standards for nursing care are set by several organizations, including the American Nurses Association (ANA), states' Nurse Practice Acts, agency policies and procedures, federal regulators, and professional nursing organizations.

### American Nursing Association's Scope and Standards of Practice

In the United States, the American Nurses Association (ANA) publishes two resources that set standards and guide professional nursing practice: *Code of Ethics for Nurses* and *Nursing: Scope and Standards of Practice*.

The *Code of Ethics for Nurses* establishes an ethical framework for nursing practice across all roles, levels, and settings.<sup>2</sup> It is discussed in greater detail in the "[Ethical Principles](#)" section of this chapter.

1. Law Office of Nicole Irmer. (2019). *Nursing standards of care issues*. <https://www.californialicensingdefense.com/practice-areas/professionals/nursing-license/standards-of-care-issues/#:~:text=Standards%20of%20care%20in%20nursing,that%20all%20nurses%20must%20follow>.
2. American Nurses Association. (2025). *Code of ethics for nurses*. American Nurses Association. <https://codeofethics.ana.org/provisions>



*Nursing: Scope and Standards of Practice* describes a professional nurse's scope of practice and defines the who, what, where, when, why, and how of nursing. This resource includes Standards of Professional Nursing Practice and Standards of Professional Performance. The **Standards of Professional Nursing Practice** are "authoritative statements of the actions and behaviors that all registered nurses, regardless of role, population, specialty, and setting, are expected to perform competently." These standards define a competent level of nursing practice based on the critical thinking model known as the nursing process and include the components of assessment, diagnosis, outcomes identification, planning, implementation, and evaluation.<sup>3</sup> Each of these standards is further discussed in the "[Application of the Nursing Process in Mental Health Care](#)" chapter of this book. The **Standards of Professional Performance** are 12 additional standards that describe a nurse's professional behavior, including activities related to ethics, advocacy, respectful and equitable practice, communication, collaboration, leadership, education, scholarly inquiry, quality of practice, professional practice evaluation, resource stewardship, and environmental health. All registered nurses are expected to engage in these professional role activities based on their level of education, position, and role. Registered nurses are accountable for their professional behaviors to themselves, health care consumers, peers, and ultimately to society.<sup>4</sup>

The ANA Standards of Professional Performance are as follows<sup>5</sup>:

- **Ethics.** The registered nurse integrates ethics in all aspects of practice.

3. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association

4. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association

5. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association

- **Advocacy.** The registered nurse demonstrates advocacy in all roles and settings.
- **Respectful and Equitable Practice.** The registered nurse practices with cultural humility and inclusiveness.
- **Communication.** The registered nurse communicates effectively in all areas of professional practice.
- **Collaboration.** The registered nurse collaborates with the health care consumer and other key stakeholders.
- **Leadership.** The registered nurse leads within the profession and practice setting.
- **Education.** The registered nurse seeks knowledge and competence that reflects current nursing practice and promotes futuristic thinking.
- **Scholarly Inquiry.** The registered nurse integrates scholarship, evidence, and research findings into practice.
- **Quality of Practice.** The registered nurse contributes to quality nursing practice.
- **Professional Practice Evaluation.** The registered nurse evaluates one's own and others' nursing practice.
- **Resource Stewardship.** The registered nurse utilizes appropriate resources to plan, provide, and sustain evidence-based nursing services that are safe, effective, financially responsible, and judiciously used.
- **Environmental Health.** The registered nurse practices in a manner that advances environmental safety and health.

## American Psychiatric Nurses Association Standards of Practice

In addition to the ANA Standards of Professional Nursing Practice, the American Psychiatric Nurses Association establishes standards of practice for psychiatric-mental health nurse specialists in *Psychiatric-Mental Health Nursing: Scope and Standards of Practice*.<sup>6</sup> These standards are built on the

6. American Nurses Association, American Psychiatric Nurses Association, and

ANA's Standards of Professional Nursing Practice, with additional activities included under the *Intervention* standard of care. These interventions are further discussed in the "[Application of the Nursing Process in Mental Health Care](#)" chapter.

► Read more about the [American Psychiatric Nurses Association](#).

## Nurse Practice Act

In addition to the professional standards of practice, nurses must legally follow regulations set by the Nurse Practice Act and enforced by the Board of Nursing in the state where they are employed. The Board of Nursing is the state-specific licensing and regulatory body that sets standards for safe nursing care and issues nursing licenses to qualified candidates, based on the Nurse Practice Act enacted by that state's legislature. The Nurse Practice Act establishes regulations for nursing practice within that state and defines the scope of nursing practice. If nurses do not follow the standards and scope of practice set forth by the Nurse Practice Act, they can have their nursing license revoked by the Board of Nursing.

Nursing students are legally accountable for the quality of care they provide to clients just as nurses are accountable. Students are expected to recognize the limits of their knowledge and experience and appropriately alert individuals in authority regarding situations that are beyond their competency. A violation of the standards of practice constitutes unprofessional conduct and can result in the Board of Nursing denying a license to a nursing graduate.

International Society of Psychiatric-Mental Health Nurses. (2014). *Psychiatric-mental health nursing: Scope and standards of practice* (2nd ed.). Nursebooks.org

## Employer Policies, Procedures, and Protocols

In addition to following professional nursing standards and the state Nurse Practice Act, nurses and nursing students must also practice according to agency policies, procedures, and protocols. For example, each agency has specific policies regarding the use of restraints. If a nurse did not follow this policy and a client was injured or died, the nurse could be held liable in a court of law.

Agencies also have their own sets of procedures and protocols. For example, each agency has specific procedural steps for performing nursing skills, such as inserting urinary catheters. Agencies also have protocols that are precisely written plans for a regimen of therapy. For example, agencies typically have a hypoglycemia protocol that nurses automatically implement when a client's blood sugar falls below a specific number and includes actions such as providing orange juice and rechecking the blood sugar. These agency-specific policies, procedures, and protocols supersede the information taught in nursing school, and nurses and nursing students can be held legally liable if they don't follow them. Therefore, it is vital for nurses and nursing students to review and follow current agency-specific procedures, policies, and protocols when providing client care.

## Federal Regulations

In addition to professional standards, state Nurse Practice Acts, and employer policies, procedures, and protocols, nursing practice is also influenced by federal regulations enacted by government agencies such as The Joint Commission and the Centers for Medicare and Medicaid.

The Joint Commission (TJC) is a national organization that accredits and certifies over 20,000 health care organizations in the United States. The mission of TJC is to continuously improve health care by setting standards for providing safe, high-quality health care. The Centers for Medicare & Medicaid Services (CMS) enforces quality standards in health care organizations that receive Medicare and Medicaid funding. Nurses must follow standards set by these agencies and implemented by their employers. For example, the

expectation that a nurse perform medication rights three times before administering medication to a client is based on a federal regulation.

Read more information at the ► [The Joint Commission](#) website and the ► [Centers for Medicare & Medicaid Services](#) website

## 5.4 Laws, Torts, Malpractice, and Disciplinary Actions

In addition to following standards of care, nurses must also follow related federal and state laws. **Criminal law** is a system of laws that punishes individuals who commit crimes. Crimes are classified as felonies, misdemeanors, and infractions. Conviction for a crime requires evidence to show the defendant is guilty beyond a shadow of doubt. This means the prosecution must convince a jury there is no reasonable explanation other than guilty that can come from the evidence presented at trial. See Figure 5.1<sup>1</sup> for an illustration of a criminal case being tried in front of a jury. **Civil law** focuses on the rights, responsibilities, and legal relationships between private citizens, and involves compensation to the injured party. A person bringing the lawsuit is called the **plaintiff**, and the parties named in the lawsuit are called **defendants**.<sup>2</sup>



Figure 5.1 Criminal Trial by Jury

1. “[Courtroom Trial with Judge, Jury – Vector Image](#)” designed by [WannaPik](#) is licensed under [CCO](#)
2. Brous, E. (2019). The elements of a nursing malpractice case, part 1: Duty. *American Journal of Nursing*, 119(7), 64–67. <https://doi.org/10.1097/01.NAJ.0000569476.17357.f5>

Civil law includes torts. A **tort** is an act of commission or omission that gives rise to injury or harm to another and amounts to a civil wrong for which courts impose liability. Tort law exists to compensate clients injured by negligent practice, provide corrective judgment, and deter negligence with consequences of action or inaction.<sup>3</sup>

Two categories of torts affecting nursing practice are intentional torts and unintentional torts. **Intentional torts** are wrongs that the defendant knew (or should have known) would be caused by their actions. Examples of intentional torts include assault, battery, false imprisonment, slander, libel, and breach of privacy or client confidentiality. **Unintentional torts** occur when the defendant's actions or inactions were unreasonably unsafe. Unintentional torts can result from acts of commission (i.e., doing something a reasonable nurse would not have done) or omission (i.e., failing to do something a reasonable nurse would do). Examples of torts affecting nursing practice are discussed in further detail in the following subsections.<sup>4</sup>

## Assault and Battery

Assault and battery are intentional torts. **Assault** is defined as intentionally putting another person in reasonable apprehension of an imminent harmful or offensive contact. For example, assault would occur by threatening to forcibly administering a medication. **Battery** is defined as intentional causation of harmful or offensive contact with another person without that person's consent. For example, battery would occur with the administration of a medication forcibly. Physical harm does not need to occur in order to be

3. Wis. JI—Civil 1005. (2016). <https://wilawlibrary.gov/jury/civil/instruction.php?n=1005>

4. Brous, E. (2019). The elements of a nursing malpractice case, part 1: Duty. *American Journal of Nursing*, 119(7), 64–67. <https://doi.org/10.1097/01.NAJ.0000569476.17357.f5>

charged with assault or battery. Battery convictions are typically misdemeanors but can be felonies if serious bodily harm occurs.<sup>5</sup>

An example related to assault and battery in health care is the client's right to refuse treatment. For example, a hospitalized client can refuse to take prescribed medication. If a nurse forcibly administers medication without a client's consent, it could be ruled assault or battery in a court of law. However, forcible administration of a medication based on a provider's order may be justified in an emergency situation to prevent imminent harm to oneself or others.<sup>6</sup>

## False Imprisonment

False imprisonment is an intentional tort. **False imprisonment** is defined as an act of restraining another person and causing that person to be confined in a bounded area. An example of possible false imprisonment in health care is the use of restraints. See Figure 5.2<sup>7</sup> for an image of a simulated client in full physical medical restraints. Restraints can be physical, chemical, or verbal. Nurses must vigilantly follow agency policies related to the use of physical and chemical restraints and monitor clients who are restrained. Chemical restraints include administration of PRN medications such as benzodiazepines and require clear documentation supporting their use. Verbal restraints consist of using speech to restrict the movement or actions of a client. Verbal threats to keep an individual in an inpatient environment is an example of a verbal restraint. This can also qualify as false imprisonment

5. Brous, E. (2019). The elements of a nursing malpractice case, part 1: Duty. *American Journal of Nursing*, 119(7), 64–67. <https://doi.org/10.1097/01.NAJ.0000569476.17357.f5>

6. Fry, S. T. (1989). The role of caring in a theory of nursing ethics. *Hypatia*, 4(2), 87-103. <https://doi.org/10.1111/j.1527-2001.1989.tb00575.x>

7. "PinelRestaint.jpg" by James Heilman, MD is licensed under [CC BY-SA 4.0](https://creativecommons.org/licenses/by-sa/4.0/)



and should be avoided. Additional information regarding the use of restraints is discussed in the “Patient Rights” section.




Figure 5.2 Full Physical Medical Restraints

## Privacy and Confidentiality

Breaching privacy and confidentiality are intentional torts. **Confidentiality** is the right of an individual to have personal, identifiable medical information, referred to as protected health information, kept private. **Protected Health Information (PHI)** is defined as individually identifiable health information, including demographic data, that relates to the individual’s past, present, or future physical or mental health or condition; the provision of health care to the individual; and the past, present, or future payment for the provision of health care to the individual.<sup>8</sup>

Confidentiality is a right protected by the **Health Insurance Portability and Accountability Act (HIPAA)**. HIPAA was enacted in 1996 and was prompted by the need to ensure privacy and protection of personal health records in an environment of electronic medical records and third-party insurance payers. There are two main sections of HIPAA law: the Privacy Rule and the Security

8. Wis. JI—Civil 1005. (2016). <https://wilawlibrary.gov/jury/civil/instruction.php?n=1005>

Rule. The Privacy Rule addresses the use and disclosure of individuals' health information. The Security Rule sets national standards for protecting the confidentiality, integrity, and availability of electronically protected health information. HIPAA regulations extend beyond medical records and apply to client information shared with others. Therefore, all types of client information and data should be shared only with health care team members who are actively providing care to them. For example, when applying HIPAA to mental health inpatient settings, nurses may not answer in the affirmative if someone calls and asks if an individual has been admitted to the unit.<sup>9</sup> See Figure 5.3<sup>10</sup> for a depiction of confidentiality.



Figure 5.3 Confidentiality

There are specific circumstances in which HIPAA does not apply. For example, nurses have a duty to warn and protect, are mandated reporters of suspected abuse or neglect, and are required to share specific information reported by minors with authorities or their parents.

## Duty to Warn and Protect

Most states have laws regarding the duty to protect third parties from

9. Wis. JI—Civil 1005. (2016). <https://wilawlibrary.gov/jury/civil/instruction.php?n=1005>

10. “[Concept of Data Privacy And Policy Illustration](#)” by [Deesign Graphics](#) at [Iconscout](#) is licensed under [CC BY 4.0](#)

potential life threats. Nurses and other mental health professionals have a duty to warn and protect third parties when they may be in danger from a client. This duty falls outside of HIPAA regulations. This includes assessing and predicting the client's threat of violence towards another person or groups of people and taking action to protect the identified victims.<sup>11</sup> An example of duty to warn and protect would be a hospitalized patient disclosing a plan to harm their coworkers upon discharge. The nurse would be obligated to report this realistic threat.

## Mandatory Reporting of Suspected Abuse or Neglect

HIPAA does not apply to reporting the suspected neglect or abuse of children, adults at risk, or older adults. Many states require health professionals to report suspected neglect or abuse. State laws vary, but they generally include a definition of abuse, a list of people required to report abuse, and the government agency designated to receive and investigate the reports. Nurses and other health professionals are referred to as **mandated reporters** because they are required by state law to report suspected neglect or abuse of children, adults at risk, and the elderly. **Adults at risk** are adults who have a physical or mental condition that impairs their ability to care for their own needs.

For example, in Wisconsin, suspected neglect or abuse is reported to Child Protective Services (CPS), Adult Protective Services, or law enforcement. Nurses should be aware of the county or state agencies to whom they should report suspected abuse. See the following box for additional information.

11. Fry, S. T. (1989). The role of caring in a theory of nursing ethics. *Hypatia*, 4(2), 87-103. <https://doi.org/10.1111/j.1527-2001.1989.tb00575.x>

- ▶ Read additional information about signs of child and elder abuse in the “[Trauma, Abuse, and Violence](#)” chapter.
- ▶ Read more about protective services in your state. Here are links to Wisconsin’s [Child Protective Services](#) and [Adult Protective Services](#).
- ▶ Find resources in your area for reporting suspected child abuse at [ChildHelp National Child Abuse Hotline](#) or elder abuse at the [National Adult Protective Services Association website](#).

## Conditional Confidentiality for Minors

“Conditional confidentiality” applies to minors under the age of 18. State laws determine what information is considered confidential and what requires reporting to law enforcement or Child Protective Services, such as child abuse, gunshot or stabbing wounds, sexually transmitted infections, abortions, suicidal ideation, and homicidal ideation. Some state laws make it optional for clinicians to inform parents/guardians if their child is seeking services related to sexual health care, substance use, or mental health care. Nurses should be aware of the state laws affecting the confidentiality of child and adolescent care in the state in which they are practicing.<sup>12</sup>

12. Nurses Service Organization and CNA Financial. (2020). *Nurse professional liability exposure claim report* (4th ed.). <https://www.nso.com/Learning/Artifacts/Claim-Reports/Minimizing-Risk-Achieving-Excellence>

- View the [Wisconsin Department of Health Services' Client Rights for Minors](#).

## Slander and Libel

Slander and libel are intentional torts. **Defamation of character** occurs when an individual makes negative, malicious, and false remarks about another person to damage their reputation. **Slander** is spoken defamation and **libel** is written defamation. Nurses must take care in their oral communication and documentation to avoid defaming clients or coworkers.<sup>13</sup>

## Fraud

**Fraud** is an intentional tort that occurs when an individual is deceived for personal gain. A nurse may be charged with fraud for documenting interventions not performed or for altering documentation to cover up an error. Fraud can result in civil and criminal charges, as well as suspension or revocation of a nurse's license.<sup>14</sup>

## Negligence and Malpractice

Negligence and malpractice are unintentional torts. **Negligence** is the failure to exercise the ordinary care a reasonable person would use in similar circumstances. Wisconsin civil jury instruction states, "A person is not using ordinary care and is negligent, if the person, without intending to do harm, does something (or fails to do something) that a reasonable person would

13. Wis. JI—Civil 1005. (2016). <https://wilawlibrary.gov/jury/civil/instruction.php?n=1005>

14. Wis. JI—Civil 1005. (2016). <https://wilawlibrary.gov/jury/civil/instruction.php?n=1005>

recognize as creating an unreasonable risk of injury or damage to a person or property.” **Malpractice** is a specific term used for negligence committed by a health professional with a license.

## Elements of Malpractice

Clients bringing a malpractice lawsuit must be able to demonstrate to the court that their interests were harmed. Most malpractice lawsuits name physicians or hospitals as defendants, although nurses can be individually named. Employers can be held liable for the actions of their employees.

Malpractice lawsuits are concerned with the legal obligations nurses have to their clients to adhere to current standards of practice. These legal obligations are referred to as the duty of reasonable care. Nurses are required to adhere to standards of practice when providing care to clients they have been assigned. This includes following organizational policies and procedures, maintaining clinical competency, and confining their activities to the authorized scope of practice as defined by their state’s Nurse Practice Act. Nurses also have a legal duty to be physically, mentally, and morally fit for practice. When nurses do not meet these professional obligations, they are said to have breached their duties to clients.<sup>15</sup>

All of the following elements must be established in a court of law to prove malpractice<sup>16</sup> :

- **Duty:** A nurse-client relationship exists.
- **Breach:** The standard of care was not met and harm was a foreseeable consequence of the action or inaction.

15. Teoli, D., & Ghassemzadeh, S. (2023). Patient self-determination act. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK538297/>

16. Teoli, D., & Ghassemzadeh, S. (2023). Patient self-determination act. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK538297/>

- **Cause:** Injury was caused by the nurse's breach.
- **Harm:** Injury resulted in damages.

## Duty

In the work environment, a duty is created when the nurse accepts responsibility for a client and establishes a nurse-client relationship. This generally occurs during inpatient care upon acceptance of a handoff report from another nurse. Outside the work environment, a nurse-client relationship is created when the nurse volunteers services. Mandatory reporting and duty to warn and protect are additional examples of a nurse's duty.<sup>17</sup>

## Breach of Duty

The second element of malpractice is breach of duty. After a plaintiff has established the first element in a malpractice suit (i.e., the nurse owed a duty to the plaintiff), the plaintiff must demonstrate that the nurse breached that duty by failing to comply with the duty of reasonable care. To demonstrate that a nurse breached their duty to a client, the plaintiff must prove the nurse deviated from acceptable standards of practice. The plaintiff must establish how a reasonably prudent nurse in the same or similar circumstances would act and then show how the defendant nurse departed from that standard of practice. The plaintiff must claim the nurse did something a reasonably prudent nurse would not have done (an act of commission) or failed to do something a reasonable nurse would have done (an act of omission).<sup>18</sup>

17. Teoli, D., & Ghassemzadeh, S. (2023). Patient self-determination act. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK538297/>

18. Teoli, D., & Ghassemzadeh, S. (2023). Patient self-determination act. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK538297/>

Experts are needed during court hearings to explain things outside the knowledge of non-nurse jurors. In reaching their opinions, experts review many materials, including the state's Nurse Practice Act and organizational policies, to determine whether the nurse adhered to them. To qualify as a nurse expert, the person testifying must have relevant experience, education, skill, and knowledge. Medical malpractice trials take place primarily in state courts, so experts are deemed qualified based on state requirements.<sup>19</sup>

## Cause

The third element of malpractice is cause. After the plaintiff has established that the nurse owed a duty to a client and then breached that duty, they must then demonstrate that damages or harm were caused by that breach. Plaintiffs cannot prevail by only demonstrating the nurse departed from acceptable standards of practice, but also must prove that such departures were the cause of any injuries. Additionally, nurses are held accountable for foreseeability, meaning a nurse of ordinary skill, care, and diligence could anticipate the risk of harm of departing from standards of practice in similar circumstances.<sup>20</sup>

Plaintiffs must be able to link the defendant's acts or omissions to the harm for which they are seeking compensation. This requires expert testimony from a physician because it requires a medical diagnosis. Unlike criminal cases, where the standard of proof is "beyond reasonable doubt," the elements of a malpractice lawsuit must be proven by a "preponderance of

19. Teoli, D., & Ghassemzadeh, S. (2023). Patient self-determination act. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK538297/>

20. Teoli, D., & Ghassemzadeh, S. (2023). Patient self-determination act. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK538297/>



evidence.” Expert testimony is required to demonstrate “medical certainty” that the nurse’s breach was the cause of an actual injury.<sup>21</sup>

## Harm

The fourth element of malpractice is harm. In a civil lawsuit, after a plaintiff has established the nurse owed a duty to the client, breached that duty, and injury was caused by the nurse’s breach, they must prove the injury resulted in damages. They request compensation for what they have lost.<sup>22</sup>

There are several types of injuries for which clients or their representatives seek compensation. Injuries can be physical, emotional, financial, professional, marital, or any combination of these. Physical injuries include loss of function, disfigurement, physical or mental impairment, exacerbation of prior medical problems, the need for additional medical care, and death. Economic injuries can include lost wages, additional medical expenses, rehabilitation, durable medical expenses, the need for architectural changes to one’s home, the loss of earning capacity, the need to hire people to perform tasks the plaintiff can no longer do, and the loss of financial support. Emotional injuries can include psychological damage, emotional distress, or other forms of mental suffering.<sup>23</sup>

Determining the specific amount a plaintiff needs can require expert witness testimony from a person known as a life care planner who is trained in analyzing and evaluating medical costs, as well as the subjective

21. Teoli, D., & Ghassemzadeh, S. (2023). Patient self-determination act. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK538297/>
22. Teoli, D., & Ghassemzadeh, S. (2023). Patient self-determination act. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK538297/>
23. Teoli, D., & Ghassemzadeh, S. (2023). Patient self-determination act. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK538297/>

determination of a jury. **Damages** are what the plaintiff requests from the court in order to remedy the injured party's situation. Damages fall into several categories, including compensatory (economic) damages, noneconomic damages, and punitive damages.<sup>24</sup> Compensatory or economic damages consist of monetary compensation for financial loss. Non economic damages consist of things like pain and suffering, but can be hard to quantify monetarily. Punitive damages are awarded to provide further punishment to the defendant in order to deter similar actions in the future.<sup>25</sup>

## Implications for Nurses

Nurses defending themselves against allegations of professional malpractice must demonstrate that their actions conformed with accepted standards of practice. They must convince a jury they acted as a reasonably prudent nurse would have in the same or similar circumstances. Nurses should follow these practices to avoid allegations of malpractice<sup>26</sup>:

- Practice according to current standards of practice.
- Carry their own professional liability insurance.
- Adhere to organizational policies and procedures. The standard of practice is to adhere to agency policy. Failing to do so creates an assumption of departure from standards.
- Document in a manner that permits accurate reconstruction of client assessments and the sequence of events, especially when notifying providers regarding clinical concerns.

24. Teoli, D., & Ghassemzadeh, S. (2023). Patient self-determination act. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK538297/>

25. Cornell Law School. (n.d.). *Legal information institute*. <https://www.law.cornell.edu/wex/damages>

26. Teoli, D., & Ghassemzadeh, S. (2023). Patient self-determination act. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK538297/>

- Maintain competence through continuing education, participation in professional conferences, membership in professional organizations, and subscriptions to professional journals.
- When using an interpreter, ensure that properly trained interpreters are used and document the name of the interpreter. The use of family, friends, or other untrained interpreters is unsafe practice and is not consistent with acceptable standards of practice.
- Maintain professional boundaries. Personal relationships with clients or their families can be red flags for juries and can be viewed as evidence of departure from professional standards.
- Engage the chain of command with client concerns and pursuing concerns to resolution.

► Read more about actual nursing malpractice cases in the [“Frequent Allegations and SBON Investigations”](#) section of the “Legal Implications” chapter in *Open RN Nursing Management and Professional Concepts*.

## Disciplinary Action by the Board of Nursing

In addition to being held liable in a court of law, nurses can have their licenses suspended or revoked by the State Board of Nursing (SBON) for unsafe nursing practice. The SBON governs nursing practice according to that state’s Nurse Practice Act to protect the public through licensure, education, legislation, and discipline. A nursing license is a contract between the state and the nurse in which the licensee agrees to provide nursing care according to that state’s Nurse Practice Act. Deviation from the Nurse Practice Act is a breach of contract that can lead to limited or revoked licensure. Nurses must

practice according to the Nurse Practice Act of the state in which they are providing client care.<sup>27</sup>

A nurse may be named in a board licensing complaint called an allegation. Allegations can be directly related to a nurse's clinical responsibilities, or they can be nonclinical (such as operating a vehicle under the influence of a substance, exhibiting unprofessional behavior, or committing billing fraud). A complaint can be filed against a nurse by anyone, such as a client, a client's family member, a colleague, or an employer. It can also be filed anonymously. After a complaint is filed, the SBON follows a disciplinary process that includes investigation, proceedings, board actions, and enforcement. The process can take months or years to resolve, and it can be costly to hire legal representation.<sup>28</sup>

Disciplinary actions by the SBON may include the following<sup>29</sup>:

- **Reprimand:** The licensee receives a public warning for a violation.
- **Limitation of License:** The licensee has conditions or requirements imposed upon their license, their scope of practice, or both.

27. American Nurses Association. (2012). *Position statement: Reduction of patient restraint and seclusion in health care settings*. <https://www.nursingworld.org/practice-policy/nursing-excellence/official-position-statements/id/reduction-of-patient-restraint-and-seclusion-in-health-care-settings/>

28. American Nurses Association. (2012). *Position statement: Reduction of patient restraint and seclusion in health care settings*. <https://www.nursingworld.org/practice-policy/nursing-excellence/official-position-statements/id/reduction-of-patient-restraint-and-seclusion-in-health-care-settings/>

29. American Nurses Association. (2012). *Position statement: Reduction of patient restraint and seclusion in health care settings*. <https://www.nursingworld.org/practice-policy/nursing-excellence/official-position-statements/id/reduction-of-patient-restraint-and-seclusion-in-health-care-settings/>

- **Suspension:** The license is completely and absolutely withdrawn and withheld for a period of time, including all rights, privileges, and authority previously conferred by the credential.
- **Revocation:** The license is completely and absolutely terminated, as well as all rights, privileges, and authority previously conferred by the credential.
- **Administrative Warning:** A warning is issued if the violation is of a minor nature, or a first occurrence and the warning will adequately protect the public. The issuance of an administrative warning is public information but the reason for issuance is not.
- **Remedial Education Order:** A remedial education order is issued when there is reason to believe that the deficiency can be corrected with remedial education, while sufficiently protecting the public.

► Find and review your state's [Nurse Practice Act](#).

## 5.5 Client Rights

When individuals with mental health disorders are admitted to a hospital, they may lose a number of abilities that we take for granted, such as the ability to come and go, schedule their time, and choose and control their activities of daily living. In many states, client's rights associated with inpatient admission to a mental health unit are spelled out in state law. Clients must be specifically informed of their rights as described in the Patient Bill of Rights document.

Clients who are determined to be legally incompetent also lose the ability to manage their financial and legal affairs and make important decisions. The Patient Self-Determination Act was passed to protect client rights.<sup>1</sup>

The Patient Self-Determination Act was passed in 1990 and is considered a landmark law for client rights. This law requires hospitals, skilled nursing facilities, home health agencies, hospice programs, and health maintenance organizations to provide clear written information for clients concerning their legal rights to make health care decisions, including the right to accept or refuse treatment. It also requires agencies to ask clients about advanced directives and to document any wishes the client has in regard to the care they do or do not want.<sup>2</sup> Interprofessional team members, including nurses, have an ethical duty to ensure that clients know and understand their health care-related rights.<sup>3</sup> See the following box for a list of client rights included in this law. These rights are further discussed in the following subsections.

1. Cady, R. F. (2010). A review of basic patient rights in psychiatric care. *JONA'S Healthcare Law, Ethics and Regulation*, 12(4), 117–127. <https://doi.org/10.1097/NHL.0b013e3181f4d357>
2. Ernstmeyer, K., & Christman, E. (Eds.). (2024). *Nursing fundamentals 2e*. Open RN | WisTech Open. <https://wtcs.pressbooks.pub/nursingfundamentals/>
3. Middleman, A. B., & Olson, K. A. (2021). Confidentiality in adolescent health care. *UpToDate*. <https://www.uptodate.com/>

## Patient Self-Determination Act<sup>4</sup>

1. The right to appropriate treatment and related services in a setting and under conditions that are the most supportive of a person's personal liberty and restrict such liberty only to the extent necessary consistent with the person's treatment needs, applicable requirements of law, and applicable judicial orders.
2. The right to an individualized, written treatment or service plan (developed promptly after admission), the right to treatment based on the plan, the right to periodic review and reassessment of treatment and related service needs, and the right to appropriate revision of the plan, including revisions necessary to provide a description of mental health services that may be needed after the person is discharged from the program or facility.
3. The right to ongoing participation, in a manner appropriate to the person's capabilities, in the planning of mental health services to be provided (including the right to participate in the development and periodic revision of the plan).
4. The right to be provided with a reasonable explanation, in terms and language appropriate to the person's mental and physical condition, the objectives of treatment, the nature and significance of possible adverse effects of recommended treatment, the reasons why the

4. Cady, R. F. (2010). A review of basic patient rights in psychiatric care. *JONA'S Healthcare Law, Ethics and Regulation*, 12(4), 117–127.  
<https://doi.org/10.1097/NHL.0b013e3181f4d357>

recommended treatment is considered appropriate, the reasons why access to certain visitors may not be appropriate, and any appropriate and available alternative treatments, services, and types of providers of mental health services.

5. The right not to receive a course of treatment in the absence of informed, voluntary, written consent to treatments except during an emergency situation or as permitted by law when the person is being treated as a result of a court order.
6. The right to not participate in experimentation in the absence of informed, voluntary, written consent.
7. The right to freedom from restraint or seclusion, other than as a course of treatment during an emergency situation with a written order by a mental health professional.
8. The right to a humane treatment environment that affords reasonable protection from harm and appropriate privacy with regard to personal needs.
9. The right to access, on request, one's mental health care records.
10. The right of a person admitted to residential or inpatient care to converse with others privately, to have convenient and reasonable access to the telephone and mail, and to see visitors during regularly scheduled hours.
11. The right to be informed promptly at the time of admission and in writing of these rights.
12. The right to assert grievances with respect to infringement of these rights.
13. The right to exercise these rights without reprisal.
14. The right of referral to other providers upon discharge.



# Informed Consent

**Informed consent** is the fundamental right of an individual to accept or reject health care. Based on the Patient Self-Determination Act, clients have the right to give informed consent before receiving medical assessment or treatment (except in emergency situations when imminent harm may occur to themselves or others). Most states allow a client to sue for battery if consent is not obtained before medical treatment is given.<sup>5</sup> However, a client must be competent and have the legal capacity to give informed consent.

**Competency** is a legal term related to the degree of cognitive ability an individual has to make decisions or carry out specific acts. Individuals are considered competent until they have been declared incompetent in a formal legal proceeding. If found incompetent, the individual is appointed a legal guardian or representative who is responsible for providing or refusing consent (while considering the individual's wishes). Guardians are typically family members such as spouses, adult children, or parents. If family members are unavailable or unwilling to serve in this role, the court may appoint a court-trained guardian.<sup>6</sup>

**Capacity** is a functional determination that an individual is or is not capable of making a medical decision within a given situation. It is outside the scope of practice for nurses to formally assess capacity, but nurses may initiate the evaluation of client capacity and contribute assessment information. Capacity

5. Cady, R. F. (2010). A review of basic patient rights in psychiatric care. *JONA'S Healthcare Law, Ethics and Regulation*, 12(4), 117–127. <https://doi.org/10.1097/NHL.0b013e3181f4d357>
6. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

may be a temporary or permanent state. The following box outlines situations where the nurse may question a client's decision-making capacity.<sup>7 8</sup>

### Triggers for Questioning a Client's Decision-Making Capacity

- Unawareness of surroundings
- Absence of questions about the treatment being offered or provided
- New inability to perform activities of daily living
- Disruptive behavior or agitation
- Labile emotions
- Hallucinations
- Intoxication

When a client does not have the capacity to provide informed consent, health care providers must obtain substituted consent for treatments. Substituted consent is authorization that another person gives on behalf of the client. For example, the activation of a client's health care power of attorney is an example of substituted consent. Substituted consent may also come from a court-appointed guardian or if state law permits, from next of kin.<sup>9</sup>

7. Cady, R. F. (2010). A review of basic patient rights in psychiatric care. *JONA'S Healthcare Law, Ethics and Regulation*, 12(4), 117–127. <https://doi.org/10.1097/NHL.0b013e3181f4d357>
8. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.
9. Cady, R. F. (2010). A review of basic patient rights in psychiatric care. *JONA'S Healthcare Law, Ethics and Regulation*, 12(4), 117–127. <https://doi.org/10.1097/NHL.0b013e3181f4d357>

- ▶ Read more about informed consent and capacity in the “[Other Legal Issues](#)” section of *Open RN Nursing Management and Professional Concepts*.

## Protective Placement

Guardianships and protective orders are legal methods in states for appointing an alternative decision-maker and identifying required services for individuals who are legally incompetent. Legally incompetent individuals may have developmental disabilities, chronic and serious mental illness, severe substance use disorders, or other conditions that limit their decision-making ability. A court can issue orders for a person who has a guardian to be protectively placed. The legal standard basically states that without the protective placement, the individual is incapable of providing for their own care and well-being that it creates a substantial risk of serious harm to themselves or others. Protective services may include case management, in-home care, nursing services, adult day care, or inpatient treatment. Protective placements must be the least restrictive setting necessary to meet the individual's needs and must be reviewed annually by the court.

## Psychiatric Advance Directive

A **Psychiatric Advance Directive (PAD)** is a legal document that describes a person's preferences for future mental health treatment or names an individual to make treatment decisions for them if they are in a crisis and unable to make decisions. Many people with mental illness, their family members, and health professionals are not familiar with PADs.<sup>10</sup>

10. The Joint Commission. (n.d.) <https://www.jointcommission.org/>

For states that do not have laws regarding PADs, an individual can still draft a PAD under the more general statutes connected to advance health care directives and living wills. However, a PAD is more beneficial because of the unique issues of mental health care and treatment, such as medication preferences and inpatient treatment considerations and the fact that a person with mental health disorders can experience recovery and wellness over time.<sup>11</sup>

- ▶ Read more about [Psychiatric Advance Directives](#) on the National Alliance on Mental Illness website.

## Restraints and Seclusion

**Restraints** are devices used in health care settings to prevent clients from causing harm to themselves or others when alternative interventions are not effective. A restraint is a device, method, or process that is used for the specific purpose of restricting a client's freedom of movement without the permission of the person. Restraints include mechanical devices such as a tie wrist device, chemical restraints, or seclusion. The Joint Commission defines a chemical restraint as a drug used to manage a client's behavior, restrict the client's freedom of movement, or impair the client's ability to appropriately interact with their surroundings that is not standard treatment or dosage for the client's condition.<sup>12</sup> It is important to note that the definition states the medication "is not standard treatment or dosage for the client's condition." For example, administering prescribed benzodiazepines as standard treatment to manage the symptoms of a diagnosed mental health disorder is

11. The Joint Commission. (n.d.) <https://www.jointcommission.org/>

12. The Joint Commission. (n.d.) <https://www.jointcommission.org>

not considered a chemical restraint, but administering benzodiazepines to limit a client's movement is considered a chemical restraint.

**Seclusion** is defined as the confinement of a client in a locked room or an area from which they cannot exit on their own. Seclusion should only be used for the management of violent or self-destructive behavior. Seclusion limits freedom of movement because, although the client is not mechanically restrained, they cannot leave the area.<sup>13,14</sup>

Although restraints are used with the intention to keep a client safe, they impact a client's psychological safety and dignity and can cause additional safety issues and death. A restrained person has a natural tendency to struggle and try to remove the restraint and can fall or become fatally entangled in the restraint. Furthermore, immobility that results from the use of restraints can cause pressure injuries, contractures, and muscle loss. Restraints take a large emotional toll on the client's self-esteem and may cause humiliation, fear, and anger.

## Restraint Guidelines

The American Nurses Association (ANA) has established evidence-based guidelines that a restraint-free environment is the standard of care. The ANA encourages the participation of nurses to reduce client restraints and seclusion in all health care settings. Restraining or secluding clients is viewed

13. Wood County, Wisconsin. (n.d.). *Guardianship and protective placements*. <https://www.co.wood.wi.us/Departments/HumanServices/GuardianshipAndProtectivePlacement.aspx>

14. Knox, D. K., & Holloman, G. H. (2012). Use and avoidance of seclusion and restraint: Consensus statement of the American Association for Emergency Psychiatry Project Beta Seclusion and Restraint Workgroup. *The Western Journal of Emergency Medicine* 13(1), 35-40. <https://doi.org/10.5811/westjem.2011.9.6867>

as contrary to the goals and ethical traditions of nursing because it violates the fundamental client rights of autonomy and dignity. However, the ANA also recognizes there are times when there is no viable option other than restraints to keep a client safe, such as during an acute psychotic episode when client and staff safety are in jeopardy due to aggression or assault. The ANA also states that restraints may be justified in some clients with severe dementia or delirium when they are at risk for serious injuries such as a hip fracture due to falling.<sup>15</sup>

The ANA provides the following guidelines: “When restraint is necessary, documentation should be done by more than one witness. Once restrained, the client should be treated with humane care that preserves human dignity. In those instances where restraint, seclusion, or therapeutic holding is determined to be clinically appropriate and adequately justified, registered nurses who possess the necessary knowledge and skills to effectively manage the situation must be actively involved in the assessment, implementation, and evaluation of the selected emergency measure, adhering to federal regulations and the standards of the The Joint Commission (2009) regarding appropriate use of restraints and seclusion.” Nursing documentation typically includes information such as client behavior necessitating the restraint, alternatives to restraints that were attempted, the type of restraint used, the time it was applied, the location of the restraint, and client education regarding the restraint.<sup>16</sup>

Any health care facility that accepts Medicare and Medicaid reimbursement

15. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. <https://www.uptodate.com/>

16. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. <https://www.uptodate.com/>

must follow federal guidelines for the use of behavioral restraints. These guidelines include the following<sup>17</sup>:

- When a restraint is the only viable option, it must be discontinued at the earliest possible time.
- Orders for the use of seclusion or restraint can never be written as a standing order or PRN (as needed).
- The treating physician must be consulted as soon as possible if the restraint or seclusion is not ordered by the client's treating physician.
- A physician or licensed independent practitioner must see and evaluate the need for the restraint or seclusion within one hour after the initiation.
- After restraints have been applied, the nurse should follow agency policy for frequent monitoring and regularly changing the client's position to prevent complications. Nurses must also ensure the client's basic needs (e.g., hydration, nutrition, and toileting) are met. Range of motion exercises and circulatory checks are typically provided hourly. Some agencies require a 1:1 client sitter or continuous monitoring when restraints are applied or seclusion is implemented.
- Each written order for a physical restraint or seclusion is limited to 4 hours for adults, 2 hours for children and adolescents ages 9 to 17, or 1 hour for clients under 9. The original order may only be renewed in accordance with these limits for up to a total of 24 hours. After the original order expires, a physician or licensed independent practitioner (if allowed under state law) must see and assess the client before issuing a new order.

▶ Review safe use of restraints and alternatives to restraints in the

17. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. <https://www.uptodate.com/>

► “Restraints” section of the “Safety” chapter in *Open RN Nursing Fundamentals*.

## Admission for Care

When clients with mental health disorders are admitted for inpatient care, the type of admission dictates certain rights and aspects of their treatment plan. Admissions may be voluntary, emergency, or involuntary.

### Voluntary Admission

Individuals over age 16 who present to a psychiatric facility and request hospitalization are considered **voluntary admissions**. Clients admitted under voluntary admission have certain rights that differ from emergency and involuntary admissions. For example, they are considered competent with the capacity to make health care decisions (unless determined otherwise). Therefore, they have the right to refuse treatment, including psychotropic medications, unless they become a danger to themselves or others.<sup>18</sup>

Clients with voluntary admission do not necessarily have an absolute right to discharge at any time but may be required to request discharge. This gives the health care team an opportunity to initiate a procedure to change the client’s admission status to involuntary if needed and associated legal requirements are met.<sup>19</sup>

18. Cady, R. F. (2010). A review of basic patient rights in psychiatric care. *JONA’S Healthcare Law, Ethics and Regulation*, 12(4), 117–127. <https://doi.org/10.1097/NHL.0b013e3181f4d357>

19. Cady, R. F. (2010). A review of basic patient rights in psychiatric care. *JONA’S*



## Emergency Admissions

Many states allow individuals to be admitted to psychiatric facilities under **emergency admission** status when they are deemed likely to harm themselves or others. State laws define the exact procedure for the initial evaluation, possible length of detainment, and treatment provided. All clients who are admitted as emergency admissions require diagnosis, evaluation, and emergency treatment according to state law. At the end of the admission period, the facility must either discharge the client, change their status to voluntary admission, or initiate a civil court hearing to determine the need for continuing treatment on an involuntary basis.<sup>20</sup>

During an emergency admission, the client's right to come and go is restricted, but they have a right to consult with an attorney and prepare for a hearing. Clients may be forced to receive psychotropic medications if they continue to be a danger to themselves or others. However, invasive procedures like electroconvulsive therapy (ECT) are not permitted unless they are ordered by the court or consented to by the client or their legal guardian.<sup>21</sup>

## Involuntary Admissions

There may be circumstances when a person becomes so mentally ill they are at risk of hurting themselves or others, and involuntary admission for care

*Healthcare Law, Ethics and Regulation*, 12(4), 117–127. <https://doi.org/10.1097/NHL.0b013e3181f4d357>

20. Cady, R. F. (2010). A review of basic patient rights in psychiatric care. *JONA'S Healthcare Law, Ethics and Regulation*, 12(4), 117–127. <https://doi.org/10.1097/NHL.0b013e3181f4d357>

21. Cady, R. F. (2010). A review of basic patient rights in psychiatric care. *JONA'S Healthcare Law, Ethics and Regulation*, 12(4), 117–127. <https://doi.org/10.1097/NHL.0b013e3181f4d357>

becomes necessary even though the individual does not desire care. An individual can have an **involuntary admission** to a psychiatric facility if they are diagnosed with a mental illness, pose a danger to themselves or others, are gravely disabled (e.g., unable to provide themselves basic necessities like food, clothing, and shelter), or are in need of treatment but their mental illness prevents voluntary help-seeking behaviors. The legal procedures are different in each state, but standards for involuntary admission are similar.<sup>22, 23</sup>

Because involuntary commitment is a serious matter, there are strict legal protections established by the U.S. Supreme Court. The standard of proof of “mentally ill and dangerous to self or others” must be based on “clear and convincing evidence.” Therefore, two physicians must certify the individual’s mental health status. Additionally, the client has the right to legal counsel and can take the case to a judge who can order release. If not released, the client can be involuntarily committed to a state-specified number of days with interim court appearances. Most states permit a 72-hour admission followed by a formal hearing. If the client feels they are being held without just cause, they can file a writ of habeas corpus (i.e., a formal written order to free the person). The court makes a decision based on the “least restrictive alternative”

22. Cady, R. F. (2010). A review of basic patient rights in psychiatric care. *JONA'S Healthcare Law, Ethics and Regulation*, 12(4), 117–127. <https://doi.org/10.1097/NHL.0b013e3181f4d357>
23. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

doctrine, meaning the least drastic action is taken to achieve the purpose of care.<sup>24,25</sup> Review your state's laws regarding involuntary admissions.

- ▶ For example, Wisconsin state law, referred to as “Chapter 51,” dictates the requirements for involuntary admissions and is further explained by [NAMI Kenosha County](#). View the PDF Document: [CHAPTER 51 State Alcohol, Drug Abuse, Developmental Disabilities and Mental Health Act](#)

## INVOLUNTARY ADMISSION OF MINORS

Special considerations apply to minors receiving psychiatric care. Many states grant minors aged 12-18 the right to provide consent for mental health treatment and to protest involuntary admission unless they are a risk to themselves or others. In many cases, a neutral mental health review officer is assigned to the case to ensure rights are upheld.<sup>26</sup> State laws are complex; therefore, nurses must be aware of legal protections of minors in the state in which they work.

24. Cady, R. F. (2010). A review of basic patient rights in psychiatric care. *JONA'S Healthcare Law, Ethics and Regulation*, 12(4), 117–127. <https://doi.org/10.1097/NHL.0b013e3181f4d357>
25. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.
26. Cady, R. F. (2010). A review of basic patient rights in psychiatric care. *JONA'S Healthcare Law, Ethics and Regulation*, 12(4), 117–127. <https://doi.org/10.1097/NHL.0b013e3181f4d357>

- Read additional information about [Clients' Rights for Minors in Wisconsin](#).

## CRIMINAL CASES AND PLEAS OF INSANITY

If a defendant pleads an insanity defense in a criminal case, they are involuntarily admitted to a mental health facility for an evaluation period determined by state law. During this time an interprofessional team of mental health professionals (including nurses) evaluates the individual's need for hospitalization and notifies the court of their treatment recommendations. This specialized mental health care is referred to as forensic psychiatry.<sup>27</sup>

## Discharge

When clients with mental health disorders are hospitalized, their admission status may impact their rights related to discharge. There are four main types of discharge<sup>28</sup>:

- **Unconditional Discharge:** Unconditional discharge refers to unconditional termination of the legal client and institution relationship. Discharge may be ordered by a psychiatrist, advanced practice provider, or the court.
- **Release Against Medical Advice (AMA):** Clients who were admitted voluntarily may elect to leave an institution against the advice of the

27. Cady, R. F. (2010). A review of basic patient rights in psychiatric care. *JONA'S Healthcare Law, Ethics and Regulation*, 12(4), 117–127. <https://doi.org/10.1097/NHL.0b013e3181f4d357>

28. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

health care provider.

- **Conditional Release:** Conditional release means the client is discharged from inpatient care but requires outpatient treatment for a specified period of time. If the client was involuntarily admitted, they can be readmitted based on the original commitment order if they don't participate in outpatient treatment.
- **Assisted Outpatient Treatment:** Assisted outpatient treatment means the conditional release is court-ordered. This treatment is tied to services and goods provided by social welfare agencies, such as disability benefits and housing.

## Reporting Unsafe or Impaired Professionals

Clients have the right to humane treatment and reasonable protection from harm. For example, if a suicidal client is admitted and left alone with the means of self-harm, the nurse has a duty to protect the client and can be held liable for injuries or death that occurs.

Nurses also have a duty to protect clients from suspected negligence by a colleague. In many states, nurses have a legal duty to intervene and report risks of harm to clients. This may include reporting concerns to a supervisor, the institution, and/or the state Board of Nursing. For example, nurses must report suspected drug diversion by colleagues because it can impact safe and humane treatment of clients. Read more about drug diversion and substance use disorder in nursing in the "[Substance Use Disorders](#)" chapter.

► View the NCSBN webpage on [Substance Use Disorder in Nursing](#).

## 5.6 Learning Activities



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=944#h5p-29>

1



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=944#h5p-30>

2



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=944#h5p-31>

1. “MH Legal/Ethical Glossary Cards ” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

2. “MH Legal Ethical Question Set 1 ” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

- ▶ Test your clinical judgment with a NCLEX Next Generation-style question: [Chapter 5, Assignment 1](#)<sup>4</sup>



- ▶ Test your clinical judgment with a NCLEX Next Generation-style question: [Chapter 5, Assignment 2](#)<sup>5</sup>



- ▶ Test your clinical judgment with a NCLEX Next Generation-style case study: [Chapter 5, Case Study 1](#)<sup>6</sup>

3. “MH Legal Ethical Question Set 2” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

4. “MH Legal Ethical Next Gen Question 1” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

5. “MH Legal Ethical Next Gen Question 2” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

6. “MH Legal Ethical Case Study” by Kellea Ewen is licensed under [CC BY-NC 4.0](#)

## V Glossary

**Adults at risk:** Adults who have a physical or mental condition that impairs their ability to care for their own needs and are at risk for neglect and/or abuse.

**Assault:** Intentionally putting another person in reasonable apprehension of an imminent harmful or offensive contact.

**Autonomy:** The capacity to determine one's own actions through independent choice, including demonstration of competence. The nurse's primary ethical obligation is client autonomy.

**Battery:** Intentional causation of harmful or offensive contact with another person without that person's consent.

**Beneficence:** Benefiting others by preventing harm, removing harmful conditions, or affirmatively acting to benefit another or others, often going beyond what is required by law.

**Board of Nursing:** The state-specific licensing and regulatory body that sets standards for safe nursing care and issues nursing licenses to qualified candidates based on the Nurse Practice Act enacted by that state's legislature.

**Capacity:** A functional determination that an individual is or is not capable of making a medical decision within a given situation.

**Competency:** A legal term related to the degree of cognitive ability an individual has to make decisions or carry out specific acts.

**Confidentiality:** The right of an individual to have personal, identifiable medical information kept private.

**Civil law:** The rights, responsibilities, and legal relationships between private citizens and involves compensation to the injured party.

**Criminal law:** A system of laws that punishes individuals who commit crimes.



**Damages:** What the plaintiff requests from the court in order to remedy the injured party's situation.

**Defamation of character:** Actions when an individual makes negative, malicious, and false remarks about another person to damage their reputation.

**Defendants:** The parties named in the lawsuit.

**Emergency admission:** Individuals are admitted to psychiatric facilities under emergency admission status when they are deemed likely to harm themselves or others.

**False imprisonment:** An act of restraining another person and causing that person to be confined in a bounded area.

**Fraud:** An intentional tort that occurs when an individual is deceived for personal gain.

**Health Insurance Portability and Accountability Act (HIPAA):** Federal regulations to ensure the privacy and protection of personal records and information.

**Informed consent:** The fundamental right of an individual to receive information about the risks, benefits, and alternatives in order to make a healthcare decision.

**Intentional tort:** A wrong that the defendant knew (or should have known) would be caused by their actions.

**Involuntary admission:** Circumstances when a person becomes so mentally ill they are at risk of hurting themselves or others, and inpatient care becomes necessary even though the individual does not desire inpatient care.

**Justice:** A moral obligation to act on the basis of equality and equity and a standard linked to fairness for all in society.

**Libel:** Written defamation.

**Malpractice:** A specific term used for negligence committed by a health professional with a license.

**Mandated reporters:** Nurses and other professionals required by state law to report suspected neglect and/or abuse of children, adults at risk, and the elderly.

**Negligence:** The failure to exercise the ordinary care a reasonable person would use in similar circumstances.

**Nonmaleficence:** The bioethical principle that specifies a duty to do no harm and balances avoidable harm with benefits of good achieved.

**Nurse Practice Act:** Law enacted by that state's legislature that establishes regulations for nursing practice within that state and defines the scope of nursing practice.

**Plaintiff:** A person bringing a lawsuit.

**Protected Health Information (PHI):** Individually identifiable health information including demographic data that relates to the individual's past, present, or future physical or mental health or condition; the provision of health care to the individual; and the past, present, or future payment for the provision of health care to the individual.

**Psychiatric Advance Directive (PAD):** A legal document that describes a person's preferences for future mental health treatment or names an individual to make treatment decisions for them if they are in a crisis and unable to make decisions.

**Restraints:** Devices used in health care settings to prevent clients from causing harm to themselves or others when alternative interventions are not effective.

**Role fidelity:** Being responsible for providing competent nursing care.

**Seclusion:** The confinement of a client in a locked room from which they

cannot exit on their own. It is generally used as a method of discipline, convenience, or coercion.

**Slander:** Spoken defamation.

**Standards of Professional Nursing Practice:** Authoritative statements from the American Nurses Association regarding the actions and behaviors that all registered nurses, regardless of role, population, specialty, and setting, are expected to perform competently.

**Standards of Professional Performance:** Twelve standards set by the American Nurses Association that describe a nurse's professional behavior, including activities related to ethics, advocacy, respectful and equitable practice, communication, collaboration, leadership, education, scholarly inquiry, quality of practice, professional practice evaluation, resource stewardship, and environmental health.

**Tort:** An act of commission or omission that gives rise to injury or harm to another and amounts to a civil wrong for which courts impose liability.

**Unintentional tort:** A wrong that occurs when the defendant's actions or inactions were unreasonably unsafe. Unintentional torts can result from acts of commission (i.e., doing something a reasonable nurse would not have done) or omission (i.e., failing to do something a reasonable nurse would do).

**Veracity:** Telling the truth.

**Voluntary admission:** An individual over age 16 who presents to a psychiatric facility and requests hospitalization. They are considered competent with the capacity to make health care decisions (unless determined otherwise).







### Learning Objectives

- Relate the anatomy and physiology of the central nervous system to mental health disorders and psychotropic medications
- Describe classes of psychotropic medications, their mechanism of action, side effects, and health teaching topics

**Psychotropic medications** are medications that affect the mind, emotions, and behavior. This chapter reviews the anatomy and physiology of the central nervous system (CNS) as it relates to mental health disorders and medications followed by an overview of several classes of psychotropic medications.

► Medication guidelines change frequently so it is vital for nurses to consult evidence-based resources for the latest drug information, warnings, and client education guidelines when administering medications. Free reference information is provided by the National Library of Medicine (NLM) on [DailyMed](#) and the [MedlinePlus](#) websites.

- DailyMed is a database containing current information from the [Food and Drug Administration](#) on prescription and over-the-counter medications. It provides essential information to health professionals for the safe and effective use of medications, including indications, dosage

and administration, contraindications, boxed warnings and precautions, adverse reactions, drug interactions, information about use in specific populations, and other important information for health care practitioners.<sup>1</sup>

- MedlinePlus is an online health information resource for clients and their loved ones with easy-to-understand medication information.<sup>2</sup>

- ▶ Additional information regarding psychotropic medications can be found on the National Alliance on Mental Illness' [Mental Health Medications](https://nami.org/About-Mental-Illness/Treatments/Mental-Health-Medications) webpage.<sup>3</sup>

1. DailyMed. (n.d.). *About us*. <https://dailymed.nlm.nih.gov/dailymed/about-dailymed.cfm>

2. MedlinePlus. (n.d.). *About us*. <https://medlineplus.gov/>

3. National Alliance on Mental Illness. (n.d.). *Mental health medications*. <https://nami.org/About-Mental-Illness/Treatments/Mental-Health-Medications>



## 6.2 Review of the Nervous System

Understanding the anatomy and physiology of the nervous system is essential to understanding how psychotropic medications work. The nervous system regulates both conscious and unconscious processes in the body, including thought, movement, sensation, and automatic functions like heart rate and digestion. It is divided into two main parts: the central nervous system (CNS) and the peripheral nervous system (PNS). See Figure 6.1<sup>1</sup> for an illustration of the central and peripheral nervous systems.




Figure 6.1 The Central and Peripheral Nervous Systems

1. “1201 Overview of Nervous System.jpg” by OpenStax is licensed under CC BY 4.0. Access for free at <https://openstax.org/books/anatomy-and-physiology/pages/12-1-basic-structure-and-function-of-the-nervous-system>

# The Central Nervous System (CNS)

The CNS consists of the brain and spinal cord, serving as the primary control center for processing and integrating information. The brain regulates cognition, emotions, and autonomic functions, while the spinal cord transmits signals between the brain and the rest of the body.

The brain's complex systems also regulate key functions such as pain perception, reward processing, and emotional regulation. For example, the opioid system in the brain plays a crucial role in pain and reward, and is integral to understanding addiction and the effects of psychoactive substances. This system, along with other neurotransmitter systems, such as dopamine and serotonin, is central to many mental health disorders. These neurotransmitters act as messengers, influencing mood, behavior, and thought processes, and are frequently targeted by psychotropic medications.

(We'll explore these systems and their role in mental health disorders and medication therapy in more detail later.)

# The Peripheral Nervous System (PNS)

The PNS consists of sensory and motor neurons that connect the CNS to the rest of the body, playing a vital role in both voluntary and involuntary processes. It functions as a bridge, relaying information between the CNS and various organs, muscles, and sensory receptors.

- Sensory neurons detect stimuli from the environment (such as light, sound, and temperature) and send signals to the brain, creating conscious perception of these stimuli.
- Motor neurons transmit signals from the brain to muscles and glands, initiating voluntary movements and autonomic responses.

Motor neurons are further divided into:

- The somatic nervous system, which controls voluntary muscle movements, such as those required for walking, talking, and writing.

- The autonomic nervous system (ANS), which regulates involuntary functions such as heart rate, digestion, and respiratory rate, maintaining homeostasis in the body.

The peripheral nervous system consists of sensory neurons and motor neurons. Sensory neurons sense the environment and conduct signals to the brain that become a person's conscious perception of that stimulus. This conscious perception may lead to a motor response that is conducted from the brain to the peripheral nervous system via motor neurons. Motor neurons are part of the somatic nervous system that stimulates voluntary movement of muscles and the autonomic nervous system that controls involuntary responses.

## Sympathetic and Parasympathetic Nervous System

The autonomic nervous system is further subdivided into the sympathetic nervous system (SNS) and the parasympathetic nervous system (PNS), each with distinct and complementary functions:

- Sympathetic Nervous System (SNS): The SNS is often associated with the “fight-or-flight” response. When activated, it prepares the body for stress or emergency situations by increasing heart rate, constricting blood vessels to raise blood pressure, and causing bronchodilation (widening of the airways) to increase oxygen intake.
- Parasympathetic Nervous System (PNS): In contrast, the PNS supports “rest-and-digest” functions, helping the body conserve energy and maintain a state of relaxation. PNS activation slows the heart rate, dilates blood vessels to reduce blood pressure, and promotes bronchoconstriction (narrowing of the airways).

The body maintains homeostasis by balancing the stimulation of the SNS and PNS. For example, during physical activity, the SNS dominates to help the body respond to the increased demand for energy, while during rest, the PNS takes over to return the body to a state of calm and relaxation.

See Figure 6.2<sup>2</sup> to compare the effects of PNS and SNS stimulation on target organs.




Figure 6.2 Effects of SNS and PNS Stimulation

## SNS Receptors

**SNS receptors** include Alpha-1, Alpha-2, Beta-1, and Beta-2 receptors which are stimulated by epinephrine and norepinephrine. Medications that activate these receptors are referred to as **adrenergic agonists** because they mimic the effects of the body's natural SNS response. For example, stimulants like methylphenidate are adrenergic agonists used to treat attention deficit hyperactivity disorder (ADHD). Conversely, **adrenergic antagonists** block SNS

2. "[Updated SNS-PNS image.png](#)" by Meredith Pomietlo for [Open RN](#) is licensed under [CC BY 4.0](#)

receptors. For example, propranolol is a Beta-2 antagonist used to manage the physical symptoms of severe anxiety (e.g., trembling, rapid heartbeat, and sweating).

## PNS Receptors

**PNS receptors** include nicotinic and muscarinic receptors that are stimulated by acetylcholine (ACh). Drugs that stimulate nicotinic and muscarinic receptors are called **cholinergics**. For example, nicotine in tobacco products stimulates nicotinic receptors. Stimulation of muscarinic receptors primarily causes smooth muscle contraction. An example of a muscarinic agonist is bethanechol used to treat urinary retention by increasing the tone of the detrusor muscle to increase bladder emptying.<sup>3</sup> Drugs that block the effects of PNS receptors are called **anticholinergics**. For example, benztropine is an anticholinergic used to treat muscle spasms associated with extrapyramidal symptoms from antipsychotic medications.<sup>4</sup> Many psychotropic medications cause anticholinergic adverse effects that can be especially hazardous for older adults. **SLUDGE** is a mnemonic for anticholinergic side effects: Salivation decreased, Lacrimation decreased, Urinary retention, Drowsiness/dizziness, GI upset, and Eyes (blurred vision/dry eyes). See Figure 6.3<sup>5</sup> for an illustration of the “SLUDGE” effects caused by anticholinergics.

3. Padda, I. S., & Derian, A. (2024). *Bethanechol*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK560587/>
4. Ahuja, A., Patel, P., & Abdijadid, S. (2024). *Benztropine*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK560633/>
5. “SLUDGE effects of Anticholinergics” by Dominic Slausen at [Chippewa Valley Technical College](#) is licensed under [CC BY 4.0](#)




Figure 6.3 SLUDGE Effects of Anticholinergics

## Opioid System

The opioid system in the brain controls pain, reward, and addictive behaviors. There are three types of **opioid receptors** called mu, delta, and kappa receptors. Opioid receptors are stimulated by endogenous peptides released by neurons (such as endorphins) and exogenous opiates. **Opiates** include powerful analgesics (such as morphine and oxycodone) prescribed to treat moderate to severe pain. Opiates also include illicit drugs (such as heroin). Chronic use of prescribed and illicit opiates can be highly addictive because of their actions on the reward system of the brain.<sup>6</sup> Read more about the addictive cycle in the “[Substance Use Disorders](#)” chapter.

## Neurotransmitters

Neurons are responsible for the communication that the nervous system

6. Dhaliwal, A., & Gupta, M. (2023). *Physiology, opioid receptor*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK546642/>

provides. **Neurotransmitters** are chemical substances released at the end of a neuron by the arrival of an electrical impulse. They diffuse across the synapse and cause the transfer of the impulse to another nerve fiber, a muscle fiber, or other structure. Neurotransmitters interact with specific receptors like a key and a lock. See Figure 6.4<sup>7</sup> for an illustration of neuron communication with neurotransmitters and receptors.

7. “Chemical synapse schema cropped.jpg” by Looie496 is licensed under Public Domain. Access for free at [https://med.libretexts.org/Bookshelves/Anatomy\\_and\\_Physiology/Book%3A\\_Anatomy\\_and\\_Physiology\\_\(Boundless\)/10%3A\\_Overview\\_of\\_the\\_Nervous\\_System/10.1%3A\\_Introduction\\_to\\_the\\_Nervous\\_System/10.1A%3A\\_Organization\\_of\\_the\\_Nervous\\_System](https://med.libretexts.org/Bookshelves/Anatomy_and_Physiology/Book%3A_Anatomy_and_Physiology_(Boundless)/10%3A_Overview_of_the_Nervous_System/10.1%3A_Introduction_to_the_Nervous_System/10.1A%3A_Organization_of_the_Nervous_System)






Figure 6.4 Neuron Communication With Neurotransmitters

There are several types of neurotransmitters associated with mental health disorders and psychoactive medications, including acetylcholine, glutamate, GABA, glycine, dopamine, serotonin, epinephrine, norepinephrine, and histamine<sup>8,9</sup>:

8. Betts, J. G., Young, K. A., Wise, J. A., Johnson, E., Poe, B., Kruse, D. H., Korol, O., Johnson, J. E., Womble, M., & DeSaix, P. (2022). *Anatomy and physiology 2e*. OpenStax. <https://openstax.org/books/anatomy-and-physiology-2e/pages/1-introduction>
9. Sheffler, Z. M., Reddy, V., & Pillarisetty, L. S. (2023). *Physiology*,



- **Acetylcholine:** Acetylcholine stimulates nicotinic and muscarinic receptors in the parasympathetic nervous system. Other substances also bind to these receptors. For instance, nicotine (in tobacco products) binds to nicotinic receptors, and muscarine (products of specific mushrooms used as a hallucinogenic) binds to muscarinic receptors.
- **Glutamate:** Glutamate is an excitatory neurotransmitter. Elevated levels of glutamate are associated with psychosis symptoms that can occur with schizophrenia, as well as with illicit drug use such as methamphetamines. Conversely, lamotrigine, a medication used to treat bipolar disorder, inhibits glutamate.
- **Gamma-Aminobutyric Acid and Glycine:** Gamma-aminobutyric acid (GABA) and glycine are inhibitory neurotransmitters that act like brakes in a car by slowing down overexcited nerve cells. Low levels of GABA are associated with seizures, anxiety, mania, and impulse control.
- **Dopamine:** Dopamine plays an essential role in several brain functions, including learning, motor control, reward, emotion, and executive functions. It is associated with several mental health disorders and is targeted by many psychotropic medications. For example, bupropion is an antidepressant that inhibits dopamine reuptake, leading to increased dopamine levels in the synapse and relieving the symptoms of depression. Conversely, chlorpromazine blocks dopamine receptors and is used to treat psychosis, but this blockade can cause extrapyramidal side effects (involuntary and uncontrolled muscle movements).
- **Serotonin:** Serotonin modulates multiple neuropsychological processes such as mood, sleep, libido, and temperature regulation. Abnormal levels of serotonin have been linked to many mental health disorders such as depression, bipolar disorder, and anxiety. Many psychotropic medications target serotonin. For example, fluoxetine belongs to a class of antidepressants called selective serotonin reuptake inhibitors (SSRIs). SSRIs prevent the reuptake of serotonin at the synapse, making more of

*neurotransmitters*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK539894/>

the chemical available in the brain and relieving depression.

- **Norepinephrine and Epinephrine:** Norepinephrine and epinephrine stimulate alpha- and beta-receptors in the sympathetic nervous system. Their release exerts effects on a variety of body processes, including stress, sleep, attention, and focus. Many psychotropic medications target these neurotransmitters. For example, venlafaxine belongs to a class of antidepressants called norepinephrine reuptake inhibitors (NRIs). NRIs are prescribed to treat depression by preventing the reuptake of norepinephrine at the synapse and boosting levels of norepinephrine in the brain.
- **Histamine:** Histamine mediates homeostatic functions in the body, promotes wakefulness, modulates feeding behavior, and controls motivational behavior. For example, diphenhydramine, a histamine antagonist, causes drowsiness and is also used to treat extrapyramidal symptoms.



View a YouTube<sup>10</sup> video called the [Neurotransmitter Anatomy](https://youtu.be/fYUpLvM5X7A) that compares the effects of neurotransmitters.

<sup>10</sup>. Khan Academy. (2014). Neurotransmitter Anatomy. [Video]. YouTube. All rights reserved. <https://youtu.be/fYUpLvM5X7A>

## 6.3 Antidepressants

Antidepressants are commonly used to treat depression and are also used to treat other conditions, such as anxiety, chronic pain, and insomnia. According to a research review by the Agency for Healthcare Research and Quality, antidepressant medications work relatively well in improving symptoms of depression and to keep depression symptoms from coming back.<sup>1</sup> For reasons not yet well understood, some people respond better to certain antidepressant medications than to others, so an individual may have to try different types of antidepressants before finding one that effectively treats their symptoms.<sup>2</sup> Additionally, it may take antidepressants two or more weeks to achieve peak effect.

There are several classes and types of antidepressants, including selective serotonin reuptake inhibitors (SSRIs), serotonin and norepinephrine reuptake inhibitors (SNRIs), norepinephrine and dopamine reuptake inhibitors (NDRIs), serotonin antagonist and reuptake inhibitors, tricyclic antidepressants (TCAs), and monoamine oxidase inhibitors (MAOIs). TCAs and MAOIs are often referred to as first-generation antidepressants because they were first marketed in the 1950s. They have many side effects and are not prescribed as frequently to treat depression as are SSRIs, SNRIs, and NDRI that have fewer side effects.

### Selective Serotonin Reuptake Inhibitors

Selective serotonin reuptake inhibitors (SSRIs) prevent the uptake of serotonin at the synapse, causing the serotonin neurotransmitter to stay in

1. Volpi-Abadie, J., Kaye, A. M., & Kaye, A. D. (2013). Serotonin syndrome. *The Ochsner Journal*, 13(4), 533–540. <https://dx.doi.org/10.1177%2F2045125311400779>
2. National Institute of Mental Health. (2023). *Mental health medications*. U.S. Department of Health & Human Services. <https://www.nimh.nih.gov/health/topics/mental-health-medications>

the synapse longer and overall raise the level of serotonin in the brain. SSRIs are primarily used to treat depression but are also used to treat bipolar disorder, obsessive-compulsive disorder, bulimia, panic disorder, post-traumatic stress disorder, anxiety, premenstrual syndrome, and migraines. Examples of common SSRIs include fluoxetine, citalopram, sertraline, paroxetine, and escitalopram.<sup>3</sup>

## Serotonin Norepinephrine Reuptake Inhibitor (SNRI)

Serotonin norepinephrine reuptake inhibitors (SNRI) prevent the reuptake of serotonin and norepinephrine, with weak inhibition of dopamine reuptake. Examples of SNRIs are venlafaxine and duloxetine.<sup>4</sup>

## Norepinephrine and Dopamine Reuptake Inhibitor (NDRI)

Bupropion is an example of a norepinephrine and dopamine reuptake inhibitor. It therefore leads to increased levels of norepinephrine and dopamine. It is used to treat depressive disorders, seasonal affective disorder, attention deficit disorder and to help people stop smoking. It is also important to note that this medication does decrease seizure threshold.<sup>5</sup>

3. National Institute of Mental Health. (2016). *Mental health medications*. U.S. Department of Health & Human Services. <https://www.nimh.nih.gov/health/topics/mental-health-medications>
4. National Institute of Mental Health. (2016). *Mental health medications*. U.S. Department of Health & Human Services. <https://www.nimh.nih.gov/health/topics/mental-health-medications>
5. MedlinePlus [Internet]. (2022). *Bupropion*. <https://medlineplus.gov/druginfo/meds/a695033.html>

## Serotonin Antagonist and Reuptake Inhibitor

Trazodone is an example of a serotonin antagonist and reuptake inhibitor. It is an antidepressant but most commonly prescribed off-label for anxiety or as a hypnotic. Trazodone reduces levels of the neurotransmitters associated with arousal effects, such as serotonin, noradrenaline, dopamine, acetylcholine, and histamine. Low-dose trazodone use exerts a sedative effect for sleep, so is typically administered in the evening.<sup>6</sup>

## Tricyclic Antidepressants

Tricyclic antidepressants (TCAs) are older first-generation antidepressants that block the reuptake of serotonin and norepinephrine in the synapse, which leads to increased concentration of these neurotransmitters in the brain. They are now more commonly used to treat neuropathic pain and insomnia. An example of a TCA is amitriptyline.<sup>7</sup>

TCAs are often administered at bedtime due to sedating effects. Older adults are particularly sensitive to the anticholinergic side effects of tricyclic antidepressants (e.g., tachycardia, urinary retention, constipation, dry mouth, blurred vision, confusion, psychomotor slowing, sedation, and delirium). Elderly clients should be started on low doses of amitriptyline and observed closely because they are at increased risk for falls. Blockage of adrenergic receptors can cause cardiac conduction disturbances and hypotension.<sup>8</sup>

Death may occur from overdose with this class of drugs. Multiple drug

6. Shin, J. J., & Saadabadi, A. (2024). *Trazodone*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK470560/>

7. National Institutes of Health. (n.d.). *Tricyclic antidepressants*. DailyMed. <https://dailymed.nlm.nih.gov/dailymed/>

8. National Institutes of Health. (n.d.). *Tricyclic antidepressants*. DailyMed. <https://dailymed.nlm.nih.gov/dailymed/>

ingestion (including alcohol) is common in deliberate tricyclic antidepressant overdose. If overdose occurs, call 911 in an outpatient setting or rapid response in an inpatient setting. Responders can consult with a Certified Poison Control Center (1-800-222-1222) or go to <sup>9</sup> <https://www.poisonhelp.org/help> for the latest treatment recommendations.

## Monoamine Oxidase Inhibitors (MAOI)

Monoamine oxidase inhibitors (MAOIs) are an older first-generation antidepressant. MAOIs are contraindicated with all other classes of antidepressants. Monoamine oxidase is an enzyme that removes the neurotransmitters norepinephrine, serotonin, and dopamine from the brain. By inhibiting this enzyme, MAOIs cause the levels of these transmitters to increase. Tranylcypromine is an example of an MAOI.<sup>10</sup>

A significant disadvantage to MAOIs is their potential to cause a hypertensive crisis when taken with stimulant medications or foods or beverages containing tyramine. Examples of foods containing tyramine are aged cheese, cured or smoked meats, alcoholic beverages, and soy sauce. Older adults are at increased risk for postural hypotension and serious adverse effects.<sup>11</sup>

Classes of antidepressant medications and their mechanisms of action are outlined in Table 6.3a.

Table 6.3. Antidepressants

9. National Institutes of Health. (n.d.). *Tricyclic antidepressants*. DailyMed. <https://dailymed.nlm.nih.gov/dailymed/>

10. National Institutes of Health. (n.d.). *Monoamine oxidase inhibitors*. DailyMed. <https://dailymed.nlm.nih.gov/dailymed/>

11. National Institutes of Health. (n.d.). *Monoamine oxidase inhibitors*. DailyMed. <https://dailymed.nlm.nih.gov/dailymed/>

Medication Class	Mechanism of Action
<b>Serotonin-Norepinephrine Reuptake Inhibitors (SNRIs)</b>  Common examples:  Venlafaxine  Duloxetine	Block the uptake of both serotonin and norepinephrine. SNRIs but with two neurotransmitters.
<b>Selective Serotonin Reuptake Inhibitors (SSRIs)</b>  Common examples:  Fluoxetine  Sertraline  Citalopram	Impact the receptors of the cell synapse. The neurotransmitter serotonin stays in the synapse.
<b>Tricyclic Antidepressants (TCAs)</b>  Common examples:  Amitriptyline  Nortriptyline	Block the presynaptic receptor for norepinephrine. The neurotransmitter norepinephrine levels are increased.
<b>Monoamine Oxidase Inhibitors (MAOIs)</b>  Common examples:  Phenelzine  Tranylcypromine	Block the enzyme that breaks down neurotransmitters serotonin and norepinephrine.
<b>Norepinephrine and Dopamine Reuptake Inhibitor (NDRI)</b>  Example: Bupropion	Block the uptake of both norepinephrine and dopamine.
<b>Serotonin Antagonist and Reuptake Inhibitor</b>  Example: Trazodone	Reduces levels of the neurotransmitters serotonin, noradrenaline, dopamine, acetylcholine.

Antidepressants commonly take four to eight weeks for noticeable effects on mood symptoms. Clients should also be counseled that if they do not feel better with the first antidepressant prescribed, the provider may need to try several different classes of medications to find one that works best for them. If a person's symptoms do not improve after trying at least two antidepressants, esketamine may be prescribed for treatment-resistant depression.

Esketamine is delivered as a nasal spray in a health care provider's office, clinic, or hospital and acts rapidly within a few hours, to relieve depression symptoms. People usually continue to take an oral antidepressant(s) to manage their symptoms.<sup>12</sup>

Clients should be instructed to never suddenly stop taking antidepressant medications or they may experience withdrawal symptoms.

## Withdrawal Symptoms

To safely stop or change antidepressants, clients must have prescribed dosage reductions with 2-6 weeks between dose reductions. Withdrawal syndrome may occur if antidepressants are stopped abruptly due to the rapid changes in levels of neurotransmitters in the brain. Withdrawal symptoms can include the following<sup>13</sup>:

- Flu-like symptoms, such as fatigue, headache, muscle aches, and sweating
- Fatigue
- Flushing
- Heart racing

12. National Institute of Mental Health. (2023). *Mental health medications*. U.S. Department of Health & Human Services. <https://www.nimh.nih.gov/health/topics/mental-health-medications>

13. Miller, K. & King, L. (2024). *Antidepressant withdrawal*. <https://www.webmd.com/depression/withdrawal-from-antidepressants>



- Trouble sleeping
- Vivid dreams or nightmares
- Nausea, diarrhea, and possibly vomiting
- Loss of appetite
- Dizziness, lightheadedness, or feeling unsteady on your feet
- Burning, tingling, or shock-like sensations
- Mood changes, such as anxiety, irritability, agitation, and aggression
- Brief shock-like feelings in the brain, short periods of blacking out, or a shock-like feeling with a buzzing sound

## Side Effects

Nurses monitor clients receiving antidepressants for side effects and report concerns to the prescribing provider. Common side effects of SSRIs and SNRIs are as follows:

- Nausea
- Diarrhea
- Weight gain
- Insomnia
- Feeling agitated, shaky or anxious
- Sexual dysfunction (reduced sex drive, difficulties achieving orgasm, or difficulties obtaining or maintaining an erection)

Side effects generally improve within a few weeks, although some can occasionally persist and require tapering or switching to a different class of antidepressant.

## Black Box Warning

A **Black Box Warning** is a significant warning from the Food and Drug Administration (FDA) that alerts the public and health care providers to serious side effects, such as injury or death. Black Box Warnings are in place for all classes of antidepressants used with children, adolescents, and young adults under age 25 due to a higher risk of suicide. All clients receiving

antidepressants should be monitored for signs of worsening depression or changing behavior, especially when the medication is started or dosages changed. Clients should be instructed to immediately call their provider if they have any of the following symptoms:

- Thoughts about suicide or dying or attempts to commit suicide
- Worsening symptoms of depression
- Anxiety or feelings of mania
- Aggression, anger, or violence

## Serotonin Syndrome

**Serotonin Syndrome** is a potentially life-threatening condition resulting from excessive serotonergic activity in the central nervous system, often triggered by the use or interaction of certain psychiatric medications. It most commonly occurs when two or more drugs that increase serotonin levels—such as selective serotonin reuptake inhibitors (SSRIs), serotonin-norepinephrine reuptake inhibitors (SNRIs), monoamine oxidase inhibitors (MAOIs), or certain over-the-counter supplements—are combined. Symptoms typically include altered mental status (e.g., agitation, confusion), autonomic instability (e.g., hyperthermia, tachycardia), and neuromuscular abnormalities (e.g., tremors, hyperreflexia, clonus). Nurses play an important role in early recognition and prompt intervention, including immediate discontinuation of serotonergic agents and supportive care.

A mnemonic commonly used to remember the symptoms of serotonin syndrome is **SHIVERS**:

- **S: Shivering:** A neuromuscular symptom similar to tremors specific to serotonin syndrome
- **H: Hyperreflexia (and myoclonus):** Hyperactive reflexes most prominent in the lower extremities.
- **I: Increased Temperature**
- **V: Vital Sign Abnormalities:** Tachycardia, tachypnea, and labile blood pressure

- **E: Encephalopathy:** Mental status changes such as agitation, delirium, and confusion
- **R: Restlessness**
- **S: Sweating**

People may get slowly worse and can become severely ill if not quickly treated. Untreated, serotonin syndrome can be deadly. With treatment, symptoms usually go away within 24 hours, but permanent kidney damage may result even with treatment. Uncontrolled muscle spasms can cause severe muscle breakdown called **rhabdomyolysis**. Myoglobin is released into the blood with muscle breakdown and clogs renal tubules, which can cause severe kidney damage if serotonin syndrome isn't recognized promptly and treated.

Treatment of serotonin syndrome may include the following<sup>14, 15</sup>:

- Stopping all serotonergic medications.
- Providing supportive care to normalize vital signs such as IV fluids, cooling measures, and medications to control heart rate and blood pressure.
- Sedating with benzodiazepines, such as diazepam or lorazepam, to decrease agitation, seizure-like movements, and muscle stiffness.
- If symptoms persist, administering cyproheptadine to block serotonin production.
- For clients with severe symptoms such as hyperthermia and muscle rigidity, more aggressive measures are required. These include sedation, intubation, neuromuscular paralysis, and active cooling techniques.<sup>16</sup>

<sup>14</sup>. Boyer, E. W. (2021). Serotonin syndrome (serotonin toxicity).

UpToDate. <https://www.uptodate.com/>

<sup>15</sup>. Tanen, D. (2021). *Serotonin syndrome*. Merck Manual Professional Version.

<https://www.merckmanuals.com/professional/injuries-poisoning/heat-illness/serotonin-syndrome>

Clients with moderate to severe serotonin syndrome should be hospitalized for close monitoring and management. Serotonin syndrome, in its most severe form, can resemble neuroleptic malignant syndrome (NMS) caused by antipsychotic medications. However, NMS develops over a period of days to weeks. Neuroleptic malignant syndrome (NMS) will be discussed further in [Chapter 11](#).

## Hypertensive Crisis

Hypertensive crisis can occur when clients taking monoamine oxidase inhibitors (MAOIs) also take medications containing pseudoephedrine or eat foods containing tyramine (aged foods; fermented foods; cured meats; alcoholic beverages such as beer or red wine; or overripe fruits such as raisins, prunes, or bananas). MAOIs inhibit the breakdown of tyramine, causing elevated tyramine levels in the body that can lead to hypertensive crisis.

**Hypertensive crisis** is a medical emergency defined as severe hypertension (blood pressure over 180/120 mm Hg) with acute end-organ damage such as stroke, myocardial infarction, or acute kidney damage. Symptoms may include a severe headache accompanied with confusion and blurred vision. Tachycardia or bradycardia may be present and associated with constricting chest pain. Other symptoms include neck stiffness or soreness, nausea or vomiting, sweating, dilated pupils, photophobia, shortness of breath, severe anxiety, and unresponsiveness. Seizures may occur, as well as intracranial bleeding in association with the increased blood pressure. Hypertensive crisis treatment involves discontinuation of the offending agent, administration of appropriate intravenous antihypertensive medications such as phentolamine

16. Spadaro, A., Scott, K. R., Koyfman, A., & Long, B. (2022). High risk and low prevalence diseases: Serotonin syndrome. *The American Journal of Emergency Medicine*, 61, 90-97. [doi: 10.1016/j.ajem.2022.08.030](https://doi.org/10.1016/j.ajem.2022.08.030).

or labetalol, and supportive care.. In severe cases, treatment in the intensive care unit may be required.<sup>17</sup><sup>[/footnote]</sup>,<sup>18</sup>

## Client Education

Clients should be instructed it may take 4 to 8 weeks for antidepressants to achieve their full effectiveness. They should not suddenly stop taking antidepressants or they may experience withdrawal symptoms. When it is time to stop the medication, the provider will slowly and safely decrease the dose. If clients stop taking the medication before the provider advises, the depression may return. They may not feel better with the first antidepressant they try, and they may need to try several different classes of medications to find one that works best for them. Education related to potential side effects and when to contact the provider, clinical worsening, avoiding alcohol, interference with cognitive or motor functioning, and potential drug interactions should also be provided.

- ▶ Review additional information in the “[Antidepressants](#)” section of the “Central Nervous System” chapter of *Open RN Nursing Pharmacology, 2e*.

17. National Institutes of Health. (n.d.). *Monoamine oxidase inhibitors*. DailyMed. <https://dailymed.nlm.nih.gov/dailymed/>

18. Sheps, S. G. (2021). *Hypertensive crisis: What are the symptoms?* Mayo Clinic. <https://www.mayoclinic.org/diseases-conditions/high-blood-pressure/expert-answers/hypertensive-crisis/faq-20058491#:~:text=A%20hypertensive%20crisis%20is%20a,higher%20%E2%80%94%20can%20damage%20blood%20vessels.>

## 6.4 Mood Stabilizer - Lithium

Mood stabilizers are used primarily to treat bipolar disorder. They are also used to treat depression (usually in combination with an antidepressant), schizoaffective disorder, and disorders of impulse control. Lithium is an example of a medication historically used as a mood stabilizer. Other medications prescribed for mood stabilizers include anticonvulsants, antipsychotic, antianxiety, and antidepressant medications.<sup>1</sup> Read more about medications used to treat bipolar disorder in the “[Treatments for Bipolar Disorder](#)” section of the “Bipolar Disorders” chapter.

### Lithium

The most commonly prescribed mood stabilizer is lithium. Lithium is primarily used to treat mania in bipolar disorder. Lithium reduces excitatory neurotransmission (dopamine and glutamate) and increases inhibitory neurotransmission (GABA). It also alters sodium transport in nerve and muscle cells and causes a shift in metabolism of catecholamines. When administered to a client experiencing a manic episode, lithium may reduce symptoms within 1 to 3 weeks. It also possesses unique antisuicidal properties that sets it apart from antidepressants. However, lithium toxicity can occur at doses close to therapeutic levels so lithium levels must be monitored regularly.<sup>2,3</sup>

1. National Institute of Mental Health. (2016). *Mental health medications*. U.S. Department of Health & Human Services. <https://www.nimh.nih.gov/health/topics/mental-health-medications>
2. This work is a derivative of [DailyMed](#) by [U.S. National Library of Medicine](#) and is available in the [Public Domain](#)
3. Malhi, G. S., Tanious, M., Das, P., Coulston, C. M., & Berk, M. (2013). Potential mechanisms of action of lithium in bipolar disorder. Current understanding. *CNS Drugs*, 27(2), 135–153. <https://doi.org/10.1007/s40263-013-0039-0>

## Side Effects

Lithium toxicity can occur at doses close to therapeutic levels, so lithium levels must be routinely monitored regularly. Signs of lithium toxicity must be promptly reported to the health care provider for dosage adjustment and treatment. Lithium blocks ADH, so symptoms of diabetes insipidus (i.e., excessive thirst and urination) should be monitored and promptly reported. Lithium's mechanism of action, nursing considerations, and side effects are summarized in Table 6.4.

Table 6.4 Lithium

Medication Class	Nursing Considerations	Common Side Effects (*Indicates more common)
Lithium	<ul style="list-style-type: none"> <li>Used as a first-line mood stabilizing agent to treat mania when symptoms are acute or as maintenance therapy</li> <li>Improved tolerance with food and better drug absorption</li> <li>Recommended water intake is 1.5 – 3 liters/day</li> <li>Given in divided doses if gastrointestinal distress occurs</li> <li>NSAIDs are not recommended because they increase lithium levels</li> <li>Therapeutic blood levels are required. Blood levels are drawn 10-12 hours after the last dose taken. The therapeutic lithium serum level is 0.6-1.2 mEq/L</li> <li>Treatments for toxicity: <ul style="list-style-type: none"> <li>Notify the health care provider regarding elevated lithium levels</li> <li>Withhold the lithium</li> <li>Encourage fluids; IV fluids may be required</li> <li>Gastric lavage</li> <li>May require urea, mannitol, aminophylline, or dialysis to hasten the excretion of the drug in severe cases</li> </ul> </li> </ul>	<p>Lithium blocks ADH (causing excessive thirst and urination)</p> <ul style="list-style-type: none"> <li>Long-term use may lead to kidney function impairment</li> </ul> <p><b>*Lithium toxicity</b></p> <ul style="list-style-type: none"> <li>Early signs: tremor, muscle weakness, ataxia</li> <li>Moderate signs: nausea, vomiting, diarrhea, complaints of blurred vision</li> <li>Severe signs: hypotension, seizures, death secondary to cardiac arrest</li> </ul> <p><b>*Lithium levels: asymptomatic</b></p>

## Client Education

Nurses teach clients that lithium must be taken as prescribed or serious side effects can occur. They reinforce the importance of adhering to regular blood tests to measure lithium levels and reporting symptoms of elevated levels of lithium, including diarrhea, vomiting, drowsiness, muscular weakness, lack of coordination, ringing in the ears (tinnitus), or large amounts of dilute urine. Driving or operating heavy machinery should be avoided when first starting lithium because it can impair mental alertness. Lithium should not be taken during pregnancy or while breastfeeding unless it is determined that the benefits to the mother outweigh the potential risks to the baby.

- ▶ Read additional information about lithium in the “[Antimaniacs](#)” section of the “Central Nervous System” chapter of *Open RN Nursing Pharmacology, 2e*.



## 6.5 Antianxiety Medications

Antianxiety medications help reduce the symptoms of anxiety, panic attacks, or extreme fear and worry. The most common class of antianxiety medications is benzodiazepines. Benzodiazepines are used to treat generalized anxiety disorder, although SSRIs or other antidepressants are typically used to treat panic disorder or social phobia (i.e., social anxiety disorder). Beta-blockers and buspirone may also be prescribed for anxiety.<sup>1</sup>

### Benzodiazepines

Benzodiazepines are used to treat anxiety and are also used for their sedation and anticonvulsant effects because they bind to GABA receptors and stimulate the effects of GABA (an inhibitory neurotransmitter).

Benzodiazepines include clonazepam, alprazolam, and lorazepam.

Benzodiazepines are a Schedule IV controlled substance because they have a potential for misuse and can cause dependence. Short-acting benzodiazepines (such as lorazepam) and beta-blockers are used to treat the short-term symptoms of anxiety. Lorazepam is available for oral, intramuscular, or intravenous routes of administration.<sup>2</sup>[/footnote]

If people suddenly stop taking benzodiazepines after taking them for a long period of time, they may have withdrawal symptoms, or their anxiety may return. Withdrawal symptoms include sleep disturbances, irritability, increased tension and anxiety, hand tremors, sweating, difficulty concentrating, nausea and vomiting, weight loss, palpitations, headache,

1. National Institute of Mental Health. (2016). *Mental health medications*. U.S. Department of Health & Human Services. <https://www.nimh.nih.gov/health/topics/mental-health-medications>

2. National Institutes of Health. (n.d.). *Benzodiazepines*. DailyMed. <https://dailymed.nlm.nih.gov/dailymed/>

muscular pain, and perceptual changes.<sup>3</sup> Therefore, benzodiazepines should be tapered off slowly.<sup>4</sup>

## Overdosage of Benzodiazepines

Overdosage of benzodiazepines is manifested by varying degrees of central nervous system depression, ranging from drowsiness to coma. If overdose occurs, call 911 or the rapid response team during inpatient care. Treatment of overdosage is mainly supportive until the drug is eliminated from the body. Vital signs and fluid balance should be carefully monitored in conjunction with close observation of the client. An adequate airway should be maintained; intubation and mechanical ventilation may be required. The benzodiazepine antagonist flumazenil may be used to manage benzodiazepine overdose. There is a risk of seizure in association with flumazenil treatment, particularly in chronic users of benzodiazepines.

## Side Effects

Overdosage of benzodiazepines causes central nervous system depression, ranging from drowsiness to coma. Children and older adults are more susceptible to the sedative and respiratory depressive effects of lorazepam and may experience paradoxical reactions such as tremors, agitation, or visual hallucinations. Benzodiazepines may cause fetal harm when administered to pregnant women. There is a boxed warning that concomitant use of benzodiazepines and opioids may result in profound sedation, respiratory

3. Pétursson, H. (1994). The benzodiazepine withdrawal syndrome. *Addiction*, 89(11), 1455-9. [doi:10.1111/j.1360-0443.1994.tb03743.x](https://doi.org/10.1111/j.1360-0443.1994.tb03743.x).
4. National Institute of Mental Health. (2018). *Anxiety disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/anxiety-disorders>

depression, coma, and death. See Table 6.5 for summarized information about benzodiazepines.

Table 6.5. Benzodiazepines

Generic	Nursing Considerations	Side/Adverse Effects
<ul style="list-style-type: none"><li>• <b>Lorazepam</b></li><li>• <b>Alprazolam</b></li><li>• <b>Clonazepam</b></li></ul>	May cause fetal harm in pregnant women and may cause paradoxical effect in children	Boxed Warning: Concomitant use of benzodiazepines and opioids may result in profound sedation, respiratory depression, coma, and death  Increased risk for falls

**BLACK BOX WARNING**

A Black Box Warning states that concurrent use of benzodiazepines and opioids may result in profound sedation, respiratory depression, coma, and death. The use of benzodiazepines exposes users to risks of misuse, substance use disorder, and addiction. Misuse of benzodiazepines commonly involves concomitant use of other medications, alcohol, and/or illicit substances, which is associated with an increased frequency of serious adverse outcomes. Additionally, the continued use of benzodiazepines may lead to clinically significant physical dependence. The risks of dependence and withdrawal increase with longer treatment duration and higher daily doses, and abrupt discontinuation or rapid dosage reduction may precipitate life-threatening withdrawal reactions. To reduce the risk of withdrawal reactions, a gradual taper should be used to stop or reduce the dosage.<sup>5</sup>

Client Education

Clients should be cautioned that driving a motor vehicle, operating machinery, or engaging in hazardous or other activities requiring attention

5. National Institutes of Health. (n.d.). *Benzodiazepines*. DailyMed.  
<https://dailymed.nlm.nih.gov/dailymed/>

and coordination should be delayed for 24 to 48 hours following administration of benzodiazepines or until the effects of the drug, such as drowsiness, have subsided. Alcoholic beverages should not be consumed for at least 24 to 48 hours after receiving lorazepam due to the additive effects on central nervous system depression. Hospitalized clients should be advised that benzodiazepines increase fall risk, and getting out of bed unassisted may result in falling and potential injury.

Read more information about benzodiazepines in the “[CNS Depressants](#)” chapter of *Open RN Nursing Pharmacology, 2e*.

## Beta-Blockers

Beta-blockers, such as propranolol, are medications that block the effects of the sympathetic nervous system by acting on Beta-1 receptors in the heart and other areas of the body. While they are most commonly prescribed to treat high blood pressure, heart rhythm disorders, and other cardiac conditions, beta-blockers may also be used off-label to help manage the physical symptoms of anxiety—such as trembling, rapid heartbeat, and sweating—especially in short-term or situational anxiety, such as public speaking or test anxiety. In these cases, they are often prescribed “as needed” rather than for continuous daily use. They may be prescribed to manage the physical symptoms of anxiety (such as trembling, rapid heartbeat, and sweating) for a short period of time or used “as needed” to reduce acute physical symptoms.<sup>6</sup>

Common side effects of beta-blockers are fatigue, hypotension, dizziness,

6. National Institute of Mental Health. (2016). *Mental health medications*. U.S. Department of Health & Human Services. <https://www.nimh.nih.gov/health/topics/mental-health-medications>

weakness, and cold hands. Beta-blockers are typically avoided in clients with asthma or diabetes.<sup>7</sup>

Read more information about propranolol in the “[Beta-2 Antagonists](#)” section of the “Autonomic Nervous System” chapter of *Open RN Nursing Pharmacology, 2e*.

## Buspirone

Buspirone is a non-benzodiazepine medication indicated for the treatment of chronic anxiety. It is included in the class of medications called anxiolytics, but it is not chemically related to benzodiazepines, barbiturates, or other sedatives. Buspirone should not be taken concurrently with a monoamine oxidase inhibitor (MAOI) due to the risk of fatal side effects. It can also cause serotonin syndrome if used in combination with MAOIs, SSRIs, or SNRIs.<sup>8</sup>[footnote]

Buspirone increases serotonin and dopamine levels in the brain. In contrast to benzodiazepines, buspirone must be taken every day for a few weeks to reach its full effect; it is not useful on an “as-needed” basis. A common side effect of buspirone is dizziness.<sup>9</sup> Buspirone is non-addictive and safer for long-term use than benzodiazepines.

7. National Institute of Mental Health. (2016). *Mental health medications*. U.S. Department of Health & Human Services. <https://www.nimh.nih.gov/health/topics/mental-health-medications>
8. National Institutes of Health. (n.d.). *Buspirone*. DailyMed. <https://dailymed.nlm.nih.gov/dailymed/>
9. National Institutes of Health. (n.d.). *Buspirone*. DailyMed. <https://dailymed.nlm.nih.gov/dailymed/>

# Hydroxyzine

Hydroxyzine is type of antihistamine which may be prescribed to alleviate anxiety for individuals for whom benzodiazepines are not appropriate. It causes sedation, so it must be used cautiously if used in combination with opioids or barbiturates. Hydroxyzine is recommended for short-term use in the treatment of anxiety and tension. The effectiveness of hydroxyzine for long-term use, defined as more than 4 months, has not been systematically assessed in clinical studies. Therefore, it is advised that physicians periodically reassess the usefulness of the drug for each individual client.<sup>10</sup>

- ▶ Read additional information about “[Treatments for Anxiety](#)” in the “Anxiety Disorders” chapter.

10. National Institutes of Health. (n.d.). *Hydroxyzine*. DailyMed. <https://dailymed.nlm.nih.gov/dailymed/>

## 6.6 Antipsychotics

Antipsychotic medicines are primarily used to treat psychosis (i.e., a loss of contact with reality that may include delusions or hallucinations). Psychosis can be a symptom of a physical condition (such as a high fever, head injury, or substance intoxication) or a mental disorder (such as schizophrenia, bipolar disorder, or severe depression).<sup>1</sup>

Antipsychotic medications reduce the intensity and frequency of psychotic symptoms by inhibiting dopamine receptors. Certain symptoms of psychosis, such as feeling agitated and having hallucinations, resolve within days of starting an antipsychotic medication. Symptoms like delusions usually resolve within a few weeks, but the full effects of the medication may not be seen for up to six weeks.<sup>2</sup>

### First-Generation (Typical) Antipsychotics

Common first-generation antipsychotic medications (also called “typical” antipsychotics) treat positive symptoms of schizophrenia. Examples include chlorpromazine, haloperidol, perphenazine, and fluphenazine.<sup>3</sup>

First-generation antipsychotics work by blocking dopamine receptors in certain areas of the CNS, such as the limbic system and the basal ganglia.

1. National Institute of Mental Health. (2023). *Mental health medications*. U.S. Department of Health & Human Services. <https://www.nimh.nih.gov/health/topics/mental-health-medications>
2. National Institute of Mental Health. (2023). *Mental health medications*. U.S. Department of Health and Human Services. [https://www.nimh.nih.gov/health/topics/mental-health-medications#part\\_2362](https://www.nimh.nih.gov/health/topics/mental-health-medications#part_2362)
3. National Institute of Mental Health. (2023). *Mental health medications*. U.S. Department of Health and Human Services. [https://www.nimh.nih.gov/health/topics/mental-health-medications#part\\_2362](https://www.nimh.nih.gov/health/topics/mental-health-medications#part_2362)

These areas are associated with emotions, cognitive function, and motor function. As a result, blockage produces a tranquilizing effect in psychotic clients. However, several adverse effects are caused by this dopamine blockade, such **extrapyramidal side effects** (e.g., involuntary or uncontrollable movements, tremors, and muscle contractions) and **tardive dyskinesia** (a syndrome of movement disorders that persists for at least one month and can last up to several years despite discontinuation of the medications).<sup>4</sup>

## Second-Generation (Atypical) Antipsychotics

Newer, second-generation antipsychotics (also called atypical antipsychotics) work by blocking specific D2 dopamine receptors and serotonin receptors. Examples of second-generation medications include risperidone, olanzapine, quetiapine, ziprasidone, aripiprazole, paliperidone, and lurasidone. Second generation antipsychotics treat both positive and negative symptoms of schizophrenia, and several are also used for treating bipolar depression or depression that has not responded to an antidepressant medication alone. Second generation antipsychotics have a significantly decreased risk of extrapyramidal side effects but are associated with weight gain and the development of metabolic syndrome.<sup>5</sup> **Metabolic syndrome** increases the risk of heart disease, stroke, and type 2 diabetes. Clinical symptoms of metabolic syndrome include high blood glucose, symptoms of diabetes (i.e., increased thirst and urination, fatigue, and blurred vision), obesity with a large abdominal girth, hypertension, elevated triglyceride, and lower levels of HDL.

See Table 6.6 for a list of common antipsychotic medications. Some are taken daily in pill or liquid form. Others can be administered as injections twice a

4. National Institute of Mental Health. (2023). *Mental health medications*. U.S. Department of Health and Human Services. [https://www.nimh.nih.gov/health/topics/mental-health-medications#part\\_2362](https://www.nimh.nih.gov/health/topics/mental-health-medications#part_2362)

5. Chokhawala, K., & Stevens, L. (2023). *Antipsychotic medications*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK519503/>



month, monthly, every three months, or every six months, which can be more convenient and improve medication adherence.

Table 6.6 Common Antipsychotic Medications<sup>6,7,8</sup>

6. National Institute of Mental Health. (2023). *Mental health medications*. U.S. Department of Health and Human Services. [https://www.nimh.nih.gov/health/topics/mental-health-medications#part\\_2362](https://www.nimh.nih.gov/health/topics/mental-health-medications#part_2362)
7. Vasan, S., & Padhy, R. K. (2023). *Tardive dyskinesia*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK448207/>
8. Jibson, M. D. (2021). Second-generation antipsychotic medications: Pharmacology, administration, and side effects. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

Medication Class	Mechanism of Action	Adverse Effects
<b>First-Generation (Typical)</b>  Examples:  Chlorpromazine  Haloperidol  Perphenazine  Fluphenazine	Postsynaptic blockade of dopamine receptors in the brain	<ul style="list-style-type: none"> <li>• Extrapyrarnidal side effects (EPS)</li> <li>• Tardive dyskinesia (TD)</li> <li>• Neuroleptic Malignant Syndrome (NMS)</li> </ul>
<b>Second-Generation (Atypical)</b>  Examples:  Risperidone  Olanzapine  Quetiapine  Ziprasidone  Aripiprazole  Paliperidone  Lurasidone  Clozapine	Postsynaptic blockade of dopamine receptors in the brain	<ul style="list-style-type: none"> <li>• Metabolic syndrome</li> <li>• Akathisia</li> <li>• Decreased risk for EPS, TD, and NMS</li> </ul>

Clients respond differently to antipsychotic medications, so it may take several trials of different medications to find the one that works best for their symptoms.<sup>9</sup>

9. National Institute of Mental Health. (2023). *Mental health medications*. U.S. Department of Health and Human Services. [https://www.nimh.nih.gov/health/topics/mental-health-medications#part\\_2362](https://www.nimh.nih.gov/health/topics/mental-health-medications#part_2362)

# Clozapine

Clients with treatment-resistant schizophrenia may be prescribed clozapine, a specific atypical antipsychotic medication that binds to both serotonin and dopamine receptors. Clozapine is often effective when other antipsychotics have failed, but it carries a complex side effect profile, including strong anticholinergic, sedative, cardiac, and hypotensive properties, as well as frequent drug-drug interactions. One of the most important risks to note is **agranulocytosis**, a rare but potentially life-threatening drop in white blood cells that significantly increases infection risk. For this reason, clients must be enrolled in a REMS (Risk Evaluation and Mitigation Strategy) program, with regular monitoring of their absolute neutrophil count (ANC)—typically weekly during the initial months of treatment. Nurses play a critical role in educating clients about the importance of lab work and recognizing early signs of infection such as fever or sore throat, which must be reported immediately.<sup>10</sup>

## Side Effects

Common side effects of both first- and second-generation antipsychotics include the following:

- Anticholinergic symptoms: dry mouth, constipation, blurred vision, or urinary retention
- Drowsiness
- Dizziness
- Restlessness
- Weight gain
- Nausea or vomiting
- Low blood pressure
- Falls related to sedation, motor instability, and postural hypotension

<sup>10</sup>. Jibson, M. D. (2021). Second-generation antipsychotic medications: Pharmacology, administration, and side effects. *UpToDate*. <https://www.uptodate.com/>

**Neuroleptic malignant syndrome (NMS)** is a rare but fatal adverse effect that can occur at any time during treatment with antipsychotics. It typically develops over a period of days to weeks and resolves in approximately nine days with treatment. Signs include increased temperature, severe muscular rigidity, confusion, agitation, hyperreflexia, elevation in white blood cell count, elevated creatinine phosphokinase, elevated liver enzymes, myoglobinuria, and acute renal failure. The antipsychotic should be immediately discontinued when signs occur. Dantrolene and bromocriptine are typically prescribed for treatment. Nursing interventions include adequate hydration, cooling, and close monitoring of vital signs and serum electrolytes.<sup>11</sup>

## Black Box Warning

A Black Box Warning states that elderly clients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death.<sup>12</sup>

## Side Effects of First-Generation Antipsychotics

First-generation antipsychotics have significant potential to cause extrapyramidal side effects and tardive dyskinesia due to their tight binding to dopamine receptors.<sup>13</sup>

Extrapyramidal (EPS) side effects refer to **akathisia** (psychomotor

11. Chokhawala, K., & Stevens, L. (2023). *Antipsychotic medications*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK519503/>

12. National Institutes of Health. (2019). *Antipsychotic drugs*. DailyMed. <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=165d01d4-a9f7-2293-e054-00144ff8d46c>

13. Jibson, M. D. (2021). Second-generation antipsychotic medications: Pharmacology, administration, and side effects. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)


restlessness), rigidity, **bradykinesia** (slowed movement), tremor, and **dystonia** (involuntary contractions of muscles of the extremities, face, neck, abdomen, pelvis, or larynx in either sustained or intermittent patterns that lead to abnormal movements or postures). Acute dystonic reactions affecting the larynx can be a medical emergency requiring intubation and mechanical ventilation. EPS symptoms usually resolve dramatically within 10 to 30 minutes of administration of parenteral anticholinergics such as diphenhydramine and benztropine.<sup>14</sup>

Tardive dyskinesia (TD) is a syndrome of movement disorders that can occur in clients taking first-generation antipsychotics. Hallmark symptoms are smacking and puckering lips, eye blinking, grimacing, and twitching. TD persists for at least one month and can last up to several years despite discontinuation of the medications. Primary treatment of TD includes discontinuation of first-generation antipsychotics and may include the addition of another medication. Second-generation VMAT2 inhibitors such as deutetrabenazine and valbenazine are considered first-line treatment for TD. Clonazepam and ginkgo biloba have also shown good effectiveness for improving symptoms of TD.<sup>15,16</sup>

14. Lewis, K., O'Day, C. S. (2025). *Dystonic reactions*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK531466/>
15. Vasan, S., & Padhy, R. K. (2023). *Tardive dyskinesia*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK448207/>
16. Pontone, G. (2020). Treating tardive dyskinesia: A clinical conundrum and new approaches. *Drug Induced Disorders: The Clinical Essentials*. Psychopharmacology Institute. <https://psychopharmacologyinstitute.com/section/treating-tardive-dyskinesia-a-clinical-conundrum-and-new-approaches-2557-4810#:~:text=Clonazepam%20probably%20improves%20tard>  
iv

## Side Effects of Second-Generation Antipsychotics

Second-generation antipsychotics have a significantly decreased risk of extrapyramidal side effects but are associated with weight gain and the development of metabolic syndrome.<sup>17</sup> Metabolic syndrome is a cluster of conditions that occur together, increasing the risk of heart disease, stroke, and type 2 diabetes. Symptoms include increased blood pressure; high blood sugar; excess body fat around the waist (also referred to as having an “apple waistline”); and abnormal cholesterol, triglyceride levels, and high-density lipoprotein (HDL) levels. Weight, glucose levels, and lipid levels should be monitored before treatment is initiated then annually.

 View a YouTube video<sup>18</sup> on metabolic syndrome: [What is Metabolic Syndrome?](#)

▶ View this PDF: [Adverse Effects of Antipsychotic Medications.](#)

## Client Education

Clients should be advised to contact their provider if and side effects occur. This includes the development of any involuntary or uncontrollable movements. They should be warned to not suddenly stop taking the

17. Chokhawala, K., & Stevens, L. (2023). *Second-generation antipsychotics*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK519503/>

18. Health Link. (2019, December 31). What is metabolic Syndrome? [Video]. YouTube. All rights reserved. [https://youtu.be/fVMvY\\_Lsqzw](https://youtu.be/fVMvY_Lsqzw)

medication because abrupt withdrawal can cause dizziness; nausea and vomiting; and uncontrolled movements of the mouth, tongue, or jaw. Clients should be warned to not consume alcohol or other CNS depressants because their ability to operate machinery or drive may be impaired.

Some people may experience relapse, meaning their psychosis symptoms come back or get worse. Relapses typically occur when people stop taking their prescribed antipsychotic medication or when they take it sporadically. Some people stop taking prescribed medications because they feel better or they feel that they don't need it anymore, but medication should never be stopped suddenly. After talking with a prescriber, clients can gradually taper their medications in some situations. However, most people with schizophrenia must stay on an antipsychotic continuously for mental wellness.<sup>19</sup>



View a supplementary YouTube video<sup>20</sup> explaining how clozapine binds to additional neuroreceptors compared to other antipsychotic medications: [The Pines, the Dones, Two Pips and a Rip](https://youtu.be/kuYGCJOcloH8)

19. National Institute of Mental Health. (2023). *Mental health medications*. U.S. Department of Health and Human Services. [https://www.nimh.nih.gov/health/topics/mental-health-medications#part\\_2362](https://www.nimh.nih.gov/health/topics/mental-health-medications#part_2362)

20. NEI Psychopharm. (2014, March 18). *The pines, the dones, two pips, and a rip* [Video]. YouTube. All rights reserved. <https://youtu.be/kuYGCJOcloH8>



View a supplementary YouTube video<sup>21</sup> exploring antipsychotics: [Pharmacology-Antipsychotics](https://youtu.be/nKklh1B2Js8?si=4s1ecekHq9v3UDWA).

- ▶ Read additional information about the mechanism of action, adverse side effects, and client education regarding antipsychotic medications in the “[Schizophrenia](#)” section of the “Psychosis and Schizophrenia” chapter.

21. Speed Pharmacology. (2018, July 18). *Pharmacology-Antipsychotics (Made Easy)*[Video]. YouTube. All rights reserved. <https://youtu.be/nKklh1B2Js8?si=4s1ecekHq9v3UDWA>



## 6.7 Stimulants

Stimulant medications are prescribed to treat children, adolescents, or adults diagnosed with attention deficit hyperactivity disorder (ADHD). Stimulants block the reuptake of norepinephrine and dopamine in the synapse and increase the overall level of these substances in the brain, but they have a paradoxical calming effect and improve the ability to focus and concentrate for individuals diagnosed with ADHD. Common stimulants used to treat ADHD include methylphenidate, amphetamine, dextroamphetamine, and lisdexamfetamine dimesylate. Stimulant medications are safe when prescribed with close supervision, but they are a Schedule II controlled substance because they have a high potential for misuse and dependence. Stimulants are available in short-, intermediate-, and long-acting formulations. See Table 6.7 for an overview of information about stimulants.

Table 6.7. Stimulants

Generic	Nursing Considerations	Side/Adverse Effects
<ul style="list-style-type: none"><li>• <b>Methylphenidate</b></li><li>• <b>Amphetamine</b></li><li>• <b>Dextroamphetamine</b></li></ul>	<ul style="list-style-type: none"><li>• High potential for misuse and drug diversion; promptly report concerns to the health care provider</li><li>• Clients should avoid alcohol</li><li>• Monitor BP and HR</li><li>• Monitor growth and weight in children</li></ul>	<ul style="list-style-type: none"><li>• Immediately report signs and symptoms of mania, psychosis, cardiac or peripheral vascular complications, and priapism</li><li>• Common side effects: headache, insomnia, upper abdominal pain, and decreased appetite</li></ul>

## Side Effects

Stimulants may cause minor side effects that resolve when dosage levels are

lowered, or a different stimulant is prescribed. The most common side effects include the following<sup>1,2</sup>:

- Difficulty falling asleep or staying asleep
- Loss of appetite and weight loss
- Stomach pain
- Headache

Less common side effects include motor or verbal tics (sudden, repetitive movements or sounds) or personality changes (such as appearing “flat” or without emotion).<sup>3</sup> Sudden death, stroke, and myocardial infarction have been reported in adults with CNS-stimulant treatment at recommended doses. Sudden death has been reported in pediatric clients with structural cardiac abnormalities and other serious heart problems taking CNS stimulants at recommended doses. If paradoxical worsening of symptoms or other adverse reactions occur, the provider should be contacted, and the dosage reduced or discontinued. Stimulants are contraindicated in clients using a monoamine oxidase inhibitor (MAOI) or using an MAOI within the preceding 14 days.<sup>4</sup>

1. National Institute of Mental Health. (2016). *Mental health medications*. U.S. Department of Health & Human Services. <https://www.nimh.nih.gov/health/topics/mental-health-medications>
2. National Institutes of Health. (n.d.). *Stimulants*. DailyMed. <https://dailymed.nlm.nih.gov/dailymed/index.cfm>
3. National Institute of Mental Health. (2016). *Mental health medications*. U.S. Department of Health & Human Services. <https://www.nimh.nih.gov/health/topics/mental-health-medications>
4. National Institutes of Health. (n.d.). *Monoamine oxidase inhibitor*. DailyMed. <https://dailymed.nlm.nih.gov/dailymed/index.cfm>

## Black Box Warning

CNS stimulants, including methylphenidate and amphetamine-like substances, have a high potential for abuse and dependence. The risk of abuse by the client or their family members should be assessed prior to prescribing stimulants, and signs of abuse and dependence should be evaluated while the client is receiving therapy.<sup>5</sup>

## Client Education

There are several important client education topics to provide to clients and/or the parents of minor children<sup>6</sup>:

- **Medication information:** Discussing the purpose, potential side effects, and the importance of adherence to prescribed treatments.
- **Controlled Substance Status/High Potential for Abuse and Dependence:** Stimulants are a controlled substance by the FDA that can be abused and lead to dependence. Stimulants should be stored in a safe (preferably locked) place to prevent misuse and should not be shared with anyone. Unused or expired stimulants should be disposed of based on state law and regulations or returned to a medicine take-back program if it is available in the community.
- **Cardiovascular Risks:** Stimulants can increase blood pressure and pulse rate. Potential serious cardiovascular risks include sudden death, cardiomyopathy, myocardial infarction, stroke, and hypertension. Instruct clients to contact a health care provider immediately if they develop symptoms, such as exertional chest pain, dizziness, or passing out.
- **Suppression of Growth:** Stimulants may cause slowing of growth in

5. National Institutes of Health. (n.d.). *CNS stimulants*. DailyMed.  
<https://dailymed.nlm.nih.gov/dailymed/index.cfm>

6. National Institutes of Health. (n.d.). *CNS stimulants*. DailyMed.  
<https://dailymed.nlm.nih.gov/dailymed/index.cfm>

children and weight loss.

- **Psychiatric Risks:** Stimulants can cause psychosis or manic symptoms, even in clients who have no prior history of these symptoms.
- **Priapism:** Painful or prolonged penile erections can occur; seek immediate medical attention.
- **Alcohol:** Alcohol should be avoided.

Nurses should reinforce with the client and their family members that the reason for the prescribed medication for ADHD is to help with self-control and the ability to focus. Possible side effects should be reviewed, and clients and their family members should be reminded it may take one to three months to determine the best pharmacological treatment, dose, and frequency of medication administration. During this time, the child's symptoms and adverse effects will be monitored weekly and the medication dose adjusted accordingly.<sup>7</sup>

- ▶ Read more information about other medications used to treat ADHD in the "[Common Disorders and Disabilities in Children and Adolescents](#)" section of the "Childhood and Adolescent Disorders" chapter.

7. Krull, K. R. (2022). Attention deficit hyperactivity disorder in children and adolescents: Treatment with medications. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

## 6.8 Psychoactive Substances and Medications to Treat Substance Use and Withdrawal

Information about the effects of substances such alcohol, cannabis, and illicit drugs is discussed in the “[Substances: Use, Intoxication, and Overdose](#)” section of the “Substance Use Disorders” chapter.

Medications to treat alcohol use disorder and opioid disorder include buprenorphine-naloxone, methadone, naltrexone, acamprosate, and disulfiram. These medications are further discussed in the “[Treatment and Recovery Services](#)” subsection of the “Substance Use Disorders” chapter.

See Table 6.8 for a list of medications commonly used to treat alcohol and opioid use disorders.

Table 6.8 Common Medications Used to Treat Alcohol and Opioid Use Disorders<sup>1</sup>

1. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

Medication	Use	DEA Schedule	Application
<b>Buprenorphine-naloxone</b>	Opioid use disorder	CIII	Used for detoxification or maintenance of abstinence.
<b>Methadone</b>	Opioid use disorder	CII	Used for withdrawal and long-term maintenance of abstinence of opioid addiction. Dispersed only at opioid treatment centers certified by SAMHSA and approved by state authority.
<b>Naltrexone</b>	Opioid use disorder and alcohol use disorder	Not scheduled under the Controlled Substances Act	Block opioid receptors, reduce cravings, and diminish rewarding effects of opioids and alcohol. Extended-release injections are recommended to prevent relapse.
<b>Acamprosate</b>	Alcohol use disorder	Not scheduled under the Controlled Substances Act	Used for maintenance of alcohol abstinence.
<b>Disulfiram</b>	Alcohol use disorder	Not scheduled under the Controlled Substances Act	Causes severe physical reactions when alcohol is ingested, such as nausea, flushing, and heart palpitations. The knowledge that the reaction will occur acts as a deterrent to drinking alcohol.

Medications used to manage symptoms of substance withdrawal/detoxification include buprenorphine, methadone, and Alpha-2 adrenergic agonists (such as clonidine and lofexidine). In the case of alcohol withdrawal, benzodiazepines (such as lorazepam or diazepam) remain the first-line treatment, particularly for preventing seizures and delirium tremens. In certain cases—especially in ICU-level care—dexmedetomidine (Precedex) may be used as an adjunct for sympathetic overactivity, though it does not treat seizures and must be used with close monitoring. Read more about these medications in the “[Withdrawal Management/Detoxification](#)” section of the “Substance Use Disorders” chapter.

## 6.9 Learning Activities



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=996#h5p-32>

1



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=996#h5p-33>

2



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=996#h5p-34>

1. “MH Psychotropic Medications Glossary Cards” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

2. “MH Psychotropic Medication Question Set 1” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

- ▶ Test your clinical judgment with a NCLEX Next Generation-style question: [Chapter 6, Assignment 1](#)<sup>4</sup>



- ▶ Test your clinical judgment with a NCLEX Next Generation-style case study: [Chapter 6, Case Study 1](#)<sup>5</sup>

3. "MH Psychotropic Medication Drag and Drop" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

4. "MH Psychotropic Medication Next Gen Question 1" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

5. "MH Psychotropic Medication Case Study" by Kellea Ewen is licensed under [CC BY-NC 4.0](#)



## VI Glossary

**Acetylcholine:** A neurotransmitter that stimulates nicotinic and muscarinic receptors in the parasympathetic nervous system.

**Adrenergic agonists:** Substances that stimulate SNS receptors and cause effects similar to epinephrine and norepinephrine.

**Adrenergic antagonists:** Substances that block SNS receptors.

**Agranulocytosis:** Extremely low white blood cell count and an adverse effect of clozapine and antipsychotic medication.

**Anticholinergics:** Substances that block the effects of PNS receptors.

**Black Box Warning:** A significant warning from the Food and Drug Administration (FDA) that alerts the public and health care providers to serious side effects, such as injury or death.

**Catecholamines:** Substances that include epinephrine, norepinephrine, and dopamine and are responsible for the body's "fight-or-flight" response.

**Central nervous system (CNS):** The brain and spinal cord.

**Cholinergics:** Substances that stimulate nicotinic and muscarinic receptors and cause effects similar to acetylcholine (ACh).

**Controlled substance:** Drugs regulated by federal law that can cause dependence and abuse.

**Dopamine:** A neurotransmitter that plays an essential role in several brain functions, including learning, motor control, reward, emotion, and executive functions.

**Extrapyramidal side effects:** Involuntary or uncontrollable movements, tremors, and muscle contractions that can occur with antipsychotic medications.

**Gamma-aminobutyric acid and Glycine:** Inhibitory neurotransmitters that

act like brakes in a car by slowing down overexcited nerve cells. Low levels of GABA are associated with seizures, anxiety, mania, and impulse control. Pregabalin is an anticonvulsant that mimics the effects of GABA and is used to treat generalized anxiety disorder.

**Glutamate:** An excitatory neurotransmitter. Elevated levels of glutamate are associated with psychosis that can occur with schizophrenia, as well as with illicit drug use such as methamphetamines. Conversely, lamotrigine, a medication used to treat bipolar disorder, inhibits glutamate.

**Histamine:** A substance that mediates homeostatic functions in the body, promotes wakefulness, modulates feeding behavior, and controls motivational behavior. For example, diphenhydramine, a histamine antagonist, causes drowsiness.

**Hypertensive crisis:** A condition that can be caused by MAOIs with severe hypertension (blood pressure greater than 180/120 mm Hg) and evidence of organ dysfunction. Symptoms may include occipital headache (which may radiate frontally), palpitations, neck stiffness or soreness, nausea or vomiting, sweating, dilated pupils, photophobia, shortness of breath, or confusion.

**Lithium toxicity:** Lithium has a narrow therapeutic range of 0.8 to 1.2 mEq/L. Levels above this range cause lithium toxicity. Signs of early lithium toxicity include diarrhea, vomiting, drowsiness, muscular weakness, and lack of coordination. At higher levels, giddiness, ataxia, blurred vision, tinnitus, and a large output of dilute urine may occur.

**Neuroleptic malignant syndrome (NMS):** A rare but fatal adverse effect that can occur at any time during treatment with antipsychotics. It typically develops over a period of days to weeks and resolves in approximately nine days with treatment. Signs include increased temperature, severe muscular rigidity, confusion, agitation, hyperreflexia, elevation in white blood cell count, elevated creatine phosphokinase, elevated liver enzymes, myoglobinuria, and acute renal failure.

**Neurotransmitters:** Chemical substances released at the end of a neuron by

the arrival of an electrical impulse. They diffuse across the synapse and cause the transfer of the impulse to another nerve fiber, a muscle fiber, or other structure. Neurotransmitters interact with specific receptors like a key and a lock.

**Norepinephrine and Epinephrine:** Substances that stimulate alpha- and beta-receptors in the sympathetic nervous system.

**Opiates:** Powerful analgesics prescribed to treat moderate to severe pain (such as morphine and oxycodone). Opiates also include illicit drugs (such as heroin).

**Opioid receptors:** Mu, delta, and kappa receptors that are stimulated by endogenous peptides released by neurons (such as endorphins) and exogenous opiates.

**Opioid system:** A system in the brain that controls pain and reward and addictive behaviors.

**Parasympathetic Nervous System (PNS) receptors:** Nicotinic and muscarinic receptors that are stimulated by acetylcholine (ACh).

**Serotonin:** A neurotransmitter that modulates multiple neuropsychological processes such as mood, sleep, libido, and temperature regulation. Abnormal levels of serotonin have been linked to many mental health disorders such as depression, bipolar disorder, and anxiety. Many psychotropic medications target serotonin.

**Serotonin syndrome:** A syndrome caused by the combination of multiple medications that affect serotonin. It typically develops within 24 hours from the combination of medication and can range from mild to a life-threatening syndrome. Signs of serotonin syndrome include mental status changes (e.g., agitation, hallucinations, coma), autonomic instability (e.g., tachycardia, labile blood pressure, hyperthermia), incoordination, or gastrointestinal symptoms (e.g., nausea, vomiting, diarrhea). Serotonin syndrome, in its most severe form, can resemble neuroleptic malignant syndrome (NMS).

**SLUDGE:** A mnemonic for anticholinergic side effects: Salivation decreased, Lacrimation decreased, Urinary retention, Drowsiness/dizziness, GI upset, Eyes (blurred vision/dry eyes).

**Sympathetic Nervous System (SNS) receptors:** Alpha-1, Alpha-2, Beta-1, and Beta-2 receptors that are stimulated by epinephrine and norepinephrine.

**Tardive dyskinesia:** A syndrome of movement disorders associated with antipsychotic medications that persists for at least one month and can last up to several years despite discontinuation of the medications.





### Learning Objectives

- Identify assessment cues of depressive disorders
- Identify nursing priorities for clients with depressive disorders
- Plan outcomes for clients with depressive disorders
- Describe safety/protective interventions for clients with depressive disorders
- Apply evidence-based practice when planning care and interventions for clients with depressive disorders
- Analyze treatments for clients with depressive disorders
- Apply the nursing process to clients with depressive disorders at risk for suicide

Depression is a common illness affecting an estimated 8% of adults in the United States annually.<sup>1</sup> Approximately 280 million people in the world have depression.<sup>2</sup>

Depressive disorders represent a group of conditions that share a core set of symptoms such as a persistent and pervasive low mood, loss of interest or pleasure in most activities, and a range of cognitive and physical symptoms

1. National Alliance on Mental Illness. (2017). *Depression*. <https://nami.org/About-Mental-Illness/Mental-Health-Conditions/Depression>
2. World Health Organization. (n.d.). *Depression*. [https://www.who.int/health-topics/depression#tab=tab\\_1](https://www.who.int/health-topics/depression#tab=tab_1)

that significantly impair daily functioning.<sup>3</sup> This chapter will discuss causes and types of depression, treatments for depression, and steps for applying the nursing process when caring for clients with depressive disorders.

3. World Health Organization. (2023). *Depressive disorder*. <https://www.who.int/news-room/fact-sheets/detail/depression>



## 7.2 Causes of Depression

There are several possible causes of depression, including faulty mood regulation by the brain, genetic vulnerability, stressful life events, medications, and medical problems. Based on current research, it is believed that several of these forces interact to bring on depression.<sup>1</sup>

### The Brain

Certain areas of the brain help regulate mood, including the hippocampus, amygdala, and hypothalamus. See Figure 7.1<sup>2</sup> for an image of these areas of the brain. Researchers believe that nerve cell connections, nerve cell growth, the functioning of nerve circuits, and levels of specific brain chemicals (called neurotransmitters) have a major impact on depression.<sup>3</sup>

1. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>
2. "1511\_The\_Limbic\_Lobe.jpg" by OpenStax College is licensed under [CC BY 3.0](https://creativecommons.org/licenses/by/3.0/). Access for free at <http://cnx.org/content/col11496/1.6/>
3. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>




Figure 7.1 Areas of the Brain Regulating Mood

The hippocampus is part of the limbic system in the brain that has a central role in processing long-term memory and recollection. For example, this part of the brain registers fear when you are confronted by a frightening situation, like an aggressive dog, and the memory of such an experience may make you wary of dogs you come across later in life. The hippocampus is smaller in some depressed people, and research suggests that ongoing exposure to stress hormones impairs the growth of nerve cells in this part of the brain.<sup>4</sup>

The amygdala is also part of the limbic system and is a group of structures deep in the brain that are associated with emotions such as anger, pleasure, sorrow, fear, and sexual arousal. The amygdala is activated when a person experiences or recalls emotionally charged memories, such as a frightening situation, and sends signals to the hypothalamus to stimulate the sympathetic fight-or-flight response. Activity in the amygdala is higher when

4. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>

a person is sad or clinically depressed. This increased activity continues even after recovery from depression.<sup>5</sup>

The hypothalamus is involved in the stress response. The stress response starts with a signal from the hypothalamus. The hypothalamus, pituitary gland, and adrenal glands form the hypothalamic-pituitary-adrenal (HPA) axis, which governs a multitude of hormonal activities in the body and also plays a role in depression. When a physical or emotional threat looms, the hypothalamus secretes corticotropin-releasing hormone (CRH) that rouses the body. CRH follows a pathway to the pituitary gland, where it stimulates the secretion of adrenocorticotrophic hormone (ACTH) into the bloodstream. When ACTH reaches the adrenal glands, it prompts the release of cortisol. The boost in cortisol readies the body to fight or flee by causing the heart to beat faster, the blood pressure to rise, and the respiratory rate to increase. CRH also affects the cerebral cortex, part of the amygdala, and the brain stem. It is thought to play a major role in coordinating one's thoughts and behaviors, emotional reactions, and involuntary responses. Working along a variety of neural pathways, it influences the concentration of neurotransmitters throughout the brain. Disturbances in hormonal systems affect neurotransmitters and vice versa.<sup>6</sup>

## Neurotransmitters

There are many types of neurotransmitters that play a role in depression<sup>7</sup>:

5. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>
6. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>
7. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical

- **Acetylcholine** enhances memory and is involved in learning and recall.<sup>8</sup> Research has also shown that increased levels have been linked to depression.<sup>9</sup>
- **Serotonin** helps regulate sleep, appetite, and mood and inhibits pain. Research supports the idea that some depressed people have reduced serotonin transmission. Low levels of a serotonin by-product have been linked to a higher risk for suicide.<sup>10</sup>
- **Norepinephrine** constricts blood vessels, raising blood pressure. It may trigger anxiety and be involved in some types of depression. It also seems to help determine motivation and reward.<sup>11</sup>
- **Dopamine** is essential to movement. It also influences motivation and plays a role in how a person perceives reality. Problems in dopamine transmission have been associated with psychosis, a severe form of distorted thinking characterized by hallucinations or delusions. It's also involved in the brain's reward system, so it is thought to play a role in

School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>

8. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>
9. Dulawa, S. C., & Janowsky, D. S. (2018). Cholinergic regulation of mood: From basic and clinical studies to emerging therapeutics, *Molecular Psychiatry*, 24(5), 694-709. doi: 10.1038/s41380-018-0219-x
10. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>
11. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>

substance abuse.<sup>12</sup> Dysregulation of dopamine pathways is also thought to contribute to anhedonia, the inability to experience pleasure, which is a common symptom in major depressive disorder.<sup>13</sup>

- **Glutamate** is a small molecule believed to act as an excitatory neurotransmitter and to play a role in bipolar disorder and schizophrenia. Animal research suggests that lithium stabilizes glutamate reuptake and smooths out the highs of mania and the lows of depression in the long-term.<sup>14</sup>
- **Gamma-aminobutyric acid (GABA)** is an amino acid that researchers believe acts as an inhibitory neurotransmitter. It is thought to help subdue anxiety.<sup>15</sup> Low levels of GABA have been associated with major depressive disorders.<sup>16</sup>

12. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>

13. Belujon, P. & Grace, A. A. (2017). Dopamine system dysregulation in major depressive disorders. *International Journal of Neuropsychopharmacology*, 20(12), 1036-1046. doi: 10.1093/ijnp/pyx056.

14. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>

15. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>

16. Luscher, B., Shen, Q., & Sahir, N. (2011). The GABAergic deficit hypothesis of major depressive disorder. *Molecular Psychiatry*, 16(4), 383-406. doi: 10.1038/mp.2010.120.

See Figure 7.2<sup>17</sup> for an illustration of neurotransmitter communication between neurons at the synapse. Antidepressants immediately boost the concentration of chemical messengers in the brain (neurotransmitters), but people typically don't begin to feel better for several weeks or longer. Experts have long wondered why people don't improve as soon as the level of neurotransmitters increases. New theories explain that antidepressants spur the growth and enhanced branching of nerve cells in the hippocampus (a process called neurogenesis), and mood improves over several weeks as nerves grow and form new connections.<sup>18</sup>

17. "[1225\\_Chemical\\_Synapse.jpg](#)" by Young, K. A., Wise, J. A., DeSaix, P., Kruse, D. H., Poe, B., Johnson, E., Johnson, J. E., Korol, O., Betts, J. G., & Womble, M. is licensed under [CC BY 4.0](#)

18. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>




Figure 7.2 Neurotransmitters at the Synapse Level

## Genes

Every part of our body, including our brain, is controlled by our genes. Humans have almost 22,000 genes in their DNA within 46 chromosomes inside the nucleus of each cell. The sequence of nitrogen-containing bases within a strand of DNA forms the genes that act as a molecular code instructing cells in the assembly of amino acids into proteins. See Figure 7.3<sup>19</sup> for an image of sequences of bases within a DNA strand that forms genes. Genes make proteins that are involved in biological processes. Throughout life, different genes turn on and off and make the right proteins at the right time. However, genes can alter biology in a way that results in a person's mood becoming unstable. In a person who is genetically vulnerable to

19. "229\_Nucleotides-01.jpg" by OpenStax College is licensed under [CC BY 3.0](https://creativecommons.org/licenses/by/3.0/). Access for free at <http://cnx.org/content/col11496/1.6/>.

depression, any stress (such as a missed deadline at work or a medical illness) can then push this system off balance.<sup>20</sup>




Figure 7.3 Sequence of Bases Within DNA That Form Genes

It is well-known that depressive and bipolar disorders run in families. Mood is affected by dozens of genes, and as our genes differ, so does depression. As researchers pinpoint specific genes involved in mood disorders and better

20. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>



understand their functions, it is hoped that treatment for depressive disorders can become more individualized and more successful as clients receive targeted medication for their specific type of depression.<sup>21</sup>

Genetics provides one perspective on how resilient an individual is in the face of difficult life events. Temperament is determined by a person's genetic inheritance and the experiences they have had in life. For example, one person may have the temperament of an introvert and tend to withdraw from social situations, whereas another person may have the temperament of an extrovert who seeks out social situations and feels energized by them. Cognitive psychologists believe that one's view of the world and assumptions about how the world works influence how a person feels. Individuals develop their assumptions about life early and automatically fall back on them when loss, disappointment, or rejection occurs. For example, a person who was continually criticized as a child may have the genetic inheritance and temperament where they become so self-critical they can't bear the slightest criticism from others, which can slow or block their career progress and make intimate relationships more difficult. Therapy and medications can shift thoughts and attitudes that have developed over time.<sup>22</sup>

## Sociocultural Influencing Factors

Many sociocultural factors play a role in depression. These factors may be related to social norms, cultural expectations, and social environments. Stressful life events—such as the death of a loved one, the loss of a job, the diagnosis of a severe illness, or the end of a significant relationship—can

21. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>
22. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>

affect anyone. In addition, family and relationship stress, including marital conflicts or domestic abuse, can contribute significantly to depressive symptoms. Cultural and societal expectations may also create psychological strain. For example, women are often expected to excel both in the workplace and as primary caregivers at home, creating conflicting demands that can lead to chronic stress and feelings of inadequacy. Socioeconomic factors such as unemployment, financial instability, and low income are strongly linked to an increased risk of depression. Gender inequality further compounds these risks, as women disproportionately experience burdens like domestic violence, caregiving responsibilities, and economic insecurity—all of which are deeply rooted in societal norms and expectations. Beyond current life stressors, many individuals have experienced traumatic childhood events that continue to affect their coping and functioning into adulthood. It is estimated that 61% of adults have experienced early adverse childhood experiences (ACEs) such as abuse, neglect, or growing up in a household with violence, mental illness, substance use, incarceration, or divorce. See Figure 7.4<sup>23</sup> for an illustration of adverse childhood experiences. Toxic stress from ACEs can change brain development and affect how the body responds to stress. ACEs are linked to chronic health problems, mental illness, and substance misuse in adulthood.<sup>24</sup>

23. “ACEs.png” by unknown author for [Centers for Disease Control and Prevention](#) is licensed in the [Public Domain](#). Access for free at [https://www.cdc.gov/injury/pdfs/priority/ACEs-Strategic-Plan\\_Final\\_508.pdf](https://www.cdc.gov/injury/pdfs/priority/ACEs-Strategic-Plan_Final_508.pdf).

24. Centers for Disease Control and Prevention. (2021). *Adverse childhood experiences (ACEs)*. <https://www.cdc.gov/violenceprevention/aces/index.html>

## Figure 1. What are Adverse Childhood Experiences?




Figure 7.4 Adverse Childhood Experiences (ACEs)

Stress triggers a chain of chemical reactions and responses in the body. If the stress is short-lived, the body usually returns to normal. But when stress is chronic or the system gets stuck in overdrive, changes in the body and brain can be long-lasting. Every real or perceived threat to one's body triggers a cascade of stress hormones that produces physiological changes called the stress response. Normally, a feedback loop allows the body to turn off "fight-or-flight" defenses when the threat passes. In some cases, though, the floodgates never close properly, and cortisol levels rise too often or simply stay high. These elevated cortisol levels can contribute to problems such as high blood pressure, immune suppression, asthma, and depression. Studies have also shown that people who have depressive disorders typically have increased levels of CRH. Antidepressants and electroconvulsive therapy are both known to reduce these high CRH levels. As CRH levels return to normal, depressive symptoms recede. Research also suggests that trauma during

childhood can negatively affect the functioning of CRH and the HPA axis throughout life.<sup>25</sup>

▶ Read the [Adverse Childhood Experiences Prevention Strategy](#).

## Medical Problems/Medication

Certain medical problems are linked to up to 10% to 15% of all depressions. For example, hypothyroidism, a condition where the body produces too little thyroid hormone, often leads to exhaustion and depression, whereas hyperthyroidism (excess thyroid hormone) can trigger manic symptoms. Heart disease has also been linked to depression, with up to half of heart attack survivors reporting feeling blue and many having significant depression. Another example of depression linked to a medical condition is postpartum depression that occurs after pregnancy.<sup>26</sup>

The following medical conditions have also been associated with depression and other mood disorders<sup>27</sup>:

- Degenerative neurological conditions, such as multiple sclerosis,

25. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>

26. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>

27. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>

Parkinson's disease, Alzheimer's disease, and Huntington's disease

- Cerebrovascular accidents (i.e., strokes)
- Some nutritional deficiencies, such as a lack of vitamin B12
- Endocrine disorders involving the parathyroid or adrenal glands
- Immune system diseases, such as lupus
- Some viruses, such as mononucleosis, hepatitis, and HIV
- Cancer
- Erectile dysfunction in men
- Chronic pain

When considering the connection between health problems and depression, an important question to address is which came first, the medical condition or the mood changes. Stress of having certain illnesses can trigger depression, whereas in other cases, depression precedes the medical illness and may even contribute to it. If depression is caused by an underlying medical problem, the mood changes should disappear after the medical condition is treated. For example, after hypothyroidism is treated, lethargy and depression often lift. In many cases, however, depression is an independent problem, which means that in order to be successful, treatment must address depression directly.<sup>28</sup>

Symptoms of depression can be a side effect of certain drugs, such as steroids or some types of blood pressure medication. Other medications associated with depressive symptoms include corticosteroids, antiepileptic drugs, oral contraceptives, and barbiturates. In addition, substances such as alcohol and cannabis can contribute to or worsen depressive symptoms, particularly with long-term or heavy use. A health care provider can help determine whether a

28. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>

new medication, a change in dosage, or interactions with other drugs or substances might be affecting an individual's mood.<sup>29</sup>

29. Harvard Health Publishing. (2019). *What causes depression?* Harvard Medical School. <https://www.health.harvard.edu/mind-and-mood/what-causes-depression>

## 7.3 Types of Depression

Depression is different from the usual mood fluctuations and short-lived emotional responses to everyday life stressors. When it is recurrent with moderate or severe intensity, depression can become a serious health condition that causes the affected person to suffer greatly and function poorly at work and school and can affect relationships with family and friends. At its worst, depression can lead to suicide. Over 700,000 people die every year due to suicide. Barriers to effective care for depression include a lack of resources, lack of trained health care providers, and social stigma associated with mental health disorders.<sup>1</sup> Depressive disorders represent a group of conditions that share a core set of symptoms but differ in important ways such as duration, timing, triggers, and presumed causes (etiology). The common feature across all depressive disorders is the presence of a sad, empty, or irritable mood that is accompanied by physical (somatic) and mental (cognitive) changes. These changes significantly impair a person's ability to function in daily life—impacting work, relationships, self-care, and overall quality of life. See Figure 7.5<sup>2</sup> for an artistic depiction of an individual experiencing depression.

1. World Health Organization. (2021). *Depression*. <https://www.who.int/news-room/fact-sheets/detail/depression>

2. “320531.png” by j4p4n at [openclipart.org](https://openclipart.org) is licensed in the [Public Domain](#).



Figure 7.5 Depression

## Major Depressive Disorder

A **major depressive episode** is a period of at least two weeks during which a person experiences a persistently depressed mood or a loss of interest or pleasure in nearly all activities, most of the day, nearly every day. This emotional state is accompanied by additional symptoms such as changes in appetite or weight, sleep disturbances (insomnia or sleeping too much), fatigue or loss of energy, feelings of worthlessness or excessive guilt, difficulty concentrating or making decisions, and recurrent thoughts of death or suicide. During a **depressive episode**, the person experiences a depressed mood (feeling sad, irritable, empty) or a loss of pleasure or interest in activities (anhedonia) for most of the day, nearly every day, for at least two weeks. Several other symptoms may also be present, which may include poor concentration, feelings of excessive guilt or low self-worth, hopelessness about the future, thoughts about dying or suicide, disrupted sleep, changes in appetite or weight, and feeling especially tired or low in energy. In some cultural contexts, some people may express their mood changes more readily in the form of bodily symptoms (such as pain, fatigue, or weakness) that are not due to another medical condition. During a depressive episode, the person experiences significant difficulty and/or impairment in important areas of personal, family, social, educational, and work functioning.<sup>3</sup>

3. World Health Organization. (2021). *Depression*. <https://www.who.int/news-room/fact-sheets/detail/depression>



A depressive episode is categorized by a provider as mild, moderate, or severe depending on the number and severity of symptoms, as well as the impact on the individual's functioning.<sup>4</sup> Mild severity indicates the symptoms to make the diagnosis are present but result in minor impairment in social or occupational functioning. Moderate severity indicates symptoms and impairment are between mild and severe. Severe indicates the intensity is seriously distressing, unmanageable, and markedly interferes with social and occupational functioning. Read the full list of criteria for Major Depressive Disorder from the *DSM-5-TR* in the following box.

### **DSM-5-TR Criteria for Major Depressive Disorder<sup>5</sup>**

1. Five or more of the following symptoms have been present during the same two-week period and represent a change from previous functioning; at least one of the symptoms is either (1) depressed mood or (2) loss of interest or pleasure.
  - Depressed mood most of the day, nearly every day, as indicated by either subjective report (e.g., feels sad, empty, hopeless) or observation made by others (e.g., appears tearful). (Note: In children and adolescents, can be irritable mood.)
  - Markedly diminished interest or pleasure in all,

4. World Health Organization. (2021). *Depression*. <https://www.who.int/news-room/fact-sheets/detail/depression>

5. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

or almost all, activities most of the day, nearly every day.

- Significant weight loss when not dieting or weight gain (i.e., a change of more than 5% of body weight in a month) or decrease or increase in appetite nearly every day. (Note: In children, consider failure to make expected weight gain.)
- Insomnia or hypersomnia nearly every day.
- Psychomotor agitation or retardation nearly every day (observable by others, not merely subjective feelings of restlessness or being slowed down).
- Fatigue or loss of energy nearly every day.
- Feelings of worthlessness or excessive or inappropriate guilt (which may be delusional) nearly every day (not merely self-reproach or guilt about being sick).
- Diminished ability to think or concentrate, or indecisiveness, nearly every day (either by subjective account or as observed by others).
- Recurrent thoughts of death (not just fear of dying), recurrent suicidal ideation without a specific plan, or a suicide attempt or a specific plan for committing suicide.

2. The symptoms cause clinically significant distress or impairment in social, occupational, or other important areas of functioning.
3. The episode is not attributable to the physiological effects of a substance or another medical condition.

4. The client has never had a manic or hypomanic episode.

## Seasonal Pattern

**Seasonal affective disorder (SAD)** is a type of major depressive disorder that affects about 5% of adults in the United States. People with SAD experience mood changes and symptoms similar to depression. The symptoms usually occur during the fall and winter months when there is less sunlight and usually improve with the arrival of spring. SAD is more than just “winter blues.” The symptoms can be distressing and overwhelming and can interfere with daily functioning.<sup>6</sup>

SAD has been linked to a biochemical imbalance in the brain prompted by shorter daylight hours and less sunlight in winter. As seasons change, people experience a shift in their biological internal clock or circadian rhythm that can cause them to be out of step with their daily schedule. SAD is more common in people living far from the equator where there are fewer daylight hours in the winter.<sup>7</sup>

SAD can be effectively treated with light therapy, antidepressant medications, cognitive behavioral therapy, or some combination of these. While symptoms will generally improve on their own with the change of season, symptoms can

6. American Psychiatric Association. (2020). *Seasonal affective disorder (SAD)*. <https://www.psychiatry.org/patients-families/depression/seasonal-affective-disorder>

7. American Psychiatric Association. (2020). *Seasonal affective disorder (SAD)*. <https://www.psychiatry.org/patients-families/depression/seasonal-affective-disorder>

improve more quickly with treatment. **Light therapy** involves sitting in front of a light therapy box that emits a very bright light (and filters out harmful ultraviolet [UV] rays). Bright light therapy typically involves exposure to a light box emitting 2,500 to 10,000 lx of light<sup>8</sup> It usually requires 20 minutes or more per day, typically first thing in the morning during the winter months. Most people see some improvements from light therapy within one or two weeks of beginning treatment. For some people, increased exposure to natural sunlight can help improve symptoms of SAD by spending time outside or arranging their home or office space to increase exposure to windows during daylight hours. General wellness activities such as performing regular exercise, eating healthfully, getting enough sleep, and staying active and connected (such as volunteering, participating in group activities and getting together with friends and family) can also help.<sup>9</sup>

## Peripartum Onset

Pregnancy and the time following childbirth represent a uniquely vulnerable period for women. This phase of life often involves intense biological changes, emotional adjustments, financial pressures, and shifting social roles. These stressors can have a profound impact on a woman's mental health, potentially resulting in a spectrum of mood disturbances—from mild and temporary feelings of sadness to more severe depressive disorders that may require clinical intervention.

8. Pjrek, E., Friedrich, M.E., Cambioli, L., Dold, M., Jäger, F., Komorowski, A., Lanzenberger, R., Kasper, S., Winkler, D. (2020). The efficacy of light therapy in the treatment of seasonal affective disorder: A meta-analysis of randomized controlled trials. *Psychotherapy Psychosomatics*, 89(1):17-24. [doi: 10.1159/000502891](https://doi.org/10.1159/000502891).
9. American Psychiatric Association. (2020). *Seasonal affective disorder (SAD)*. <https://www.psychiatry.org/patients-families/depression/seasonal-affective-disorder>

## BABY BLUES

It is common for new mothers to experience what is known as the “baby blues,” which affects up to 70% of women after childbirth. The baby blues typically begin within a few days of delivery and include emotional lability, tearfulness without an obvious cause, mild irritability, restlessness, and a general sense of anxiety or being overwhelmed. These symptoms are usually mild, self-limited, and do not interfere significantly with a mother’s ability to care for herself or her baby. Most cases resolve spontaneously within about two weeks, and medical treatment is generally not necessary.

## PERINATAL DEPRESSION

Perinatal depression represents a more serious mental health condition.

**Perinatal depression** is a broad term that refers to depressive symptoms that occur any time during pregnancy and up to one year after childbirth. This includes both antenatal (or prenatal) depression, which develops while the person is still pregnant, and postpartum depression, which occurs after the baby is born. Perinatal depression can affect emotional well-being, physical health, bonding with the baby, and overall family functioning. It is more than just occasional sadness or stress. Symptoms of perinatal depression often include persistent feelings of sadness, anxiety, fatigue, irritability, changes in sleep and appetite, and difficulty concentrating or feeling connected to the baby.<sup>10</sup>

### Postpartum Depression

Postpartum depression is a subset of perinatal depression and refers to a major depressive episode that begins after childbirth. It is much more serious mental health condition than the “baby blues”. **Postpartum depression** is characterized by persistent feelings of sadness, hopelessness, anxiety, and

10. National Institute of Mental Health (2023). *Perinatal Depression*.  
<https://www.nimh.nih.gov/health/publications/perinatal-depression>

exhaustion that interfere with a mother's ability to function and care for her infant. These symptoms go beyond the scope of the baby blues in both intensity and duration, often requiring professional treatment such as psychotherapy, medication, or a combination of both. If left untreated, postpartum depression can significantly impair the mother–infant bond, contribute to difficulties with infant feeding and sleeping, and may have long-term consequences for the child's emotional, cognitive, and social development. Symptoms of postpartum depression usually start between one to three weeks after birth<sup>11</sup> These symptoms reflect the same characteristics as those in the perinatal depression group, yet the time of onset is different. Many healthcare providers recognize that postpartum depression can emerge at any point within the first year following childbirth, even though this extended timeframe is not explicitly captured in DSM-5-TR criteria.

In rare and severe cases, postpartum depression may escalate into a condition called postpartum psychosis. This is a psychiatric emergency that requires immediate medical intervention. Postpartum psychosis can involve hallucinations, delusions, disorganized thinking, mania, severe agitation, and confusion. Because of the potential risk for self-harm or harm to the infant, individuals experiencing symptoms of postpartum psychosis should be brought to the nearest emergency room or emergency services should be contacted immediately.


Understanding the range and severity of mood disorders related to pregnancy and childbirth is essential for early identification and intervention. Prompt recognition and appropriate care can significantly improve outcomes for both the mother and her child.

Pregnancy and the period after delivery can be a vulnerable time for women. Mothers can experience significant biological, emotional, financial, and social

11. National Library of Medicine. (2022). Postpartum depression screening. <https://medlineplus.gov/lab-tests/postpartum-depression-screening/>

changes during this time. Up to 70 percent of new mothers experience the “baby blues,” a short-term condition that does not interfere with their daily activities and does not require medical attention. Symptoms include crying for no reason, irritability, restlessness, and anxiety. These symptoms generally last up to two weeks and resolve on their own.<sup>12</sup>

See Figure 7.6<sup>13</sup> for an illustration of perinatal depression from the National Institutes of Health.



12. American Psychiatric Association (2020). *What is peripartum depression (formerly postpartum)?* <https://www.psychiatry.org/patients-families/postpartum-depression/what-is-postpartum-depression>
13. This image is a derivative of “20-mh-8116-perinataldepression.pdf” by [National Institute of Mental Health](https://www.nimh.nih.gov/health/publications/perinatal-depression) and is licensed in the [Public Domain](https://www.nimh.nih.gov/health/publications/perinatal-depression). Access for free at <https://www.nimh.nih.gov/health/publications/perinatal-depression>.

Figure 7.6 Perinatal Depression

Untreated peripartum depression is not only a problem for the mother's health and quality of life but can also affect the well-being of the baby. Postpartum depression can cause bonding issues with the baby and also contribute to sleeping and feeding problems for the baby. In the long-term, children of mothers with peripartum depression are at greater risk for cognitive, emotional, developmental and verbal deficits, and impaired social skills.<sup>14</sup>

- ▶ Read more information about perinatal depression on the National Institute of Mental Health's [Perinatal Depression](#) webpage.

## Persistent Depressive Disorder

**Persistent Depressive Disorder**, previously known as dysthymia, is a chronic form of depression that is typically less severe in intensity than Major Depressive Disorder but lasts much longer. In adults, symptoms must be present for most of the day, more days than not, for at least two years. In children and adolescents, the required duration is at least one year, and the mood may be irritable rather than depressed. Individuals with this disorder often describe themselves as feeling “down in the dumps,” “low,” or “just not themselves.” Because the symptoms are milder and may develop gradually, they can become part of a person's normal daily experience, leading many to

14. American Psychiatric Association (2020). *What is peripartum depression (formerly postpartum)?* <https://www.psychiatry.org/patients-families/postpartum-depression/what-is-postpartum-depression>



believe their persistent low mood is simply part of their personality rather than a treatable mental health condition.<sup>15</sup>

Common features of Persistent Depressive Disorder include poor appetite or overeating, insomnia or hypersomnia, low energy or fatigue, low self-esteem, poor concentration or difficulty making decisions, and feelings of hopelessness. While these symptoms may not be as intense as those seen in a major depressive episode, they can still significantly affect a person's functioning and quality of life over time.<sup>16</sup>

Because the condition is long-lasting, it can interfere with relationships, work or school responsibilities, and the ability to enjoy life. Treatment often includes a combination of psychotherapy, such as cognitive behavioral therapy, and antidepressant medications like SSRIs or SNRIs. Ongoing support and follow-up care are important, as the chronic nature of the disorder requires long-term management to help individuals regain and maintain emotional well-being.<sup>17</sup>

## Premenstrual Dysphoric Disorder

**Premenstrual Dysphoric Disorder (PMDD)** is a severe and disabling form of premenstrual syndrome (PMS) that involves significant mood disturbances and physical symptoms occurring in the final week before menstruation. While many individuals experience some degree of emotional or physical discomfort during their menstrual cycle, PMDD is much more intense and

15. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.
16. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.
17. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

can seriously interfere with daily life, relationships, and functioning. The symptoms become minimal or absent in the week post-menses.<sup>18</sup>

Individuals with PMDD must have one or more of the following symptoms<sup>19</sup> :

- Extreme affective lability (mood swings, feeling suddenly sad or tearful, or increased sensitivity to rejection)
- Significant irritability or anger or increased interpersonal conflicts
- Significant depressed mood, feelings of hopelessness, or self-deprecating thoughts
- Significant anxiety, tension, or feelings of being keyed up or on edge.

One or more of the following symptoms must be additionally present to reach a total of five symptoms when combined with the symptoms in the preceding paragraph<sup>20</sup> :

- Decreased interest in usual activities (e.g., work, school, friends, hobbies)
- Difficulty concentrating
- Lethargy, easy fatigability, or significant lack of energy
- Significant change in appetite, overeating, or specific food cravings
- Hypersomnia or insomnia
- A sense of being overwhelmed or out of control
- Physical symptoms such as breast tenderness or swelling, joint or muscle pain, a sense of bloating, or weight gain.

18. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

19. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

20. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

A key feature of PMDD is that these symptoms are not merely uncomfortable—they are severe enough to disrupt work, school, and personal relationships, and they go beyond what is typically expected in PMS.<sup>21</sup>

The exact cause of PMDD is not fully understood, but it is believed to be related to sensitivity to normal hormonal fluctuations during the menstrual cycle, particularly in individuals with an underlying vulnerability to mood disorders. Treatment may involve a combination of antidepressant medications, birth control hormonal medications, psychotherapy (especially cognitive behavioral therapy), and lifestyle modifications such as regular exercise, vitamin supplementation with B-6 and magnesium, and adaptive coping strategies such as yoga and meditation.<sup>22</sup>

## Substance/Medication-Induced Depressive Disorder

**Substance/Medication-Induced Depressive Disorder** is a type of depression that occurs as a direct result of using, misusing, or withdrawing from certain substances or medications. This disorder is diagnosed when depressive symptoms such as a persistently low mood, lack of interest or pleasure, fatigue, difficulty concentrating, and feelings of worthlessness are closely linked to the use of a particular drug or substance and not better explained by another mental health condition. The depressive symptoms typically develop during or soon after substance use or withdrawal and must cause significant distress or impairment in daily functioning.<sup>23</sup>

21. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.
22. Cleveland Clinic. (2023). *Premenstrual dysphoric disorder*.  
<https://my.clevelandclinic.org/health/diseases/9132-premenstrual-dysphoric-disorder-pmdd>
23. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

A wide range of substances can trigger this disorder. Common examples include alcohol, cannabis, opioids, sedatives, stimulants, and hallucinogens, as well as prescription medications such as corticosteroids, antihypertensives, hormonal contraceptives, barbiturates, and certain antiepileptic drugs. The effects may occur in people who are taking the substance as prescribed or in those who are misusing it. In some cases, individuals may not realize that their depressive symptoms are linked to a medication or drug, especially if the symptoms have developed gradually.<sup>24</sup>

An important part of diagnosing this condition is establishing the temporal relationship between substance use and the onset of depression. Once the substance is discontinued or the medication is adjusted, symptoms may improve—though in some cases, clinical treatment is still necessary to help manage mood changes. Treatment involves identifying and stopping the responsible substance, providing emotional support, and in some cases, using antidepressant medications or therapy to help stabilize mood during recovery.<sup>25</sup>

## Depressive Disorder Due to Another Medical Condition

**Depressive Disorder Due to Another Medical Condition** is diagnosed when a person experiences a persistent depressed mood or a significant loss of interest or pleasure in activities, and these symptoms are directly caused by a medical illness or condition. Unlike other types of depression, this disorder is not primarily psychological in origin—it results from the physiological effects

24. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

25. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

of a medical issue. The depressive symptoms must cause significant distress or impair the person's ability to function in daily life.<sup>26</sup>

Common medical conditions associated with this type of depression include neurological disorders (such as stroke, Parkinson's disease, multiple sclerosis), endocrine disorders (like hypothyroidism or Cushing's disease), and chronic illnesses (such as cancer, diabetes, or heart disease). The timing of symptom onset is important for diagnosis, as the depressive symptoms must clearly follow the development of the medical condition and not be better explained by another mental health disorder.<sup>27</sup>

Treatment involves addressing both the underlying medical condition and the depressive symptoms, which may include psychotherapy, medication, or supportive care depending on the individual's needs.<sup>28</sup>

## Mixed Features

Depressive episodes can also be classified as "mixed," meaning symptoms are present but do not predominate.<sup>29</sup> Mixed features associated with a major depressive episode have been found to be a significant risk factor for the development of bipolar I or bipolar II disorder. Characteristics and treatment of bipolar disorder are further discussed in the "[Bipolar Disorders](#)" chapter.

26. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

27. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

28. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

29. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

See Figure 7.7<sup>30</sup> for a comparison of the features of different types of mood disorders.





Figure 7.7 Mood Disorders. Used under Fair Use.

- ▶ Read more information at the American Psychiatric Association's [What is Depression?](#) webpage.
- ▶ Listen to [The Dark Place](#) podcast on the Podbean website to hear personal perspectives on depression and other mental health topics.

30. "mood-disorders.jpg" by Dr. Vipul Rastogi, MBBS; DCP (Ireland); MRCPsych (UK) Speciality Registrar, Hampshire Partnership Trust, UK for [www.clinicaljunior.com](http://www.clinicaljunior.com). Image used under Fair Use. Access for free at <http://www.clinicaljunior.com/psychmoodvipul.html>

View the following YouTube video of an advocate  
 explaining her experiences with depression and her  
journey to recovery<sup>31</sup> :



One or more interactive elements has been excluded from this version of the text. You can view them online here: <https://wtcs.pressbooks.pub/nursingmhcc/?p=326#oembed-1>

## Grief

Individuals often have difficulty coping with the death or loss of a loved one, job, or relationship. It is normal for feelings of sadness or grief to develop in response to such situations, but these symptoms are not the same as having a major depressive disorder. The grieving process is natural and unique to each individual and shares some of the same features of depression. Both grief and depression can involve intense sadness and withdrawal from usual activities with increased risk for suicide.

According to the American Psychiatric Association, grief and depression are different in these ways<sup>32</sup> :

- In grief, painful feelings come in waves, often intermixed with positive

31. World Health Organization (WHO). (2018, September 19). *Youth voices: Kylie Verzosa on depression* [Video]. YouTube. Licensed in the [Public Domain](#). <https://youtu.be/tRc4XZXYTgY>

32. American Psychiatric Association. (2020). *What is depression?* <https://www.psychiatry.org/patients-families/depression/what-is-depression>

memories of the deceased. In major depression, mood and/or interest (pleasure) are decreased for most of two weeks.

- In grief, one's self-esteem is usually maintained. In major depression, feelings of worthlessness and self-loathing are common.
- In grief, thoughts of death may surface when thinking of or fantasizing about "joining" the deceased loved one. In major depression, thoughts are focused on ending one's life due to feeling worthless or undeserving of living or being unable to cope with the pain of depression.

## Prolonged Grief Disorder

The DSM-5-TR introduced a new diagnosis called **Prolonged Grief Disorder** that is characterized by intense longing and preoccupation with the deceased person that persists beyond 12 months and causes significant impairment. Additionally, since the death, at least three of the following symptoms are present<sup>33</sup> :

- Identity disruption (e.g., feeling as though part of oneself has died) since the death
- Significant sense of disbelief about the death
- Avoidance of reminders that the person is dead
- Intense emotional pain related to the death
- Difficulty reintegrating into one's relationships and activities after the death
- Emotional numbness as a result of the death
- Feeling that life is meaningless as a result of the death
- Intense loneliness as a result of the death

33. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.



- ▶ Read more information about grief in the “[Grief and Loss](#)” chapter in *Open RN Nursing Fundamentals, 2e*.

## 7.4 Treatments for Depression

Depression is one of the most treatable mental disorders. Between 80% and 90% percent of people with depression eventually respond well to treatment. Almost all patients gain some relief from their symptoms with effective treatment. Before a diagnosis is made or treatment planned, a mental health care provider should conduct a thorough diagnostic evaluation, including an interview, mental status examination, psychosocial assessment, and a physical examination performed by a primary care provider. In some cases, a blood test might be done to make sure the depression is not due to a medical condition like a thyroid problem or a vitamin deficiency. The evaluation also explores medical and family histories, cultural beliefs, and environmental factors as part of the psychosocial assessment with the goal of arriving at a diagnosis and planning a course of action. Treatments for depression include medications, psychotherapy, electroconvulsive therapy, transcranial magnetic stimulation, and encouraging self-care and effective coping strategies. Nurses should keep patient-centered care and client preferences in mind when implementing a client's treatment plan. For example, the client should be educated about the possible adverse effects associated with antidepressant medication, and their ability to pay for and obtain transportation to treatments like psychotherapy should be considered.<sup>1</sup>

### Antidepressant Medications

Brain chemistry may contribute to an individual's depression and may factor into their treatment. Read more about the causes of depression in the [“Causes of Depression”](#) section of this chapter. For this reason, providers may prescribe antidepressants to help modify an individual's brain chemistry.

Antidepressants are used to regulate and increase neurotransmitters in the brain to improve mood. There are several classes of antidepressants with

1. American Psychiatric Association. (2020). *What is depression?*  
<https://www.psychiatry.org/patients-families/depression/what-is-depression>

different mechanisms of action and potential adverse side effects. Classes of antidepressants, their mechanism of action, and common side effects are outlined in Table 7.4.<sup>2</sup> Common side effects are listed in this table by frequency of their occurrence. Some medications can cause QTc prolongation, which refers to delayed cardiac electrical conduction that can lead to dysrhythmias. Tricyclic antidepressants (TCAs) and monoamine oxidase inhibitors (MAOIs) often cause anticholinergic side effects (e.g., tachycardia, urinary retention, constipation, dry mouth, blurred vision, confusion, psychomotor slowing, sedation, and delirium). A specific medication class is typically initiated by the provider based on the individual's symptoms, potential side effects, cost, and family history of success with certain medications.

Review neurotransmitter actions and related central nervous system physiology in the “[Psychotropics Medications](#)” chapter.

Table 7.4 Antidepressants

2. Rush, J. A. (2025). Side effects of antidepressant medication. UpToDate. <https://www.uptodate.com/>

Medication Class	Mechanism of Action	Common Side Effects  (*Indicates medical emergency)
<b>Serotonin-Norepinephrine Reuptake Inhibitors (SNRIs)</b>  Common examples:  Venlafaxine  Duloxetine	Block the uptake of both serotonin and norepinephrine from the cell synapse. Similar to SSRIs but with two neurotransmitters.	<ul style="list-style-type: none"> <li>• Gastrointestinal upset</li> <li>• Sexual dysfunction</li> <li>• Drowsiness</li> <li>• Insomnia</li> <li>• Agitation</li> <li>• Weight gain</li> </ul>
<b>Selective Serotonin Reuptake Inhibitors (SSRIs)</b>  Common examples:  Fluoxetine  Sertraline  Citalopram	Impact the receptors of the cell synapse to inhibit or prevent the uptake of serotonin, making the neurotransmitter serotonin stay in the synapse longer.	<ul style="list-style-type: none"> <li>• Sexual dysfunction</li> <li>• Gastrointestinal upset</li> <li>• QTc prolongation</li> <li>• Weight gain</li> <li>• Insomnia</li> <li>• Agitation</li> <li>• Drowsiness</li> <li>• Dizziness</li> <li>• Headache</li> <li>• Dry mouth</li> <li>• Constipation</li> <li>• Tremor</li> </ul> <b>*Serotonin syndrome</b>

<p><b>Tricyclic Antidepressants (TCAs)</b></p> <p>Common examples:</p> <p>Amitriptyline</p> <p>Nortriptyline</p>	<p>Block the presynaptic receptor for norepinephrine and partially serotonin. This makes the neurotransmitter norepinephrine level increase in the synapse.</p>	<ul style="list-style-type: none"> <li>• Anticholinergic side effects</li> <li>• Orthostatic hypotension</li> <li>• Drowsiness</li> <li>• QTc prolongation</li> <li>• Insomnia</li> <li>• Agitation</li> <li>• Weight gain</li> <li>• Sexual dysfunction</li> </ul>
<p><b>Monoamine Oxidase Inhibitors (MAOIs)</b></p> <p>Common examples:</p> <p>Phenelzine</p> <p>Tranylcypromine</p>	<p>Block the enzyme that breaks down monoamine, which causes an increase in the level of neurotransmitters serotonin and norepinephrine.</p>	<ul style="list-style-type: none"> <li>• Sexual dysfunction</li> <li>• Orthostatic hypotension</li> <li>• Insomnia</li> <li>• Agitation</li> <li>• Anticholinergic side effects</li> <li>• Drowsiness</li> <li>• Gastrointestinal upset</li> <li>• Weight gain</li> </ul> <p><b>*Hypertensive crisis</b></p>

<b>Norepinephrine and Dopamine Reuptake Inhibitor (NDRI)</b>  Bupropion	Blocks the uptake of both norepinephrine and dopamine from the cell synapse.	<ul style="list-style-type: none"> <li>• Tachycardia</li> <li>• Diaphoresis</li> <li>• Weight loss</li> <li>• Gastrointestinal upset</li> <li>• Agitation</li> <li>• Dizziness,</li> <li>• Headache</li> <li>• Insomnia</li> <li>• Tremor</li> <li>• Blurred vision</li> </ul>
---	--	--

Antidepressants may produce some improvement within the first week or two, but full benefits may not be seen for two to three months. If a patient feels little or no improvement after several weeks, the mental health care provider can alter the dose of the medication, add another medication, or substitute another antidepressant. In a similar manner, if a client develops ongoing bothersome side effects, the provider can switch them to a different medication. Nurses should encourage clients to contact their mental health care prescriber if a medication is not working within the expected time frame or if they are experiencing ongoing bothersome side effects. It is typically recommended that clients continue to take medication(s) for six or more months after the symptoms have improved. Long-term maintenance therapy may be suggested to decrease the risk of future episodes for people at high risk for recurrence.<sup>3</sup>

## Black Box Warning

A **Black Box Warning** is a significant warning from the Food and Drug

3. American Psychiatric Association. (2020). *What is depression?*

<https://www.psychiatry.org/patients-families/depression/what-is-depression>

Administration (FDA) that alerts the public and health care providers to serious side effects, such as injury or death. Black Box Warnings are in place for all classes of antidepressants used with children, adolescents, and young adults due to a higher risk of suicide. All clients receiving antidepressants should be monitored for signs of worsening depression or changing behavior, especially when the medication is started or dosages changed.

## Adverse/Side Effects

Nurses must monitor clients receiving antidepressants for side effects and report concerns to the prescribing provider. Clients should be instructed to immediately call their provider if they have any of the following symptoms, especially if they are new or worsening<sup>4</sup>:

- Thoughts about suicide or dying
- Attempts to commit suicide
- Worsening depression
- Anxiety
- Agitation or restlessness
- Panic attacks
- Trouble sleeping (insomnia)
- Irritability
- Aggression, anger, or violence
- Dangerous impulses
- Increased activity and talking (i.e., signs of mania)
- Other unusual changes in behavior or mood

4. National Institute of Mental Health. (2016). *Mental health medications*. U.S. Department of Health & Human Services. <https://www.nimh.nih.gov/health/topics/mental-health-medications>

## SEROTONIN SYNDROME

High doses of antidepressants or a combination of medications that affect serotonin, such as antidepressants or triptans (used to treat migraine headaches) can cause a medical emergency called serotonin syndrome. The presentation of **serotonin syndrome** is extremely variable, ranging from mild symptoms to a life-threatening syndrome. It typically develops within 24 hours from the increased dosage or combination of medications and can be fatal.

Symptoms of serotonin syndrome can be classified into three categories<sup>5 6 7</sup>:

- Mental status changes: Agitation, restlessness, or delirium
- Autonomic hyperactivity: Tachycardia, hypertension, hyperthermia, diaphoresis, shivering, vomiting, or diarrhea
- Neuromuscular hyperactivity: Tremor, muscle hypertonia or rigidity, myoclonus, hyperreflexia, or clonus (including rapid, horizontal eye movements)

A mnemonic commonly used to remember the symptoms of serotonin syndrome is **SHIVERS**:

- **S**: Shivering: A neuromuscular symptom similar to tremors specific to serotonin syndrome
- **H**: Hyperreflexia (and myoclonus): Hyperactive reflexes most prominent in

5. A.D.A.M. Medical Encyclopedia [Internet]. (2021). *Serotonin syndrome*. <https://medlineplus.gov/ency/article/007272.htm>

6. Boyer, E. W. (2021). Serotonin syndrome (serotonin toxicity). *UpToDate*. <https://www.uptodate.com/>

7. Tanen, D. (2021). *Serotonin syndrome*. Merck Manual Professional Version. <https://www.merckmanuals.com/professional/injuries-poisoning/heat-illness/serotonin-syndrome>



the lower extremities.

- **I:** Increased Temperature
- **V:** Vital Sign Abnormalities: Tachycardia, tachypnea, and labile blood pressure
- **E:** Encephalopathy: Mental status changes such as agitation, delirium, and confusion
- **R:** Restlessness
- **S:** Sweating

People may get slowly worse and can become severely ill if not quickly treated. Untreated, serotonin syndrome can be deadly. With treatment, symptoms usually go away within 24 hours, but permanent kidney damage may result even with treatment. Uncontrolled muscle spasms can cause severe muscle breakdown called **rhabdomyolysis**. Myoglobin is released into the blood with muscle breakdown and clogs renal tubules, which can cause severe kidney damage if serotonin syndrome isn't recognized promptly and treated.

Treatment of serotonin syndrome may include the following<sup>8,9</sup>:

- Stopping all serotonergic medications.
- Providing supportive care to normalize vital signs such as IV fluids, cooling measures, and medications to control heart rate and blood pressure.
- Sedating with benzodiazepines, such as diazepam or lorazepam, to decrease agitation, seizure-like movements, and muscle stiffness.
- If symptoms persist, administering cyproheptadine to block serotonin production.

8. Boyer, E. W. (2021). Serotonin syndrome (serotonin toxicity).

UpToDate. <https://www.uptodate.com/>

9. Tanen, D. (2021). *Serotonin syndrome*. Merck Manual Professional Version.

<https://www.merckmanuals.com/professional/injuries-poisoning/heat-illness/serotonin-syndrome>

- For patients with severe symptoms such as hyperthermia and muscle rigidity, more aggressive measures are required. These include sedation, intubation, neuromuscular paralysis, and active cooling techniques.<sup>10</sup>

Clients with moderate to severe serotonin syndrome should be hospitalized for close monitoring and management. Serotonin syndrome, in its most severe form, can resemble neuroleptic malignant syndrome (NMS) caused by antipsychotic medications. However, NMS develops over a period of days to weeks. Neuroleptic malignant syndrome (NMS) will be discussed further in [Chapter 11](#).

## HYPERTENSIVE CRISIS

Hypertensive crisis can occur when clients taking monoamine oxidase inhibitors (MAOIs) also take medications containing pseudoephedrine or eat foods containing tyramine (aged foods; fermented foods; cured meats; alcoholic beverages such as beer or red wine; or overripe fruits such as raisins, prunes, or bananas). MAOIs inhibit the breakdown of tyramine, causing elevated tyramine levels in the body that can lead to hypertensive crisis.

**Hypertensive crisis** is a medical emergency defined as severe hypertension (blood pressure over 180/120 mm Hg) with acute end-organ damage such as stroke, myocardial infarction, or acute kidney damage. Symptoms may include a severe headache accompanied with confusion and blurred vision. Tachycardia or bradycardia may be present and associated with constricting chest pain. Other symptoms include neck stiffness or soreness, nausea or vomiting, sweating, dilated pupils, photophobia, shortness of breath, severe anxiety, and unresponsiveness. Seizures may occur, as well as intracranial bleeding in association with the increased blood pressure. Hypertensive crisis treatment involves discontinuation of the offending agent, administration of

10. Spadaro, A., Scott, K. R., Koyfman, A., & Long, B. (2022). High risk and low prevalence diseases: Serotonin syndrome. *The American Journal of Emergency Medicine*, 61, 90-97. [doi: 10.1016/j.ajem.2022.08.030](https://doi.org/10.1016/j.ajem.2022.08.030).

appropriate intravenous antihypertensive medications such as phentolamine or labetalol, and supportive care.. In severe cases, treatment in the intensive care unit may be required.<sup>11,12</sup>

## Client Education

Clients should be instructed it may take 4 to 6 weeks for antidepressants to achieve their full effectiveness. They should not suddenly stop taking antidepressants or they may experience withdrawal symptoms. When it is time to stop the medication, the provider will slowly and safely decrease the dose. If clients stop taking the medication before the provider advises, the depression may return. They may not feel better with the first antidepressant they try, and they may need to try several different classes of medications to find one that works best for them. Education related to potential side effects and when to contact the provider, clinical worsening, avoiding alcohol, interference with cognitive or motor functioning, and potential drug interactions should also be provided.

Nurses also teach clients about coping strategies to reduce symptoms of depression. For example, regular exercise helps create positive feelings and improves mood. Getting enough quality sleep on a regular basis, eating a healthy diet, and avoiding alcohol (a depressant) can also reduce symptoms

11. U.S. National Library of Medicine. (2024). *Hypertensive crisis*. MedlinePlus. <https://medlineplus.gov/>
12. Sheps, S. G. (2021). *Hypertensive crisis: What are the symptoms?* Mayo Clinic. <https://www.mayoclinic.org/diseases-conditions/high-blood-pressure/expert-answers/hypertensive-crisis/faq-20058491#:~:text=A%20hypertensive%20crisis%20is%20a,higher%20%E2%80%94%20can%20damage%20blood%20vessels.>

of depression.<sup>13</sup> Read more about coping strategies in the “[Stress, Coping, and Crisis Intervention](#)” chapter.

## Psychotherapy

Psychotherapy may be used alone for treatment of mild depression or in combination with antidepressant medications for moderate to severe depression. Psychotherapy may involve only the individual, but it can include others such as family members or couples therapy to help address issues within these close relationships. Depending on the severity of the depression, significant improvement can be made in 10 to 15 sessions. **Group therapy** brings people with similar disorders together in a supportive environment to learn how others cope in similar situations.<sup>14</sup>

## Cognitive Behavioral Therapy

**Cognitive behavioral therapy (CBT)** is a form of psychotherapy that is effective for a range of problems including depression, anxiety disorders, alcohol and drug use problems, marital conflict, eating disorders, and severe mental illness. CBT helps a person to recognize distorted/negative thinking with the goal of changing thoughts and behaviors to respond to changes in a more positive manner.<sup>15</sup> Numerous research studies suggest that CBT leads to significant improvement in functioning and quality of life. Studies show that

13. American Psychiatric Association. (2020). *What is depression?*  
<https://www.psychiatry.org/patients-families/depression/what-is-depression>
14. American Psychiatric Association. (2020). *What is depression?*  
<https://www.psychiatry.org/patients-families/depression/what-is-depression>
15. American Psychiatric Association. (2020). *What is depression?*  
<https://www.psychiatry.org/patients-families/depression/what-is-depression>

CBT has been demonstrated to be as effective as, or more effective than, other forms of psychological therapy or psychiatric medications.<sup>16</sup>

CBT is based on these core principles<sup>17</sup>:

- Psychological problems are based, in part, on faulty or unhelpful ways of thinking.
- Psychological problems are based, in part, on learned patterns of unhelpful behavior.
- People suffering from psychological problems can learn better ways of coping with them, thereby relieving their symptoms and increasing quality of life.

CBT treatment involves efforts to change thinking patterns. These strategies might include the following<sup>18</sup>:

- Learning to recognize one's distortions in thinking that are creating problems, and then reevaluating them in light of reality.
- Gaining a better understanding of the behavior and motivation of others.

16. Clinical Practice Guideline for the Treatment of Posttraumatic Stress Disorder. (2017). *What is cognitive behavioral treatment?* American Psychological Association. <https://www.apa.org/ptsd-guideline/patients-and-families/cognitive-behavioral>

17. Clinical Practice Guideline for the Treatment of Posttraumatic Stress Disorder. (2017). *What is cognitive behavioral treatment?* American Psychological Association. <https://www.apa.org/ptsd-guideline/patients-and-families/cognitive-behavioral>

18. Clinical Practice Guideline for the Treatment of Posttraumatic Stress Disorder. (2017). *What is cognitive behavioral treatment?* American Psychological Association. <https://www.apa.org/ptsd-guideline/patients-and-families/cognitive-behavioral>

- Using problem-solving skills to cope with difficult situations.
- Learning to develop a greater sense of confidence in one's own abilities.

CBT treatment also usually involves efforts to change behavioral patterns. These strategies might include facing one's fears instead of avoiding them, using role-playing to prepare for potentially problematic interactions with others, and learning to calm one's mind and relax one's body.<sup>19</sup>

CBT aims to help clients develop skills to manage their feelings in healthy ways. Through in-session exercises and "homework" between sessions, clients develop coping skills, whereby they learn ways to change their own thinking and behavior, ultimately changing how they feel. CBT therapists focus on current situations, thought patterns, and behaviors rather than past events. A certain amount of information about one's history is needed, but the focus is primarily on developing more effective ways of coping with life moving forward.<sup>20</sup>

## DIALECTICAL BEHAVIOR THERAPY

**Dialectical behavior therapy (DBT)** is a type of cognitive behavioral therapy that provides clients with new skills to manage painful emotions and decrease conflict in relationships. It has been used successfully to treat people experiencing depression, bulimia, binge-eating, bipolar disorder, post-

19. Clinical Practice Guideline for the Treatment of Posttraumatic Stress Disorder. (2017). *What is cognitive behavioral treatment?* American Psychological Association. <https://www.apa.org/ptsd-guideline/patients-and-families/cognitive-behavioral>
20. Clinical Practice Guideline for the Treatment of Posttraumatic Stress Disorder. (2017). *What is cognitive behavioral treatment?* American Psychological Association. <https://www.apa.org/ptsd-guideline/patients-and-families/cognitive-behavioral>

traumatic stress disorder, borderline personality disorder, and substance abuse. DBT focuses on providing therapeutic skills in four key areas<sup>21</sup>:

- Mindfulness focuses on improving an individual's ability to accept and be present in the current moment.
- Distress tolerance is geared toward increasing a person's tolerance of negative emotion, rather than trying to escape from it.
- Emotion regulation strategies are used to manage and change intense emotions that are causing problems in a person's life.
- Interpersonal effectiveness techniques allow a person to communicate with others in a way that is assertive, maintains self-respect, and strengthens relationships.

## Electroconvulsive Therapy

**Electroconvulsive therapy (ECT)** is a medical treatment reserved for clients with severe major depression who have not responded to medications, psychotherapy, or other treatments. It involves a brief electrical stimulation of the brain while the client is under anesthesia. A client typically receives ECT two to three times a week for a total of 6 to 12 treatments. It is usually managed by a team of trained medical professionals, including a psychiatrist, an anesthesiologist, and a nurse.<sup>22</sup> See Figure 7.8<sup>23</sup> for an image showing ECT electrode placement.

21. Psychology Today. (n.d.). *Dialectical behavior therapy*.

<https://www.psychologytoday.com/us/therapy-types/dialectical-behavior-therapy>

22. American Psychiatric Association. (2020). *What is depression?*

<https://www.psychiatry.org/patients-families/depression/what-is-depression>

23. “[Electroconvulsive Therapy.png](#)” by [BruceBlaus](#) is licensed under [CC BY-SA 4.0](#)





Figure 7.8 Electroconvulsive Therapy

Nursing considerations regarding ECT include the following:

- Pre-procedure education
- Securement of informed consent
- Pre-procedure preparation
- Screening tools used pre- and post-ECT to evaluate side effects, including memory loss
- Medication administration
- Post-ECT care, such as monitoring vital signs and for changes in airway, breathing, and circulation, and implementing fall risk precautions post-anesthesia

A client must provide written informed consent before ECT is administered. In situations where a client is too ill to make decisions for themselves, the consent process is governed by state law (for example, a court-appointed guardian).<sup>24</sup>

Clients and their families should discuss all options for treatment with the psychiatrist before making a specific treatment decision. They should be provided with sufficient information to fully understand the procedure and

24. American Psychiatric Association. (2019). *What is electroconvulsive therapy (ECT)?* <https://www.psychiatry.org/patients-families/ect>



the potential benefits, risks, and side effects of each treatment option before providing written consent.<sup>25</sup>

General anesthesia is provided during ECT, so presurgical preparation is provided with typical dietary restrictions before the procedure. Typically, this means no food or water after midnight and only a sip of water with morning medications. An intravenous line is inserted, and electrode pads are placed on the head.


A client typically receives ECT two or three times a week for a total of 6 to 12 treatments, depending on the severity of symptoms and the response to treatment. At the time of each treatment, a client is given general anesthesia and a muscle relaxant, and electrodes are attached to the scalp at precise locations. The client's brain is stimulated with a brief controlled series of electrical pulses. This causes a seizure within the brain that lasts for approximately one minute. The patient is asleep for the procedure and awakens after 5-10 minutes, much as from minor surgery.<sup>26</sup>

ECT treatment has been associated with some risks such as short-term memory loss and difficulty learning. Some people have trouble remembering events that occurred in the weeks before the treatment or earlier. In most cases, memory problems improve within a couple of months. Some clients may experience longer-lasting problems, including permanent gaps in memory.<sup>27</sup>

25. American Psychiatric Association. (2019). *What is electroconvulsive therapy (ECT)?* <https://www.psychiatry.org/patients-families/ect>

26. American Psychiatric Association. (2019). *What is electroconvulsive therapy (ECT)?* <https://www.psychiatry.org/patients-families/ect>

27. American Psychiatric Association. (2019). *What is electroconvulsive therapy (ECT)?* <https://www.psychiatry.org/patients-families/ect>

 View the following YouTube video of ECT therapy<sup>28</sup>

## Transcranial Magnetic Stimulation

**Transcranial Magnetic Stimulation (TMS)** is a noninvasive procedure that uses magnetic fields to stimulate nerve cells in the brain to improve symptoms of depression. TMS is typically used when other depression treatments haven't been effective. It uses a magnet to activate the brain. Unlike electroconvulsive therapy (ECT), in which electrical stimulation is more generalized, TMS can be targeted to a specific site in the brain. A typical TMS session lasts 30 to 60 minutes and does not require anesthesia. During the procedure, an electromagnetic coil is held against the forehead near an area of the brain that is thought to be involved in mood regulation. Short electromagnetic pulses are administered through the coil. The magnetic pulses easily pass through the skull and cause small electrical currents that stimulate nerve cells in the targeted brain region. The magnetic field is about the same strength as that of a magnetic resonance imaging (MRI) scan. The person generally feels a slight knocking or tapping on the head as the pulses are administered. The muscles of the scalp, jaw, or face may contract or tingle during the procedure, and mild headaches or brief light-headedness may result after the procedure. It is also possible that the procedure could cause a seizure, although this adverse effect is uncommon. Because the treatment is relatively new, long-term side effects are unknown.<sup>29</sup>

28. Michigan Medicine. (2022). *ECT: Disrupting the Stigma Around an Essential Treatment Option* [Video]. YouTube. [https://youtu.be/qk\\_FjhitKDI?si=hwi32ffhJGvOGM-a](https://youtu.be/qk_FjhitKDI?si=hwi32ffhJGvOGM-a)

29. National Institute of Mental Health. (2016). Brain stimulation therapies. U.S.

Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/brain-stimulation-therapies/brain-stimulation-therapies>

## 7.5 Applying the Nursing Process to Depressive Disorders

The following subsections will describe how a nurse applies the nursing process and the National Council of State Board of Nursing Clinical Judgment Measurement Model (NCJMM) to caring for clients with depression. The stages of the NCJMM are listed in parentheses next to each stage of the nursing process.

### Assessment (Recognize Cues)

The initial assessment should determine if clinical depression exists, consider other conditions that might account for the client's symptoms, and assess for risk of harm to self or others.<sup>1</sup> Assessing a client with a depressive disorder focuses on both verbal and nonverbal assessments. As the nurse conducts follow-up assessments, findings are compared to baseline admission assessments. Assessment includes several components, such as a mental status examination, psychosocial assessment, cultural assessment, spiritual assessment, screening with validated tools, and review of laboratory testing results while also considering lifespan considerations.



The role of the nurse in caring for clients with depression is related to primary nursing care, as well as collaboration with interprofessional team members. As a team member, the nurse may collaborate with psychiatrists, psychologists, licensed social workers, and other health care providers. The scope and practice of each team member is clearly defined within their professional licensure.

1. Gaynes, B.N. (2024). Depression in adults: Clinical features and diagnosis. *UpToDate*. <https://www.uptodate.com/>

## Mental Status Examination

A psychiatric interview aims to collect information from the client as well as create a therapeutic relationship between the nurse and the client. The registered nurse uses specific questions during the client's admission process based on agency policy. See Table 7.5a for common findings when assessing a client with a depressive disorder. See expected findings for these components of a mental status examination in the "[Assessment](#)" section in Chapter 4. Critical findings that require immediate notification of the provider are bolded with an asterisk.

Table 7.5a Common Findings During A Mental Status Examination Of A Client With A Depressive Disorder

Mental Status Examination Component	Common Findings in Depressive Disorders (*Indicates immediately notify provider)
Level of Consciousness and Orientation	<ul style="list-style-type: none"> <li>• Disoriented/confused. The client may be very self-focused and may appear disoriented when others initiate conversation or talk about current events because they are unaware of what is occurring around them due to their depression.</li> </ul>
Appearance and General Behavior	<ul style="list-style-type: none"> <li>• Disheveled. The client's hair may not be combed, and they may be unwashed with poor dental care. They may be wearing dirty clothing with food stains and have body odor with little or no attention to self-care.</li> <li>• Sleep disturbances. The client may exhibit too much sleep (i.e., 14-18 hours daily) or have insomnia (i.e., less than 4 hours of sleep or in intervals of sleep).</li> <li>• Psychomotor retardation. The client may have a slow response with walking, talking, and reacting; may tend to stay put on the couch or in bed.</li> <li>• Vegetative signs. The client may exhibit weight loss, insomnia, constipation, and self-care deficits.</li> <li>• Avolition (reduced motivation or goal-directed behavior). For example, the client states, "I just don't want to shower today."</li> <li>• Anergia (low energy). For example, the client states, "I have no energy; I cannot get out of bed."</li> <li>• Social isolation.</li> <li>• Aggressiveness and agitation.</li> <li>• <b>*Verbal and nonverbal threats of harm.</b></li> <li>• <b>*Self-harming behaviors</b> such as cutting, picking at skin, knocking head against the wall, tightening string or items on wrists, or stabbing self with anything fashioned into a weapon.</li> <li>• <b>*"Cheeking pills"</b> (i.e., holding pills in mouth and not swallowing them). Clients may save medications for use in a later suicide attempt.)</li> </ul>

Speech	<ul style="list-style-type: none"> <li>• May speak slowly in a monotone.</li> <li>• May exhibit <b>latency</b> (i.e., a delayed response to a question or comment). However, be aware of the client's baseline because this may be their normal if they have a history of traumatic brain injuries, dementia, or are English language learners.</li> </ul>
Motor Activity	<ul style="list-style-type: none"> <li>• Psychomotor retardation.</li> <li>• Motor restlessness and anxiety.</li> <li>• Inactive or not initiating self-activity or care.</li> </ul>

## Mood and Affect

- **Apathy** (a lack of interest in events that one previously found enjoyable). For example, the client is no longer interested in participating in pleasurable activities. This may be verbalized or observed.
- Anxious, irritable, angry, euphoric, tearful, or depressed.
- **Labile mood** (rapid, exaggerated changes in mood). For example, a client is crying uncontrollably but when asked about the reason for crying, they state, "I don't know, I have no reason but can't stop crying."
- **Dysphoria** (a state of unease or dissatisfaction). For example, the client states, "I don't like it here."
- Sadness.
- Crying episodes.
- Flat or blunted affect.
- Constricted/restricted affect. For example, a client usually laughs at jokes and giggles at funny comments but now does neither.
- **Incongruency** (a lack of alignment between response and actions). For example, a client responds that they are fine, yet their body language is curled in a fetal position with no eye contact, and they are mumbling.
- Mood is inappropriate for the current situation.
- **\*Hopelessness**. For example, a client may feel there is no longer any hope for them getting better or for life improving.
- **\*Worthlessness**. For example, a client may have feelings of guilt regarding themselves and their depression. They may feel like a burden to others and are unable to recognize their own value.
- **\*Helplessness**. For example, a client does not feel in control of life events.

**\*Note: Hopelessness, worthlessness, and helplessness are related to an increased risk of self-injury behavior and suicide and must be reported to provider. Do not leave clients alone if statements such as these are being made.**



<b>Thought and Perception</b>	<ul style="list-style-type: none"> <li>• <b>Poverty of content</b> (responds without saying anything substantive or says much more than is necessary to convey a message). For example, if a client is asked, “How did you sleep last night?,” the client answers with a long response about different brands of bedsheets without answering the question.</li> <li>• Decreased attention span.</li> <li>• Obsessions/preoccupations/ruminations. For example, a client may dwell on negative aspects of self-concept or faults and failures because they are unable to focus on anything else and thus repeat these thoughts often.</li> <li>• <b>*Suicidal ideation.</b></li> <li>• <b>*Homicidal ideation.</b></li> <li>• <b>*Violence ideation.</b></li> </ul> <p><b>*Note: Suicidal, homicidal, and violence ideations are characteristics of depression with recurring thoughts of death. These types of comments indicate increased risk for self-injury, suicide, or injury to others and must be reported to provider. Do not leave clients alone if statements such as these are being made.</b></p>
<b>Attitude and Insight</b>	<ul style="list-style-type: none"> <li>• Unaware of the current situation.</li> <li>• Unable to relate to current trends or settings.</li> <li>• Lack of perception.</li> <li>• Low awareness outside of self.</li> <li>• Irritability.</li> </ul>
<b>Cognitive Abilities and Level of Judgment</b>	<ul style="list-style-type: none"> <li>• The inability to make decisions, think clearly, or solve problems is common in clients with depression.</li> </ul>

## Psychosocial Assessment

As previously discussed in the “[Application of the Nursing Process in Mental](#)

Health Care” chapter, a psychosocial assessment obtains additional subjective data that detects risks and identifies treatment opportunities and resources.<sup>23</sup>

- Reason for seeking health care (i.e., “chief complaint”)
- Thoughts of self-harm or suicide (both current and historical)
- Cultural assessment
- Spiritual assessment
- Family dynamics
- Current and past medical history
- Current medications
- History of previously diagnosed mental health disorders
- Previous hospitalizations
- Educational background
- Occupational background
- History of exposure to psychological trauma, violence, and domestic abuse
- Substance use (tobacco, alcohol, recreational drugs, misused prescription drugs)
- Family history of mental illness
- Coping mechanisms
- Functional ability/Activities of daily living

After identifying the reason the client is seeking health care, additional focused questions are used to obtain detailed information used to plan care. The mnemonic PQRSTU can be used to ask questions in an organized

2. Clinical Practice Guideline for the Treatment of Posttraumatic Stress Disorder. (2017). *What is cognitive behavioral treatment?* American Psychological Association. <https://www.apa.org/ptsd-guideline/patients-and-families/cognitive-behavioral>
3. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

fashion. See Table 7.5b for a sample PQRST assessment for depression and sample responses by a client.

Table 7.5b Sample PQRSTU Questions for Assessing Depression

PQRSTU	Sample Questions	Sample Client Response
<b>Provocation/ Palliation</b>	“What makes your low mood or sadness better or worse?”	“It gets worse when I’m alone or when I think about things I haven’t done right. Sometimes walking outside or talking to a friend helps, but I rarely have the energy to try.”
<b>Quality</b>	“Can you describe what your low mood or sadness feels like?”	“I feel numb, tired, and like I’m just going through the motions. Nothing feels enjoyable anymore.”
<b>Region</b>	“Do you feel any physical symptoms associated with your low mood?”	“Sometimes it feels like a heavy weight on my chest and shoulders. I get a lot of body aches, especially in my back and shoulders. My legs feel tired and my head feels cloudy.”
<b>Severity</b>	“On a scale of 0 to 10, how would you rate how bad your sadness or low mood feels right now?”	“It’s about a 7 on most days. On bad days it’s a 9 and I feel like I can’t pull myself out of bed.”
<b>Timing/ Treatment</b>	“When did this sadness or low mood start? How often do you feel this way? Does it come and go or stay constant?”	“It started about six months ago after I lost my job. I feel like this nearly every day. Mornings are the worst, but some days it never really lifts.”
<b>Understanding</b>	“What do you think is causing your sadness or low mood? How do you make sense of these feelings?”	“I think it’s because I lost my job and haven’t felt useful since. I feel like a failure at everything I do.”

## SUICIDE AND SELF INJURY SCREENING

Clients being evaluated or treated for depression often have suicidal ideation. It is important for the nurse to introduce suicide screening in a way that helps the client understand its purpose and normalize questions that might otherwise seem intrusive. The Patient Safety Screener (PSS-3) is an example of a brief screening tool to detect suicide risk in all client presenting to acute care settings.<sup>4</sup>

**Non-suicidal self-injury (NSSI)** refers to intentional self-inflicted destruction of body tissue without suicidal intention and for purposes not socially sanctioned. Common forms of NSSI include behaviors such as cutting, burning, scratching, and self-hitting. It is considered a maladaptive coping strategy without the desire to die. NSSI is a common finding among adolescents and young adults in psychiatric inpatient settings.<sup>5</sup>

Review information about suicide screening, the Patient Safety Screener, and screening for non-suicidal self-injury (NSSI) in the “[Assessment](#)” section of the *Applying the Nursing Process to Mental Health Care* chapter.

## CULTURAL ASSESSMENT

Cultural Formulation Interview (CFI) questions help nurses understand a client’s cultural background and how it influences their experience of mental health symptoms, including depression.<sup>6</sup> Sample CFI questions focused

4. Suicide Prevention Resource Center. (n.d.). *The patient safety screener: A brief tool to detect suicide risk*. <https://sprc.org/micro-learning/patientsafetyscreener>
5. Cipriano, A., Cella, S., & Cotrufo, P. (2017). Nonsuicidal self-injury: A systematic review. *Frontiers in Psychology*, 8. <https://www.frontiersin.org/articles/10.3389/fpsyg.2017.01946/full>
6. DeSilva, R., Aggarwall, N. K., & Lewis-Fernandez, R. (2015). The DSM-5 cultural formulation interview and the evolution of cultural assessment in psychiatry.

specifically on understanding depression within a cultural context include the following:

- Cultural Definition of the Problem
  - “Can you describe what you’re experiencing emotionally? How would you name or label this problem?”
  - “What do people in your family or community call these kinds of feelings or mood?”
  - “How does your culture or family view feeling sad or losing interest in life?”
- Cultural Perceptions of Cause, Context, and Support
  - “What do you think caused your sadness or low mood?”
  - “Are there events in your life, such as stress or loss, that you think may have contributed to these feelings of sadness or low mood?”
  - “Do you have any cultural or spiritual beliefs that help explain what you’re going through?”
- Cultural Factors Affecting Coping and Help-Seeking
  - “What kinds of things have you done to cope with these feelings so far?”
  - “Are there any traditional remedies, rituals, or religious practices you use to feel better?”
  - “Have you tried to talk to anyone about these feelings, like family members, friends, religious leaders, or traditional healers?”
- Cultural Features of the Nurse–Client Relationship
  - ““Is there anything I should know about your background or beliefs that would help me better understand you?”
  - “Do you have any concerns or hesitations you have about seeing a mental health professional?”

*Psychiatric Times*, 32(6). <https://www.psychiatrictimes.com/view/dsm-5-cultural-formulation-interview-and-evolution-cultural-assessment-psychiatry>

- “What kind of help do you think would work best for you?”

Findings from the cultural formulation interview are used to individualize a client’s plan of care to their preferences, values, beliefs, and goals.

### **Reflective Questions**

Reflect on how a client’s cultural values, beliefs, or preferences impact presenting symptoms, the nursing care plan, or treatment modalities. Consider the following components:

- Religious/Spiritual beliefs
- Language/Communication
- Nutritional preferences
- Fasting
- Potential drug-food interactions
- Rituals/Customs/Practices
- Gender dysphoria

## **SPIRITUAL ASSESSMENT**

The FICA Spiritual History Tool is a widely used assessment model for evaluating a client’s spiritual beliefs and how they may influence health, illness, and coping. FICA© is a mnemonic for the domains of Faith, Importance, Community, and Address in Care.<sup>7</sup> The data obtained from a FICA assessment can be helpful in understanding how clients with depression draw on spirituality or religion for support—or how spiritual distress may be contributing to feelings of depression. Spiritual distress is very common for

7. GW School of Medicine & Health Sciences. (n.d.). *Clinical FICA tool*.  
<https://smhs.gwu.edu/spirituality-health/program/transforming-practice-health-settings/clinical-fica-tool>

clients experiencing serious illness, injury, loss, or the dying process, and nurses assist clients to adopt healthy coping strategies to deal with these life events. Addressing a client's spirituality and advocating spiritual care have been shown to improve clients' health and quality of life.<sup>8,9</sup>

Table 7.5c summarizes a sample spiritual assessment questions and sample responses from a client experiencing depression.

Table 7.5c Sample FICA Spiritual Assessment Questions for Clients with Depression

8. Pilger, C., Molzahn, A. E., de Oliveira, M. P., & Kusumota, L. (2016). The relationship of the spiritual and religious dimensions with quality of life and health of patients with chronic kidney disease: An integrative literature review. *Nephrology Nursing Journal: Journal of the American Nephrology Nurses' Association*, 43(5), 411–426. <https://pubmed.ncbi.nlm.nih.gov/30550069/>
9. Puchalski, C., Jafari, N., Buller, H., Haythorn, T., Jacobs, C., & Ferrell, B. (2020). Interprofessional spiritual care education curriculum: A milestone toward the provision of spiritual care. *Journal of Palliative Medicine*, 23(6), 777–784. <https://doi.org/10.1089/jpm.2019.0375>

Domain	Sample Assessment Question	Sample Client Response
<b>Faith</b>	“Do you consider yourself spiritual or religious? What gives your life meaning?”	“I was raised in a religious family that went to church every Sunday, but lately I’ve felt disconnected. I used to find comfort in prayer, but now I don’t even feel like trying.”
<b>Importance</b>	“What importance does your faith or belief have in your life? Has it influenced how you cope with sadness or low mood?”	“My faith used to mean a lot to me, but lately it’s been hard to feel hopeful that things will get better.”
<b>Community</b>	“Are you part of a spiritual or religious community? Does participation in this community provide support when you’re feeling sad or in a low mood?”	“I used to go to church regularly and even went to a prayer group, but I’ve stopped attending. I don’t want people to see me like this.”
<b>Address in Care</b>	“How would you like me (or the health care team) to address spiritual issues during your care? Would you like to speak with a chaplain?”	“I would talk with a chaplain if they will just listen and not impose their faith on me.”

Feelings of guilt, abandonment, and spiritual distress are common in individuals experiencing depression and may compound their symptoms. Nurses may recognize cues of spiritual distress and offer to connect the client with a chaplain or spiritual care services. Spiritual goals may be included in the nursing care plan if the client finds them valuable.

## FAMILY DYNAMICS

Family dynamics are included in a psychosocial assessment, especially for children, adolescents, and older adults. **Family dynamics** refers to the patterns of interactions among relatives, their roles and relationships, and the various factors that shape their interactions. Because family members rely on each other for emotional, physical, and economic support, they are primary sources of relationship security or stress. Family dynamics and the quality of family relationships can have either a positive or negative impact on an individual’s health. For example, secure and supportive family relationships



can provide love, advice, and care, whereas stressful family relationships can be burdened with arguments, unhealthy relationships, and a lack of support.<sup>10</sup>

Unhealthy family dynamics can cause children to experience trauma and stress as they grow up. This type of exposure, known as adverse childhood experiences (ACEs), is linked to an increased risk of developing physical and mental health problems such as heart, lung, and liver disease, depression, and anxiety. Unhealthy family dynamics also correlate with an increased risk of substance use and addiction among adolescents.<sup>11</sup> Review information about adverse childhood experiences (ACEs) in the “[Mental Health and Mental Illness](#)” section of Chapter 1.

## Screening Tools

Screening tools assess characteristics of specific mental health disorders. The screening tools listed below are examples of screenings, assessments, and question/answer prompts designed to address depressive disorders. These screening tools may be used on admission and at different times throughout the hospital or treatment stay. The findings may be used to compare and contrast client progress within the hospital stay, from a previous admission, or periodically on an outpatient basis. The registered nurse often conducts these tools as a collaborative member of the health care team.

### **Links to Common Screening Tools for Depressive Disorders**

10. Jabbari, B., Schoo, C., & Rouster, A. S. (2023). *Family dynamics*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK560487/>

11. Jabbari, B., Schoo, C., & Rouster, A. S. (2023). *Family dynamics*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK560487/>

- ▶ **Columbia-Suicide Severity Rating Scale (C-SSRS) PDF:** A rating scale for suicidal ideation and behaviors that rates the degree of risk or intent of harm. It can be a self-assessment or administered by the health care professional.
- ▶ **Patient Health Questionnaire-9 (PHQ-9):** A quick screening tool with nine criteria for assessing a client's risk of depression.
- ▶ **Beck Depression Inventory (BDI):** A 21-item self-assessment questionnaire that determines the severity of depression from none to severe.
- ▶ **Hamilton Depression Scale (HDRS) PDF:** A 17-item questionnaire used to rate the severity of one's depression.
- ▶ **Geriatric Depression Scale (GDS) PDF:** A self-report of depressive symptoms for older adults; the new version is 15 questions.
- ▶ **Edinburgh Postnatal Depression Scale (EPDS):** A self-report of ten statements by mothers to screen for postpartum depression.

## Laboratory Testing

There is no specific blood test to diagnose depression, but health care providers often order laboratory tests to rule out other conditions that can mimic depression symptoms, like anemia or thyroid disease. Testing may also be performed to assess overall health and kidney/liver function, especially if medication will be prescribed. Common laboratory tests for new onset depression include complete blood count, chemistry panels, kidney and liver function tests, Vitamin B12 levels, urinalysis, thyroid stimulating hormone,

rapid plasma reagin (RPR) for syphilis, human chorionic gonadotropin (HCG) for pregnancy, and toxicology screening for drug use.<sup>12</sup>

## Life Span Considerations

Life span considerations influence how the client is assessed, as well as the selection of appropriate nursing interventions. Depressive disorders can be found across the life span from the very young to the very old. It is important to individualize all interventions to the age and developmental level of the client. Review developmental stages in the “[Application of the Nursing Process in Mental Health Care](#)” chapter.

## CHILDREN AND ADOLESCENTS

Children may not verbalize sadness. Instead, depression often presents as irritability, frequent temper tantrums, somatic complaints (e.g., stomachaches, headaches), withdrawal from school, friends, or family, or behavioral changes like clinginess, acting out, or poor academic performance. It is important to ask children and adolescents who are withdrawn or sad about thoughts of suicide or self-harm. Adolescents may perceive a single disappointment (such as a relationship break-up) as so catastrophic they feel suicidal or begin to hurt themselves. Nurses can use play-based or art-based therapeutic techniques to facilitate a therapeutic nurse-client relationship. If a school-aged child or adolescent is suspected to have undiagnosed depression, a nurse or school counselor can refer them to a mental health professional to conduct a comprehensive assessment and plan effective treatments.

12. Chand, S. P., & Arif, H. (2023). *Depression*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK430847/>

## OLDER ADULTS

Depression in older adults may manifest differently as somatic symptoms (e.g., fatigue, pain, GI issues), anxiety, irritability, reduced verbal expression of sadness, or cognitive changes that may be mistaken for dementia.

Furthermore, individuals with dementia are at higher risk for depression.

### Depression Associated With Dementia

**Dementia** refers to a group of symptoms that lead to a progressive, irreversible decline in mental function severe enough to disrupt daily life caused by a group of conditions including Alzheimer's disease, vascular dementia, frontal-temporal dementia, and Lewy body disease. Alzheimer's disease is one of the most common forms of dementia. Alzheimer's disease causes impaired memory and the ability to learn, reason, make judgments, communicate, and carry out daily activities. An early symptom of Alzheimer's disease can be subtle memory loss and personality changes that differ from normal age-related memory problems. They seem to tire or become upset or anxious more easily. They do not cope well with change. For example, they can follow familiar routes, but traveling to a new place confuses them, and they can easily become lost. In the early stages of the illness, people with Alzheimer's disease are particularly susceptible to depression.<sup>13</sup>

While changes in the brain that cause dementia are permanent and worsen over time, thinking and memory problems can be

13. American Psychiatric Association. (2019). *What is Alzheimer's disease?* <https://www.psychiatry.org/patients-families/alzheimers/what-is-alzheimers-disease>

aggravated by untreated depression.<sup>14</sup> Nurses should report new symptoms of depression in clients who have been diagnosed with dementia.

- ▶ Read more about dementia at the [Alzheimer's Association's](https://www.alz.org/alzheimers-dementia/what-is-dementia) webpage.

### Reflective Questions

How does a nurse differentiate between symptoms that could indicate depression, delirium, dementia, or psychosis?

1. What are some common underlying medical conditions that could potentially mimic the symptoms of depression or mania in those who are elderly?
2. What other symptoms might a client who is a child/adolescent display that would indicate the need to assess for disorders other than depression?

## Diagnosis (Analyze Cues)

Mental health disorders are diagnosed by health providers using the *DSM-5-TR*, similar to how medical conditions are diagnosed by trained medical professionals. Nurses create individualized nursing care plans based

<sup>14</sup>. Alzheimer's Association. (n.d.) *What is dementia?* <https://www.alz.org/alzheimers-dementia/what-is-dementia>

on the client’s response to mental health disorders. See common nursing diagnoses related to mental health disorders in the “[Diagnosis](#)” section of the “Application of the Nursing Process in Mental Health Care” chapter.

Risk for suicide is always evaluated for clients with depressive disorders because suicidal ideation is a symptom of depression. Other common nursing diagnoses with sample expected outcomes for clients with depressive disorders are discussed in the following section in Table 7.5d.

## Outcome Identification (Generate Solutions)

SMART outcomes are identified in relation to the established nursing diagnoses for each client. SMART is an acronym for Specific, Measurable, Attainable/Actionable, Relevant, and Timely. Read more about outcomes identification in the “[Application of the Nursing Process in Mental Health Care](#)” chapter. Table 7.5d provides common nursing diagnoses and sample expected outcomes for each nursing diagnosis.

Table 7.5d. Common Expected Outcomes for Nursing Diagnoses Related to Depressive Disorders<sup>15</sup>

15. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

Nursing Diagnosis <sup>16,17</sup>	Sample Expected Outcomes
<b>Risk for Suicide</b>	<p>The client will communicate feelings and thoughts of suicide to the health care team, prior to acting on thoughts, during their inpatient stay.</p> <p><b>*Note: Clients with depression are at higher risk of suicide when experiencing sudden euphoric recovery from major depression.</b><sup>18</sup></p>
<b>Ineffective Coping/Readiness for Enhanced Coping</b>	<p>The client will identify effective coping strategies within 24 hours of admission.</p> <p>The client will engage in preferred stress management techniques by Day 3 of admission.</p>
<b>Self-Neglect</b>	The client will increase participation in baseline personal care each day during their stay.
<b>Fatigue/Sleep Deprivation</b>	The client will, within one week, report feeling rested upon awakening.
<b>Imbalanced Nutrition: Less than Body Requirements</b>	The client will eat 50% or more of their meal tray at each meal.
<b>Constipation</b>	The client will have a soft, formed stool at least every three days during their inpatient stay.
<b>Social Isolation</b>	The client will communicate with others during their inpatient stay by participating in daily group offerings within the milieu.
<b>Chronic Low Self-Esteem</b>	The client will verbalize at least three personal strengths within three days of admission.
<b>Hopelessness</b>	The client will describe plans for a positive future by discharge.
<b>Spiritual Distress</b>	The client will identify a meaning and purpose in life within two weeks.
<b>Readiness for Enhanced Knowledge</b>	The client will verbalize three common side effects of their medications by the end of the shift.

16. Ackley, B., Ladwig, G., Makic, M. B., Martinez-Kratz, M., & Zanotti, M. (2020). *Nursing diagnosis handbook: An evidence-based guide to planning care* (12th ed.). Elsevier.

# Planning (Generate Solutions)

## Safety

Safety receives top priority when planning and implementing interventions for clients with depression. Clients with depressive disorders are monitored closely for risk of suicide, and interventions are planned according to their level of risk. Review interventions for clients at risk of suicide in the [“Application of the Nursing Process in Mental Health Care”](#) chapter.

## Phases of Treatment and Recovery

As discussed earlier in this chapter, a combination of pharmacological treatments and psychotherapies are often an effective approach to treating depressive disorders. There are three phases in treatment and recovery from major depression<sup>19</sup>:

- The active phase (6 to 12 weeks) is directed at reduction of depressive symptoms and restoration of psychosocial and work function. Hospitalization may be required, and medication and other biological treatments may be initiated.
- The continuation phase (4 to 9 months) is directed at prevention of relapse through pharmacotherapy, education, and depression-specific psychotherapy. This phase focuses on maintaining the client as a functional and contributing member of the community after recovery

17. Herdman, T. H., & Kamitsuru, S. (Eds.). (2018). *Nursing diagnoses: Definitions and classification, 2018-2020*. Thieme Publishers New York.

18. Ackley, B., Ladwig, G., Makic, M. B., Martinez-Kratz, M., & Zanotti, M. (2020). *Nursing diagnosis handbook: An evidence-based guide to planning care* (12th ed.). Elsevier.

19. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.



from the acute phase.

- The maintenance phase (1 year or more) is directed at preventing future episodes of depression. Medication may be phased out or continued.

Nurses target interventions based on the client's current phase of treatment and recovery, their current nursing diagnoses, and established expected outcomes.

## Implementation (Take Action)

### Nursing Interventions for Depression Based on Categories of the APNA Implementation Standard

Nursing interventions for clients with depressive disorders can be categorized based on the American Psychiatric Nurses Association (APNA) standard for *Implementation* that includes the *Coordination of Care; Health Teaching and Health Promotion; Pharmacological, Biological, and Integrative Therapies; Milieu Therapy; and Therapeutic Relationship and Counseling*. Read more about these subcategories in the "[Application of the Nursing Process in Mental Health Care](#)" chapter. See examples of interventions for each of these categories for clients with depressive disorders in Table 7.5e.

Table 7.5e Examples of Nursing Interventions for Clients with Depressive Disorders Based on Subcategories of APNA Implementation Standard

Subcategory of the APNA Standard of Implementation	The nurse will ...	Rationale
<b>Coordination of Care</b>	<ul style="list-style-type: none"> <li>– Collaborate with mental health professionals (e.g., therapists, psychiatrists) to ensure continuity of care.</li> <li>– Communicate client assessment data, such as suicidal ideation or checking of medications, with interprofessional team members.</li> <li>– Facilitate referrals to community support resources such as support groups or home health.</li> <li>– Assist with care transitions (e.g., hospital discharge to long-term care or other community-based facilities).</li> </ul>	Interdisciplinary collaboration ensures a comprehensive treatment plan and helps reduce relapse risk through continued support. All team members providing care must be aware of the client's suicide risk to maintain a safe environment.
<b>Health Teaching and Health Promotion</b>	<ul style="list-style-type: none"> <li>-Educate the client and family members about depression, symptoms, and treatment options.</li> <li>– Promote lifestyle modifications such as regular sleep, exercise, and healthy nutrition.</li> <li>– Address stigma and normalize help-seeking.</li> </ul>	Empowering clients with knowledge improves self-management and treatment adherence. Health promotion activities can help alleviate mild to moderate depression symptoms and reduce recurrence.

<b>Pharmacological, Biological, and Integrative Therapies</b>	<ul style="list-style-type: none"> <li>-Administer prescribed antidepressants and monitor for side effects (e.g., GI distress, suicidality, serotonin syndrome). Open all medications in front of the client.</li> <li>– Provide health teaching on medication purpose, effects, adherence, and expected timeframes for improved symptoms.</li> <li>-Promote positive coping strategies such as journaling, meditation, and yoga.</li> <li>– Encourage adherence to treatment (therapy, medications).</li> </ul>	<p>Client understanding of their medications and potential side effects can increase medication compliance. Opening all medications in front of the client may decrease paranoia, if present. Nurses encourage resilience by adopting positive coping strategies.</p>
<b>Milieu Therapy</b>	<ul style="list-style-type: none"> <li>-Provide a structured, safe, and supportive environment.</li> <li>– Encourage participation in group therapy or therapeutic recreation.</li> <li>– Use daily routines and posted schedules to increase stability and reduce decision fatigue.</li> <li>-Perform and document intentional rounding every 15 to 60 minutes on a varied schedule.</li> </ul>	<p>A therapeutic milieu fosters a sense of safety, belonging, and predictability, which helps stabilize mood and reduce isolation. Visually rounding on every client in the milieu creates a strong safety plan for all clients and staff. Following a varied schedule prevents the client from anticipating when staff will check on them, which can reduce the ability to plan for suicide/risky behaviors.</p>

<b>Therapeutic Relationship and Counseling</b>	<ul style="list-style-type: none"> <li>– Use active listening, empathy, and validation to build trust and a supportive nurse-client relationship.</li> <li>– Explore cognitive distortions through supportive dialogue.</li> <li>– Facilitate goal-setting and positive affirmations.</li> <li>– Encourage verbalization of feelings, especially guilt, hopelessness, or suicidal thoughts.</li> <li>– Refer to spiritual supports such as chaplains, based on client preferences.</li> <li>– Review additional communication interventions in the “Communication Tips” subsection below.</li> </ul>	Providing effective therapeutic techniques for clients with depression can promote hope and positive self-esteem.
--	--	---

## Nursing Interventions for Physiological Signs of Depression

Nursing interventions are also planned that target common physiological signs of depression and associated self-care deficits. See common interventions for these conditions in Table 7.5f.

Table 7.5f Nursing Interventions Targeting Physiological Signs of Depression and Self-Care Deficit<sup>20</sup>

20. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

Problem/Intervention	Rationale
<p><b>Nutrition</b></p> <ul style="list-style-type: none"> <li>• Offer small, high-calorie, and high-protein snacks and fluids frequently.</li> <li>• When possible, encourage family or friends to join the client during meals.</li> <li>• Encourage the client to participate in selecting food and drinks.</li> <li>• Teach about nutrient-dense foods that support mood (e.g., omega-3s, B vitamins).</li> <li>• Refer the client to a dietician if necessary.</li> <li>• Weight the client weekly and monitor trends.</li> <li>• Observe the client's eating patterns, maintain safety. <i>(In the hospital setting, client's that are suicidal are given their food on a tray with plastic silverware, food in Styrofoam containers, etc. to reduce the risk of injury to self/others).</i></li> </ul>	<p>Poor nutrition increases the risk for physical illness and improving nutrition can help stabilize energy and mood. Small, frequent snacks are more easily tolerated than large portions of food if the client has a loss of appetite. Fluids prevent dehydration and minimize constipation.</p> <p>Eating is a social event. Eating with loved ones reinforces the idea that someone cares about them and can serve as an incentive to eat. The client is more likely to eat foods they prefer. A dietician can help create an individualized diet plan. Monitoring the client's status provides data for evaluating effectiveness.</p>

<p><b>Sleep</b></p> <ul style="list-style-type: none"> <li>• <b>Assess sleep patterns and disturbances.</b></li> <li>• <b>Encourage consistent sleep-wake cycles and limit daytime naps. Provide periods of rest after activities, if needed.</b></li> <li>• <b>Encourage the client to get up and dress and stay out of bed during the day.</b></li> <li>• <b>Encourage relaxation measures in the evening, such as a warm bath, warm milk, or soothing music.</b></li> <li>• <b>Avoid caffeinated beverages.</b></li> </ul>	<p>Sleep disruption is a hallmark of depression and worsens emotional regulation. Promoting good sleep hygiene helps restore circadian rhythm and reduce fatigue. Minimizing sleep during the day and establishing routines increase the likelihood of restful sleep at night. Relaxation techniques induce sleep. Decreasing caffeine intake increases the possibility of sleep.</p>
<p><b>Elimination (Constipation)</b></p> <ul style="list-style-type: none"> <li>• <b>Monitor frequency of bowel movements.</b></li> <li>• <b>Encourage fluids and foods high in fiber.</b></li> <li>• <b>Provide periods of exercise.</b></li> <li>• <b>Evaluate medications for side effects affecting elimination (e.g., TCAs, SSRIs).</b></li> <li>• <b>Support toileting routines and privacy.</b></li> <li>• <b>Evaluate the need for a bowel management program with stool softeners and laxatives.</b></li> </ul>	<p>Many depressed clients are constipated, so frequency of bowel movements should be monitored. Fluids, fiber, and exercise stimulate peristalsis and soften stools. Bowel management programs may be needed to avoid constipation or fecal impaction. Addressing elimination improves physical comfort and self-esteem.</p>

<p><b>Fatigue/Energy Deficit</b></p> <ul style="list-style-type: none"> <li>• <b>Schedule activities during periods of higher energy (usually morning).</b></li> <li>• <b>Promote light physical activity, such as walking or stretching.</b></li> <li>• <b>Provide rest periods without encouraging excessive sleeping.</b></li> <li>• <b>Use motivational interviewing to support gradual engagement according to the treatment plan.</b></li> </ul>	<p>Depression often causes feelings of extreme fatigue. Light exercise improves endorphin levels and sleep quality. Motivational interviewing helps encourage clients to set personal goals and participate in the treatment plan.</p>
<p><b>Self-Care Deficits</b></p> <ul style="list-style-type: none"> <li>• <b>Assess ability to perform ADLs (e.g., bathing, grooming, dressing).</b></li> <li>• <b>Offer verbal prompts, encouragement, and step-by-step support. When appropriate, give step-by-step reminders, such as “Wash the right side of your face and now your left.”</b></li> <li>• <b>Establish a simple daily hygiene routine.</b></li> <li>• <b>Reinforce successes positively.</b></li> </ul>	<p>Depression impairs motivation and concentration, leading to neglect of self-care. Slowed thinking and difficulty concentrating make organizing simple tasks difficult. Supporting ADLs restores self-worth and promotes dignity. Being clean and well-groomed can improve self-esteem.</p>

## Communication Tips for Clients with Depressive Disorders

Some clients with depression may be so withdrawn they are unwilling or unable to speak. Sitting with them in silence may feel like a waste of time, but nurses should be aware that providing therapeutic presence can be meaningful in supporting the client with depression. Helpful communication techniques for clients with depression and their rationale are described in the following box.

## Communication Tips for Clients with Depressive Disorders<sup>21</sup>

- Use a calm, soft, and patient tone.
  - **Rationale:** Depressed individuals may feel overwhelmed or hopeless; a soft tone reduces pressure and invites connection.
- Allow extra time for responses.
  - **Rationale:** Psychomotor retardation (slowed thinking or movement) is common in depression. Waiting without rushing encourages the client to participate without feeling inadequate.
- Use simple, concrete words and allow the client time to respond.
  - **Rationale:** Depression can impair concentration and processing. Short, straightforward phrases are easier to understand and less cognitively demanding.
- Acknowledge and validate feelings.
  - **Rationale:** Statements like “It sounds like things have been really difficult for you” show empathy and help the client feel seen and understood,

21. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.



rather than judged.

- Be alert for signs of suicidal ideation and ask directly.
  - **Rationale:** Use nonjudgmental phrasing like, “Sometimes people with depression have thoughts of not wanting to live. Have you felt that way?” Asking does not increase risk for suicide and opens a vital safety dialogue. People often experience relief and decreased feelings of isolation when they share thoughts of suicide.
- Avoid platitudes such as, “Just think positive,” “Everyone feels down once in a while,” or “Just snap out of it.”
  - **Rationale:** Platitudes invalidate the individual’s feelings and can increase feelings of guilt or worthlessness because they cannot “snap out of it.”
- Normalize their experience.
  - **Rationale:** Letting clients know that “many people with depression feel this way” helps reduce shame and stigma, and reassures them they are not alone.
- When a client is silent, use the technique of making observations, such as “There are new pictures on the wall,” or “You are wearing new shoes.”
  - **Rationale:** When an individual is not ready to talk, direct questions can raise their anxiety

levels. Respect the client's silence or limited responses. Pointing out objects in the environment can draw the person into reality. Being present without pressuring them to talk builds trust and shows unconditional support.

- Offer hope in small, realistic ways.
  - **Rationale:** Gently reinforce that treatment is available and recovery is possible, e.g., “There are things we can try that have helped others feel better.”

Nurses counsel individuals with depression to help them explore positive coping strategies<sup>22</sup>:

- Encourage stress management techniques such as exercise, good sleep, and healthy food choices.
- Promote the formation of supportive relationships such as peer support and support groups to reduce social isolation and enable the individual to work on personal goals and relationship needs.
- Provide information about spiritual support as the individual defines it, such as chaplain or pastoral visits or spending time in nature; many people find strength and comfort in spiritual and/or religious activities.
- Help the client reconstruct a healthier and more hopeful attitude about the future (without providing false reassurance).

22. Halter, M. (2022). *Varc Carolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

# Collaborative Mental Health Treatments

Nurses assist in implementing collaborative interventions based on the client's treatment plan. Review collaborative mental health treatments and common medications used to treat depression in the "[Treatments for Depression](#)" section of this chapter.

## Client Education Regarding Antidepressant Medications

Nurses educate clients about their medications, including the manner in which they work, common side effects, and issues to report to their provider. Clients taking antidepressants should also be educated regarding the following considerations<sup>23</sup>:

- When taking antidepressants, it is important to follow the instructions on how much to take. Some people start to feel better a few days after starting the medication, but it can take four to eight weeks to feel the most benefit. Antidepressants work well and are safe for most people, but it is still important to talk with your mental health care provider if you have side effects such as sexual dysfunction, weight gain, dizziness, nausea, palpitations, drowsiness, insomnia, or anxiety. Side effects may go away as your body adjusts to the medication, but in some cases, switching to a different medication may be required.
- Don't stop taking an antidepressant without first talking to your health care provider. Stopping your medicine suddenly can cause symptoms or worsen depression.
- Antidepressants cannot solve all of your problems. Antidepressants work best when combined with psychotherapy and healthy coping strategies. If you notice that your mood is getting worse or if you have thoughts

23. Centers for Disease Control and Prevention. (2021). *Mental health conditions: Depression and anxiety*. <https://www.cdc.gov/tobacco/campaign/tips/diseases/depression-anxiety.html>

about hurting yourself, it is important to call your provider right away.

- Some people who are depressed may think about hurting themselves or committing suicide (taking their own life). If you are having thoughts about committing suicide, please seek immediate help by calling your provider, 911, or 1-800-273-TALK to reach a 24-hour crisis center that provides free, confidential help to people in crisis.
- Some antidepressants may cause risks to the baby during pregnancy. Talk with your provider if you are pregnant or might be pregnant or if you are planning to become pregnant.
- For individuals who are very depressed or suicidal, it is important to provide close monitoring when the individual first starts taking an antidepressant medication. Often an individual may have increased energy to make a suicidal attempt when they first begin a medication, whereas previously they may have had suicidal thoughts but lacked the energy to make an attempt.

## Supporting Family Members

It is important to support the family members and significant others who are living with an individual with a depressive disorder. Read tips on living with someone with depression in Figure 7.9.<sup>24</sup>

24. “2\_living\_with\_someone\_with\_depression.png” by unknown author for [World Health Organization](https://www.who.int/campaigns/world-mental-health-day/2021/campaign-materials) is in the [Public Domain](https://www.who.int/campaigns/world-mental-health-day/2021/campaign-materials). Access for free at <https://www.who.int/campaigns/world-mental-health-day/2021/campaign-materials>.



Figure 7.9 Supporting Family Members and Significant Others

## Evaluation (Evaluate Outcomes)

Evaluation of the client's progress towards meeting expected outcomes occurs continuously throughout the treatment phase. Evaluation includes comparing results from screening tools, reviewing laboratory results, and monitoring the effectiveness of prescribed medications, treatments, and nursing interventions. Based on the evaluation findings, the nursing care plan may be modified, or new interventions or outcomes may be added.

## 7.6 Spotlight Application

### Postpartum Depression<sup>1</sup>

Maya is a healthy 32-year-old woman who has been married for over two years and is expecting her first child, a baby boy. She had a history of depression and generalized anxiety disorder.

She has been doing well with a combination of medication and cognitive behavioral therapy (CBT) for many years. Maya had decided in the months leading up to getting pregnant that she wanted to be off medication and worked with her psychiatrist to carefully accomplish this. She continued weekly therapy. She was mostly active, upbeat, and cheerful during her pregnancy. She gave birth to a healthy 7.3-pound baby boy. After the delivery, she started to feel sad, overwhelmed, and consistently tearful. She frequently felt irritable and on edge. This feeling persisted for the first ten weeks after the baby was born. She had limited support—her parents were divorced, and her mother was living in another state. Her in-laws were much older with numerous health complications and couldn't help regularly.

Maya and her husband went to see her psychiatrist. She was quite tearful and felt she was a “failure as a mom.” Her baby cried incessantly, and she could barely get any sleep. Maya felt utterly incapable of soothing her baby and would get frustrated and tearful. She was so afraid of what she had learned about sudden infant death syndrome (SIDS) that she would barely allow herself to sleep. She felt that it was a constant race against the clock—with nursing, pumping, and changing. She was always cleaning bottles and diapers. She felt horrified with how she looked. She had expected to wear pre-pregnancy clothes immediately after childbirth. She hadn't had a meal in peace or gotten her hair or nails done and couldn't even think about having

1. American Psychiatric Association. (n.d.). *Patient story: Postpartum depression*. <https://www.psychiatry.org/patients-families/postpartum-depression/patient-story>

sex with her husband. He tried to be supportive, but also felt overwhelmed by it all. He felt she was inconsolable, and they both felt at a loss.

The psychiatrist talked about a variety of tools, including participating in cognitive behavioral therapy, incorporating 15-20 minutes of daily relaxation, practicing mindfulness skills, hiring help, getting her mom to stay with her for a few weeks, and seeking other support. Her husband understood the urgency of the situation and offered to take time off work and to do some of the overnight feedings. Maya decided to restart on her previous antidepressant and also joined a new moms' support group and continued CBT weekly therapy.

Over the next few months, she was exercising more and getting more sleep and support and had significant improvement in mood and energy. She received some sleep training tips from her pediatrician as well. Maya and her husband shared with her psychiatrist that they were feeling significantly better. They were excited to share that they found a series of self-help parenting books to be particularly helpful and had gotten some helpful tips from others in the moms' group.

"Wow, it really does take a village to raise a child, doesn't it?" Maya commented to her psychiatrist. They spoke about how in previous generations new couples could rely on extended family support and how that support often doesn't exist now. Also, inaccurate beliefs, such as babies are easy and infancy should be a happy time for parents, add to stress, conflict, and guilt. Being able to normalize the stress of adjusting to parenthood was extremely helpful for Maya and her husband.

**Reflective Questions:**

1. What educational tips and strategies may have been of benefit to Maya as part of her prenatal care?
2. How have cultural and societal norms impacted families in relation to child rearing?

3. What support strategies and education might be offered to birth partners to help them anticipate challenges and recognize early signs of post-partum depression?



## 7.7 Learning Activities



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=313#h5p-17>

1



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=313#h5p-18>

2



*An interactive H5P element has been excluded from this version of the text. You can view it*

1. “MH Depression Glossary Cards” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
2. “MH Depression Drag and Drop” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

online here:

<https://wtcs.pressbooks.pub/nursingmhcc/?p=313#h5p-19>

3

## Case Study

Kinsey Valdez, age 26 years, is brought into the urgent care clinic today by her roommate, Sara, who reports a recent change in Kinsey's behavior and difficulty concentrating. Kinsey states she let her roommate bring her to the outpatient clinic because she was not feeling well, and "I just do not know what's wrong." During the initial visit with Kinsey, the nurse notes that Kinsey turns away, does not make eye contact, and speaks quietly.

Sarah reports that Kinsey has had a lack of energy for several weeks. She says that Kinsey frequently talks about her life being "hopeless," doesn't get out of bed in the morning, and has no interest in sleeping or eating.

Sarah further explains that when Kinsey was 16, her best friend died in a car accident and she experienced severe depression at that time. She received family support but did not take any medication and was not involved in traditional counseling services. Additionally, Kinsey recently discovered that she is HIV

positive and this occurred roughly on the anniversary of her best friend's passing.

Kinsey has been living with Sarah for 3 years in an apartment and they also work together as nurses on the night shift in the Med-Tele unit at a local hospital. Kinsey's closest family member, her younger brother, lives 3 hours away and they rarely speak. Kinsey admits that she has few friends and states, "I'm usually not much fun to hang out with, but I'm okay with that." She has also been picking up many extra shifts at work due to short staffing in recent weeks. Kinsey states, "I'm worried that this positive HIV test will cause me to no longer be able to work as a nurse."

Kinsey is instructed to go to the nearest Emergency Department (ED) for further assessment and follow-up. Sara takes her to the ED. The nurse from the urgent care clinic called the ED nurse with brief report. When Kinsey and Sarah arrive, they are placed in a private ED room immediately upon arrival.

### **Reflective Questions:**

1. What CUES do you recognize as relevant in planning and providing care for Kinsey?
2. What is Kinsey's priority nursing problem?
3. What are your first steps in caring for Kinsey?

### **ASSESSMENT**

The ED nurse performs an assessment and collects the following data.

**Vital signs:** BP 110/64, P 48, T 35.8, O2 sat 97% on room air, R 16

**Pain (PQRST):** Pain location: knees and ankles bilaterally.

Provocation (P): "Always there, does feel a bit better after

sleeping for a while." Quality (Q): "Achy." Radiation (R): None,

Severity (S): 3/10. Timing (T): Started about 3 weeks ago after falling on the ice at work when leaving after a night shift.

### **Focused Assessment Findings**

**Neuro:** Reports a history of migraine headaches but has had none in recent years. Wears contacts most of the time, sometimes wears glasses. Denies dizziness, changes in vision, or any numbness in limbs.

**Respiratory:** Breathing is unlabored and regular, but occasionally shallow when resting.

**Cardiac:** Denies any complaints.

**GI:** Weight: 52.3 kg, a 3 kg decrease from previous visit to the hospital 1 year ago.

Eating 1 meal daily but only eats if roommate makes food and brings it to her in bed. "I'm just too tired to eat most of the time." Nibbles on snacks and drinks Gatorade and protein shakes at work. Last BM yesterday, pattern is regular. Denies nausea or vomiting.

**GU:** Denies any complaints.

**Skin:** Three small scratches are present on her left forearm that are evenly spaced and superficial. No redness, tenderness, or warmth noted. Face is pale. Small rash across chest that is non-raised, light pink, not warm or itching.

### **Psychosocial Assessment Findings**

**General appearance and behavior:** Wearing pajamas, hair is in loose bun, no makeup. Apprehensive and detached, avoiding eye contact. Slouched in chair, with feet up in a modified fetal position. Cooperative, but passive.

**Affect and mood:** Sad facial expression, blunted affect, slow movements, and quiet monotone voice

**Speech pattern:** Slow, minimal speech. Quiet, logical sentences.

**Thought content:** Depressive thought content, without evidence

of delusions or hallucinations.

**Cognition:** Alert and oriented x 3. Has Bachelor's degree in Nursing and an RN license and is articulate and intelligent.

**Judgment:** No evidence of changes in judgment.

### **Labs Results**

Results are unremarkable, negative for ETOH and cannabis. CD4 is 641 u/L

### **Additional Reflective Questions**

4. What additional CUES are relevant for planning Kinsey's care based on the nurse assessment?
5. What is your hypothesis for Kinsey after analyzing the nursing assessment data?
6. Write a SMART goal for Kinsey's care based on priority nursing problem(s).
7. What priority nursing interventions are important to implement when caring for Kinsey?

- ▶ Test your clinical judgment related to the case study provided with a NCLEX Next Generation-style question: [Chapter 7, Assignment 1](#)<sup>4</sup>



4. "Depression Next Gen Question 1" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

- ▶ Test your clinical judgment related to the case study provided with a NCLEX Next Generation-style case study: [Chapter 7, Case Study 1](#)<sup>5</sup>



5. “Depression Next Gen Case Study” by Kellea Ewen is licensed under [CC BY-NC 4.0](#)

## VII Glossary

**Cognitive behavioral therapy (CBT):** A type of psychotherapy that helps a person recognize distorted/negative thinking with the goal of changing thought and behaviors to respond to changes in a more positive manner.

**Depressive episode:** An episode where the person experiences a depressed mood (feeling sad, irritable, empty) or a loss of pleasure or interest in activities for most of the day, nearly every day, for at least two weeks. Several other symptoms are also present, which may include poor concentration, feelings of excessive guilt or low self-worth, hopelessness about the future, thoughts about dying or suicide, disrupted sleep, changes in appetite or weight, and feeling especially tired or low in energy.

**Depressive Disorder Due to Another Medical Condition:** When a person experiences a persistent depressed mood or a significant loss of interest or pleasure in activities, and these symptoms are directly caused by a medical illness or condition.

**Electroconvulsive therapy (ECT):** A medical treatment reserved for clients with severe major depression who have not responded to medications, psychotherapy, or other treatments. It involves a brief electrical stimulation of the brain while the client is under anesthesia.

**Group therapy:** A type of psychotherapy that brings people with similar disorders together in a supportive environment to learn how others cope in similar situations.

**Hypertensive crisis:** An acute rise and significantly elevated blood pressure, typically over 180/120 mm Hg, that causes acute end-organ damage such as stroke, myocardial infarction, or acute kidney damage. It can be caused by MAOIs, a class of antidepressants.

**Latency:** A delayed response to a question or comment.

**Light therapy:** Therapy for seasonal affective disorder (SAD) that involves sitting in front of a light therapy box that emits a very bright light. It usually

requires 20 minutes or more per day, typically first thing in the morning during the winter months. Most people see some improvements from light therapy within one or two weeks of beginning treatment.

**Major depressive episode:** A period of at least two weeks during which a person experiences a persistently depressed mood or a loss of interest or pleasure in nearly all activities, most of the day, nearly every day.

**Perinatal depression:** Depressive disorder that occurs during pregnancy.

**Peristent depressive disorder:** A chronic form of depression that is typically less severe in intensity than Major Depressive Disorder but lasts much longer.

**Postpartum depression:** Feelings of extreme sadness, anxiety, and exhaustion that may make it difficult for mothers of newborns to complete daily care activities for themselves and/or for their babies. Severe postpartum depression can lead to postpartum psychosis.

**Postpartum psychosis:** Severe postpartum depression can cause delusions (thoughts or beliefs that are not true), hallucinations (seeing, hearing, or smelling things that are not there), mania (a high, elated mood that often seems out of touch with reality), paranoia, and confusion. Women who have postpartum psychosis are at risk for harming themselves or their child and should receive help as soon as possible by calling 911 or taking the mother to the emergency room.

**Premenstrual Dysphoric Disorder (PMDD):** A severe and disabling form of premenstrual syndrome (PMS) that involves significant mood disturbances and physical symptoms occurring in the week or two before menstruation.

**Prolonged Grief Disorder:** A disorder that is characterized by intense longing and preoccupation with the deceased person that persists beyond 12 months and causes significant impairment.

**Rhabdomyolysis:** Severe muscle breakdown which releases myoglobin into the bloodstream and subsequently clogging renal filtrations and causing kidney damage.



**Seasonal affective disorder (SAD):** A type of depression causing symptoms during the fall and winter months when there is less sunlight and usually improves with the arrival of spring. SAD is more than just “winter blues.” The symptoms can be distressing and overwhelming and can interfere with daily functioning.

**Serotonin syndrome:** A medical emergency that can occur in clients taking medications that affect serotonin levels.

**Substance/Medication-Induced Depressive Disorder:** A type of depression that occurs as a direct result of using, misusing, or withdrawing from certain substances or medications.

**Transcranial Magnetic Stimulation (TMS):** A noninvasive procedure that uses magnetic fields to stimulate nerve cells in the brain to improve symptoms of depression when other depression treatments haven’t been effective.







### Learning Objectives

- Identify assessment cues of bipolar disorders
- Identify nursing priorities for clients with bipolar disorders
- Plan outcomes for clients with bipolar disorders
- Describe safety/protective interventions for clients with bipolar disorders
- Apply evidence-based practice when planning care and interventions for clients with bipolar disorders
- Analyze treatments for clients with bipolar disorders
- Apply the nursing process to clients with bipolar disorders at risk for suicide

Bipolar disorder is a mental illness that causes dramatic shifts in a person's mood, energy, and ability to think clearly. Moods shift from abnormally elevated moods called manic episodes to abnormal low moods of depression. See Figure 8.1<sup>1</sup> for a depiction of the shifts in mood that occur with bipolar disorder. Severe bipolar episodes of mania can also include hallucinations or delusions, which can be confused with symptoms of schizophrenia.<sup>2</sup> This chapter will discuss the signs, symptoms, and treatments for bipolar disorder

1. "P\_culture.svg" by [he:משתמש:נועמה](#) is licensed under [CC BY-SA 3.0](#)

2. National Alliance on Mental Illness. (2017). *Bipolar disorder*.  
<https://www.nami.org/About-Mental-Illness/Mental-Health-Conditions/Bipolar-Disorder>

and explain how to apply the nursing process when caring for clients with bipolar disorder.



*Figure 8.1 Bipolar Disorder*

## 8.2 Basic Concepts of Bipolar Disorders

### Manic Episodes

Bipolar disorders include shifts in mood from abnormal highs (called manic episodes) to abnormal lows (i.e., depressive episodes). A **manic episode** is a persistently elevated or irritable mood with abnormally increased energy lasting at least one week. The mood disturbance is severe and causes marked impairment in social or occupational function. Severe episodes often require hospitalization to prevent harm to self or others. As the manic episode intensifies, the individual may become psychotic with hallucinations, delusions, and disturbed thoughts. The episode is not caused by the physiological effects of a substance (such as drug abuse, prescribed medication, or other treatment) or by another medical condition.<sup>1</sup>

According to the *DSM-5-TR*, three or more of the following symptoms are present during a manic episode<sup>2</sup>:

- Inflated self-esteem or grandiosity
- Decreased need for sleep (i.e., feels rested after only three hours of sleep)
- More talkative than usual or pressure to keep talking
- Flight of ideas or subjective experience that thoughts are racing
- Distractibility (i.e., attention is too easily drawn to unimportant or irrelevant stimuli)
- Increase in goal-directed activity (either socially, at work or school, or sexually) or psychomotor agitation
- Excessive involvement in activities that have a high potential for painful consequences (e.g., engaging in unrestrained buying sprees, sexual

1. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

2. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

indiscretions, or foolish business investments)

People experiencing a manic episode may become physically exhausted.

## Hypomanic episodes

Hypomanic episodes have similar symptoms to a manic episode but are less severe and do not cause significant impairment in social or occupational functioning or require hospitalization. A **hypomanic episode** is defined as a distinct period of abnormally and persistently elevated, expansive, or irritable mood, accompanied by increased activity or energy, lasting at least four consecutive days and present most of the day, nearly every day.<sup>3</sup>

During this period, three or more of the following symptoms (or four if the mood is only irritable) must be present to a significant degree, representing a noticeable change from the individual's typical behavior<sup>4</sup>:

- Inflated self-esteem or grandiosity, decreased need for sleep (e.g., feeling rested after only three hours)
- Being more talkative than usual or feeling pressure to keep talking
- Experiencing flight of ideas or racing thoughts, distractibility
- Increased goal-directed activity or psychomotor agitation
- Excessive involvement in activities with a high potential for painful consequences (such as reckless spending, risky sexual behavior, or unwise business investments).

The episode must be associated with a clear change in functioning that is uncharacteristic of the person when asymptomatic, and this change must be

3. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

4. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.



observable by others. However, unlike manic episodes, hypomanic episodes are not severe enough to cause marked impairment in social or occupational functioning or to require hospitalization. If psychotic features are present, the episode is considered manic by definition. The symptoms must not be attributable to the physiological effects of a substance or another medical condition.<sup>5</sup>

## Depressive Episodes

A **depressive episode**[\[pb\\_glossary\]](#) is characterized by a persistently low or sad mood and a significant loss of interest or pleasure in most activities. These symptoms must last for at least two weeks and represent a noticeable change from the individual's typical functioning. Depressive episodes in bipolar disorder are severe and debilitating, often causing marked impairment in social, academic, or occupational performance. Depressive episodes associated with bipolar disorder can lead to suicide. The mortality ratio due to suicide for people with bipolar disorder is 20 times above the general population rate and exceeds rates for other mental health disorders.[\[footnote\]](#)Baldessarini, R.J., Vázquez, G.H. & Tondo, L. (2020, January 6). Bipolar depression: A major unsolved challenge. *International Journal of Bipolar Disorders* 8, (1). <https://doi.org/10.1186/s40345-019-0160-1>[\[footnote\]](#)

Like manic episodes, depressive episodes are not attributable to the physiological effects of a substance or another medical condition.

According to the DSM-5-TR, five (or more) of the following symptoms must be present during the same two-week period, with at least one of the symptoms being either (1) depressed mood or (2) loss of interest or pleasure.[\[footnote\]](#)American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.[\[footnote\]](#).

5. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

- Depressed mood most of the day, nearly every day (e.g., feels sad, empty, hopeless, or appears tearful)
- Markedly diminished interest or pleasure in all, or almost all, activities
- Significant weight loss when not dieting, weight gain, or a decrease or increase in appetite
- Insomnia or hypersomnia nearly every day
- Psychomotor agitation or retardation observable by others
- Fatigue or loss of energy
- Feelings of worthlessness or excessive, inappropriate guilt
- Diminished ability to think or concentrate, or indecisiveness
- Recurrent thoughts of death, suicidal ideation, or suicide attempt

Unlike the increased energy and impulsive behaviors of a manic episode, depressive episodes are characterized by slowed thinking, physical fatigue, and feelings of hopelessness. Individuals may withdraw from others and experience difficulty performing basic daily tasks. In some cases, symptoms become so intense that individuals may experience delusions or hallucinations with depressive content (e.g., believing they are responsible for catastrophic events or hearing voices that reinforce feelings of worthlessness).

## Bipolar Disorders

There are three major types of bipolar and related disorders called Bipolar I, Bipolar II, and Cyclothymia. See Figure 8.2<sup>[footnote]</sup> [“Bipolar\\_mood\\_shifts.png”](#) by Osmosis is licensed under [CC BY-SA 4.0](#)<sup>[/footnote]</sup> for an illustration comparing these three types of bipolar disorders.

[caption id="attachment\_1467" align="aligncenter" width="457"]




Figure 8.3 Bipolar

Disorders[/caption]

**[pb\_glossary id="363"]Bipolar I Disorder** is the most severe bipolar disorder. Individuals with Bipolar I Disorder have had at least one manic episode and often experience additional hypomanic and depressive episodes. (Read about the symptoms of depressive episodes in the “[Depressive Disorders](#)” chapter of this book.) One manic episode in the course of an individual's life can change an individual's diagnosis from depression to bipolar disorder. Manic episodes last at least one week and are present for most of the day, nearly every day. They can be so severe that the person requires hospitalization. Depressive episodes typically last at least two weeks. Episodes of depression with mixed

features (having depressive symptoms and manic symptoms at the same time) are also possible.<sup>6,7</sup>

**Bipolar II Disorder** is defined by a pattern of depressive episodes and hypomanic episodes. The hypomania episodes last at least 4 consecutive days and present most of the day, nearly every day, but individuals do not experience the full-blown manic episodes typical of Bipolar I disorder. Individuals with Bipolar II Disorder often have higher productivity when they are hypomanic and may exhibit increased irritability.<sup>8,9</sup>

**Cyclothymia** is defined by periods of hypomanic symptoms and depressive symptoms lasting for at least two years (1 year in children and adolescents). However, the symptoms do not meet the diagnostic requirements for hypomanic episodes or depressive episodes.<sup>10,11</sup>

Some people with Bipolar I or Bipolar II disorders experience rapid cycling with at least four mood episodes in a 12-month period. These mood episodes

6. National Institute of Mental Health. (2024). *Bipolar disorder*. National Institutes of Health. <https://www.nimh.nih.gov/health/topics/bipolar-disorder>
7. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.
8. National Institute of Mental Health. (2024). *Bipolar disorder*. National Institutes of Health. <https://www.nimh.nih.gov/health/topics/bipolar-disorder>
9. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.
10. National Institute of Mental Health. (2024). *Bipolar disorder*. National Institutes of Health. <https://www.nimh.nih.gov/health/topics/bipolar-disorder>
11. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

can be manic episodes, hypomanic episodes, or major depressive episodes. Cycling can also occur within a month or even a 24-hour period. **Rapid cycling** is associated with severe symptoms and impaired functioning and is more difficult to treat.<sup>12</sup>

## Coexisting Disorders

It is common for people with bipolar disorder to also have anxiety disorders, attention deficit hyperactivity disorder, or substance use disorders<sup>13</sup>.

Sometimes, a person with severe episodes of mania or depression may experience psychotic symptoms, such as hallucinations or delusions, resulting in an incorrect diagnosis of schizophrenia. People with bipolar disorder may misuse alcohol or drugs and engage in other high-risk behaviors in times of impaired judgment during manic episodes. In some cases, people with bipolar disorder also have an eating disorder, such as binge eating or bulimia.<sup>14</sup>

Physiological causes can also cause mania-like symptoms. For example, hyperthyroidism can cause difficulty sleeping, irritability, anxiety, and unintentional weight loss. Individuals can also experience substance-induced bipolar symptoms that develop during intoxication by a substance or withdrawal from a substance. For example, alcohol, sedatives, cocaine, methamphetamines, and phencyclidine (PCP) can cause bipolar-like

12. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

13. Hunt, G.E., Malhi, G.S., Cleary, M., Lai, H.M., & Sitharthan, T. (2016). Prevalence of comorbid bipolar and substance use disorders in clinical settings, 1990-2015: Systematic review and meta-analysis, *Journal of Affective Disorders*, 206, 331-349. doi [10.1016/j.jad.2016.07.011](https://doi.org/10.1016/j.jad.2016.07.011)

14. National Institute of Mental Health. (2023). *Bipolar disorder*. National Institutes of Health. <https://www.nimh.nih.gov/health/topics/bipolar-disorder>

symptoms.<sup>15</sup> For these reasons, on initial evaluation of a client experiencing a manic episode, screening is often typically performed for thyroid disorders and substance use.

## Causes of Bipolar Disorder

Researchers continue to study the possible causes of bipolar disorder. Similar to depressive disorders, most experts agree there is no single cause, and there are many factors that contribute to bipolar disorder. Research shows that people who have a parent or sibling with bipolar disorder have genetic predisposition toward having the disorder. Some studies indicate that neurobiological factors may also predispose individuals to bipolar disorder. The brains of people with bipolar disorder may differ from the brains of people who do not have bipolar disorder. People with certain genes are more likely to develop bipolar disorder. Newer research indicates altered intracellular calcium signaling occurs in people with bipolar disorders, and antiseizure medications can provide effective treatment.<sup>16,17</sup>

Environmental factors may also causative role in the development of bipolar disorders. Stress is a common trigger for mania and depression in adults, and previous adverse childhood events (ACEs) are significantly associated with bipolar disorder.<sup>18</sup> For example, a person with an unstable, chaotic childhood may experience bipolar disorder later in adulthood that is triggered by

15. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.
16. National Institute of Mental Health. (2024). *Bipolar disorder*. National Institutes of Health. <https://www.nimh.nih.gov/health/topics/bipolar-disorder>
17. [The Emerging Neurobiology of Bipolar Disorder](#) by Harrison, Geddes, & Tunbridge is licensed under [CC BY 4.0](#)
18. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

extreme stress. Read more about ACEs in the "[Mental Health and Mental Illness](#)" section in Chapter 1.

- ▶ Read more on the National Institute of Mental Health's [Bipolar Disorder](#) webpage.

## 8.3 Treatments for Bipolar Disorders

Bipolar disorder is often a lifelong illness, and episodes of mania and depression typically recur. Between episodes, many people with bipolar disorder are free of mood disruption, but some people have lingering symptoms. Long-term, continuous treatment can help people manage symptoms and prevent relapse. An effective treatment plan for bipolar disorder typically includes a combination of pharmacologic, psychotherapy, lifestyle changes, and other medical treatments.<sup>1</sup>

### Pharmacologic Treatments

Pharmacologic treatments for bipolar disorder include mood stabilizers, anti-seizure medications, and second-generation (“atypical”) antipsychotics. Treatment plans may also include medications that target sleep disruption or anxiety. Antidepressant medication may be used to treat depressive episodes in bipolar disorder in combination with a mood stabilizer and/or an antipsychotic to prevent precipitating a manic episode.<sup>2</sup> See Table 8.3 for a list of common medications used to treat bipolar disorders.<sup>3</sup> Review information about the associated neurotransmitters in the “[Psychotropic Medications](#)” chapter.

Table 8.3 Common Medications Used to Treat Bipolar Disorders

1. National Institute of Mental Health. (2024). *Bipolar disorder*. National Institutes of Health. <https://www.nimh.nih.gov/health/topics/bipolar-disorder>
2. National Institute of Mental Health. (2024). *Bipolar disorder*. National Institutes of Health. <https://www.nimh.nih.gov/health/topics/bipolar-disorder>
3. Post, R. M. (2022). Bipolar treatment in adults: Choosing maintenance treatment. *UpToDate*. <https://www.uptodate.com/>



Medication Class	Nursing Considerations	Common Side Effects  (*Indicates medical emergency)
Mood Stabilizers	Used to treat symptoms of mania and mood lability.	
Lithium	<p>Used as a first-line mood stabilizing agent to treat mania when symptoms are acute or as maintenance therapy</p> <p>Improved tolerance with food and better drug absorption</p> <p>Recommended water intake is 1.5 – 3 liters/day</p> <p>Given in divided doses if gastrointestinal distress occurs</p> <p>NSAIDs are not recommended because they increase lithium levels</p> <p>Therapeutic blood levels are required. Blood levels are drawn 10-12 hours after the last dose taken. The therapeutic lithium serum level is 0.6-1.2 mEq/L</p>	<p>Lithium blocks antidiuretic hormone (ADH), so monitor for symptoms of diabetes insipidus (i.e., excessive thirst and urination)</p> <p>Long-term use increases risk for hypothyroidism, hyperparathyroidism, and impaired kidney functioning</p> <p><b>*Lithium toxicity</b> (notify the health care provider)</p> <p>Early signs (&lt;1.5 mEq/L): nausea, vomiting, diarrhea, thirst, polyuria, slurred speech, muscle weakness, or fine tremors</p> <p>Moderate signs (1.6-1.9 mEq/L): coarse hand tremors, mental confusion, persistent GI complaints, muscle hyperirritability, EEG changes, or uncoordinated movements</p> <p>Severe signs (&gt;2.0 mEq/L): ataxia, blurred vision, large output of dilute urine, severe hypotension, clonic movements, overt confusion, cardiac dysrhythmias, proteinuria, or death secondary to pulmonary complications</p> <p><b>*Lithium levels &gt; 2.5 mEq/L constitute a medical emergency, even if the client is asymptomatic.</b></p>

Anti-seizure Medications	Used to treat mania and other symptoms.	
<b>Valproic Acid (Depakote)</b>	<p>Used for rapid cycling in acute manic phase when not responding to other mood stabilizers</p> <p>Requires periodic therapeutic serum valproic acid blood levels. Therapeutic range is 50-125 mcg/mL</p> <p>Requires laboratory monitoring, including liver function tests, amylase, and lipase and platelet levels due to higher risk for blood dyscrasias and pancreatitis</p>	<ul style="list-style-type: none"> <li>• Weight gain</li> <li>• Sedation</li> <li>• Nausea and/or vomiting</li> <li>• Hair loss</li> <li>• Tremors</li> <li>• Gastrointestinal discomfort</li> </ul> <p><b>*Signs of toxicity</b> (notify the health care provider):</p> <ul style="list-style-type: none"> <li>• <b>Abdominal pain</b></li> <li>• <b>Dark colored urine</b></li> <li>• <b>Jaundice</b></li> </ul>
<b>Carbamazepine (Tegretol)</b>	<p>Used in combination with lithium and antipsychotic drugs for resistive symptoms</p> <p>Used when treatment resistant due to rapid cycling, paranoia, and extreme hyperactivity</p> <p>Requires routine laboratory monitoring, including white blood cells and liver function tests because it can cause bone marrow suppression and liver damage</p> <p>Therapeutic serum blood levels are 4-12 mcg/mL</p>	<ul style="list-style-type: none"> <li>• Hyponatremia</li> <li>• Fatigue</li> <li>• Blurred vision</li> <li>• Nausea</li> <li>• Ataxia</li> </ul> <p>Risk for toxicity if tegretol blood level is &gt;20 mcg/mL. If suspect toxicity, hold the drug and notify the health care provider.</p>

<b>Lamotrigine (Lamictal)</b>	<p>Used with acute mania or as maintenance therapy</p> <p>Therapeutic serum blood level is 2.5 to 15 mcg/mL</p> <p>Improved tolerance in divided doses</p>	<ul style="list-style-type: none"> <li>• Stevens-Johnson syndrome (A rare but potentially life-threatening reaction to medication that starts with flu-like symptoms, followed by a painful rash that spreads and blisters.)</li> <li>• Other common side effects: nausea/vomiting, headache, dizziness, photosensitivity, rash</li> </ul>
<b>Topiramate (Topamax)</b>  <b>Oxcarbazepine (Oxtellar)</b>	<p>Used for treatment resistant mania</p>	<p>Topiramate</p> <ul style="list-style-type: none"> <li>• Weight loss</li> <li>• Fatigue</li> <li>• Visual disturbances</li> </ul> <p>Oxcarbazepine</p> <ul style="list-style-type: none"> <li>• Hyponatremia</li> </ul>
<b>Antipsychotics Medications</b>	<b>Used to treat symptoms of psychosis, agitation, insomnia, or anxiety.</b>	

<b>Olanzapine (Zyprexa)</b>  Risperidone (Risperdal)  Quetiapine (Seroquel)  Ziprasidone (Geodon)  Aripiprazole (Abilify)  Clozapine (Clozaril)	Help stabilize mood and slow down hyperactive motor activity  Several drugs come in oral, parental, liquid, oral disintegrating tablets, and long-acting injections	Review common adverse effects in the “ <a href="#">Schizophrenia</a> ” section of the “Psychosis and Schizophrenia” chapter.
<b>Antianxiety Medications</b>	<b>Used to treat symptoms of anxiety and irritability.</b>	
<b>Benzodiazepines</b>  Lorazepam (Ativan)  Alprazolam (Xanax)  <b>Other</b>  Gabapentin (Neurontin)  Buspirone (Buspar)	Help reduce the level of hyperactivity during an acute manic phase as a supplement with mood stabilizing drugs  Avoid benzodiazepines in clients with substance use disorders due to high risk of addiction	Review common adverse effects in the “ <a href="#">Treatments for Anxiety</a> ” section of the “Anxiety Disorders” chapter.
<b>Antidepressants Medications</b>	<b>Used to treat symptoms of depressive episodes.</b>	

<b>Medication Classes:</b>  SSRIs  SNRIs  TCAs  MAOIs	Can precipitate a manic episode if not used in combination with a mood stabilizer and/or antipsychotic.	Review common adverse effects in the “ <a href="#">Treatments for Depression</a> ” section of the “Depressive Disorders” chapter.
---	---	---

## Lithium

When administered to a client experiencing a manic episode, lithium may reduce symptoms within 1 to 3 weeks. It also possesses unique antisuicidal properties that sets it apart from antidepressants.<sup>4,5</sup> Lithium levels must be monitored closely during treatment to ensure clients remain within therapeutic and safe parameters.

### BLACK BOX WARNING

A Black Box Warning for lithium states that lithium toxicity is closely related to serum lithium levels and can occur at doses close to therapeutic levels. Before initiating therapy, ensure that lithium levels will be able to be measured promptly and accurately.<sup>6</sup>

4. National Institutes of Health. (2024). *Antisuicidal properties*. DailyMed. <https://www.nlm.nih.gov/>
5. Malhi, G. S., Tanious, M., Das, P., Coulston, C. M., & Berk, M. (2013). Potential mechanisms of action of lithium in bipolar disorder. Current understanding. *CNS Drugs*, 27(2), 135–153. <https://doi.org/10.1007/s40263-013-0039-0>
6. National Institutes of Health. (2024). *Lithium*. DailyMed. <https://www.nlm.nih.gov/>

## ADVERSE/SIDE EFFECTS

Lithium must be closely monitored with routine blood work because it has a narrow therapeutic range of 0.8 to 1.2 mEq/L. Levels above this range cause **lithium toxicity**. Lithium levels > 2.5 mEq/L constitute a medical emergency, even if the client is asymptomatic. Signs of early lithium toxicity include diarrhea, vomiting, drowsiness, muscular weakness, and lack of coordination. At higher levels giddiness, ataxia, blurred vision, tinnitus, and a large output of dilute urine may be seen. Treatment of lithium toxicity includes withholding the lithium and pushing fluids; IV fluids may be required. In severe cases, gastric lavage, mannitol, urea, aminophylline, or dialysis may be used to hasten the excretion of the drug. Lithium is contraindicated in clients with renal or cardiovascular disease, severe dehydration or sodium depletion, and those receiving diuretics because these conditions increase the risk of lithium toxicity.<sup>7,8</sup>

Fine hand tremor, polyuria, and mild thirst may persist throughout treatment. Lithium can cause abnormal electrocardiographic (ECG) findings and risk of sudden death. Clients should be advised to seek immediate emergency assistance if they experience fainting, light-headedness, abnormal heartbeats, or shortness of breath.<sup>9</sup>

Renal function is adversely affected by lithium and requires routine laboratory testing including urinalysis, blood urea nitrogen, and creatinine. Thyroid and parathyroid functioning can also be adversely affected by lithium, requiring

7. National Institutes of Health. (2024). *Lithium*. DailyMed.

<https://www.nlm.nih.gov/>

8. Janicak, P. G. (2022). Bipolar disorder in adults and lithium: Pharmacology, administration, and management of adverse effects. *UpToDate*.

<https://www.uptodate.com>

9. National Institutes of Health. (2024). *Lithium*. DailyMed.

<https://www.nlm.nih.gov/>

routine laboratory testing including thyroid function studies and calcium levels.<sup>10</sup>

Lithium can cause fetal harm in pregnant women. Safety has not been established for children under 12 and is not recommended.<sup>11</sup>

## CLIENT EDUCATION FOR LITHIUM

Lithium must be taken as prescribed or serious side effects can occur. Blood tests to measure lithium levels will be ordered regularly by the provider. The provider should be immediately notified of symptoms of elevated levels of lithium, including diarrhea, vomiting, drowsiness, muscular weakness, lack of coordination, ringing in the ears (tinnitus), or large amounts of dilute urine. Driving or operating heavy machinery should be avoided when first starting lithium because it can impair mental alertness. Lithium should not be taken during pregnancy or while breastfeeding unless it is determined that the benefits to the mother outweigh the potential risks to the baby. Clients taking lithium must also be taught to monitor their dietary intake of sodium consistent as drastic reduction in sodium can lead to lithium toxicity and a drastic increase in sodium intake can lead to sub-therapeutic lithium<sup>12</sup>

## Psychotherapy

**Psychotherapy** is a term for a variety of treatment techniques that help an individual identify and change troubling emotions, thoughts, and behaviors.

10. Janicak, P. G. (2021). Bipolar disorder in adults and lithium pharmacology, administration, and management of adverse effects. *UpToDate*.

<https://www.uptodate.com/>

11. National Institutes of Health. (2024). *Lithium*. DailyMed.

<https://www.nlm.nih.gov/>

12. National Institutes of Health. (2024). *Lithium*. DailyMed.

<https://www.nlm.nih.gov/>

Treatment may include therapies such as cognitive-behavioral therapy (CBT), dialectical behavior therapy (DBT), and psychoeducation. Read more information about CBT and DBT in the [“Treatments for Depression”](#) section in the “Depressive Disorders” chapter.

Treatment may also include newer therapies designed specifically for the treatment of bipolar disorder, including **Interpersonal and Social Rhythm Therapy (IPSRT)** and **family-focused therapy**. IPSRT emphasizes the importance of establishing stable daily routines such as sleeping, waking up, working, and eating meals. Family-focused therapy focuses on psychoeducation, communication enhancement training, and problem-solving skills. It includes attention to family dynamics and relationships as contributing factors to the client’s mood.

## Electroconvulsive Therapy

Electroconvulsive therapy (ECT) is used to treat bipolar disorders. ECT is a brain stimulation procedure delivered under general anesthesia. It can be effective in treating severe depressive and manic episodes when medication and psychotherapy are not effective. ECT can also be effective when a rapid response is needed, as in the case of suicide risk or **catatonia** (a state of unresponsiveness). Read more about electroconvulsive therapy (ECT) in the [“Depressive Disorders”](#) chapter.

## Repetitive Transcranial Magnetic Stimulation

Repetitive transcranial magnetic stimulation (rTMS) is a brain stimulation procedure that uses magnetic waves to relieve depressive episodes associated with bipolar disorders and does not require general anesthesia.

## Lifestyle Changes and Coping

Lifestyle changes can play a significant role in managing symptoms of bipolar disorder. Implementing these lifestyle changes can complement pharmacotherapy and psychotherapy, leading to better management of bipolar disorder symptoms and overall quality of life.



Regular aerobic exercise, such as jogging, brisk walking, swimming, or bicycling, helps with depression and anxiety, promotes better sleep, and is healthy for the heart and brain. There is also some evidence that anaerobic exercise such as weightlifting, yoga, and Pilates can be helpful.<sup>13</sup> Adopting a balanced diet, such as the Mediterranean diet, can positively impact mood and metabolic health. A vegetarian diet has been associated with reduced depression scores, better psychosocial functioning, and improved metabolic parameters. Individuals should also maintain a regular sleep schedule and practice mindfulness or meditation to help manage stress. Alcohol, tobacco products, and recreational drug use should be avoided. Individuals should build and maintain strong social connections to help develop emotional support systems for symptom management.<sup>14</sup>

Even with proper treatment, mood changes can occur. Treatment is more effective when a client and health care provider work together and talk openly about concerns and choices. Keeping a life chart that records daily mood symptoms, treatments, sleep patterns, and life events can help clients and health care providers track and treat bipolar disorder over time. Clients can easily share data collected via smartphone apps – including self-reports, self-ratings, and activity data – with their health care providers and therapists.<sup>15</sup>

13. National Institute of Mental Health. (2024). *Bipolar disorder*. National Institutes of Health. <https://www.nimh.nih.gov/health/topics/bipolar-disorder>
14. Simjanoski, M., Patel, S., Boni, R., Balanzá-Martínez, V., Frey, B. N., Minuzzi, L., Kapczinski, F., Cardoso, T. A. (2023). Lifestyle interventions for bipolar disorders: A systematic review and meta-analysis. *Neuroscience and Biobehavioral Reviews*. 152, 105257. [doi:10.1016/j.neubiorev.2023.105257](https://doi.org/10.1016/j.neubiorev.2023.105257).
15. National Institute of Mental Health. (2024). *Bipolar disorder*. National Institutes of Health. <https://www.nimh.nih.gov/health/topics/bipolar-disorder>

- ▶ Take the American Psychiatric Association's free course: [The Treatment of Bipolar Depression: From Pills to Words.](#)

## 8.4 Applying the Nursing Process to Bipolar Disorders

### Assessment (Recognize Cues)

Assessment of a client with a mood disorder focuses on both verbal and nonverbal cues. People with bipolar disorder experience periods of unusually intense emotion, grandiose delusions, changes in sleep patterns and activity levels, and impulsive behaviors, often without recognizing potential harmful effects.<sup>1</sup> See Figure 8.3<sup>2</sup> for an artistic depiction of grandiose delusions when a cat looking in a mirror sees a lion.

1. National Institute of Mental Health. (2024). *Bipolar disorder*. National Institutes of Health. <https://www.nimh.nih.gov/health/topics/bipolar-disorder>
2. “[Cat and lion in mirror illustration.svg](#)” by [Arlo Barnes](#) is licensed under [CC BY 3.0](#)




Figure 8.3 Grandiose Delusions

It is often helpful to interview family members or significant others of clients with mood disorders. Clients with mania, hypomania, or psychosis often have poor insight and may have difficulty providing an accurate history.<sup>3</sup>

Safety guidelines for assessing a client with a bipolar disorder include the following:<sup>4</sup>

- Assess if the client is a danger to self or others. The client may have suicidal or homicidal ideation. Poor impulse control may result in harm to

3. Suppes, T. (2021). Bipolar disorder in adults: Assessment and diagnosis.

*UpToDate*. <https://www.uptodate.com/>

4. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

self or others. Assess the need for protection from uninhibited behaviors. For example, external controls may be needed to protect the client from consequences such as bankruptcy.

- Assess physiological stability while obtaining vital signs and lab results including electrolytes. The client may not eat or sleep for days at a time with potential physiological consequences.

## Mental Status Examination

Table 8.4a outlines typical assessment findings a nurse may observe in a client experiencing a manic episode. Typical findings relate to mood, behavior, thought processes, speech patterns, and cognitive function.

Table 8.4a Typical Mental Status Examination Findings for a Client Experiencing a Manic Episode<sup>5,6,7</sup>

5. National Institute of Mental Health. (2024). *Bipolar disorder*. National Institutes of Health. <https://www.nimh.nih.gov/health/topics/bipolar-disorder>
6. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.
7. Suppes, T. (2021). Bipolar disorder in adults: Assessment and diagnosis. *UpToDate*. <https://www.uptodate.com/>

Assessment	Typical Findings During a Manic Episode
<b>Level of Consciousness and Orientation</b>	May be disoriented/confused, but can be oriented to person, place, and time.
<b>Mood and Affect</b>	Exhibits an unstable, euphoric mood. Client may state they feel “up,” “high,” “jumpy,” or “wired,” but mood can quickly change to irritation and anger.
<b>Appearance and General Behavior</b>	<p>Typically exhibits a decreased need for sleep and a loss of appetite that may result in dehydration or poor nutritional status.</p> <p>May exhibit inappropriate dress or grooming or dress provocatively, sloppily, flamboyantly, or bizarrely. May change clothes frequently throughout the day. May use excessive makeup or demonstrate little attention to grooming.</p> <p>May demonstrate risky behaviors with poor impulse control and poor judgment, such as eating and drinking excessively, spending or giving away a lot of money, or having reckless sex. Excessive spending can lead to financial hardship from credit card debt from buying items they don't need.</p>
<b>Speech</b>	Typically talk very fast (e.g., pressured speech) about many different topics (hyperv verbal). May have difficulty in accurately communicating needs due to flight of ideas or slurred or garbled speech.
<b>Motor Activity</b>	Typically hyperactive with an inability to recognize need for rest or sleep.
<b>Thought and Perception</b>	<p>Client may state they feel as if their “thoughts are racing.”</p> <p>May feel as if they are unusually important, talented, or powerful.</p> <p>May describe hallucinations, illusions, or paranoia.</p> <p>May exhibit flight of ideas, loose associations, and clang associations. (See definitions of terms in the “<a href="#">Application of the Nursing Process in Mental Health Care</a>” chapter.)</p> <p>May exhibit suicidal, homicidal, or violence ideation.</p>
<b>Attitude and Insight</b>	Typically exhibit limited insight with an inability to make sound decisions impacting their adherence to taking prescribed medications.

<b>Cognitive Abilities</b>	Typically exhibit decreased attention span, distraction, and impaired judgement.
----------------------------	--

## Psychosocial Assessment

It is helpful to begin the psychosocial assessment by obtaining the reason why the client is seeking health care in their own words. For example, the client may identify a problem such as a relationship issue, stressful job, or school challenges that could be addressed by counseling.<sup>8</sup>

A comprehensive psychosocial assessment includes the following components:

Reason for seeking health care (i.e., “chief complaint”)

Thoughts of suicide or self injury

Cultural assessment

Spiritual assessment

Family dynamics

Current and past medical history

Current medications

History of previously diagnosed mental health disorders

Previous hospitalizations

Educational background

Occupational background

History of exposure to psychological trauma, violence, and domestic abuse

Substance use (tobacco, alcohol, recreational drugs, misused prescription drugs)

Family history of mental illness

Coping mechanisms

Functional ability/Activities of daily living

8. Halter, M. (2022). Varcarolis' foundations of psychiatric-mental health nursing (9th ed.). Saunders.

After identifying the reason the client is seeking health care, additional focused questions are used to obtain detailed information. The mnemonic PQRSTU can be used to ask questions in an organized fashion. See Table 8.4b for a sample PQRSTU assessment for assessing mania.

Table 8.4b Sample PQRSTU Questions for Assessing Mania<sup>9</sup>

PQRSTU	Sample Questions
Provocation/ Palliation	“What tends to trigger or worsen your high energy or mood? What helps
Quality	“How would you describe what you’re feeling when you’re in a high or elev
Region	“Do you experience physical symptoms or sensations when you’re in a ma
Severity	“On a scale of 0 to 10, how intense do your manic symptoms feel when the
Timing/ Treatment	“When did this elevated mood or energy begin? How long does it usually
Understanding	“What do you think is happening when you feel this way? Do you think yo

9. National Institute of Mental Health. (2024). *Bipolar disorder*. National Institutes of Health. <https://www.nimh.nih.gov/health/topics/bipolar-disorder>



## SUICIDE AND SELF INJURY SCREENING

Clients being evaluated or treated for bipolar disorder may have suicidal ideation. The Patient Safety Screener (PSS-3) is an example of a brief screening tool to detect suicide risk in all client presenting to acute care settings.<sup>10</sup>

Non-suicidal self-injury (NSSI) refers to intentional self-inflicted destruction of body tissue without suicidal intention and for purposes not socially sanctioned. Common forms of NSSI include behaviors such as cutting, burning, scratching, and self-hitting. It is considered a maladaptive coping strategy without the desire to die. NSSI is a common finding among adolescents and young adults in psychiatric inpatient settings.<sup>11</sup>

Review information about suicide screening, the Patient Safety Screener, and screening for non-suicidal self-injury (NSSI) in the “[Assessment](#)” section of the Applying the Nursing Process to Mental Health Care” chapter.

## CULTURAL ASSESSMENT

Cultural Formulation Interview (CFI) questions help nurses understand a client’s cultural background and how it influences their experience of mental health symptoms, including manic and depressive episodes.<sup>12</sup> Sample CFI

10. Suicide Prevention Resource Center. (n.d.). The patient safety screener: A brief tool to detect suicide risk. <https://sprc.org/micro-learning/patientsafetyscreener>
11. Cipriano, A., Cella, S., & Cotrufo, P. (2017). Nonsuicidal self-injury: A systematic review. *Frontiers in Psychology*, 8. <https://www.frontiersin.org/articles/10.3389/fpsyg.2017.01946/full>
12. DeSilva, R., Aggarwall, N. K., & Lewis-Fernandez, R. (2015). The DSM-5 cultural formulation interview and the evolution of cultural assessment in psychiatry. *Psychiatric Times*, 32(6). <https://www.psychiatrictimes.com/view/>

questions focused specifically on understanding mania and bipolar disorders within a cultural context include the following:

- Cultural Definition of the Problem
  - “How would you describe the changes in your mood or energy that you’ve been experiencing?”
  - “In your community or family, what do people call this kind of problem—feeling very ‘up’ or energetic at times, and very ‘down’ at others?”
  - Do you feel this is as a medical issue, a spiritual issue, or something else?”
- Cultural Perceptions of Cause, Context, and Support
  - “What do you think has caused these shifts in your mood or energy?”
  - “Have there been any stressors, life events, or spiritual changes that you believe might explain this?”
  - “Do other people in your family or community view these mood changes as a strength, a weakness, or something else?”
- Cultural Factors Affecting Coping and Help-Seeking
  - “When you feel very energetic or very low, how do you usually cope with those feelings?”
  - “Have you sought help from anyone—such as a spiritual leader, traditional healer, or family member?”
  - “Are there treatments or practices you trust or prefer when managing your health?”
- Cultural Features of the Nurse–Client Relationship
  - “Are there any concerns you have about talking to a mental health professional?”
  - “Would you feel more comfortable speaking with someone of a similar gender, cultural, or religious background?”

[dsm-5-cultural-formulation-interview-and-evolution-cultural-assessment-psychiatry](#)

- “What would help you feel more supported or understood during treatment?”

Findings from the cultural formulation interview are used to individualize a client’s plan of care to their preferences, values, beliefs, and goals.

## SPIRITUAL ASSESSMENT

The FICA Spiritual History Tool is a widely used assessment model for evaluating a client’s spiritual beliefs and how they may influence health, illness, and coping. It’s especially helpful in understanding how clients with bipolar disorder draw on spirituality or religion for support. Addressing a client’s spirituality and advocating spiritual care have been shown to improve clients’ health and quality of life.<sup>13 14</sup>

The FICA Spiritual History Tool© is a common tool used to gather information about a client’s spiritual history and preferences. FICA© is a mnemonic for the domains of Faith, Importance, Community, and Address in Care.<sup>15</sup> Table 8.4c

13. Pilger, C., Molzahn, A. E., de Oliveira, M. P., & Kusumota, L. (2016). The relationship of the spiritual and religious dimensions with quality of life and health of patients with chronic kidney disease: An integrative literature review. *Nephrology Nursing Journal: Journal of the American Nephrology Nurses’ Association*, 43(5), 411–426. <https://pubmed.ncbi.nlm.nih.gov/30550069/>
14. Puchalski, C., Jafari, N., Buller, H., Haythorn, T., Jacobs, C., & Ferrell, B. (2020). Interprofessional spiritual care education curriculum: A milestone toward the provision of spiritual care. *Journal of Palliative Medicine*, 23(6), 777–784. <https://doi.org/10.1089/jpm.2019.0375>
15. GW School of Medicine & Health Sciences. (n.d.). *Clinical FICA tool*. <https://smhs.gwu.edu/spirituality-health/program/transforming-practice-health-settings/clinical-fica-tool>

summarizes a sample FICA Spiritual Assessment and sample responses from a client experiencing bipolar disorder.

Table 8.4c Sample FICA Spiritual Assessment Questions For Clients With Bipolar Disorder

Domain	Sample Assessment Question	Sample Client Response
<b>Faith</b>	“Do you consider yourself spiritual or religious? What gives your life meaning?”	“During my high periods, I sometimes feel like I’m on a mission from God. When I crash, I feel abandoned or punished.”
<b>Importance</b>	“What importance does your faith or belief have in your life? Has it influenced how you cope with your mood changes?”	“My faith can be confusing at times depending on how I’m feeling. When I’m up, it can feel like a blessing because I am so productive and can get so much done. When I’m down, I question everything and wonder why I am being punished with these up and down periods.”
<b>Community</b>	“Are you part of a spiritual or religious community? Do they provide support to help manage your health or emotions?”	“When I’m feeling depressed, I don’t feel like going to church. I haven’t talked to anyone about what happens during my up periods.”
<b>Address in Care</b>	“How would you like me (or the health care team) to address spiritual issues during your care? Would you like to speak with a chaplain or contact your pastor?”	“Yes I would be interested in speaking to a chaplain.”

Nurses may recognize cues of spiritual distress related to feelings or behaviors that occur during manic and depressive episodes. Nurses can offer to connect the client with a chaplain or spiritual care services. Spiritual goals may be included in the nursing care plan if the client finds them valuable.

## FAMILY DYNAMICS

Family dynamics are included in a psychosocial assessment, especially for children, adolescents, and older adults. **Family dynamics** refers to the patterns of interactions among relatives, their roles and relationships, and the

various factors that shape their interactions. Because family members rely on each other for emotional, physical, and economic support, they are primary sources of relationship security or stress. Family dynamics and the quality of family relationships can have either a positive or negative impact on an individual's mental health. For example, secure and supportive family relationships can provide love, advice, and care, whereas stressful family relationships can be burdened with arguments, unhealthy relationships, and a lack of support.<sup>16</sup>

## Screening Tools

Many screening tools exist to assess mood disorders. Common examples include the following:

- ▶ **Mood Disorder Questionnaire (MDQ) PDF:** Thirteen questions with yes/no responses for assessing mania.
- ▶ **Young Mania Rating Scale (YMRS) PDF:** An 11-item assessment based on the client's subjective report of behaviors over the past 24 hours regarding manic symptoms. It is useful to evaluate baseline functioning and progress being made. There is also a parent version for assessing children and adolescents.
- ▶ **Altman Self-Rating Mania Scale:** A self-assessment questionnaire to evaluate the severity of mania or hypomania.

## Laboratory Testing

Initial medical evaluation of clients with a possible or established diagnosis of bipolar disorder typically includes the following:

16. Jabbari, B., Schoo, C., & Rouster, A. S. (2023). Family dynamics. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK560487/>

- Thyroid stimulating hormone
- Complete blood count
- Chemistry panel
- Urine toxicology to screen for substances of abuse

Thereafter, routine laboratory testing for clients with bipolar disorders can include these items:

- Therapeutic Medication Levels: Medication dosages may need adjustment based on blood levels to avoid toxicity and ensure they are in therapeutic range. Read more about therapeutic drug levels under the “Medications” subsection of the [“Treatments for Bipolar Disorders”](#) section of this chapter.
- Kidney or liver function tests, based on medications prescribed.
- Thyroid function studies and calcium levels.
- Nutrition or hydration status, such as serum sodium levels, hematocrit, albumin, and prealbumin levels, which can become impaired during manic episodes due to poor intake.

### Reflective Questions

1. What are some common underlying medical conditions that could potentially mimic the symptoms of mania in older adults?
2. Why are some individuals with bipolar disorder misdiagnosed with schizophrenia?

## Life Span Considerations

Life span considerations influence how the client is assessed, as well as the selection of appropriate nursing interventions.

## CHILDREN AND ADOLESCENTS

Most cases of bipolar disorder are diagnosed in adolescence or adulthood, but the symptoms can appear earlier in childhood. While less common than in adults, it is possible for children to experience manic and depressive episodes, especially if there is a family history of bipolar disorder. Research also indicates that trauma and adverse childhood experiences (ACEs) may increase the chances of developing bipolar disorder in people with a genetic risk of having the condition. The symptoms of manic episodes may look different in children than in adults, often presenting as significant irritability, mood swings, changes in sleep patterns, and destructive outbursts. Depressive episodes may present as somatic complaints or withdrawal. Bipolar disorder may be misdiagnosed as ADHD, ODD, or conduct disorder.<sup>17</sup> Review information about adverse childhood experiences (ACEs) in the “[Mental Health and Mental Illness](#)” section of Chapter 1.

## OLDER ADULTS

Onset of bipolar disorder after age 60 is rare. Individuals often present with depression as the primary symptom rather than hypomania or mania. New symptoms of mania and should be communicated to the health care provider for investigation of medical causes such as stroke, thyroid dysfunction, or dementia.<sup>18</sup>

## Diagnosis (Analyze Cues)

Mental health disorders are diagnosed by trained mental health professionals

17. Cleveland Clinic. (2022). Bipolar disorder in children.

<https://my.clevelandclinic.org/health/diseases/14669-bipolar-disorder-in-children>

18. Tampi, R. (2023). Assessment and management of bipolar disorder in older adults. *Psychiatric Times*, 40(12). <https://www.psychiatrictimes.com/view/assessment-and-management-of-bipolar-disorder-in-older-adults>

using the *Diagnostic and Statistical Manual of Mental Disorders (DSM-5-TR)*. Nurses create individualized nursing care plans using nursing diagnoses based on the client's response to their mental health disorders. Examples of common nursing diagnoses associated with bipolar disorders are listed in Table 8.4d.

Table 8.4d Common Nursing Diagnoses Related to Bipolar Disorder<sup>19, 20</sup>

19. Ackley, B., Ladwig, G., Makic, M. B., Martinez-Kratz, M., & Zanotti, M. (2020). *Nursing diagnosis handbook: An evidence-based guide to planning care* (12th ed.). Elsevier.
20. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.



<b>Nursing Diagnosis</b>	<b>Associated Behaviors and Characteristics</b>
<b>Safety: Risk for Injury Risk for Suicide Risk for Violence</b>	Impulsive, risky behaviors with poor personal boundaries. Lack of insight into illness. May exhibit agitation, self-harm, or threatening behaviors.
<b>Communication: Impaired Cognition Impaired Communication</b>	Grandiose thinking with poor judgment, flight of ideas, or pressured speech with loose associations.
<b>Self-Care Deficit: Bathing, Grooming, Hygiene, Dressing</b>	Poor hygiene and distracted from tasks.
<b>Impaired Nutrition</b>	Unable to sit long enough to eat; poor appetite. May eat excessive amounts of food during hypomanic episodes.
<b>Disturbed Sleep Patterns</b>	Inability to rest or sleep without frequent awakenings; often hyperactive at night.
<b>Fatigue r/t Psychological Demands</b>	Hyperactive and restless.
<b>Social Isolation r/t Ineffective Coping or Intrusive Behaviors</b>	Feel different from others and preoccupied with own thoughts. Social behavior may be incongruent with norms. May demonstrate an excessive amount of verbal exchange or violation of personal boundaries. May engage in inappropriate sexual language or behavior.
<b>Risk for Spiritual Distress</b>	Demonstrate ineffective coping strategies, separation from support system, or hopelessness.

## Outcome Identification (Generate Solutions)

During an acute manic episode, the overall goals are symptom management, achieving remission of symptoms, preventing injury, and supporting

physiological integrity. Examples of goals during the acute phase include the following<sup>21</sup>:

- Maintain stable cardiac status.
- Be well-hydrated.
- Get sufficient sleep and rest.
- Make no attempt at self-harm.
- Demonstrate thought control with the aid of staff and/or medication.
- Maintain tissue integrity.

The maintenance phase occurs after acute symptoms have been controlled and the goals become focused on preventing future exacerbations of manic episodes through education, support, and problem-solving skills. The following are examples of goals during the maintenance phase<sup>22</sup>:

- Identifying and avoiding triggers for developing acute mania
- Attending therapy sessions
- Developing new coping skills

Outcome criteria are identified based on the phase of bipolar illness the client is experiencing, either acute or maintenance phase.<sup>23</sup> SMART outcomes are Specific, Measurable, Attainable/Actionable, Relevant, and Timely. Read more about SMART outcomes in the “[Application of the Nursing Process in Mental Health Care](#)” chapter. The following are sample SMART outcomes for clients with bipolar disorders:

21. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.
22. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.
23. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

- The client will communicate feelings and thoughts of self-harm (self-injury) to the health care team, prior to acting on thoughts, during this shift.
- The client will eat breakfast within one hour of the arrival of the breakfast tray.
- The client will attend one or more group meetings each day while in the outpatient setting.

## Planning (Generate Solutions)

### Safety

Clients with bipolar disorders are monitored closely for risk of suicide, and interventions are planned according to their level of risk. Review interventions for clients at risk of suicide in the “[Application of the Nursing Process in Mental Health Care](#)” chapter.

### Planning Interventions During Acute Manic Episodes

When a client is hospitalized during an acute manic episode, planning focuses on stabilizing the client while maintaining safety. Nursing care focuses on managing medications, decreasing physical activity, increasing food and fluid intake, reinforcing a minimum of 4 to 6 hours of sleep per night, and ensuring self-care needs are met.<sup>24</sup>

### Planning Interventions During the Maintenance Phase

During the maintenance phase, planning focuses on preventing relapse and limiting the severity and duration of future episodes. During this period, individuals with bipolar disorders often face multiple hardships resulting from

24. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

their behaviors during previous acute manic episodes. Interpersonal, occupational, educational, and financial consequences may occur. Clients need support as they recover from acute illness and repair their lives.<sup>25</sup>

Individuals are often ambivalent about treatment, but bipolar disorders typically require medications to be taken over long periods of time or for a lifetime to prevent relapse. Self-medication through alcohol or other substances often complicates recovery and treatment. Nurses must establish a therapeutic nurse-client relationship to support continued treatment. See [Chapter 2](#) for additional information on establishing a therapeutic nurse-client relationship. Individuals are typically referred to community resources and outpatient mental health care settings. In addition to medication management, outpatient services provide structure and decrease social isolation.<sup>26</sup>

## Implementation (Take Action)

### Nursing Interventions for Clients With Bipolar Disorder Based on Categories of the APNA Implementation Standard

Nursing interventions for clients with bipolar disorders can be categorized based on the American Psychiatric Nurses Association (APNA) standard for *Implementation* that includes the *Coordination of Care; Health Teaching and Health Promotion; Pharmacological, Biological, and Integrative Therapies; Milieu Therapy; and Therapeutic Relationship and Counseling*. Read more about these subcategories in the “[Application of the Nursing Process in](#)

25. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

26. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

Mental Health Care” chapter. See examples of interventions for each of these categories for clients with bipolar disorders in Table 8.4e.

Table 8.4e Nursing Interventions for Mania Based on the Categories of the APNA Implementation Standard<sup>27</sup>

27. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

Categories of Interventions Based on the APNA Standard of Implementation	What the nurse will do...	Rationale
--	---------------------------	-----------

<b>Coordination of care</b>	<ul style="list-style-type: none"> <li>• Collaborate with psychiatrists, social workers, and therapists for integrated care.</li> <li>• Maintain safety by communicating safety precautions with interprofessional team members as needed to prevent self-harm, suicide, or homicide risks.</li> <li>• Ensure consistency of behavioral expectations among all staff on the unit by including expectations in the nursing care plan.</li> <li>• Assist in care transitions (e.g., hospitalization, discharge planning).</li> <li>• Provide referrals to outpatient mental health care settings, community resources, support groups, peer programs, and housing resources.</li> </ul>	<p>Bipolar disorder requires a multidisciplinary approach to address mood stabilization, medication adherence, psychosocial needs, and relapse prevention.</p> <p>The client may exhibit high risk or impulsive behaviors that could pose a risk of harm to self/others. They may experience altered thought processes with poor insight and judgment.</p> <p>Consistent expectations help prevent manipulative behaviors and pushing of limits.</p> <p>The nurse assists in coordinating care delivery during inpatient care and for after discharge.</p>
-----------------------------	---	--

	Plan for quality of life, independence, and optimal recovery.	
<b>Health Teaching and Health Promotion</b>	<ul style="list-style-type: none"> <li>• Educate about bipolar disorder, medication side effects, early warning signs, and mood tracking.</li> <li>• Promote healthy lifestyle behaviors such as good sleep hygiene, healthy diet, and avoiding the use of alcohol and illicit drugs.</li> <li>• Address stigma and support self-advocacy.</li> <li>• See “Client Education” topics for bipolar disorder in the box following these tables.</li> </ul>	Teaching promotes insight, helps prevent relapse, and empowers the client to participate actively in their care and safety planning. Nurses encourage resilience by promoting positive coping strategies.



<b>Pharmacological, Biological, and Integrative Therapies</b>	<ul style="list-style-type: none"> <li>• Provide health teaching about prescribed mood stabilizers and other medications with expected time frames for improvement.</li> <li>• Monitor for side effects of medications and signs of medication toxicity by reviewing lab results (e.g., lithium levels and liver function tests)</li> <li>• Teach about integrative therapies like journaling, mindfulness or yoga, as appropriate.</li> <li>• Open all medications in front of the client.</li> </ul>	<p>Medication adherence is critical in bipolar management. The client's understanding of their medications and potential side effects can increase medication adherence. Monitoring lab results helps prevent the development of complications. There is a small margin of safety between therapeutic and toxic doses of lithium. Holistic therapies enhance emotional regulation and stability. Opening all medications in front of the client may decrease paranoia that may occur with bipolar disorder.</p>
---	--	---

<b>Milieu Therapy</b>	<ul style="list-style-type: none"> <li>• Maintain a structured, low-stimulation, and safe environment. Avoid competitive activities or games. Encourage frequent rest periods and “down time.” Redirect aggressive behavior with structured activities to help decrease tension and provide focus.</li> <li>• Limit environmental triggers (e.g., excessive noise, stimulation). The client may require a private room.</li> <li>• Provide cues for orientation and routine (e.g., activity schedules).</li> <li>• During acute mania, use prescribed medications, seclusion, or restraint to minimize physical harm.</li> </ul>	<p>A therapeutic milieu supports emotional containment during mania and reduces confusion or distress during depressive phases. Reducing stimuli may prevent escalation of anxiety and agitation. Structured activities provide security and focus, but competitive activities should be avoided because they may be too stimulating and can cause escalation of anxiety and agitation. Rest can prevent exhaustion that can result from constant physical activity. The nurse’s priority is to protect the client and others from harm. Storing valued items protects the client from giving away money and possessions. Group therapy can encourage effective coping skills.</p>
-----------------------	--	--

	<ul style="list-style-type: none"> <li>• Store valuables in the hospital safe until safe judgment returns.</li> <li>• Encourage participation in group therapy after acute manic episode has resolved, addressing social skills, personal grooming, mindfulness, and stress management.</li> </ul>	
--	--	--

<b>Therapeutic Relationship and Counseling</b>	<ul style="list-style-type: none"> <li>• Build trust with clear boundaries, active listening, and nonjudgmental support.</li> <li>• Use limit setting to address risky or intrusive behavior during manic episodes.</li> <li>• Encourage exploration of thoughts and feelings during depressive episodes.</li> <li>• Assist in developing insight and realistic goals.</li> <li>• Be consistent in approach and expectations. Use a firm and calm approach with short and concise statements. For example, “John, come with me. Eat this sandwich.” Identify expectations in simple, concrete terms with consequences. For example, “John, do</li> </ul>	<p>A therapeutic relationship helps clients feel safe, fosters self-awareness, and facilitates engagement. Limit setting and empathy help stabilize mood and improve self-control.</p> <p>Structure and control can improve feelings of security for a client who is feeling out of control. Consistent limits and expectations minimize the potential for the client’s manipulation of staff and provide feelings of security. Clear expectations help the client experience outside controls and understand reasons for medication, seclusion, or restraints if they are not able to control their behaviors.</p> <p>Listening to legitimate complaints can reduce underlying feelings of helplessness and can minimize acting-out behaviors. Clients may be impulsive and hyperv verbal and interrupt, blame, ridicule, or manipulate others, so personal boundaries must be set.</p>
--	--	--

	<p>not yell at or hit Peter. If you cannot control your behaviors, you will be escorted to the seclusion room to prevent harm to yourself and others.”</p> <ul style="list-style-type: none"> <li>• Listen to and act on legitimate complaints.</li> <li>• Redirect energy into appropriate and constructive channels.</li> <li>• Set limits with personal boundaries.</li> <li>• See additional “Communication Tips” in the box below.</li> </ul>	
--	--	--

## Nursing Interventions for Physiological Symptoms of Manic and Depressive Episodes

Nurses implement interventions for common physiological symptoms that occur during manic and depressive episodes. Table 8.4f summarizes these interventions.

Table 8.4f Nursing Interventions to Promote Physiological Integrity During  
Manic Episodes<sup>28</sup>

28. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

Problem/ Intervention	Rationale
<p><b>Nutrition</b></p> <ul style="list-style-type: none"> <li>• <b>Monitor weight, dietary and fluid intake, and hydration.</b></li> <li>• <b>Encourage regular meals/ snacks, especially in clients with mania who may forget to eat. Offer frequent, high-calorie, high-protein snacks, drinks, and finger foods.</b></li> <li>• <b>Limit caffeine and excessive sugar intake.</b></li> <li>• <b>Collaborate with a dietician as indicated.</b></li> </ul>	<p>During manic phases, clients may experience hyperactivity and decreased appetite, whereas during depressive phases, they may have overeating or loss of appetite. Medication can also cause weight gain or metabolic syndrome, requiring dietary management. Ensure adequate dietary and fluid intake and minimize development of dehydration. “Finger foods” allow for “eating on the run.”</p>

<p><b>Sleep/Rest</b></p> <ul style="list-style-type: none"> <li>• <b>Promote a calm, quiet environment with consistent bedtime routines.</b></li> <li>• <b>Limit evening stimulation and screen time. Encourage relaxation techniques and avoidance of caffeine.</b></li> <li>• <b>Monitor for symptoms of insomnia or hypersomnia.</b></li> <li>• <b>Administer sleep aids as prescribed and monitor for side effects.</b></li> <li>• <b>During manic episodes, set limits on excessive activity to prevent exhaustion. Encourage frequent rest periods during the day and adequate sleep.</b></li> </ul>	<p>Manic episodes often cause reduced need for sleep or insomnia, which can worsen symptoms. Depressive episodes may cause excessive sleep or poor sleep quality, impacting recovery and energy. Sleep regulation supports mood stabilization. Bedtime routines and relaxation techniques induce sleep. Encouraging bedtime routines and decreasing caffeine intake increases the possibility of sleep.<sup>29</sup></p>
--	--



29. Centers for Disease Control and Prevention. (2024). *About sleep*. [https://www.cdc.gov/sleep/about/index.html#cdc\\_behavioral\\_basics\\_steps-what-to-do](https://www.cdc.gov/sleep/about/index.html#cdc_behavioral_basics_steps-what-to-do)

<p><b>Elimination (Constipation)</b></p> <ul style="list-style-type: none"> <li>• <b>Assess bowel and bladder patterns</b></li> <li>• <b>Encourage regular toileting.</b></li> <li>• <b>Encourage fluids and foods high in fiber.</b></li> <li>• <b>Evaluate the need for a bowel management program with stool softeners and laxatives.</b></li> </ul>	<p>Psychotropic medications commonly cause constipation or urinary retention, which may be exacerbated by poor diet or decreased mobility during depressive episodes. Regular monitoring ensures comfort and safety. Fluids and fiber stimulate peristalsis and soft stools. The client experiencing acute mania is easily distracted and not aware of bodily needs. Bowel management programs may be needed to avoid fecal impaction.</p>
---	--

<p><b>Self-Care Deficits</b></p> <ul style="list-style-type: none"> <li>• <b>Assess independence in ADL performance: bathing, grooming, dressing.</b></li> <li>• <b>Encourage the use of a toothbrush, washcloth, soap, and makeup or shaving supplies with concrete instructions as needed.</b></li> <li>• <b>Encourage appropriate clothing choices based on environmental conditions.</b></li> <li>• <b>Use visual reminders, schedules, or prompts to encourage hygiene.</b></li> <li>• <b>Assist with tasks as needed while promoting independence.</b></li> </ul>	<p>Both manic and depressive episodes can lead to neglect of hygiene and daily routines due to distractibility, low energy, or impaired insight. Supporting ADLs helps maintain dignity, identity, and structure. Distractibility and poor concentration are countered through simple, concrete instructions.</p>
---	---

# Communication Tips for Clients with Bipolar Disorder

Clients experiencing mania may be easily agitated or hyperv verbal. Communication tips are summarized in the following box.

## Effective Communication Tips for Clients with Manic Episode of Bipolar Disorder

- Maintain a calm, non-reactive demeanor.
  - **Rationale:** A calm and steady tone helps de-escalate emotional intensity and provides a model of self-regulation.
- Use short, clear, and concrete statements.
  - **Rationale:** During mania, attention span and focus are impaired. Direct, simple communication prevents confusion and helps the client process instructions more effectively.
- Avoid arguing or challenging grandiose ideas.
  - **Rationale:** Debating delusional or inflated beliefs can increase defensiveness and agitation. Instead, acknowledge the emotion behind the belief and gently redirect without confrontation.
- Set firm but respectful personal boundaries and limits on behaviors.
  - **Rationale:** Clients may become intrusive, impulsive, or overstimulated. Clear, consistent boundaries help ensure safety for the client and

others.

- Validate emotions. For example, say, “It sounds like you’re feeling really energized today” instead of reinforcing behaviors by saying, “Wow, you’ve done so much!”).
  - Validating emotions promotes insight into behaviors.
- Reinforce reality gently and consistently.
  - During manic episodes, reality orientation may be impaired. Use clear reminders such as, “Today is Monday, and the group therapy session is this afternoon” to ground the conversation.
- Monitor for rapid topic changes and redirect.
  - Clients experiencing mania often exhibit flight of ideas. Respectfully bring the focus back to the established topic to gather accurate information. For example, “Earlier you mentioned you were having problems sleeping – let’s talk more about that.”
- Acknowledge progress and small successes.
  - Positive reinforcement builds self-esteem. Celebrate even small steps toward recovery.
- Assess safety and ask directly about suicidal or homicidal risk, such as, “Have you had any thoughts of hurting yourself or others?”

- Direct questioning ensures timely intervention to reduce risk of harm. If a client demonstrates agitation with escalation of manic behavior, review interventions described in the “[Crisis and Crisis Intervention](#)” section of the “Stress, Coping, and Crisis Intervention” chapter. Additional de-escalation techniques to maintain safety are described in the “[Workplace Violence](#)” section of the “Trauma, Abuse, and Violence” chapter.

## Client Education for Bipolar Disorder

Living with bipolar disorder can be challenging, but there are ways to control symptoms and live a healthy life. The client may be resistant to teaching during the acute phase of a manic episode, so it is beneficial to wait until manic symptoms begin to resolve. Client education topics are described in the following box.

### **Client Education: Bipolar Disorder**

Client education regarding bipolar disorder includes the following guidelines<sup>30</sup>:

30. National Institute of Mental Health. (2020). *Bipolar disorder*. National Institutes of Health. <https://www.nimh.nih.gov/health/topics/bipolar-disorder>

- Get treatment and stay committed to it. Recovery from a manic episode takes time and it's not easy, but treatment is the best way to start feeling better.
- Keep medical and therapy appointments and talk with the provider about treatment options.
- Take all medicines as directed.
- Maintain a structure for daily activities; keep a routine for eating, getting enough sleep, and exercising.
- Learn to recognize mood swings and warning signs for manic episodes such as decreased need for sleep.
- Ask for help when experiencing barriers or challenges for treatment. Social support helps coping.
- Be patient; improvement takes time.
- Avoid using alcohol and illicit drugs.
- Encourage participation in cognitive behavioral therapy (CBT) or dialectical behavioral therapy (DBT).
- Use stress management, relaxation techniques, and coping strategies to minimize anxiety.
- Participate in support groups

Read more information about the following support groups using these links:

- ▶ [National Alliance on Mental Illness \(NAMI\)](#)
- ▶ [Depression and Bipolar Support Alliance \(DBSA\)](#).

# Implementing Seclusion or Restraints

De-escalation techniques should be attempted at early signs of escalating agitation to avoid the need for seclusion or restraints. However, if a client is escalating out of control to a point where they pose an immediate risk of injury to themselves or others, the use of medications, seclusion, or restraints may become necessary to maintain a safe environment. Most state laws prevent the use of unnecessary restraint or seclusion, so their use is associated with complex ethical, legal, and therapeutic issues.<sup>31</sup> Review ANA guidelines on using restraints in the “[Client Rights](#)” section of the “Legal and Ethical Considerations in Mental Health Care” chapter and information on safely implementing restraints in the “[Workplace Violence](#)” section of the “Trauma, Abuse, and Violence” chapter.

Controlling escalating agitation during the acute phase of a manic episode may include immediate administration of a prescribed antipsychotic and/or benzodiazepine. A combination of haloperidol (Haldol) and lorazepam (Ativan) that can be injected for rapid onset of action is commonly used. The nurse must monitor for respiratory depression, hypotension, and oversedation after administering these types of medication.

## Evaluation (Evaluate Outcomes)

Evaluation occurs continuously throughout the treatment of bipolar disorders. The registered nurse individualizes assessments based on the established goals and SMART outcomes for each client. The effectiveness of nursing and collaborative interventions is evaluated and revised as needed. Questions used to guide the evaluation process include the following:

- Is the client medically stable with nutritional intake, sleep patterns, labs, or activity levels?

31. Halter, M. (2022). *Varc Carolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.



- Is the client engaging in self-care measures?
- Is the client safe with no self-harm behaviors, statements, gestures, or threats of harm towards others?
- Is the client engaging appropriately in unit-based activities and the therapeutic milieu?
- Is the client able to maintain appropriate personal boundaries with others?
- Is the client able to engage appropriately in verbal conversations and interactions with others?
- Is the client able to demonstrate insight into own illness?
- Is the client tolerating prescriptive medications at therapeutic serum levels and without key side effects?
- Is the client at or near baseline optimal functioning without manic symptoms?
- Is the client able to participate in their own plan of care including discharge planning?

## 8.5 Spotlight Application

### Sample Nursing Documentation

Here is a sample nursing narrative note for a client recently admitted who is currently experiencing an acute phase of a manic episode.

*S: Client reports feeling “very happy” with “high energy” and “no need for sleep.” She states, “I’ve got lots to accomplish today; now move out of my way.”*

*O: Euphoric mood with full range affect. Denies suicidal or homicidal ideation. Is skipping in the hallways and singing loudly with rapid, pressured speech. Is easily distracted and unable to focus on the current conversation. Is intrusive with others with a poor sense of personal boundaries; she is giving away clothing items and going into other clients’ rooms and taking their belongings. Dressing in multiple layers of clothes and unable to sit to eat or finish meals. She has experienced weight loss of four pounds in the past three days.*

*A: Acute mania nearing exhaustion with disruptive behaviors.*

*P: Reduce stimuli with calming milieu and redirect firmly to maintain appropriate personal boundaries with others. Encourage nutritional intake with hydration and finger foods, along with rest periods twice a day. Refrain from large group activities with short 1:1 time for unit programming needs.*

## 8.6 Learning Activities



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=380#h5p-20>

1



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=380#h5p-21>

2



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=380#h5p-23>

1. “MH Bipolar Disorders Glossary Cards” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
2. “MH Bipolar Disorders Question Set 1” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

## Case Study

Jennifer Rogers is a 31-year-old female with bipolar I disorder who was admitted to the hospital last night for surgical repair of a left tibial fracture sustained in a fall during a manic episode. She was diagnosed with bipolar disorder in her mid-twenties, shortly after she was married and graduated from law school. Her most recent manic episode occurred 6 months after her wedding and required hospitalization at that time. She recently stopped taking her medication.

Staff initially had difficulty communicating with Jennifer due to her profound hearing loss and deafness. Jennifer is able to read lips and communicates using American Sign Language. She went to surgery early this morning, shortly after arriving at the ED via ambulance. Post-surgery, she has been admitted to the orthopedic unit and has been there for a couple of hours. She is currently manic, expressing grandiose ideas that she is urgently needed in Washington, D.C., to solve current political issues.

Jennifer is married and has no children. Her sister lives in the area and is supportive. Jennifer works as an attorney who specializes in tax law.

Her husband, Joe Rogers, is in the room with her and is serving as her interpreter at this time.

3. “MH Bipolar Disorders Drag and Drop” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

Jennifer's injuries were sustained from a fall and altercation at a bar. Paramedics who responded to the 911 call at the bar stated that she was ranting about the government and talking about how her destiny is to be "an angel," who can help solve all of our government's problems. She climbed onto a table in the bar, and in the struggle that ensued with security, she fell off of the table and sustained a tibial fracture.

Jennifer's husband had contacted the emergency room when he discovered that she was missing from their home last evening, and he was here prior to her surgery. Her sister and he had been searching for her for hours before the accident when they were informed that she had left work early that day. Joe reported that she had not been sleeping or eating much for the past 2 weeks. The last time she stopped taking her medications was 2 years ago, resulting in a manic episode. At that time, she accumulated over \$12,000 in credit card debt on excessive shopping.

### **Reflective Questions**

1. What CUES do you recognize as relevant in providing nursing care for Jennifer?
2. What is your hypothesis for Jennifer's priority nursing problem(s)?
3. What initial steps should be taken to provide client-centered care to Jennifer?

- ▶ Test your clinical judgment with a NCLEX Next Generation-style question: [Chapter 8, Assignment 1](#)<sup>4</sup>



- ▶ Test your clinical judgment with a NCLEX Next Generation-style question: [Chapter 8, Assignment 2](#)<sup>5</sup>



- ▶ Test your clinical judgment with a NCLEX Next Generation-style case study: [Chapter 8, Case Study 1](#)<sup>6</sup>

4. "MH Bipolar Disorders Next Gen Question 1" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

5. "MH Bipolar Disorders Next Gen Question 2" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

6. "MH Bipolar Disorders Next Gen Case Study" by Kellea Ewen is licensed under [CC BY-NC 4.0](#)

## VIII Glossary

**Bipolar I Disorder:** The most severe bipolar disorder with at least one manic episode; most individuals experience additional hypomanic and depressive episodes.

**Bipolar II Disorder:** A pattern of depressive episodes and hypomanic episodes, but individuals have never experienced a full-blown manic episode typical of Bipolar I Disorder.

**Catatonia:** A state of unresponsiveness due to a person's mental state.

**Cyclothymia:** A disorder defined by periods of hypomanic symptoms and periods of depressive symptoms lasting for at least two years (1 year in children and adolescents). However, the symptoms do not meet the diagnostic requirements for hypomanic episodes or depressive episodes.

**Family-focused therapy:** Psychotherapy that focuses on psychoeducation, communication enhancement training, and problem-solving skills. It includes attention to family dynamics and relationships as contributing factors to the client's mood.

**Grandiose delusions:** A symptom associated with bipolar disorders of feeling unusually important, talented, or powerful.

**Hypomanic episode:** Episodes similar to symptoms of a manic episode, but they are less severe and do not cause significant impairment in social or occupational functioning or require hospitalization.

**Interpersonal and social rhythm therapy (IPSRT):** Psychotherapy that emphasizes the importance of establishing stable daily routines such as sleeping, waking up, working, and eating meals.

**Manic episode:** A persistently elevated or irritable mood with abnormally increased energy lasting at least one week. The mood disturbance is severe and causes marked impairment in social or occupational function. Severe episodes often require hospitalization to prevent harm to self or others.

**Psychotherapy:** A variety of treatment techniques that help an individual identify and change troubling emotions, thoughts, and behaviors.

**Rapid cycling:** At least four mood episodes associated with bipolar disorder occurring in a 12-month period.







### Learning Objectives

- Identify assessment cues of anxiety disorder behaviors, obsessive-compulsive disorder, and post-traumatic stress disorder
- Identify nursing priorities for clients with anxiety disorders, obsessive-compulsive disorder, and post-traumatic stress disorder
- Plan outcomes for clients with anxiety disorders, obsessive-compulsive disorder, and post-traumatic stress disorder
- Differentiate safety/protective interventions for clients with anxiety disorders, obsessive-compulsive disorder, and post-traumatic stress disorder
- Apply evidence-based practice when planning care and interventions for clients with anxiety disorders, obsessive-compulsive disorder, and post-traumatic stress disorder
- Analyze treatments for clients with anxiety disorders, obsessive-compulsive disorder, and post-traumatic stress disorder
- Apply the nursing process to clients with anxiety disorders, obsessive-compulsive disorder, and post-traumatic stress disorder at risk for suicide

Anxiety is a natural part of everyday life and can range from helpful to harmful, depending on the situation and intensity. It is the body's response to stress, whether the stressor is perceived as positive or negative. For example, positive stressors—such as cleaning the house before a holiday gathering or

studying for an exam—can create a sense of urgency that enhances focus and productivity. In contrast, negative stressors—like losing one’s car keys or getting into a disagreement with a loved one—may trigger distress and frustration. Mild anxiety can be beneficial by providing the energy and concentration needed to complete important tasks or motivating healthy behavioral changes. However, excessive anxiety can become overwhelming, leading to significant distress and impairing a person’s ability to function in social, educational, occupational, or other areas of life.

This chapter will explore the levels of anxiety and common anxiety disorders, such as generalized anxiety disorder, panic disorder, and various phobias. It will also examine how anxiety is related to conditions like obsessive-compulsive disorder (OCD), post-traumatic stress disorder (PTSD), and certain medical conditions. Finally, the chapter will demonstrate how the nursing process can be applied to caring for individuals experiencing anxiety.

## 9.2 Basic Concepts

**Anxiety** is a universal human experience that often includes feelings of apprehension, uneasiness, uncertainty, or dread in response to a real or perceived threat.<sup>1</sup> While fear is a reaction to a specific and present danger, anxiety often involves a vague sense of dread to a specific or unknown danger. However, the body reacts physiologically with the stress response to both anxiety and fear.<sup>2</sup> See an artistic depiction of a person facing their feelings of anxiety from a perceived threat in Figure 9.1.<sup>3</sup> Review the stress response in the “[Stress, Coping, and Crisis Intervention](#)” chapter.



Figure 9.1 Anxiety

1. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.
2. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.
3. “[fear-g84d457182\\_1280](#)” by [mohamed\\_hassan](#) on [Pixabay.com](#) is licensed under [CC0](#)

# Levels of Anxiety

Hildegard Peplau, a psychiatric mental health nurse theorist, developed a model describing four levels of anxiety which are categorized as mild, moderate, severe, and panic. Behaviors and characteristics can overlap across these levels, but it can be helpful to tailor interventions based on the level of anxiety the client is experiencing.<sup>4</sup>

## Mild

Mild anxiety is a natural part of everyday life and can help an individual use their senses to perceive reality in sharp focus. Symptoms of mild anxiety include restlessness, irritability, or mild tension-relieving behaviors such as finger tapping, fidgeting, or nail biting.<sup>5</sup> These symptoms are generally manageable and do not significantly impair daily functioning.

## Moderate Anxiety

As anxiety increases, the perceptual field narrows, and the ability of the individual to fully observe their surroundings is diminished/reduced. The person experiencing moderate anxiety may demonstrate selective inattention where only certain things in the environment are seen or heard unless they are pointed out. The individual's ability to think clearly, learn, and problem solve is hampered, but can still take place. The physiological stress response kicks in with symptoms such as perspiration, elevated heart rate, and elevated respiratory rate. The individual may also experience headaches, gastric

4. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

5. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

discomfort, urinary urgency, voice tremors, and shakiness; however, they may not be aware these symptoms are related to their level of stress and anxiety.<sup>6</sup>

## Severe Anxiety

In severe anxiety, the perceptual field is significantly diminished, causing the individual to focus on either one specific detail or many scattered details. They often have difficulty noticing what is going on in their environment, even if it is pointed out, and may appear dazed or confused, exhibiting automatic behavior. Learning, problem-solving, and critical thinking become impossible at this level. Symptoms of the stress response intensify and may include hyperventilation, a pounding heart, insomnia, and/or a sense of impending doom.<sup>7</sup> See Figure 9.2<sup>8</sup> for an artist's rendition of severe anxiety.

6. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.
7. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.
8. "[Walter Gramatté, Die Grosse Angst \(The Great Anxiety\), 1918, NGA 71292.jpg](#)" by Walter Gramatté is licensed in the [Public Domain](#)



Figure 9.2 Severe Anxiety

## Panic

**Panic** is the most extreme level of anxiety that results in significantly dysregulated behavior. The individual is unable to process information from the environment and may lose touch with reality. They may demonstrate behavior such as pacing, running, shouting, screaming, or withdrawal, and hallucinations may occur. Acute panic can lead to exhaustion.<sup>9</sup> It is important to distinguish between the broad concept of panic and a panic attack. While panic refers to an intense experience of fear, a panic attack is a specific, acute episode marked by defined symptoms and diagnostic criteria. Panic attack is discussed in [Chapter 9.3 Anxiety-Related Disorders](#).

9. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.



## Coping with Anxiety

Individuals may use several strategies to cope with anxiety. **Coping strategies** are an action, a series of actions, or a thought process used to address a stressful or unpleasant situation or modify one's reaction to such a situation. Coping strategies are classified as adaptive or maladaptive. Adaptive coping strategies include problem-focused coping and emotion-focused coping. Problem-focused coping typically focuses on seeking treatment such as counseling or cognitive behavioral therapy. Emotion-focused coping includes strategies such as engaging in mindfulness, meditation, or yoga; using humor and jokes; seeking spiritual or religious pursuits; engaging in physical activity or breathing exercises; and seeking social support. Maladaptive coping responses include responses such as avoidance of the stressful condition, withdrawal from a stressful environment, disengagement from stressful relationships, and misuse of alcohol or other substances.

**Defense mechanisms** are reaction patterns used by individuals to protect themselves from anxiety that arises from stress and conflict. Adaptive use of defense mechanisms can help people achieve their goals, but excessive or maladaptive use of defense mechanisms can be unhealthy.<sup>10</sup> Excessive use of defense mechanisms are associated with specific mental health disorders.

Read more about coping strategies and defense mechanisms in the "[Stress, Coping, and Crisis Intervention](#)" chapter.

10. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

## 9.3 Anxiety - Related Disorders

Anxiety disorders are one of the most common mental health disorders in the United States, with about one in five adults reporting symptoms of anxiety. In 2022, of those reporting symptoms of anxiety, 11.4% were categorized as having mild anxiety symptoms, 3.9% were categorized as having moderate anxiety symptoms, and 2.8% were categorized as having severe anxiety. A higher percentage of women report anxiety symptoms compared to men, and anxiety is more prevalent among younger adults.<sup>1</sup>

Anxiety disorders often co-occur with other psychiatric conditions such as depression and substance use disorders.<sup>2</sup> An **anxiety disorder** is characterized by persistent and excessive anxiety that is disproportionate to the actual threat and leads to significant distress or impairment in social, occupational, or other important areas of functioning.<sup>3</sup> According to the DSM-5-TR, an anxiety disorder is diagnosed when specific criteria are met, including the frequency, duration, and intensity of symptoms, as well as the presence of functional impairment and distress<sup>4</sup>

There are several types of anxiety-related disorders, including generalized anxiety disorder, social anxiety disorder, panic disorder, phobia-related disorders, separation anxiety, and selective mutism.

1. Terlizzi, E. & Zablotsky, B. (2024). Symptoms of anxiety and depression among adults: United States, 2019 and 2022 [PDF]. *National Health Statistics Report*, 213. <https://www.cdc.gov/nchs/data/nhsr/nhsr213.pdf>
2. Penninx, B. W., Pine, D. S., Holmes, E. A., & Reif, A. (2021). Anxiety disorders. *Lancet*, 397(10277), 914-927. [doi: 10.1016/S0140-6736\(21\)00359-7](https://doi.org/10.1016/S0140-6736(21)00359-7)
3. Penninx, B. W., Pine, D. S., Holmes, E. A., & Reif, A. (2021). Anxiety disorders. *Lancet*, 397(10277), 914-927. [doi: 10.1016/S0140-6736\(21\)00359-7](https://doi.org/10.1016/S0140-6736(21)00359-7)
4. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing..

# Generalized Anxiety Disorder

The *Diagnostic and Statistical Manual of Mental Disorders (DSM-5-TR)* defines **generalized anxiety disorder (GAD)** as excessive anxiety and worry, occurring on more days than not for at least six months, about a number of events or activities (such as work or school performance). The individual finds it difficult to control the anxiety and worry, and it is associated with at least three of the following symptoms<sup>5</sup>:

- Restless or feeling keyed up or on edge
- Being easily fatigued
- Difficulty concentrating or going mentally blank
- Irritability
- Muscle tension
- Sleep disturbance (difficulty falling or staying asleep or restless, unsatisfying sleep)

The anxiety, worry, or physical symptoms cause clinically significant distress or impairment in social, occupational, or other areas of functioning and are not attributable to the physiological effects of a substance, medical condition, or other mental disorder.<sup>6</sup> See Figure 9.3<sup>7</sup> for an artist's depiction of feelings of anxiety.

5. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.
6. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.
7. "[Edvard Munch - Anxiety - Google Art Project.jpg](#)" by [Edvard Munch](#) is licensed in the [Public Domain](#)




Figure 9.3 Anxiety

## Social Anxiety Disorder

The *DSM-5* defines **social anxiety disorder** as marked fear or anxiety about one or more social situations in which the individual is exposed to possible scrutiny by others. Examples include social interactions (e.g., having a conversation, meeting unfamiliar people), being observed (e.g., eating or drinking), and performing in front of others (e.g., giving a speech). The individual fears they will act in a way or show anxiety symptoms that will be negatively evaluated by others, so social situations are avoided or endured with intense fear or anxiety. This fear, anxiety, or avoidance is persistent and typically lasts for six months or more and is not related to a substance, another mental health disorder, or medical condition. It results in clinically significant impairment in social, occupational, or other important areas of

functioning. In children, the fear or anxiety may be expressed by crying, tantrums, freezing, clinging, shrinking, or failing to speak in social situations.<sup>8</sup>

## Panic Disorder

People with panic disorder have recurrent unexpected panic attacks. **Panic attacks** are sudden periods of intense fear that come on quickly and reach their peak within minutes. Attacks can occur unexpectedly or can be brought on by a trigger, such as a feared object or situation. A panic attack can be caused solely by the fear of having a panic attack.<sup>9</sup> See Figure 9.4.<sup>10</sup>



Figure 9.4 Panic Attack

8. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.
9. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.
10. "[A subjective impression of a panic attack.png](#)" by [Yitzilitt](#) is licensed under [CC BY-SA 4.0](#)

The *DSM-5* defines a panic attack when four or more of the following symptoms occur<sup>11</sup>:

- Palpitations, a pounding heartbeat, or an accelerated heart rate
- Sweating
- Trembling or shaking
- Sensations of shortness of breath or smothering
- Feelings of choking
- Chest pain or discomfort
- Nausea or abdominal distress
- Feeling dizzy, unsteady, light-headed, or faint
- Chills or heat sensations
- Paresthesia (numbness or tingling sensations)
- Derealization (feelings of unreality) or depersonalization (being detached from oneself)
- Feelings of losing control or “going crazy”
- Fear of dying

A panic attack is not related to the physiological effects of a substance, medical condition, or another mental disorder. Culture-specific symptoms may also be seen, such as physical symptoms of anxiety of tinnitus, soreness, headache, and uncontrollable screaming or crying, but these should not be counted as one of the four symptoms.<sup>12</sup>

To be diagnosed as a panic disorder, at least one of the panic attacks has

11. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

12. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.



been followed by one month (or more) of one or both of the following characteristics<sup>13</sup>:

- Persistent concern or worry about additional panic attacks or their consequences
- A significant maladaptive change in behavior related to the attacks (such as avoiding unfamiliar situations)

People with panic disorder often worry about when the next attack will happen and actively try to prevent future attacks by avoiding places, situations, or behaviors they associate with panic attacks. Worry about panic attacks and the effort spent trying to avoid attacks cause significant problems in various areas of the person's life, including the potential development of agoraphobia.<sup>14</sup> Read more about agoraphobia in the following “Phobia-Related Disorders” section.

## Phobia-Specific Disorders

A **phobia** is an intense fear or aversion to specific objects or situations (e.g., flying, heights, animals, receiving an injection, or seeing blood). Although it can be realistic to feel anxious about certain objects or circumstances, the fear felt by people with phobias is out of proportion to the actual danger caused by the situation or object.<sup>15</sup> Although individuals with specific phobias often recognize their reactions as disproportionate, they tend to overestimate

13. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

14. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

15. National Institute of Mental Health. (n.d.). *Any anxiety disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/statistics/any-anxiety-disorder>

the danger in their feared situations and thus the judgment of being out of proportion is made by the clinician.<sup>16</sup> See Figure 9.5<sup>17</sup> for an illustration of a phobia called arachnophobia (fear of spiders).



Figure 9.5 Arachnophobia

The phobic object almost always provokes immediate fear or anxiety and is actively avoided or endured with intense fear or anxiety. This fear, anxiety, or avoidance is persistent and typically lasts for six months or more.<sup>18</sup> Physical symptoms can include palpitations, sweating, trembling, shortness of breath, dizziness, and gastrointestinal distress.

<sup>16</sup>. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

<sup>17</sup>. “[Depiction of a person living with a phobia.png](https://www.myupchar.com/en)” by <https://www.myupchar.com/en> is licensed under [CC BY-SA 4.0](https://creativecommons.org/licenses/by-sa/4.0/)

<sup>18</sup>. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.



Three common phobias are social anxiety disorder, agoraphobia, and separation anxiety disorder.<sup>19</sup>

## Agoraphobia

The *DSM-5* defines **agoraphobia** as intense fear of two or more of the following situations:

- Using public transportation
- Being in open spaces (e.g., parking lots, marketplaces, or bridges)
- Being in enclosed spaces (e.g., shops or theaters)
- Standing in line or being in a crowd
- Being outside of the home alone

People with agoraphobia often avoid these situations because they think it may be difficult or impossible to leave in the event they have a panic-like reaction or other embarrassing symptoms such as incontinence. In the most severe form of agoraphobia, the individual can become housebound. Physical symptoms can include palpitations, sweating, trembling, shortness of breath, dizziness, and gastrointestinal distress when exposed to or anticipating the feared situations. Agoraphobia can occur with or without a panic disorder.<sup>20</sup> See Figure 9.6<sup>21</sup> for an artistic rendition of agoraphobia.

19. National Institute of Mental Health. (n.d.). *Any anxiety disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/statistics/any-anxiety-disorder>
20. National Institute of Mental Health. (n.d.). *Any anxiety disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/statistics/any-anxiety-disorder>
21. “[Agoraphobia](#)” by [Ali Emad](#) is licensed under [CC BY-NC-ND 4.0](#)



Figure 9.6 Agoraphobia

## Separation Anxiety Disorder

Separation anxiety is often thought of as something that occurs in children, but adults can also be diagnosed with separation anxiety disorder. People who have **separation anxiety disorder** have fears about being separated from people to whom they are attached. They often worry that some sort of harm will happen to those to whom they are attached while they are separated. This fear leads them to avoid being separated from their attachment figures and to avoid being alone. People with separation anxiety may have nightmares about being separated from attachment figures or experience physical symptoms when separation occurs or is anticipated. The fear, anxiety, or avoidance is persistent, lasting at least four weeks in children and typically

six months or more in adults, causing significant distress or impairment to social, occupational, or other important areas of functioning.<sup>22,23</sup>

## Selective Mutism

Selective mutism is a rare disorder associated with anxiety. **Selective mutism** occurs when people fail to speak in specific social situations despite having normal language skills. Selective mutism usually occurs before the age of five and is often associated with extreme shyness and fear of social embarrassment.

- **Freezing:** A common response where the child becomes physically immobile or rigid when expected to speak.
- **Avoidance and security behaviors:** These behaviors include avoiding eye contact, hiding behind objects or people, and using gestures or nonverbal communication to avoid speaking.
- **Physical symptoms:** These can include blushing, sweating, trembling, and gastrointestinal distress when faced with speaking situations.

It can also be a symptom of post-traumatic stress syndrome. Individuals who are diagnosed with selective mutism are often also diagnosed with other anxiety disorders, and their symptoms are not related to a communication disorder, autism, schizophrenia, or other psychotic disorder.<sup>24,25</sup>

22. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

23. National Institute of Mental Health. (n.d.). *Any anxiety disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/statistics/any-anxiety-disorder>

24. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

25. National Institute of Mental Health. (n.d.). *Any anxiety disorder*. U.S.



View the following YouTube video on anxiety disorders:

[Mental Health Minute: Anxiety Disorders in Adults.](https://youtu.be/UjPRVKS4OBg)<sup>26</sup>

## Risk Factors for Anxiety Disorders

Occasional anxiety is an expected part of life. It is common to feel anxious when faced with a problem at work, before taking a test, or before making an important decision. However, anxiety disorders involve more than temporary worry or fear. For a person with an anxiety disorder, the anxiety does not go away and can worsen over time, and their symptoms can interfere with daily activities such as job performance, schoolwork, and relationships.<sup>27</sup>

Researchers have found that both genetic and environmental factors contribute to the risk of developing an anxiety disorder. Although the risk factors for each type of anxiety disorder can vary, some general risk factors for all types of anxiety disorders include the following<sup>28</sup> :

Department of Health and Human Services. <https://www.nimh.nih.gov/health/statistics/any-anxiety-disorder>

26. National Institute of Mental Health (NIMH). (2021, December 20). *Mental health minute: Anxiety disorders in adults* [Video]. YouTube. All rights reserved. <https://youtu.be/UjPRVKS4OBg>

27. Herdman, T. H., & Kamitsuru, S. (Eds.). (2018). *Nursing diagnoses: Definitions and classification, 2018-2020*. Thieme Publishers New York.

28. National Institute of Mental Health. (n.d.). *Any anxiety disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/statistics/any-anxiety-disorder>

- Temperamental traits of shyness or behavioral inhibition in childhood
- Exposure to traumatic life or environmental events in early childhood or adulthood
- A history of anxiety or other mental health disorders in biological relatives

▶ Review information about adverse childhood events in the [“Mental Health and Mental Illness”](#) section in Chapter 1.

Some medical conditions (such as thyroid problems, chronic obstructive pulmonary disease, and angina) or substances (such as caffeine, certain prescribed medications, or illicit drugs) can also cause symptoms of anxiety. When a client is initially evaluated for anxiety, a physical health examination is performed to rule out other potential causes of their anxiety symptoms.

## 9.4 Treatments for Anxiety

Anxiety disorders are generally treated with psychotherapy, medication, or a combination of both treatments.<sup>1</sup> Support groups, coping strategies, and psychoeducation can also help individuals manage their anxiety.

### Psychotherapy

**Psychotherapy** or “talk therapy” can help people with anxiety disorders. To be effective, psychotherapy must be directed at the person’s specific anxieties and tailored to their needs. Examples of psychotherapy include **cognitive behavioral therapy (CBT)** and **dialectal behavior therapy (DBT)**. CBT teaches people different ways of thinking, behaving, and reacting to anxiety-producing situations and fearful objects. It can also help people learn and practice social skills, which is vital for treating social anxiety disorder. DBT assists individuals to develop distress tolerance skills and emotional regulation skills in managing their anxiety. CBT and DBT can be conducted individually or with a group of people who have similar difficulties.<sup>2</sup> Read more about CBT and DBT in the “[Depressive Disorders](#)” chapter.

**Exposure therapy** is a form of CBT which may be used alone or with other types of CBT to treat social anxiety disorder. Exposure therapy focuses on confronting the fears underlying an anxiety disorder to help people engage in activities they have been avoiding. Exposure therapy can be used in combination with relaxation exercises and/or imagery.<sup>3</sup> A similar therapy that

1. National Institute of Mental Health. (2024). *Anxiety disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/anxiety-disorders>
2. National Institute of Mental Health. (2024). *Anxiety disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/anxiety-disorders>
3. National Institute of Mental Health. (2024). *Anxiety disorders*. U.S. Department

is particularly helpful with phobias is systematic desensitization. Systematic desensitization is a graduated exposure therapy conducted at a very slow pace used in combination with relaxation techniques and imagery.<sup>4</sup>

## Medications

Medications do not cure anxiety disorders but are used to help relieve symptoms of anxiety, panic attacks, extreme fear, and worry. The most common classes of medications used to combat anxiety disorders are antianxiety (anxiolytic) drugs (such as benzodiazepines), antidepressants, and beta-blockers.<sup>5</sup>

- ▶ Review information regarding the effects of benzodiazepines and beta-blockers on neurotransmitters in the “[Antianxiety Medications](#)” section of the “Psychotropic Medications” chapter.

of Health and Human Services. <https://www.nimh.nih.gov/health/topics/anxiety-disorders>

4. Udayangani, S. (2021). *What is the difference between systematic desensitization and exposure therapy*. Difference Between. <https://www.differencebetween.com/what-is-the-difference-between-systematic-desensitization-and-exposure-therapy/#Systematic%20Desensitization>
5. National Institute of Mental Health. (2024). *Anxiety disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/anxiety-disorders>



# Benzodiazepines

Benzodiazepines are prescribed for their immediate effect in relieving anxiety. For long-term use, benzodiazepines are considered a second-line treatment for anxiety (with antidepressants or buspirone considered first-line treatment), as well as an “as-needed” treatment for any distressing flare-ups of symptoms. These drugs are known for their rapid onset. However, people can build up a tolerance if taken over a long period of time, and they may need higher and higher doses to get the same effect. Benzodiazepines are a Schedule IV controlled substance because they have a potential for misuse and can cause dependence. To avoid these problems, benzodiazepines are typically prescribed for short periods of time, especially for people who have a history of substance use disorders. Short-acting benzodiazepines (such as lorazepam) and beta-blockers are used to treat the short-term symptoms of anxiety. Lorazepam is available for oral, intramuscular, or intravenous routes of administration.<sup>6,7</sup>

If people suddenly stop taking benzodiazepines after taking them for a long period of time, they may have withdrawal symptoms, or their anxiety may return. Withdrawal symptoms include sleep disturbances, irritability, increased tension and anxiety, hand tremors, sweating, difficulty concentrating, nausea and vomiting, weight loss, palpitations, headache,

6. National Institute of Mental Health. (2024). *Anxiety disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/anxiety-disorders>

7. National Institutes of Health. (2023). Lorazepam. DailyMed. <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=73bfaeab-94db-48c2-a194-8b173025de78>



muscular pain, and perceptual changes.<sup>8</sup> Therefore, benzodiazepines should be tapered off slowly.<sup>9</sup>

## BLACK BOX WARNING

A Black Box Warning states that concurrent use of benzodiazepines and opioids may result in profound sedation, respiratory depression, coma, and death. The use of benzodiazepines exposes users to risks of misuse, substance use disorder, and addiction. Misuse of benzodiazepines commonly involves concomitant use of other medications, alcohol, and/or illicit substances, which is associated with an increased frequency of serious adverse outcomes. Additionally, the continued use of benzodiazepines may lead to clinically significant physical dependence. The risks of dependence and withdrawal increase with longer treatment duration and higher daily doses, and abrupt discontinuation or rapid dosage reduction may precipitate life-threatening withdrawal reactions. To reduce the risk of withdrawal reactions, a gradual taper should be used to stop or reduce the dosage.<sup>10</sup>

## ADVERSE/SIDE EFFECTS

Children and older adults are more susceptible to the sedative and respiratory depressive effects of benzodiazepines and may experience paradoxical reactions such as tremors, agitation, or visual hallucinations. Debilitated

8. Pétursson, H. (1994). The benzodiazepine withdrawal syndrome. *Addiction*, 89(11):1455-9. [doi:10.1111/j.1360-0443.1994.tb03743.x](https://doi.org/10.1111/j.1360-0443.1994.tb03743.x).
9. National Institute of Mental Health. (2024). *Anxiety disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/anxiety-disorders>
10. National Institutes of Health. (2023). *Lorazepam*. DailyMed. <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=73bfaeab-94db-48c2-a194-8b173025de78>

clients should be monitored frequently and have their dosage adjusted carefully according to their response; the initial dosage should not exceed 2 mg. Dosage for clients with severe hepatic insufficiency should be adjusted carefully according to client response. Benzodiazepines may cause fetal harm when administered to pregnant women.<sup>11</sup>

## OVERDOSAGE

Overdosage of benzodiazepines is manifested by varying degrees of central nervous system depression, ranging from drowsiness to coma. If overdose occurs, call 911 or the rapid response team during inpatient care. Treatment of overdosage is mainly supportive until the drug is eliminated from the body. Vital signs and fluid balance should be carefully monitored in conjunction with close observation of the client. An adequate airway should be maintained; intubation and mechanical ventilation may be required. The benzodiazepine antagonist flumazenil may be used to manage benzodiazepine overdose. There is a risk of seizure in association with flumazenil treatment, particularly in chronic users of benzodiazepines.

## CLIENT EDUCATION

Clients should be cautioned that driving a motor vehicle, operating machinery, or engaging in hazardous or other activities requiring attention and coordination should be delayed for 24 to 48 hours following administration of benzodiazepines or until the effects of the drug, such as drowsiness, have subsided. Alcoholic beverages should not be consumed for at least 24 to 48 hours after receiving benzodiazepines due to the additive effects on central nervous system depression. Hospitalized clients should be advised that benzodiazepines increase fall risk, and getting out of bed

11. National Institutes of Health. (2023). *Lorazepam*. DailyMed. <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=73bfaeab-94db-48c2-a194-8b173025de78>

unassisted may result in falling and potential injury if undertaken within eight hours of taking benzodiazepines.

## Antidepressants

Selective serotonin reuptake inhibitors (SSRIs) and serotonin-norepinephrine reuptake inhibitors (SNRIs) are commonly used as first-line treatments for anxiety. Less commonly used treatments for anxiety disorders are older classes of antidepressants, such as tricyclic antidepressants and monoamine oxidase inhibitors (MAOIs).<sup>12</sup> Read more about antidepressants in the “[Depressive Disorders](#)” chapter. If antidepressants are prescribed, they can take several weeks to reach their optimal effectiveness, so it is important to teach clients to give the medication appropriate time before reaching a conclusion about its effectiveness. They should be advised to avoid abruptly stopping the medication or without talking to their prescribing provider. Antidepressants should be tapered off slowly to safely decrease the dose because stopping them abruptly can cause withdrawal symptoms.

## Buspirone

Buspirone is a non-benzodiazepine medication indicated for the treatment of chronic anxiety. It is included in the class of medications called anxiolytics, but it is not chemically related to benzodiazepines, barbiturates, or other sedatives. Buspirone should not be taken concurrently with a monoamine oxidase inhibitor (MAOI) due to the risk of fatal side effects. It can also cause serotonin syndrome if used in combination with MAOIs, SSRIs, or SNRIs.<sup>13</sup>

12. National Institute of Mental Health. (2024). *Anxiety disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/anxiety-disorders>

13. National Institutes of Health. (2019). *Buspirone*. DailyMed. <https://www.nlm.nih.gov/>

Buspirone increases serotonin and dopamine levels in the brain. In contrast to benzodiazepines, buspirone must be taken every day for a few weeks to reach its full effect; it is not useful on an “as-needed” basis. A common side effect of buspirone is dizziness.<sup>14</sup> Buspirone is non-addictive and safer for long-term use than benzodiazepines.

## Beta-Blockers

Although beta-blockers are typically used to treat high blood pressure and other cardiac conditions, they can also be used to help relieve the physical symptoms of anxiety, such as rapid heartbeat, shaking, trembling, and flushing. These medications, when taken for a short period of time, can help people keep their physical symptoms under control. Beta-blockers can also be used “as needed” to reduce acute anxiety or as a preventive intervention for predictable forms of performance anxieties.<sup>15</sup> For example, some students who experience severe test anxiety that impairs their exam performance may take prescribed beta-blockers before their exams.

Common side effects of beta-blockers are fatigue, hypotension, dizziness, weakness, and cold hands. Beta-blockers are typically avoided in clients with asthma or diabetes.<sup>16</sup>

14. Wilson, T. K., Tripp, J. (2023). *Buspirone*. StatPearls [Internet].<https://www.ncbi.nlm.nih.gov/books/NBK531477/>
15. National Institute of Mental Health. (2024). *Anxiety disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/anxiety-disorders>
16. National Institute of Mental Health. (2023). *Mental health medications*. U.S. Department of Health & Human Services. <https://www.nimh.nih.gov/health/topics/mental-health-medications>

# Hydroxyzine

Hydroxyzine is type of antihistamine which may be prescribed to alleviate anxiety for individuals for whom benzodiazepines are not appropriate. It causes sedation, so it must be used cautiously if used in combination with opioids or barbiturates. Hydroxyzine is recommended for short-term use in the treatment of anxiety and tension. The effectiveness of hydroxyzine for long-term use, defined as more than 4 months, has not been systematically assessed in clinical studies. Therefore, it is advised that physicians periodically reassess the usefulness of the drug for each individual client.<sup>17</sup>

## Client Education

Clients should be educated about symptoms of their diagnosed anxiety disorder and techniques to manage it. For some individuals, even being aware that something is a symptom of anxiety, naming it, and connecting it to anxiety can help reduce the intensity of the anxiety.


Stress management techniques and coping strategies can help people with anxiety disorders calm themselves and enhance the effects of therapy. Research suggests that aerobic exercise can help some people manage their anxiety. Other coping strategies include meditation, yoga, journaling, prayer, or spending time in nature. Read more about stress management and coping strategies in the “[Stress, Coping, and Crisis Intervention](#)” chapter.

Support groups can be helpful for individuals experiencing anxiety disorders by sharing their problems and achievements with others experiencing similar symptoms. Talking with a trusted family member, friend, chaplain, or clergy member can also provide support.

Certain substances such as caffeine, some over-the-counter cold medicines, illicit drugs, and herbal supplements may aggravate the symptoms of anxiety

17. National Institutes of Health. (2019). *Hydroxyzine*. DailyMed.  
<https://www.nlm.nih.gov/>

disorders or interact with prescribed medications. Clients should be advised to avoid these substances.

View the following YouTube video on teaching adolescents  
 how to deal with anxiety disorders: [Mental Health Minute: Stress and Anxiety in Adolescents.](#)<sup>18</sup>

<sup>18</sup>. National Institute of Mental Health (NIMH). (2021, September 21). *Mental health minute: Stress and anxiety in adolescents* [Video]. YouTube. All rights reserved. <https://youtu.be/wr4N-SdekqY>

## 9.5 Obsessive - Compulsive Disorder

**Obsessive-compulsive disorder (OCD)** is a common chronic disorder in which a person has uncontrollable, reoccurring thoughts (obsessions) and/or behaviors (compulsions) they feel the urge to repeat over and over. These compulsions often temporarily relieve the stress/tension of the obsession.<sup>1</sup>

Historically, the relationship of OCD with anxiety disorders was strongly emphasized. However, the *Diagnostic and Statistical Manual of Mental Disorders, 5th edition* classified OCD under a separate group of disorders called “Obsessive-Compulsive and Related Disorders” due to the presence of obsessions and compulsions.<sup>2</sup> A systematic review found that lifetime psychiatric comorbidities were present in 69% of individuals with OCD, with anxiety disorders the most common comorbidity in children, and depressive disorders being the most common comorbidity in adults.<sup>3</sup>

### Signs and Symptoms

Symptoms of OCD can interfere with all aspects of life, such as work, school, and personal relationships. Symptoms may come and go, ease over time, or

1. National Institute of Mental Health. (2024). *Obsessive-compulsive disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/obsessive-compulsive-disorder-ocd>
2. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.
3. Sharma, E, Sharma, L. P., Balachander, S., Lin, B., Manohar, H., Khanna, P., Lu, C., Garg, K., Thomas, T. L., Au, A. C. L., Selles, R. R., Højgaard, D. R. M. A., Skarphedinsson, G, Stewart, S. E. (2021). *Comorbidities in obsessive-compulsive disorder across the lifespan: A systematic review and meta-analysis*. *Frontiers in Psychiatry*, 12:703701. <https://doi.org/10.3389%2Ffpsy.2021.703701>

worsen.<sup>4</sup> See Figure 9.7<sup>5</sup> for an artist's depiction of obsessive thoughts related to OCD.



Figure 9.7 Obsessive Compulsive Disorder

**Obsessions** are recurrent and persistent thoughts, urges, or images that are

4. National Institute of Mental Health. (2024). *Obsessive-compulsive disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/obsessive-compulsive-disorder-ocd>
5. “[Cope-With-Obsessive-Compulsive-Disorder-Step-15.jpg](#)” by unknown author at [wikihow.com](https://www.wikihow.com) is licensed under [CC BY-NC-SA 3.0](#)



experienced as intrusive and unwanted, and in most individuals cause anxiety.<sup>6</sup> Common obsessions are as follows<sup>7</sup>:

- Fear of germs or contamination
- Unwanted forbidden or taboo thoughts involving sex, religion, or harm
- Aggressive thoughts towards self or others
- Having things symmetrical or in a perfect order

**Compulsions** are repetitive behaviors or mental acts that a person feels driven to perform in response to an obsession or according to their rules that must be applied rigidly. The behaviors or mental acts are aimed at preventing or reducing anxiety or distress, but are excessive and are not connected realistically with what they are intended to neutralize or prevent.<sup>8</sup> Examples of compulsions include the following<sup>9</sup>:

- Excessive cleaning and/or handwashing
- Ordering and arranging things in a particular, precise way
- Repeatedly checking on things, such as repeatedly checking to see if the door is locked or that the oven is off
- Compulsive counting

6. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

7. National Institute of Mental Health. (2024). *Obsessive-compulsive disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/obsessive-compulsive-disorder-ocd>

8. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

9. National Institute of Mental Health. (2024). *Obsessive-compulsive disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/obsessive-compulsive-disorder-ocd>

Not all rituals or habits are compulsions; everyone double-checks things sometimes. For example, many people double-check that their doors are locked as they exit the vehicle. However, a person with OCD generally exhibits the following characteristics<sup>10</sup>:

- Spends at least one hour a day on these thoughts
- Can't control their thoughts or behaviors, even when those thoughts or behaviors are recognized as excessive
- Does not experience pleasure when performing the behaviors or rituals, but may feel relief from the anxiety the obsessive thoughts cause
- Experiences significant problems in their daily life due to these thoughts or behaviors

Sometimes compulsions are accompanied by a fear of potential consequences if they are not carried out. For this reason, an individual with OCD may become distressed if not able to complete a compulsive act.

Some individuals with OCD also have a tic disorder. Motor tics are sudden, brief, repetitive movements, such as eye blinking and other eye movements, facial grimacing, shoulder shrugging, and head or shoulder jerking. Vocal tics include repetitive throat-clearing, sniffing, or grunting sounds.<sup>11</sup>

People with OCD may try to cope by avoiding situations that trigger their obsessions or use alcohol or drugs to calm themselves. Although most adults with OCD have good insight and recognize what they are doing doesn't make sense, some may not realize that their behavior is out of the ordinary (i.e., they

10. National Institute of Mental Health. (2024). *Obsessive-compulsive disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/obsessive-compulsive-disorder-ocd>

11. National Institute of Mental Health. (2024). *Obsessive-compulsive disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/obsessive-compulsive-disorder-ocd>

demonstrate “poor insight”). Most children do not have good insight into their thoughts and behaviors, so parents or teachers typically recognize OCD symptoms in children.

## Risk Factors

The causes of OCD are unknown, but risk factors include genetics, brain structure and functioning, and environmental factors such as adverse childhood events (ACEs).<sup>12</sup>

## Genetics

Twin and family studies have shown that people with first-degree relatives (such as a parent, sibling, or child) who have OCD are at a higher risk for developing OCD. The risk is higher if the first-degree relative developed OCD as a child or teen.<sup>13</sup>

## Brain Structure and Functioning

Imaging studies have shown differences in the frontal cortex and subcortical structures of the brain in clients with OCD, but the connection with symptoms is not clear. Research is still underway because understanding potential causes may help determine specific, personalized treatments to treat OCD.<sup>14</sup>

12. National Institute of Mental Health. (2024). *Obsessive-compulsive disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/obsessive-compulsive-disorder-ocd>

13. National Institute of Mental Health. (2024). *Obsessive-compulsive disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/obsessive-compulsive-disorder-ocd>

14. National Institute of Mental Health. (2024). *Obsessive-compulsive disorder*.

## Environment

Research has found an association between childhood trauma (otherwise known as adverse childhood events (ACEs) and obsessive-compulsive symptoms. Some studies also found that children may develop OCD symptoms following a streptococcal infection, referred to as Pediatric Autoimmune Neuropsychiatric Disorders Associated with Streptococcal Infections (PANDAS). Children with PANDAS have a very sudden onset or worsening of their symptoms after a streptococcal infection, followed by a slow, gradual improvement.<sup>15</sup>

- ▶ Review information about adverse childhood events (ACEs) in the “[Mental Health and Mental Illness](#)” section in Chapter 1.

- ▶ Read more at the National Institute of Mental Health (NIMH) [PANDAS – Questions and Answers](#) webpage.

## Treatment

OCD is typically treated with medication, psychotherapy, or a combination of both. Although most clients with OCD improve with treatment, some clients continue to experience symptoms. Individuals with OCD may also have other

U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/obsessive-compulsive-disorder-ocd>

15. National Institute of Mental Health. (2024). *Obsessive-compulsive disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/obsessive-compulsive-disorder-ocd>

mental health disorders, such as anxiety, depression, and body dysmorphic disorder (a disorder in which someone mistakenly believes that a part of their body is abnormal). It is important to consider these other comorbid disorders when planning interventions related to treatment.<sup>16</sup>

## Medication

Selective serotonin reuptake inhibitors (SSRIs) are used to help reduce OCD symptoms. SSRIs often require higher daily doses in the treatment of OCD than of depression and may take 8 to 12 weeks to start working. If symptoms do not improve with SSRIs, research shows that some clients may respond well to an antipsychotic medication, especially if they also have a tic disorder.<sup>17</sup>

► Read more about SSRIs and antipsychotics in the “[Psychotropic Medications](#)” chapter.

## Psychotherapy

Psychotherapy can be an effective treatment for adults and children with OCD. Research shows that certain types of psychotherapy, including cognitive behavior therapy (CBT) and other related therapies (such as habit reversal training), can be as effective as medication for many individuals. Research

16. National Institute of Mental Health. (2024). *Obsessive-compulsive disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/obsessive-compulsive-disorder-ocd>

17. National Institute of Mental Health. (2024). *Obsessive-compulsive disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/obsessive-compulsive-disorder-ocd>

also shows that a type of CBT called **Exposure and Response Prevention (EX/RP)** is effective in reducing compulsive behaviors in clients with OCD. EX/RP includes spending time in the very situation that triggers compulsions (for example, touching dirty objects) but then being prevented from undertaking the usual resulting compulsion (handwashing). For many clients, EX/RP is an add-on treatment when SSRIs do not effectively treat OCD symptoms.<sup>18</sup>

## Other Treatment Options

In 2018 the FDA approved transcranial magnetic stimulation as an adjunct in the treatment of OCD in adults. **Repetitive Transcranial Magnetic Stimulation (rTMS)** uses a magnet to activate the brain. Unlike electroconvulsive therapy (ECT), in which electrical stimulation is more generalized, rTMS can be targeted to a specific site in the brain. A typical rTMS session lasts 30 to 60 minutes and does not require anesthesia. During the procedure, an electromagnetic coil is held against the forehead near an area of the brain that is thought to be involved in mood regulation. Short electromagnetic pulses are administered through the coil. The magnetic pulses easily pass through the skull and cause small electrical currents that stimulate nerve cells in the targeted brain region. The magnetic field is about the same strength as that of a magnetic resonance imaging (MRI) scan. The person generally feels a slight knocking or tapping on the head as the pulses are administered. The muscles of the scalp, jaw, or face may contract or tingle during the procedure, and mild headaches or brief light-headedness may result after the procedure. It is also possible that the procedure could cause a seizure, although this adverse effect is uncommon. Because the treatment is relatively new, long-term side effects are unknown.<sup>19</sup>

18. National Institute of Mental Health. (2024). *Obsessive-compulsive disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/obsessive-compulsive-disorder-ocd>

19. National Institute of Mental Health. (2024). *Brain stimulation therapies*. U.S.

## Client Education

In addition to teaching clients about the symptoms of OCD, prescribed medications, and other treatments, nurses should teach clients how to manage stress and anxiety associated with OCD:

- Create a consistent sleep schedule
- Make regular exercise a part of your routine
- Eat a healthy, balanced diet
- Seek support from capable family and friends

View a supplementary YouTube video<sup>20</sup> on OCD: [Obsessive compulsive disorder \(OCD\) – causes, symptoms & pathology](https://www.youtube.com/watch?v=l8Jofzx_8p4)

Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/brain-stimulation-therapies/brain-stimulation-therapies>

20. Osmosis for Elsevier. (2016, March 7). *Obsessive compulsive disorder (OCD) – causes, symptoms & pathology* [Video]. YouTube. All rights reserved. [https://www.youtube.com/watch?v=l8Jofzx\\_8p4](https://www.youtube.com/watch?v=l8Jofzx_8p4)

## 9.6 Post-Traumatic Stress Disorder

**Post-traumatic stress disorder (PTSD)** is diagnosed in individuals who have been exposed to a traumatic event with chronic stress symptoms lasting more than one month that are so severe they interfere with relationships, school, or work. PTSD was formerly classified as an anxiety disorder but was placed in a new diagnostic category in the DSM-5 called “Trauma and Stressor-Related Disorders.”

Post-traumatic stress disorder has similar characteristics to severe anxiety and phobia-related disorders because of the physiological stress response that occurs. Post-traumatic stress disorder (PTSD) can develop in some people who have experienced a shocking, frightening, or dangerous event. It is natural to feel afraid during and after a traumatic situation, and the “fight-or-flight” stress response is a physiological reaction intended to protect a person from harm. Most people recover from the range of reactions that can occur after experiencing trauma. However, people who do not recover from these reactions and continue to experience problems are diagnosed with PTSD. People who have PTSD may feel stressed or frightened, even when they are not in danger.<sup>1</sup>

### Symptoms

Symptoms of PTSD typically begin three months after the traumatic incident, but they may also begin years afterward. If symptoms occur within one month of the traumatic event, it is diagnosed as acute stress disorder. Symptoms must last more than a month and be severe enough to interfere with social or occupational functioning to be considered PTSD. The course of the illness varies; some people recover within six months, while others have

1. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>



symptoms that last much longer. In some people, the condition becomes chronic.<sup>2,3</sup>

To be diagnosed with PTSD, an adult must have the following types of symptoms for at least one month<sup>4,5</sup>:

- At least one “re-experiencing” symptom
- At least one “avoidance” symptom
- At least two “arousal and reactivity” symptoms
- At least two “cognition and mood” symptoms

## Re-Experiencing Symptoms

Re-experiencing symptoms include the following<sup>6</sup>:

- Flashbacks—reliving the trauma over and over, including physical

2. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>
3. American Psychiatric Association. (2013). *Desk reference to the diagnostic criteria from DSM-5*.
4. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>
5. American Psychiatric Association. (2013). *Desk reference to the diagnostic criteria from DSM-5*.
6. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>

symptoms like a racing heart or sweating

- Bad dreams
- Frightening thoughts

Re-experiencing symptoms can start from the person's own thoughts and feelings. Words, objects, or situations that are reminders of the event can also trigger re-experiencing symptoms. Re-experiencing symptoms may cause problems in a person's everyday routine and relationships.<sup>7</sup>

## Avoidance Symptoms

Avoidance symptoms are as follows<sup>8</sup>:

- Staying away from places, events, or objects that are reminders of the traumatic experience
- Avoiding thoughts or feelings related to the traumatic event

These symptoms may cause a person to change their personal routine. For example, after a car accident, a person who usually drives may avoid driving or riding in a car.

7. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>

8. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>

## Arousal and Reactivity Symptoms

Arousal and reactivity symptoms include the following<sup>9</sup>:

- Being easily startled
- Feeling tense or “on edge”
- Having difficulty sleeping
- Having angry outbursts

Arousal symptoms are usually constant instead of being triggered by things that remind one of the traumatic events. These symptoms can make the person feel stressed and angry and can make it hard to do daily tasks, such as sleeping, eating, or concentrating.<sup>10</sup>

## Cognition and Mood Symptoms

Cognition and mood symptoms are as follows<sup>11</sup>:

- Trouble remembering key features of the traumatic event
- Negative thoughts about oneself or the world
- Distorted feelings like guilt or blame
- Loss of interest in enjoyable activities

9. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>

10. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>

11. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>

Cognition and mood symptoms can begin or worsen after the traumatic event and make the person feel alienated or detached from friends or family members.<sup>12</sup>

It is natural to have some of these types of symptoms for a few weeks after a traumatic event. However, when the symptoms last more than a month, seriously affect a person's functioning, and are not related to substance use, medical illness, or anything except the event itself, they can be symptoms of PTSD. PTSD is also often accompanied by depression, substance abuse, or other anxiety disorders.<sup>13</sup>

## Life Span Considerations

Children and teens can have extreme reactions to trauma, but they may exhibit different symptoms than adults. Symptoms of PTSD can be seen in young children (less than six years old) and may include the following:

- Bedwetting after having learned to use the toilet
- Forgetting how to talk or being unable to talk (i.e., selective mutism)
- Acting out the scary event during playtime
- Being unusually clingy with a parent or other adult

Older children and teens are more likely to show symptoms similar to those seen in adults. They may also develop disruptive, disrespectful, or destructive behaviors. Hypersexual behavior may occur if the trauma was related to a sexual assault. Older children and teens may also feel guilty for not preventing

12. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>

13. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>

injury or death in certain traumatic situations and may have thoughts of revenge.<sup>14</sup>

## Risk and Resilience Factors

Anyone can develop PTSD at any age, including war veterans; children who have experienced trauma; or adults who have experienced a physical or sexual assault, abuse, accident, disaster, being a refugee, or some other serious event. According to the National Center for PTSD, about 6 out of every 100 people will experience PTSD symptoms at some point in their lives. Women are more likely to develop PTSD than men, and genes may make some people more likely to develop PTSD than others.<sup>15</sup>

Not everyone with PTSD has directly experienced a dangerous event. Some people develop PTSD after a friend or family member experiences danger or harm. The sudden, unexpected death of a loved one can also lead to PTSD.<sup>16</sup> PTSD is triggered by events that are perceived to be life-threatening, and this can vary from individual to individual. It disrupts the general sense of safety that allows individuals to function in the world.

It is important to remember that not everyone who lives through a dangerous event develops PTSD. In fact, most people will not develop the disorder. Many factors play a part in whether a person will develop PTSD. Risk factors make a

14. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>

15. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>

16. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>

person more likely to develop PTSD, but other factors, called resilience factors, can help reduce the risk of developing the disorder or promote recovery from the disorder.<sup>17</sup>

These factors increase the risk for developing PTSD<sup>18</sup>:

- Living through dangerous events and traumas
- Being injured from a traumatic event
- Seeing another person hurt or seeing a dead body
- Experiencing childhood trauma or adverse childhood events (ACEs)
- Feeling horror, helplessness, or extreme fear
- Having little or no social support after the event
- Dealing with extra stress after the event, such as loss of a loved one, pain and injury, or loss of a job or home
- Having a history of mental illness or substance abuse

▶ Review information about adverse childhood events (ACEs) in the “[Mental Health and Mental Illness](#)” section of Chapter 1.

Resilience factors that may promote recovery after trauma include the following<sup>19</sup>:

17. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>
18. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>
19. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S.

- Receiving support from other people, such as friends and family
- Finding a support group after a traumatic event
- Learning to feel good about one's own actions in the face of danger, recognizing that we controlled what we could in an uncontrollable situation
- Having a positive coping strategy or a way of getting through the bad event and learning from it
- Being able to act and respond effectively despite feeling fear

If a child or adolescent discloses traumatic events to caregivers, teachers, or other adults, it is important for them to feel their concerns are validated by the adult in order to develop resilience.

Researchers are studying the importance of risk and resilience factors, as well as the impact of genetics and neurobiology. With more research, it may be possible to someday predict who is likely to develop PTSD and how to prevent it from occurring.<sup>20</sup> See Figure 9.8<sup>21</sup> for an image of a veteran with PTSD using a service dog as an effective coping strategy.

Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>

20. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>

21. "[Local veterans group gets national exposure 141127-F-ZZ999-001.jpg](#)" by unknown author is licensed in the [Public Domain](#)





Figure 9.8 A Veteran with PTSD Using a Service Dog as a Coping Strategy

## Treatments

For people with PTSD, treatments include medications, psychotherapy, or a combination of both. Everyone is different, and PTSD affects people differently, so a treatment that works for one person may not work for another. It is important for anyone with PTSD to be treated by a mental health provider who has experience treating PTSD.<sup>22</sup>

If someone with PTSD is also experiencing an ongoing trauma, such as an abusive relationship, both of the problems need to be addressed. Other ongoing problems can include panic disorder, depression, substance use disorder, and suicidal ideation.<sup>23</sup>

22. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>

23. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S.



## Medications

Antidepressants can help control PTSD symptoms such as sadness, worry, anger, and feeling numb inside. For example, two FDA-approved medications for PTSD are the antidepressants sertraline and paroxetine. Other medications may be helpful for treating specific PTSD symptoms, such as sleep problems and nightmares.<sup>24</sup> Read more information about antidepressants in the “[Antidepressants](#)” section of the “Psychotropic Medications” chapter.

## Psychotherapy

Psychotherapy can occur one-on-one or in a group setting. It typically lasts 6 to 12 weeks, but it can continue for as long as the individual finds it helpful. Research shows that additional support from family and friends can be an important part of recovery.<sup>25</sup>

Many types of psychotherapy can help people with PTSD. Some target the symptoms of PTSD directly, whereas other therapies focus on social, family, or job-related problems. Effective psychotherapies emphasize key components such as education about symptoms, identification of triggers or symptoms, and skills to manage the symptoms. Examples of psychotherapies used to treat PTSD are cognitive behavioral therapy, exposure therapy, eye movement

Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>

24. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>

25. National Institute of Mental Health. (2024). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>

desensitization and reprocessing, animal therapy programs, and MDMA-assisted psychotherapy.

Cognitive behavioral therapy (CBT) combined with exposure therapy helps people face and control their fear by gradually exposing them to the trauma they experienced in a safe way. It uses imagining, writing, or visiting the place where the event happened to help reduce the intensity of PTSD symptoms. Read more about CBT in the “[Depressive Disorders](#)” chapter.


Cognitive restructuring helps individuals create new thought patterns about the trauma. Sometimes events are remembered differently than how they truly happened, and individuals may experience feelings of guilt or shame in relation to the trauma, regardless of whether they played an active role or not. They may feel guilt or shame about something that is not their fault. Therapists can help individuals assess the trauma through a variety of lenses to process and continue the healing journey.


Eye movement desensitization and reprocessing (EMDR) is a psychotherapy treatment that was originally designed to alleviate the distress associated with traumatic memories. During EMDR therapy, the client attends to emotionally disturbing material in brief sequential doses while simultaneously focusing on an external stimulus. Therapist-directed lateral eye movements are the most commonly used external stimulus, but a variety of other stimuli such as hand-tapping and audio stimulation are also used. It is hypothesized that EMDR therapy facilitates the individual’s access of their traumatic memory network so that new associations can be forged between the traumatic memory and more adaptive memories or information. These new associations are thought to result in elimination of emotional distress and development of cognitive insights.<sup>26,27</sup>

26. EDMR Institute. (n.d.) *What is EDMR?* <https://www.emdr.com/what-is-emdr/>

27. American Psychological Association. (2025). *Clinical practice guidelines for the treatment of post-traumatic stress disorder: Eye Movement*

Animal assisted intervention (AAI), also referred to as animal therapy, is a commonly used complementary treatment for PTSD. It most often includes dogs or horses. A systematic review examined the outcomes in studies using AAI with trauma survivors. Although the reviewed studies were diverse and limited, all reported positive outcomes of AAI, such as reduced depression, PTSD symptoms, and anxiety.<sup>28</sup>

View a supplementary YouTube video<sup>29</sup> on EMDR therapy:  
 [What is Eye Movement Desensitization Reprocessing Therapy?](#)

 View a supplementary YouTube video on PTSD<sup>30</sup>: [NIMH-](#)

*Desensitization and Reprocessing (EMDR) Therapy.* <https://www.apa.org/ptsd-guideline/treatments/eye-movement-reprocessing>

28. O'Haire, M., Guerin, N., & Kirkham, A. (2015). Animal-Assisted Intervention for trauma: a systematic literature review. *Frontiers in Psychology*, 6. <https://doi.org/10.3389/fpsyg.2015.01121>

29. Pysch Hub. (2019, April 16). *What is eye movement desensitization reprocessing therapy?* [Video]. YouTube. All rights reserved. <https://www.youtube.com/watch?v=1IPsBPH2M1U>

30. National Institute of Mental Health (NIMH). (2021, June 21). *NIMH-funded researcher Dr. Barbara Rothbaum discusses post-traumatic stress*



Funded Researcher Dr. Barbara Rothbaum Discusses Post-Traumatic Stress Disorder.

*disorder* [Video]. YouTube. All rights reserved. <https://youtu.be/wlcWlbM4hLE>

## 9.7 Applying the Nursing Process to Anxiety Disorders

People with anxiety disorders rarely require hospitalization unless they are suicidal, although anxiety can occur with other mental disorders requiring hospitalization. As a nurse working with individuals with diagnosed anxiety disorders, be aware of your self-reaction. It is not uncommon to have feelings of frustration, especially if you feel as if the symptoms are a matter of choice or under the client's control. The client often acknowledges the fear is unrealistic or exaggerated but continues to engage in avoidant behavior. Recall that avoidant behavior is a symptom, and behavioral changes are accomplished slowly with treatment.<sup>1</sup>

It is also important to be aware that hospitalized clients may develop anxiety in association with other medical conditions (i.e., chronic obstructive pulmonary disease [COPD], angina, or hyperthyroidism) or medical procedures. Anxiety is a nursing diagnosis, as well as a potential mental health disorder. While implementing interventions that address medical conditions, often the nurse must also implement interventions that address associated anxiety.

### Assessment (Recognize Cues)

#### Mental Status Examination

See Table 9.7a for common findings when assessing a client with an anxiety disorder. (See expected findings for these components of a mental status examination in the “[Assessment](#)” section in Chapter 4.) Critical findings that require immediate notification of the provider are bolded with an asterisk.

1. Halter, M. (2022). *Varc Carolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

Table 9. 7a Common Findings During Mental Status Examinations for Clients With Anxiety Disorders<sup>2</sup>

2. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

Mental Status Examination Component	Common Findings in Anxiety Disorders (*Indicates immediately notify provider)
<b>Signs of Distress</b>	<p>Determine the client's current level of anxiety (mild, moderate, severe, or panic) and assess for risk of suicide or self-harm. If a client is experiencing a panic attack, they may exhibit <b>*shortness of breath</b>, <b>*chest pain</b> or tightness, a choking sensation, dizziness, <b>*palpitations</b>, nausea, abdominal pain, diaphoresis, or a fear of dying or losing control.</p> <p>Anxiety often coexists with depression and can lead to feelings of being overwhelmed and unable to cope, which can be a significant risk factor for suicidal ideation.</p> <p><b>*Note: Suicidal ideations indicate increased risk for self-injury, suicide, or injury to others and must be reported to provider. Do not leave clients alone if statements such as these are being made.</b></p>
<b>Level of Consciousness and Orientation</b>	Typically alert and oriented to person, place, time, and situation. However, during acute panic episodes, transient confusion or disorientation may occur.
<b>Appearance and General Behavior</b>	May appear nervous and restless with fidgeting, wringing hands, foot tapping, facial tension, and tremors. Posture may be rigid or defensive. May avoid eye contact, appear guarded, or display frequent checking behaviors (e.g., looking around the environment for threats). May experience urinary frequency. May appear disheveled. May report altered sleep patterns. <b>*Verbal and nonverbal threats of harm</b> or <b>*self-harming behaviors</b> such as cutting, picking at skin, knocking head against the wall, tightening string or items on wrists, or stabbing self with anything fashioned into a weapon should be immediately reported to the provider.
<b>Speech</b>	Speech may be rapid, pressured, or hesitant; may include stammering, especially in social anxiety; often driven by urgency to express worries or fears.
<b>Motor Activity</b>	Psychomotor agitation is common (e.g., pacing, fidgeting, tapping fingers) with restless movements and muscle tension evident. May display shakiness or tremors.

<b>Mood and Affect</b>	Affect is often irritable. Mood is often reported as “worried,” “on edge,” “nervous,” or “panicked.” Mood and affect are generally congruent.
<b>Thought and Perception</b>	Thought content is dominated by worry, anticipatory fear, or intrusive thoughts (especially in clients with OCD). Thought process may be circumstantial with excessive unnecessary details, tangents, and related ideas included before addressing the core topic. Perceptions are generally intact with no hallucinations.
<b>Attitude and Insight</b>	Often cooperative but may be guarded or overly compliant. May recognize thoughts as excessive or irrational (especially in clients with GAD or OCD), but still struggles to control them.
<b>Cognitive Abilities and Level of Judgment</b>	Usually intact, although attention and concentration may be impaired by excessive worry. Working memory and decision-making may be affected with severe anxiety or panic.

## Psychosocial Assessment

It is helpful to begin the psychosocial assessment by obtaining the reason why the client is seeking health care in their own words and focus on what factors could be contributing to their anxiety. For example, the client may identify a problem such as a relationship issue, stressful job, or school challenges that could be addressed by counseling.<sup>3</sup>

A comprehensive psychosocial assessment includes the following components:

- Reason for seeking health care (i.e., “chief complaint”)

3. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.



- Thoughts of suicide or self injury
- Cultural assessment
- Spiritual assessment
- Family dynamics
- Current and past medical history
- Current medications
- History of previously diagnosed mental health disorders
- Previous hospitalizations
- Educational background
- Occupational background
- History of exposure to psychological trauma, violence, and domestic abuse
- Substance use (tobacco, alcohol, recreational drugs, misused prescription drugs)
- Family history of mental illness
- Coping mechanisms
- Functional ability/Activities of daily living

After identifying the reason the client is seeking health care, additional focused questions are used to obtain detailed information. The mnemonic PQRSTU can be used to ask questions in an organized fashion. See Table 9.7b for a sample PQRST assessment for anxiety.

Table 9.7b Sample PQRSTU Questions for Assessing Anxiety

PQRSTU	Sample Questions
<b>Provocation/ Palliation</b>	“What makes your anxiety get worse?”  “What helps calm your anxiety?”
<b>Quality</b>	“Can you describe what it feels like when you have anxiety?”
<b>Region</b>	“Do you feel the anxiety in specific places in your body?”
<b>Severity</b>	“On a scale of 0 to 10, how bad is your anxiety right now?”
<b>Timing/ Treatment</b>	“When did the anxiety start? How long does it usually last? Is it constant or intermittent?”
<b>Understanding</b>	“What do you think is causing your anxiety?”

## SUICIDE AND SELF INJURY SCREENING

Clients being evaluated or treated for anxiety may have suicidal ideation. It is important for the nurse to introduce suicide screening in a way that helps the client understand its purpose and normalize questions that might otherwise seem intrusive. The Patient Safety Screener (PSS-3) is an example of a brief screening tool to detect suicide risk in all client presenting to acute care settings.<sup>4</sup>

4. Suicide Prevention Resource Center. (n.d.). *The patient safety screener: A brief*

**Non-suicidal self-injury (NSSI)** refers to intentional self-inflicted destruction of body tissue without suicidal intention and for purposes not socially sanctioned. Common forms of NSSI include behaviors such as cutting, burning, scratching, and self-hitting. It is considered a maladaptive coping strategy without the desire to die. NSSI is a common finding among adolescents and young adults in psychiatric inpatient settings.<sup>5</sup>

Review information about suicide screening, the Patient Safety Screener, and screening for non-suicidal self-injury (NSSI) in the “[Assessment](#)” section of the Applying the Nursing Process to Mental Health Care” chapter.

## CULTURAL ASSESSMENT

Cultural Formulation Interview (CFI) questions help nurses understand a client’s cultural background and how it influences their experience of mental health symptoms, including anxiety.<sup>6</sup> Sample CFI questions focused specifically on understanding anxiety within a cultural context include the following:

- Cultural Definition of the Problem
  - “Can you tell me more about your experience of feeling anxious or

*tool to detect suicide risk.* <https://sprc.org/micro-learning/patientsafetyscreener>

5. Cipriano, A., Cella, S., & Cotrufo, P. (2017). Nonsuicidal self-injury: A systematic review. *Frontiers in Psychology*, 8. <https://www.frontiersin.org/articles/10.3389/fpsyg.2017.01946/full>
6. DeSilva, R., Aggarwall, N. K., & Lewis-Fernandez, R. (2015). The DSM-5 cultural formulation interview and the evolution of cultural assessment in psychiatry. *Psychiatric Times*, 32(6). <https://www.psychiatrictimes.com/view/dsm-5-cultural-formulation-interview-and-evolution-cultural-assessment-psychiatry>

nervous?”

- “What do you call these feelings or symptoms in your own words or language?”
- “What do people in your family or community think is going on when someone feels like this?”
- Cultural Perceptions of Cause, Context, and Support
  - “Why do you think this anxiety started?”
  - “Are there any stressors in your life or community that you believe contribute to these feelings?”
  - “Do you believe your anxiety is related to physical health, spiritual issues, or emotional distress?”
  - “What do others in your family or community (parents, elders, spiritual leaders) believe causes anxiety?”
- Cultural Factors Affecting Coping and Help-Seeking
  - “How have you tried to deal with your anxiety so far?”
  - “Are there any traditional practices, herbs, or dietary supplements you have tried to use to manage it?”
  - “Have you sought help from community leaders, healers, or religious figures?”
  - “What kinds of treatment or help do you think would be most useful or acceptable to you?”
- Cultural Features of the Nurse–Client Relationship
  - “Are there any concerns you have about talking to a mental health professional?”
  - “Would you feel more comfortable speaking with someone of a similar gender, cultural, or religious background?”
  - “What would help you feel more supported or understood during treatment?”

Findings from the cultural formulation interview are used to individualize a client’s plan of care to their preferences, values, beliefs, and goals.

Cultural beliefs can affect an individual's expression of their feelings of anxiety. An example of a culture-mediated response related to anxiety and panic disorder is ataque de nervios (ADN) or “attack of the nerves” that may be exhibited in Hispanic populations. Symptoms of ADNs can vary widely but are typically described as an experience of distress characterized by a general sense of being out of control. The most common symptoms include uncontrollable shouting, attacks of crying, trembling, and heat in the chest rising into the head. Suicidal gestures, seizures, or fainting episodes may be observed. These symptoms are reported to typically occur following a distressing event such as an interpersonal conflict or the death of a loved one.<sup>7</sup>

## SPIRITUAL ASSESSMENT

The FICA Spiritual History Tool is a widely used assessment model for evaluating a client's spiritual beliefs and how they may influence health, illness, and coping. It's especially helpful in understanding how clients with anxiety draw on spirituality or religion for support—or how spiritual distress may be contributing to feelings of anxiety. Spiritual distress is very common for clients experiencing serious illness, and nurses assist clients to adopt healthy coping strategies to deal with these life events. Addressing a client's

7. Keough, M. E., Timpano, K. R., & Schmidt, N. B. (2009). Ataques de nervios: Culturally bound and distinct from panic attacks? *Depression & Anxiety*, 26(1), 16-21. <https://onlinelibrary.wiley.com/doi/10.1002/da.20498>

spirituality and advocating spiritual care have been shown to improve clients' health and quality of life.<sup>8,9</sup>

The FICA Spiritual History Tool© is a common tool used to gather information about a client's spiritual history and preferences. FICA© is a mnemonic for the domains of Faith, Importance, Community, and Address in Care.<sup>10</sup> Table 9.7c summarizes a sample FICA Spiritual Assessment for a client experiencing anxiety.

Table 9.7c Sample FICA Spiritual Assessment Questions for Clients with Anxiety

8. Pilger, C., Molzahn, A. E., de Oliveira, M. P., & Kusumota, L. (2016). The relationship of the spiritual and religious dimensions with quality of life and health of patients with chronic kidney disease: An integrative literature review. *Nephrology Nursing Journal: Journal of the American Nephrology Nurses' Association*, 43(5), 411–426. <https://pubmed.ncbi.nlm.nih.gov/30550069/>
9. Puchalski, C., Jafari, N., Buller, H., Haythorn, T., Jacobs, C., & Ferrell, B. (2020). Interprofessional spiritual care education curriculum: A milestone toward the provision of spiritual care. *Journal of Palliative Medicine*, 23(6), 777–784. <https://doi.org/10.1089/jpm.2019.0375>
10. GW School of Medicine & Health Sciences. (n.d.). *Clinical FICA tool*. <https://smhs.gwu.edu/spirituality-health/program/transforming-practice-health-settings/clinical-fica-tool>

Domain	Sample Assessment Question	Sample Client Response
<b>Faith</b>	“Do you consider yourself spiritual or religious? What gives your life meaning?”	“Yes, I believe in God. My faith gives me strength when I feel overwhelmed.”
<b>Importance</b>	“What importance does your faith or belief have in your life? Has it influenced how you cope with stress or your anxiety?”	“I believe that everything happens for a reason. I try to trust in God when I feel anxious, but sometimes I wonder if I’m being punished for something I have done.”
<b>Community</b>	“Are you part of a spiritual or religious community? Does participation in this community provide support when you’re feeling anxious or stressed?”	“Sometimes I go to church on Sundays. Being with others who have similar beliefs helps me feel less alone.”
<b>Address in Care</b>	“How would you like me (or the health care team) to address spiritual issues during your care? Would you like to speak with a chaplain?”	“Yes I would be interested in speaking to a chaplain.”

Nurses may recognize cues of spiritual distress or beliefs in divine punishment that may exacerbate feelings of anxiety. Nurses can offer to connect the client with a chaplain or spiritual care services. Spiritual goals may be included in the nursing care plan if the client finds them valuable.

## FAMILY DYNAMICS

Family dynamics are included in a psychosocial assessment, especially for children, adolescents, and older adults. **Family dynamics** refers to the patterns of interactions among relatives, their roles and relationships, and the various factors that shape their interactions. Because family members rely on each other for emotional, physical, and economic support, they are primary sources of relationship security or stress. Family dynamics and the quality of family relationships can have either a positive or negative impact on an individual’s mental health. For example, secure and supportive family relationships can provide love, advice, and care, whereas stressful family

relationships can be burdened with arguments, unhealthy relationships, and a lack of support.<sup>11</sup>

Unhealthy family dynamics can cause children to experience trauma and stress as they grow up. This type of exposure, known as adverse childhood experiences (ACEs), is linked to an increased risk of developing physical and mental health problems such as heart, lung, and liver disease, depression, and anxiety. Unhealthy family dynamics also correlate with an increased risk of substance use and addiction among adolescents.<sup>12</sup>

- ▶ Review information about adverse childhood experiences (ACEs) in the “[Mental Health and Mental Illness](#)” section of Chapter 1.

## Screening Tools

The Severity Measure for Generalized Anxiety Disorder in Adults is a common tool for measuring anxiety. High scores may indicate generalized anxiety disorder or panic disorder, although it can also be associated with major depressive disorder.

11. Jabbari, B., Schoo, C., & Rouster, A. S. (2023). *Family dynamics*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK560487/>
12. Jabbari, B., Schoo, C., & Rouster, A. S. (2023). *Family dynamics*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK560487/>



▶ View the [Severity Measure for Generalized Anxiety Disorder—Adult](#) PDF screening tool.

## Diagnostic and Lab Work

When assessing for anxiety disorders, the provider will typically order lab work to rule out common medical causes of anxiety, such as hyperthyroidism, hypoglycemia, hypercalcemia, hyperkalemia, hyponatremia, or hypoxia. Toxicology screen to identify potential substance abuse. Review and/or monitor the results of these tests as part of the nursing assessment.

## Lifespan Considerations

### CHILDREN AND ADOLESCENTS

All children experience some anxiety, and anxiety is expected at specific times of a child's development. For example, from approximately age eight months through the preschool years, healthy children may show anxiety when separated from their parents or caregivers, called separation anxiety. Young children also commonly have fears, such as fear of the dark, storms, animals, or strangers. Anxiety is considered normal when situational. Consider the fear of dangerous situations such as approaching a rattlesnake or standing on a steep cliff; at crucial times such as these, anxiety is important because it provides safety. Anxiety can also be motivational if it drives clients to accomplish goals.

When a child is overly worried or anxious, a nurse's initial assessment should determine if conditions in the child's environment are causing this feeling. For example, is the anxiety resulting from being bullied or from adverse childhood experiences (ACEs)? If so, protective interventions should be put into place. If no realistic threat exists and the anxiety causes significant life

dysfunction, then the child should be referred to a mental health provider to determine if an anxiety disorder exists.

Children with anxiety disorders are overly tense or fearful and their worries interfere with daily activities. Some children may require significant amounts of reassurance. Because anxious children may also be quiet, compliant, and eager to please, their feelings of anxiety can be easily missed. When a child does not outgrow the typical fears and anxieties in childhood or when there are so many fears and worries they interfere with school, home, or play activities, the child may be diagnosed with an anxiety disorder.

Anxiety can also make children irritable and angry and can include physical symptoms like fatigue, headaches, stomachaches, or trouble sleeping. Early treatment of anxiety disorders in children and adolescents can enhance friendships, social and academic potential, and self-esteem.

## **OLDER ADULTS**

Anxiety disorders in older adults may be impacted by age-related physical, psychological, and social factors. Older adults may not use the word “anxiety”, but may instead describe feeling “worried,” “nervous,” “or “tense.” Some older adults may view anxiety or expression of their emotions as a personal weakness due to generational beliefs. Nurses can normalize mental health conversations and reduce stigma by framing care in terms of well-being, quality of life, and stress reduction. Hearing or memory issues may affect how an older adult responds to questions, so nurses should allow more time for responses. Death of a spouse, family members, or friends can lead to increased feelings of loss and isolation. It is often beneficial to include social engagement and reminiscence therapy in treatment of anxiety in older adults. Chronic illnesses like chronic obstructive pulmonary disease (COPD), cardiovascular disease, or diabetes mellitus can increase levels of anxiety due to medical concerns and may be compounded by issues like fear of falling, losing autonomy, or being placed in a long-term care setting. Symptoms of anxiety may be mistaken for signs of early dementia, such as restlessness and difficulty concentrating. Side effects from certain medications can also cause

or worsen feelings of anxiety, such corticosteroids, bronchodilators, or stimulants. Benzodiazepines are not recommended for use by older adults to treat anxiety due to side effects of oversedation, confusion, and increased fall risk.

## Diagnosis (Analyze Cues)

Anxiety is a NANDA-I nursing diagnosis and described as “vague, uneasy feeling of discomfort or dread accompanied by an autonomic response; a feeling of apprehension caused by anticipation of danger. It is an alerting sign that warns of impending danger and enables the individual to take measures to deal with the threat.”<sup>13</sup> Review signs and symptoms of anxiety in the preceding “Assessment” subsection or consult an evidence-based nursing care plan resource for defining characteristics of this nursing diagnosis.

## Outcome Identification (Generate Solutions)

The overall goal for clients experiencing anxiety is to reduce the frequency and intensity of the anxiety symptoms. SMART outcomes are individualized to the client’s specific diagnosed conditions, situational factors, and current status. Planning outcomes in small, attainable steps can help a client gain a sense of control over their anxiety.<sup>14</sup>

Examples of SMART outcomes include:

- The client’s vital signs will return to baseline within one hour.
- The client will identify and verbalize symptoms of anxiety by the end of the shift.

13. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

14. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

- The client will verbalize three preferred stress management and coping strategies for controlling their anxiety by the end of Week 1.

## Planning (Generate Solutions)

The client should be encouraged to participate in planning outcomes and interventions tailored to their situation and needs. This will increase the likelihood that the interventions will be successful. Keep in mind that clients with severe anxiety or panic may not be able to participate in planning and rely on the nurse to take a directive role.<sup>15</sup>

## Safety

Immediately planning and implementing interventions to maintain client safety receive priority. Review interventions for clients with a risk for suicide in the “[Application of the Nursing Process in Mental Health Care](#)” chapter.

If a client’s anxiety continues to escalate and they become severely agitated, measures must be taken to keep them and others safe. The nurse may find that administering prescribed medications, initiating time in a quiet room, seclusion, or restraints is required. Review crisis intervention in the “[Stress, Coping, and Crisis Intervention](#)” chapter. Review information regarding the use of seclusion and restraints in the “[Psychosis and Schizophrenia](#)” chapter. Review ANA guidelines on using restraints in the “[Client Rights](#)” section of the “Legal and Ethical Considerations in Mental Health Care” chapter and information on safely implementing restraints in the “[Workplace Violence](#)” section of the “Trauma, Abuse, and Violence” chapter.

## Mild to Moderate Anxiety

The nurse can reduce a client’s anxiety level and prevent escalation by

15. Halter, M. (2022). *Varc Carolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

providing a calm presence in a quiet environment, acknowledging their feelings of distress, and actively listening. Using therapeutic techniques like open-ended questions, distraction, exploring, and seeking clarification can be used to relieve the client's feelings of tension and focus on previously successful coping strategies.<sup>16</sup> Review therapeutic communication techniques in the "[Therapeutic Communication and the Nurse-Client Relationship](#)" chapter.

It may be helpful to encourage the client to participate in physical activities that may provide relief from tension and increase endorphin levels. For example, the nurse can encourage the mildly anxious client to walk or play ping-pong.<sup>17</sup>

## Severe Anxiety to Panic

A person experiencing severe anxiety to panic is often unable to solve problems or grasp what is going on in the environment. The nurse should also remain with a client experiencing acute, severe, or panic levels of anxiety. Therapeutic communication should focus on helping the client feel safe. Firm, short, simple statements using a slow, low-pitched voice are helpful.<sup>18</sup>

In addition to keeping the client and others safe, priority nursing interventions for a client experiencing severe anxiety focus on the client's physical needs, such as fluids to prevent dehydration, blankets for warmth, and rest to prevent exhaustion. If a person continues to constantly move or

16. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

17. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

18. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

pace despite interventions, high-calorie finger foods may be offered to maintain their nutrition.<sup>19</sup> Read additional interventions related to crisis intervention in the “[Stress, Coping, and Crisis Intervention](#)” chapter.

## Implementation (Take Action)

### Nursing Interventions for Anxiety Based on Categories of the APNA Implementation Standard

Nursing interventions for anxiety disorders can be categorized based on the American Psychiatric Nurses Association (APNA) standard for *Implementation* that includes the *Coordination of Care; Health Teaching and Health Promotion; Pharmacological, Biological, and Integrative Therapies; Milieu Therapy; and Therapeutic Relationship and Counseling*. Read more about these subcategories in the “[Application of the Nursing Process in Mental Health Care](#)” chapter. See examples of interventions for each of these categories for clients with anxiety disorders in Table 9.7d.

Table 9.7d Examples of Nursing Interventions for Anxiety by APNA Subcategories<sup>20, 21</sup>

19. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.
20. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.
21. Ackley, B., Ladwig, G., Makic, M. B., Martinez-Kratz, M., & Zanoliti, M. (2020). *Nursing diagnosis handbook: An evidence-based guide to planning care* (12th ed.). Elsevier.

Subcategory of the APNA Standard of Implementation	The nurse will ...	Rationale
<b>Coordination of Care</b>	<ul style="list-style-type: none"> <li>– Collaborate with interdisciplinary team members to develop a comprehensive anxiety management plan.</li> <li>– Ensure continuity of care during transitions.</li> <li>– Facilitate referrals to specialists or community programs.</li> </ul>	Anxiety is often multifactorial, requiring input from various disciplines. Coordinated care ensures consistent, holistic support across settings and improves outcomes.
<b>Health Teaching and Health Promotion</b>	<ul style="list-style-type: none"> <li>– Teach about the physiological signs of anxiety.</li> <li>– Teach stress management techniques (eating a healthy diet, avoiding caffeine and other stimulants, participating in regular physical activity, obtaining adequate sleep, using mindfulness activities).</li> <li>– Teach positive coping strategies (meditation, prayer, deep breathing exercises, grounding, yoga, journaling, mindfulness activities).</li> </ul>	Increasing the client's understanding of the symptoms of anxiety helps empower self-management. Lifestyle changes and coping strategies reduce physiological arousal and improve resilience.
<b>Pharmacological, Biological, and Integrative Therapies</b>	<ul style="list-style-type: none"> <li>– Monitor effectiveness prescribed medications and for side effects.</li> <li>– Educate client and family members on medication purpose and adherence.</li> <li>– Support safe use of integrative therapies (e.g., mindfulness, aromatherapy).</li> <li>– Monitor for misuse of benzodiazepines.</li> </ul>	Pharmacologic treatment can reduce symptom severity. Client education improves adherence and safety. Integrative therapies can enhance relaxation and overall well-being. Monitoring reduces the risk of adverse effects or dependence.

<b>Milieu Therapy</b>	<ul style="list-style-type: none"> <li>– Maintain a calm and structured unit with minimal overstimulation.</li> <li>– Provide quiet, safe spaces for de-escalation.</li> <li>– Encourage participation in group therapy or activities.</li> <li>– Use environmental cues to reduce uncertainty.</li> </ul>	A structured, low-stimulation environment reduces external stressors and promotes emotional stability. Group participation fosters connection, reduces isolation, and models healthy coping behaviors.
<b>Therapeutic Relationship and Counseling</b>	<ul style="list-style-type: none"> <li>– Establish trust and rapport using active listening and empathy.</li> <li>– Use therapeutic communication techniques.</li> <li>– Encourage expression of thoughts and fears.</li> <li>– Support realistic goal-setting and problem-solving.</li> </ul>	A strong nurse–client relationship and therapeutic communication provides emotional safety, fosters insight, and motivates engagement in treatment.

## Nursing Interventions for Physiological Signs of Anxiety

Nursing interventions also target common physiological signs of anxiety and associated self-care deficits. See common interventions for these conditions in Table 9.7e.

Table 9.7e Nursing Interventions Targeting Physiological Signs of Anxiety and Self-Care Deficit<sup>22</sup>

22. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.



Problem/Intervention	Rationale
<p><b>Nutrition</b></p> <ul style="list-style-type: none"> <li>• Monitor nutritional intake and weight regularly.</li> <li>• Encourage small, frequent meals if appetite is decreased.</li> <li>• Teach about avoiding caffeinated beverages and excessive sugar intake.</li> <li>• Refer the client to a dietician if necessary.</li> </ul>	<p>Anxiety may reduce appetite or lead to emotional eating (e.g., consumption of high-sugar foods can exacerbate physiological symptoms such as restlessness). Balanced nutrition supports energy levels.</p>
<p><b>Sleep</b></p> <ul style="list-style-type: none"> <li>• Assess sleep patterns and quality using sleep diary.</li> <li>• Promote sleep hygiene: establish bedtime routine, limit screen time, avoid stimulants.</li> <li>• Teach relaxation techniques such as deep breathing or guided imagery before bed.</li> <li>• Consult with health care provider for sleep medications or referral for CBT if insomnia persists.</li> </ul>	<p>Anxiety can cause difficulty falling asleep or staying asleep, which worsens anxiety and impairs coping. Non-pharmacological interventions promote better rest without dependence.</p>

<p><b>Elimination</b></p> <ul style="list-style-type: none"> <li>• <b>-Monitor for constipation, diarrhea, or urinary frequency, which may be exacerbated by anxiety.</b></li> <li>• <b>Encourage hydration and high-fiber foods.</b></li> <li>• <b>Promote regular toileting schedule and privacy.</b></li> <li>• <b>Evaluate for side effects of anxiety medications affecting elimination.</b></li> </ul>	<p>Autonomic arousal due to feelings of anxiety can affect urinary patterns. Supporting elimination helps manage increased levels of anxiety. Some medications can affect function.</p>
<p><b>Self-Care Deficits</b></p> <ul style="list-style-type: none"> <li>• <b>Assess for performance of Activities of Daily Living (ADL) including bathing, grooming, and dressing.</b></li> <li>• <b>Provide structure and gentle prompts for hygiene tasks as needed.</b></li> <li>• <b>Encourage participation in morning routines and setting daily goals.</b></li> <li>• <b>Promote positive reinforcement for completing self-care tasks.</b></li> </ul>	<p>Anxiety can cause fatigue, low motivation, or affect hygiene and self-care. Structured support helps with functioning.</p>

## Communication Tips for Clients With Anxiety

Communicating effectively with someone experiencing severe anxiety must foster feelings of trust, safety, and therapeutic alliance. People in states of heightened anxiety may struggle to process information, express themselves clearly, or feel emotionally overwhelmed. Table 9.7f provides communication tips when speaking with clients with severe anxiety.

Table 9.7f Communication Tips for Clients With Severe Anxiety<sup>23</sup>

Tip	Explanation
<b>Use a Calm, Reassuring Tone</b>	Speak slowly and in a steady, gentle voice. A calm tone can reduce the client's arousal level and model a sense of control.
<b>Keep Language Simple and Clear</b>	Avoid complex instructions and medical jargon. Use short sentences and one idea at a time. Anxiety impairs concentration and memory, so information must be simplified.
<b>Validate Feelings Without Dismissing Them</b>	For example, validate feelings by saying things like "I can see this is really overwhelming for you," or "It's okay to feel scared." Avoid saying "Calm down" or "You're overreacting," which can feel invalidating to the client.
<b>Avoid Rapid-Fire Questions</b>	Ask one question at a time and give the person time to respond. Allow silence for processing. Over-questioning can escalate anxiety and create cognitive overload.
<b>Offer Grounding Techniques</b>	Gently guide the person to the present: "Can you tell me 3 things you see in the room?" or "Let's take a slow breath together." This helps redirect their focus away from anxious thoughts.
<b>Be Patient and Nonjudgmental</b>	Avoid correcting or challenging irrational fears if a person is experiencing severe anxiety or panic. Focus on providing support and safety rather than using logic to try to dispel fear. Use therapeutic presence.
<b>Respect Personal Space</b>	Stay at a comfortable distance to avoid making the person feel trapped. Maintain non-threatening body language with a relaxed posture and uncrossed arms.
<b>Provide Structure and Predictability</b>	Explain what you're doing step by step: "I'm going to put this cuff on your arm to take your blood pressure now." Uncertainty increases anxiety.
<b>Encourage Autonomy and Choices</b>	Offer choices when possible (e.g., "Would you prefer to sit here or over by the window?"). Choices restore a sense of control that is often lost during severe anxiety.

23. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

## Client Education

Nurses teach clients how to recognize signs of anxiety and panic attacks and how to use coping strategies to help manage symptoms.

- ▶ Read additional information about anxiety triggers and coping strategies on the “[Anxiety Checklist University](#)” website.

## Evaluation (Evaluate Outcomes)

Nurses refer to the individualized SMART outcomes established for each client when evaluating the effectiveness of interventions in the care plan. In general, evaluation of goals for clients with anxiety disorders includes the following questions<sup>24</sup>:

- Is the client experiencing a reduced level of anxiety?
- Does the client recognize their symptoms are related to anxiety?
- Is the client successfully implementing adaptive coping strategies to manage their anxiety?
- Is the client adequately performing self-care activities (e.g., hygiene, eating, and elimination)?
- Is the client able to maintain satisfying interpersonal relationships?
- Is the client able to successfully function socially, occupationally, or in other important areas of functioning?

24. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

If the client outcomes or goals are not met or only partially met, the nursing care plan should be revised and reimplemented.

## 9.8 Spotlight Application

Melissa is a 20-year-old nursing student who visits her primary care provider for a refill on her birth control pills. During the admission assessment, Melissa shares with the intake nurse that her “anxiety has gotten much worse” during nursing school. She states, “I always feel tired, but I can’t sit down and relax. I feel like there is always more studying that I should be doing for my classes. Even though I’m tired at night, I don’t sleep well and wake up frequently throughout the night. If I can’t go back to sleep, I turn on my computer and study. I have also been getting headaches every day, and I think it is from being on the computer so much. My boyfriend keeps asking me why I am so crabby. Lately I have had a lot of test anxiety, and sometimes my mind goes blank during an exam even though I have memorized all of the material.”

The primary care provider diagnoses Melissa with Generalized Anxiety Disorder and encourages her to attend a local support group with other people experiencing anxiety. A beta-blocker is prescribed on an “as needed” basis for before exams or other types of performance assessments, and a referral is made to a psychotherapist for cognitive behavioral therapy. Melissa is encouraged to follow up with her primary health care provider in two weeks for evaluation of her progress.

1. What priority assessment data should the nurse collect at this time?  
(Assessment/Recognizing Cues)

*The frequency, duration and triggers of her anxiety; including how her anxiety affects her daily life, academic performance and relationship with others. The number of hours Melissa sleeps at night, the quality of her sleep and how often she wakes up during the night. Further assessment of Melissa’s headaches, including frequency, severity, location and possible triggers. Coping mechanisms that Melissa has tried to use to manage her stress. Signs of fatigue or nutritional imbalances. Presence of support system. Signs and symptoms associated with the “flight or fight” stress response. The levels of anxiety Melissa has been experiencing (mild, moderate, severe or panic). Screening tools, such as the Severity Measure for Generalized Anxiety Disorder. Lab work to assess for physiological causes of Melissa’s anxiety.*

*Cultural beliefs that may affect Melissa's perception of anxiety or treatment options.*

2. Based on the assessment data provided, which nursing diagnoses would be appropriate at this time? (Diagnosis/Analyzing Cues)

- *Anxiety r/t academic demands*
- *Disturbed sleep pattern r/t anxiety*
- *Chronic pain r/t excessive screen use*
- *Impaired social interaction r/t irritability*
- *Ineffective coping r/t inability to manage anxiety*

3. Provide a sample of expected outcomes that would be appropriate for Melissa. (Outcome Identification/Generate Solutions)

- *Melissa will report a decrease in anxiety symptoms within two weeks.*
- *Melissa will sleep 6-8 hours per night without frequent awakenings within two weeks.*
- *Melissa will experience a reduction in headache frequency and severity within two weeks.*
- *Melissa will maintain healthy relationships by managing her irritability within two weeks.*
- *Melissa will list three different stress management techniques prior to leaving the clinic.*

4. What nursing interventions would be appropriate for this client? (Planning & Implementation/Generate Solutions & Take Action)

- *Use therapeutic communication techniques such as actively listening, providing a calm environment and validating Melissa's feelings.*
- *Teach Melissa how to create a realistic study schedule that balances study time with self-care activities.*
- *Provide Melissa with resources on stress management techniques such as exercise, guided imagery, journaling or practicing mindfulness.*
- *Encourage Melissa to limit screen time before bed and avoid studying*

*in bed to help improve quality of sleep.*

- *Provide education on how a well-balanced diet can improve anxiety and overall wellbeing.*
- *Reinforce the importance of support groups and following through with psychotherapy referral.*
- *Teach Melissa to recognize the early signs of anxiety so it can be managed effectively.*
- *Educate Melissa on her prescribed beta-blocker, including proper timing and potential side effects.*
- *Encourage Melissa to avoid caffeine as this can exacerbate anxiety as well as interfere with sleep.*

5. How would you evaluate if Melissa's outcomes were met? (Evaluation/Evaluate Outcomes)

*Every time the nurse interacts with Melissa, outcomes should be evaluated. Continued assessment of Melissa's anxiety levels, sleep patterns, headaches, relationship status and ability to apply stress management techniques will help the nurse determine if outcomes are met, partially met, or not met. If outcomes are partially met or not met, the nurse may need to continue monitoring Melissa's progress or the care plan may need revision.*



## 9.9 Learning Activities



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=492#h5p-24>

1



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=492#h5p-25>

2



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=492#h5p-27>

1. “MH Anxiety Glossary Cards” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
2. “MH Anxiety Drag and Drop” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

- ▶ Test your clinical judgment with a NCLEX Next Generation-style question: [Chapter 9, Assignment 1](#)<sup>4</sup>



- ▶ Test your clinical judgment with a NCLEX Next Generation-style case study: [Chapter 9, Case Study 1](#)<sup>5</sup>



3. "MH Anxiety Question Set 1" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

4. "MH Anxiety Next Gen Question by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

5. "MH Anxiety Next Gen Case Study by Kellea Ewen is licensed under [CC BY-NC 4.0](#)

## IX Glossary

**Agoraphobia:** Intense fear of two or more of the following situations: using public transportation, being in open spaces (e.g., parking lots, marketplaces, or bridges), being in enclosed spaces (e.g., shops or theaters), standing in line or being in a crowd, or being outside of the home alone.

**Anxiety:** A universal human experience that includes feelings of apprehension, uneasiness, uncertainty, or dread resulting from a real or perceived threat.

**Compulsions:** Repetitive behaviors that a person with OCD feels the urge to do in response to an obsessive thought.

**Coping strategies:** An action, a series of actions, or a thought process used to address a stressful or unpleasant situation or modify one's reaction to such a situation.

**Defense mechanisms:** Reaction patterns used by individuals to protect themselves from anxiety that arises from stress and conflict.

**Dialectal behavioral therapy (DBT):** A structured, evidence-based psychotherapy that combines cognitive-behavioral techniques for emotion regulation and reality-testing with concepts of mindfulness, acceptance, and dialectics (the idea that two seemingly opposite things can both be true).

**Exposure and response prevention (EX/RP):** A type of psychotherapy effective in reducing compulsive behaviors in clients with OCD. EX/RP includes spending time in the very situation that triggers compulsions (for example, touching dirty objects) but then being prevented from undertaking the usual resulting compulsion (handwashing).

**Exposure therapy:** A type of psychotherapy that focuses on confronting the fears underlying an anxiety disorder to help people engage in activities they have been avoiding.

**Generalized anxiety disorder (GAD):** Excessive anxiety and worry occurring

for at least six months about a number of events or activities (such as work or school performance).

**Obsessions:** Repeated thoughts, urges, or mental images that cause anxiety.

**Obsessive-compulsive disorder (OCD):** A common chronic disorder in which a person has uncontrollable, reoccurring thoughts (obsessions) and/or behaviors (compulsions) they feel the urge to repeat over and over. Performing the compulsive behaviors often brings the person brief relief from the anxiety the obsessive thoughts cause them.

**Panic:** The most extreme level of anxiety that results in significantly dysregulated behavior. The individual is unable to process information from the environment and may lose touch with reality.

**Panic attacks:** Sudden periods of intense fear that come on quickly and reach their peak within minutes. Attacks can occur unexpectedly or can be brought on by a trigger, such as a feared object or situation.

**Phobia:** An intense fear or aversion to specific objects or situations (e.g., flying, heights, animals, receiving an injection, or seeing blood).

**Post-traumatic stress disorder (PTSD):** A disorder that develops in some people who have experienced a shocking, frightening, or dangerous event where they feel stressed or frightened even when they are not in danger.

**Repetitive Transcranial Magnetic Stimulation (rTMS):** Treatment that uses a magnet to activate specific sites in the brain.

**Selective mutism:** A condition when people fail to speak in specific social situations despite having normal language skills.

**Separation anxiety disorder:** A condition where an individual has a fear about being separated from people to whom they are attached.

**Social anxiety disorder:** Significant fear or anxiety about one or more social situations in which the individual is exposed to possible scrutiny by others.





### Learning Objectives

- Identify assessment cues of personality disorder behaviors
- Identify nursing priorities for clients with personality disorders
- Plan outcomes for clients with personality disorders
- Differentiate safety/protective interventions for clients with personality disorders
- Apply evidence-based practice when planning care and interventions for clients with personality disorders
- Analyze treatments for clients with personality disorders
- Apply the nursing process to clients with personality disorders at risk for suicide

Research studies have demonstrated that nine percent of Americans have a personality disorder. A large proportion of this population is also diagnosed with one or more other mental illnesses.<sup>1</sup> This chapter will describe the signs and symptoms of ten personality disorders and associated treatments.

1. National Institute of Mental Health. (n.d.). *Personality disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/statistics/personality-disorders>

## 10.2 Basic Concepts

A person's **personality** is a relatively stable pattern of thinking, feeling, and behaving that evolves over their lifetime. It is unique to each individual and influenced by their experiences, environment (surroundings and life situations), and inherited characteristics. **Personality traits** are characteristics, whether considered positive or negative, that make up one's personality. Healthy personality traits include characteristics such as the following:

- Demonstrating healthy personal boundaries
- Accepting responsibility for personal actions
- Communicating in a healthy and effective manner
- Obeying laws
- Showing mutual respect in relationships
- Being independent
- Displaying confidence
- Behaving in a non-impulsive manner

Review healthy personal boundaries in the “[Foundational Mental Health Concepts](#)” chapter. An individual's personality is considered unhealthy and classified as a disorder when it impacts their interpersonal relationships and results in impaired functioning in social, occupational, or other important areas of their life.<sup>1</sup>

The *Diagnostic and Statistical Manual of Mental Disorders (DSM-5-TR)* defines a **personality disorder** as an enduring pattern of inner experience and behavior that deviates significantly from the expectations of the individual's culture. Its onset can be traced back to adolescence or early

1. American Psychiatric Association. (2013). *Desk reference to the diagnostic criteria from DSM-5*.



adulthood and is present in a variety of contexts. This abnormal pattern of behavior is manifested in two or more of the following areas<sup>2</sup>:

- Cognition (i.e., ways of perceiving and interpreting self, other people, and events)
- Affect (i.e., the range, intensity, lability, and appropriateness of emotional response)
- Interpersonal functioning
- Impulse control

There are ten different personality disorders that are categorized into three clusters (A, B, and C) in the *DSM-5-TR*. Personality disorders within each cluster have similar patterns of behavior. The ten disorders include Cluster A (paranoid, schizoid, schizotypal), Cluster B (antisocial, narcissistic, borderline, histrionic), and Cluster C (dependent, avoidant, and obsessive-compulsive personality disorder).<sup>3</sup> These disorders are often **ego-syntonic**, meaning that the behaviors and thoughts are consistent with the individual's self-perception and are not seen as problematic by the individual, despite causing significant issues in their life.

Each personality disorder is further described in the following sections. As you read through each section, keep in mind that these disorders are more than just personality traits; they are diagnosed based on patterns of behaviors that significantly impair a person's functioning.

## Cluster A Personality Disorders

**Cluster A personality disorders** include paranoid personality disorder,

2. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

3. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

schizoid personality disorder, and schizotypal personality disorder. Cluster A is characterized as the odd, eccentric cluster. Individuals with these types of disorders often experience social awkwardness.

## Paranoid Personality Disorder

The *DSM-5-TR* defines **paranoid personality disorder** as a pervasive distrust and suspiciousness of others such that their motives are interpreted as malevolent, beginning in early childhood and present in a variety of contexts. It is diagnosed in individuals with four or more of the following characteristics<sup>4</sup>:

- Suspects without evidence that others are exploiting, harming, or deceiving them
- Preoccupied with unjustified doubts about the loyalty or trustworthiness of friends or associates
- Reluctant to confide in others because of unwarranted fear that the information will be used maliciously against them
- Reads hidden meaning or threatening meanings into benign remarks or events
- Persistently bears grudges (i.e., is unforgiving of insults, injuries, or slights)
- Perceives attacks on their character or reputation that are not apparent to others and is quick to react angrily or to counterattack
- Has recurrent suspicions without justification regarding fidelity of spouse or sexual partner

See Figure 10.1<sup>5</sup> for a word cloud image representing paranoid personality disorder.

4. American Psychiatric Association. (2013). *Desk reference to the diagnostic criteria from DSM-5*.

5. “[Word\\_Cloud\\_PPD.png](#)” by [MissLunaRose12](#) is licensed under [CC BY-SA 4.0](#)




Figure 10.1 Paranoid Personality Disorder

## Schizoid Personality Disorder

The *DSM-5-TR* defines **schizoid personality disorder** as a pervasive pattern of detachment from social relationships and a restricted range of expression of emotions in interpersonal settings, beginning in early childhood and present in a variety of contexts. It is diagnosed in individuals with four or more of the following characteristics<sup>6</sup>:

- Neither desires nor enjoys close relationships including being part of a family
- Almost always chooses solitary activities
- Has little, if any, interest in having sexual experiences with another person
- Takes pleasure in few, if any, activities
- Lacks close friends other than first-degree relatives

6. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

- Appears indifferent to the praise or criticism of others
- Shows emotional coldness, detachment, or flat affect

See Figure 10.2<sup>7</sup> for a word cloud image representing schizoid personality disorder.




Figure 10.2 Schizoid Personality Disorder

## Schizotypal Personality Disorder

The *DSM-5* defines **schizotypal personality disorder** as a pervasive pattern of social and interpersonal deficits marked by acute discomfort with and reduced capacity for close relationships, as well as by cognitive or perceptual distortions and eccentricities of behavior, beginning in early adulthood and

7. “[Word\\_Cloud\\_SzPD.png](#)” by [MissLunaRose12](#) is licensed under [CC BY-SA 4.0](#)

present in a variety of contexts. It is diagnosed in individuals with five or more of the following characteristics<sup>8</sup>:

- **Ideas of reference** (i.e., the false belief that coincidental events relate to oneself). For example, a person shopping in a store sees two strangers laughing and believes that they are laughing at them, when, in reality, the other two people do not even notice them.
- Odd beliefs or magical thinking that influence behavior and are inconsistent with cultural norms (**Magical thinking** refers to the idea that one can influence the outcome of specific events by doing something that has no bearing on the circumstances. For example, a person watching a baseball game exhibits magical thinking when believing that holding the remote control in a certain position caused their favorite player to hit a home run.)
- Unusual perceptual experiences including bodily illusions (A **body illusion** refers to a perception that one's body is significantly different from its actual configuration. For example, a person lying in bed feels as if they are levitating.)
- Odd thinking and speech
- Suspiciousness or paranoid ideation
- Inappropriate or constricted affect
- Behavior or appearance that is odd, eccentric, or peculiar
- Lack of close friends or confidants other than first-degree relatives
- Excessive social anxiety that does not diminish with familiarity and tends to be associated with paranoid fears rather than negative judgments about self

Please Note: There are key differences between the schizotypal and schizoid. Schizoid is marked by social detachment and emotional coldness without

8. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

cognitive or perceptual distortions, whereas schizotypal includes these social deficits along with cognitive or perceptual distortions and eccentric behavior.

See Figure 10.3<sup>9</sup> for a word cloud image representing schizotypal personality disorder.



Figure 10.3 Schizotypal Personality Disorder

## Cluster B Personality Disorders

**Cluster B personality disorders** include antisocial, borderline, histrionic, and narcissistic personality disorders. Cluster B personality disorders are characterized by dramatic, overly emotional, or unpredictable thinking or behavior.

### Antisocial Personality Disorder

The *DSM-5-TR* defines **antisocial personality disorder** as a pervasive pattern

9. “[Word\\_Cloud\\_StPD.png](#)” by [MissLunaRose12](#) is licensed under [CC BY-SA 4.0](#)

of disregard for and violation of the rights of others since age 15. It is diagnosed in individuals with three or more of the following characteristics<sup>10]</sup>:

- Failure to conform to social norms with respect to lawful behaviors, as indicated by repeatedly performing acts that are grounds for arrest
- Deceitfulness, as indicated by repeated lying, use of aliases, or conning others for personal profit or pleasure
- Impulsivity or failure to plan ahead
- Irritability and aggressiveness, as indicated by repeated physical fights and assaults
- Reckless disregard for the safety of self or others
- Consistent irresponsibility, as indicated by repeated failure to sustain consistent work behavior or honor financial obligations
- Lack of remorse, as indicated by being indifferent to or rationalizing having hurt, mistreated, or stolen from another person

See Figure 10.4<sup>11]</sup> for a word cloud image representing antisocial personality disorder.

10. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

11. “[Word\\_Cloud\\_AsPD.png](#)” by [MissLunaRose12](#) is licensed under [CC BY-SA 4.0](#)



Figure 10.4 Antisocial Personality Disorder

## Borderline Personality Disorder

The *DSM-5-TR* defines **borderline personality disorder** as a pervasive pattern of instability of personal relationships, self-image, and affect with significant impulsivity, beginning in early adulthood and present in a variety of contexts. It is diagnosed in individuals with five or more of the following characteristics<sup>12</sup>:

- Frantic efforts to avoid real or imagined abandonment
- A pattern of unstable and intense personal relationships characterized by alternating between extremes of idealization and devaluation (referred to as **splitting**)
- Identity disturbance with significantly and persistently unstable self-image or sense of self
- Impulsivity in at least two areas that are potentially self-damaging (e.g.,

12. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.



- spending, sex, substance misuse, reckless driving, or binge eating)
- Recurrent suicidal behavior or self-mutilating behavior
- Unstable affect with significant mood reactivity (i.e., intense anxiety or irritability usually lasting only a few hours)
- Chronic feelings of emptiness
- Inappropriate, intense anger or difficulty controlling anger
- Transient, stress-related paranoid ideation or severe dissociative symptoms (**Dissociative symptoms** include the experience of detachment or feeling as if one is outside one's body. It is often associated with loss of memory of the experience. Dissociative disorders are associated with an individual's previous experience of trauma.)

See Figure 10.5<sup>13</sup> for a word cloud image representing borderline personality disorder.



Figure 10.5 Borderline Personality Disorder

13. "Word\_Cloud\_BPD.png" by [MissLunRose12](#) is licensed under [CC BY-SA 4.0](#)

# Histrionic Personality Disorder

The *DSM-5-TR* defines **histrionic personality disorder** as a pervasive pattern of excessive emotionality and attention seeking, beginning in early adulthood and present in a variety of contexts. It is diagnosed in individuals with five or more of the following characteristics<sup>14</sup>:

- Uncomfortable in situations in which they are not the center of attention
- Interaction with others is characterized by inappropriate sexually seductive or provocative behavior
- Rapidly shifting and shallow expression of emotion
- Consistently uses physical appearance to draw attention to oneself
- Excessively impressionistic speech that is lacking in detail
- Shows self-dramatization, theatricality, and exaggerated expression of emotion
- Suggestible (i.e., easily influenced by others)
- Considers relationships to be more intimate than they actually are

See Figure 10.6<sup>15</sup> for a word cloud image representing histrionic personality disorder.

14. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

15. “[Word\\_Cloud\\_HPD.png](#)” by [MissLunRose12](#) is licensed under [CC BY-SA 4.0](#)




Figure 10.6 Histrionic Personality Disorder

## Narcissistic Personality Disorder

The *DSM-5-TR* defines **narcissistic personality disorder** as a pervasive pattern of grandiosity (in fantasy or behavior), need for attention, and lack of empathy, beginning in early adulthood and present in a variety of contexts. It is diagnosed in individuals with five or more of the following characteristics<sup>16</sup>:

- Has a grandiose sense of self-importance (i.e., exaggerates achievements and talents)
- Is preoccupied with fantasies of unlimited success, power, brilliance, beauty, or ideal love
- Believes they are “special” and can only be understood by, or should only associate with, other “special” or high-status people (or institutions)
- Requires excessive admiration

16. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

- Has a sense of **entitlement** (i.e., unreasonable expectations of especially favorable treatment)
- Exploits others to achieve their own goals
- Lacks empathy and the ability to identify with the feelings and needs of others
- Is often envious of others or believes that others are envious of them
- Shows arrogant or haughty behaviors or attitudes

See Figure 10.7<sup>17</sup> for a word cloud image representing narcissistic personality disorder.



Figure 10.7 Narcissistic Personality Disorder

## Cluster C Personality Disorders

**Cluster C personality disorders** include avoidant, dependent, and obsessive-

17. “[Word\\_Cloud\\_NPD.png](#)” by [MissLunRose12](#) is licensed under [CC BY-SA 4.0](#)

compulsive personality disorders. Cluster C personality disorders are characterized by anxious, fearful thinking or behavior.

## Avoidant Personality Disorder

The *DSM-5-TR* defines **avoidant personality disorder** as a pervasive pattern of social inhibition, feelings of inadequacy, and hypersensitivity to negative evaluation, beginning in early adulthood and present in a variety of contexts. It is diagnosed in individuals with four or more of the following characteristics<sup>18</sup>:

- Avoids occupational activities that involve significant interpersonal contact because of fears or criticism, disapproval, or rejection
- Is unwilling to get involved with people unless certain of being liked
- Shows restraint within intimate relationships because of the fear of being shamed or ridiculed
- Is preoccupied with being criticized or rejected in social situations
- Is inhibited in new interpersonal situations because of feelings of inadequacy
- Views self as socially inept, personally unappealing, or inferior to others
- Is unusually reluctant to take personal risks or to engage in any new activities because they may prove embarrassing

See Figure 10.8<sup>19</sup> for a word cloud image representing avoidant personality disorders.

18. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

19. “[Word\\_Cloud\\_AvPD.png](#)” by [MissLunRose12](#) is licensed under [CC BY-SA 4.0](#)




Figure 10.8 Avoidant Personality Disorder


## Dependent Personality Disorder

The *DSM-5-TR* defines **dependent personality disorder** as a pervasive and excessive need to be taken care of that leads to submission and clinging behavior and fears of separation, beginning in early childhood and present in a variety of contexts. It is diagnosed by five or more of the following characteristics<sup>20</sup>:

- Has difficulty making everyday decisions without an excessive amount of advice and reassurance from others
- Needs others to assume responsibility for most major areas of their life
- Has difficulty expressing disagreement with others because of fear of loss or support or approval
- Has difficulty initiating projects or doing things on their own because of lack of self-confidence in judgment or abilities

20. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

- See Figure 10.9<sup>21</sup> for a word cloud image representing dependent personality disorder.



The *DSM-5-TR* defines **obsessive-compulsive personality disorder** as a

10.2 Basic Concepts | 707

pervasive pattern or preoccupation with orderliness, perfectionism, and mental and interpersonal control at the expense of flexibility, openness, and efficiency, beginning in early childhood and present in a variety of contexts. It is diagnosed in individuals with four or more of the following characteristics<sup>22</sup>:

- Is preoccupied with details, rules, lists, order, organization, or schedules to the extent that the major point of the activity is lost
- Shows perfectionism that interferes with task completion (i.e., is unable to complete a project because their overly strict standards are not met)
- Is excessively devoted to work and productivity to the exclusion of leisurely activities and friendships
- Is overconscientious, scrupulous, and inflexible about matters of morality, ethics, or values (not accounted for by cultural or religious identification)
- Is unable to discard worn-out or worthless objects even when they have no sentimental value
- Is reluctant to delegate tasks or work with others unless they submit to exactly their way of doing things
- Adopts a miserly spending style towards self and others; money is viewed as something to be hoarded for future catastrophes
- Shows rigidity and stubbornness

It is important to note that obsessive-compulsive personality disorder (OCPD) is a different disorder than obsessive-compulsive disorder (OCD). OCPD includes long-term personality traits characterized by extreme perfectionism, rigidity, and adherence to rules. A person with OCPD is often proud of these personality traits. Conversely, OCD includes uncontrollable, recurring thoughts (obsessions) and/or behaviors (compulsions) that cause the individual significant emotional distress.

22. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.



- ▶ Review the “[Obsessive-Compulsive Disorder](#)” section of the “Anxiety Disorders” chapter for more details about obsessive-compulsive disorder.

See Figure 10.10<sup>23</sup> for a word cloud image representing obsessive-compulsive personality disorder.




Figure 10.10 Obsessive-Compulsive Personality Disorder

23. “[Word\\_Cloud\\_OCPD.png](#)” by [MissLunRose12](#) is licensed under [CC BY-SA 4.0](#)



View the following Khan Academy YouTube video explaining personality disorders<sup>24</sup> : [Personality Disorders](#).

### Reflective Question:

1. When does one's personality become a disorder versus just personality traits?

## Risk Factors for Personality Disorders

Research suggests that genetics, childhood trauma, peer influences, and other factors can contribute to the development of personality disorders.<sup>25,26</sup>

- **Adverse Childhood Experiences (ACEs):** Multiple studies have shown that childhood adversities such as emotional, physical, and sexual abuse,

24. khanacademymedicine. (2015, July 20). *Personality disorders | Behavior | MCAT | Khan Academy* [Video]. YouTube. All rights reserved.  
<https://youtu.be/-Xv593jgyJ4>

25. American Psychological Association (2010). *What causes personality disorders?* <https://www.apa.org/topics/personality-disorders/causes>

26. Solmi, M., Dragioti, E., Croatto, G., Radua, J., Borgwardt, S., Carvalho, A. F., Demurtas, J., Mosina, A., Kurotschka, P., Thompson, T., Cortese, S., Shin, J. I., & Fusar-Poli, P. (2021). Risk and protective factors for personality disorders: An umbrella review of published meta-analyses of case-control and cohort studies. *Front Psychiatry*, 12, 679379. [doi:10.3389/fpsy.2021.679379](https://doi.org/10.3389/fpsy.2021.679379).

as well as neglect, significantly increase the risk of developing personality disorders. For instance, childhood emotional abuse and neglect have been strongly associated with borderline personality disorder (BPD).<sup>27</sup>

- **Perinatal Complications:** Factors such as prematurity, asphyxia, and complicated deliveries have been linked to an increased risk of personality disorders. These perinatal complications may lead to abnormal brain development, which can predispose individuals to personality pathology.<sup>28</sup>
- **Prenatal Stress:** Exposure to maternal stress during pregnancy has been associated with a higher likelihood of developing personality disorders in offspring. This association remains significant even after adjusting for other factors such as parental psychiatric history and prenatal smoking.<sup>29</sup>
- **Childhood Behavioral Problems:** Early conduct problems, depressive symptoms, anxiety, and immaturity have been identified as predictors of personality disorders in adolescence and adulthood. Conduct problems, in particular, have been shown to be an independent predictor across all clusters of personality disorders.<sup>30</sup>

27. Broekhof, R., Nordahl, H. M., Eikenæs, I. U., & Selvik, S. G. (2024). Adverse childhood experiences are associated with personality disorder: A prospective, longitudinal study. *Journal of Personality Disorders*, 38(1), 19-33. [doi: 10.1521/pedi.2024.38.1.19](https://doi.org/10.1521/pedi.2024.38.1.19).
28. Fazel, S., Bakiyeva, L., Cnattingius, S., Grann, M., Hultman, C. M., Lichtenstein, P., & Geddes, J. R. (2012). Perinatal risk factors in offenders with severe personality disorder: A population-based investigation. *Journal of Personality Disorders*, 26(5), 737-50. [doi: 10.1521/pedi.2012.26.5.737](https://doi.org/10.1521/pedi.2012.26.5.737).
29. Brannigan, R., Tanskanen, A., Huttunen, M. O., Cannon, M., Leacy, F. P., & Clarke, M. C. (2020). The role of prenatal stress as a pathway to personality disorder: Longitudinal birth cohort study. *The British Journal of Psychiatry*, 216(2), 85-89. [doi: 10.1192/bjp.2019.190](https://doi.org/10.1192/bjp.2019.190).
30. Bernstein, D. P., Cohen, P., Skodol, A., Bezirgianian, S., & Brook, J. S. (1996).

- **Genetic and Temperamental Factors:** Certain temperamental traits, such as high neuroticism and low agreeableness, have been implicated in the development of personality disorders. These traits can interact with environmental factors to increase the risk.<sup>31,32</sup>
- **Invalidating Relational Experiences:** Maladaptive parenting, problematic peer relationships, and other invalidating experiences during childhood and adolescence can contribute to the development of personality disorders, particularly.<sup>33</sup>

Certain positive factors can help prevent children from developing personality disorders. For example, a single strong relationship with a relative, teacher, or friend can offset negative influences. Strong resiliency factors contribute to the health and development of a child.<sup>34</sup>

Childhood antecedents of adolescent personality disorders. *American Journal of Psychiatry*, 153(7), 907-13. doi: 10.1176/ajp.153.7.907.

31. American Psychological Association (2010). *What causes personality disorders?* <https://www.apa.org/topics/personality-disorders/causes>
32. Junewicz, A., & Billick, S. B. (2021). Preempting the development of antisocial behavior and psychopathic traits, *The Journal of American Academy of Psychiatry Law*, 49(1), 66-76. <https://doi10.29158/JAAPL.200060-20>
33. Solmi, M., Dragioti, E., Croatto ,G., Radua, J., Borgwardt, S., Carvalho, A.F., Demurtas, J., Mosina, A., Kurotschka, P., Thompson, T., Cortese, S., Shin, J. I., & Fusar-Poli, P. (2021). Risk and protective factors for personality disorders: An umbrella review of published meta-analyses of case-control and cohort studies. *Front Psychiatry*, 12, 679379. doi: 10.3389/fpsyt.2021.679379.
34. American Psychological Association (2010). *What causes personality disorders?* <https://www.apa.org/topics/personality-disorders/causes>

## 10.3 Treatment for Personality Disorders

Clients with some types of personality disorders do not exhibit insight for their condition and are unlikely to seek treatment. They often go undiagnosed unless they seek treatment for another psychiatric or medical diagnosis. Some may also be forced to seek treatment at the influence of family members or as required by law if legal infractions have occurred. Personality disorders can be challenging for mental health professionals to treat. Individuals with personality disorders struggle to recognize that their difficulties in life are related to their personalities. They may truly believe their problems are a result of other people or outside factors which makes treatment difficult. It is very common for clients with personality disorders to also have substance abuse, anxiety, depression, or eating disorders.<sup>1</sup>

Treatment for personality disorders is tailored to the specific disorder and individual client needs, and it is a gradual process. Psychotherapy is the first line of treatment for personality disorders. Medications may also be prescribed to treat underlying co-occurring conditions such as anxiety or depression.

### Psychotherapy

During psychotherapy, an individual can gain insight and knowledge about the personality disorder and what is contributing to their symptoms. They talk about their thoughts, feelings, and behaviors and ideally will develop an understanding of the impact of their thoughts and behaviors on themselves and others. They learn strategies for managing and coping with their symptoms. This treatment can help reduce problematic behaviors that impact an individual's relationships and functioning.<sup>2</sup>

1. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.
2. American Psychiatric Association. (n.d.). *What are personality disorders?*

Common types of psychotherapy used to treat personality disorders are as follows<sup>3</sup>:

- Cognitive behavior therapy
- Dialectical behavior therapy
- Interpersonal therapy
- Psychoanalytic/psychodynamic therapy
- Psychoeducation

Cognitive behavior therapy (CBT) teaches the client to become more aware of the way they think so they can ultimately change the way they behave.

**Dialectical behavior therapy (DBT)** is a type of cognitive behavioral therapy that was originally created for clients with borderline personality disorder to help them cope with stress, control emotions, and establish healthy relationships. It is considered the gold standard for treating borderline personality disorder and is also used for other types of disorders. The client learns how to be aware of how thoughts, feelings, behaviors link together. They learn how to use their senses to be aware of what is happening around them and how to use strategies (such as mindfulness, distress tolerance, interpersonal effectiveness, and emotion regulation) to react calmly in a crisis, avoid negative impulsive behavior, and improve relationships.<sup>4</sup> See Figure

<https://www.psychiatry.org/patients-families/personality-disorders/what-are-personality-disorders>

3. American Psychiatric Association. (n.d.). *What are personality disorders?* <https://www.psychiatry.org/patients-families/personality-disorders/what-are-personality-disorders>
4. Schimelpfening, N. (2021). What is dialectical behavior therapy (DBT)? *Verywell Mind*. <https://www.verywellmind.com/dialectical-behavior-therapy-1067402>

10.11<sup>5</sup> for an illustration of how thoughts, feelings and behaviors are linked together.




Figure 10.11 Dialectical Behavior Therapy

**Interpersonal therapy** focuses on developing healthy relationships with others. Negative aspects of relationships, such as social isolation and aggression, are identified, and strategies to improve relationships are discussed and planned.

**Psychodynamic therapy** promotes self-reflection and self-examination of problematic relationship patterns and unresolved conflicts. It can bring awareness of how one's past has an influence on one's present behavior.

**Mentalization-Based Therapy:** A type of psychotherapy designed to help individuals improve their ability to understand and interpret their own and others' mental states—such as thoughts, feelings, and intentions. The goals

5. "Dialectical Behavior Therapy by Kim Ernstmeier.png" by Kim Ernstmeier for [Chippewa Valley Technical College](#) is licensed under [CC BY 4.0](#)

are to improve self-awareness and understanding of one's own emotions, enhance empathy and understanding of others' perspectives. This subsequently helps individuals develop better interpersonal skills and communication.

**Psychoeducation** teaches the client about their medications, psychotherapies, and support groups available in the community. Wraparound services may be planned with an individualized plan of care that brings multiple services/supports together for the client.

## Pharmacotherapy

There are no specific medications approved to treat personality disorders. However, clients may be prescribed medications by their health care provider as adjunctive treatment to psychotherapy. Medications that can be fatal in overdose should be avoided or used with caution. Benzodiazepines can be especially toxic when combined with alcohol or opioids, and they can also cause behavioral disinhibition in clients with personality disorders.<sup>6</sup>

The following medications may be prescribed for symptoms associated with personality disorders: commonly used to treat them include the following<sup>7</sup>:

- Cognitive-perceptual symptoms (e.g., hallucinations and paranoid ideation): Treated with low-dose antipsychotics such as aripiprazole, risperidone, and quetiapine.
- Impulsive behaviors (e.g., self-injury, theft, interpersonal conflict): Impulsivity and behavioral dysregulation may be treated with mood stabilizers (such as valproate and lamotrigine).
- Affective dysregulation (e.g., depressed mood, mood lability, anxiety,

6. Nelson, K. J. (2024). Personality disorders: Overview of pharmacotherapy. *UpToDate*. <https://www.uptodate.com>

7. Nelson, K. J. (2024). Personality disorders: Overview of pharmacotherapy. *UpToDate*. <https://www.uptodate.com>



anger): Treated with mood stabilizers and low dose antipsychotics, which are more effective than antidepressants in this population.

## Support Groups

Support groups and supportive relationships can be helpful for sharing experiences with others who are going through similar situations. Talking with trusted friends or family members can also provide emotional support and encouragement.

## 10.4 Applying the Nursing Process to Personality Disorders

The nursing process is crucial in the care of patients with personality disorders due to its structured and systematic approach that ensures comprehensive and individualized care.

### Assessment (Recognize Cues)

As previously discussed in this chapter, there are ten different personality disorders that are categorized into three clusters (A, B, and C) in the DSM-5. Personality disorders within each cluster have similar patterns of behavior. Cluster A personality disorders include paranoid personality disorder, schizoid personality disorder, and schizotypal personality disorder. Cluster A is characterized as the odd, eccentric cluster. Individuals with these types of disorders often experience social awkwardness. Cluster B personality disorders include antisocial, borderline, histrionic, and narcissistic personality disorders. Cluster B personality disorders are characterized by dramatic, overly emotional, or unpredictable thinking or behavior. Cluster C personality disorders include avoidant, dependent, and obsessive-compulsive personality disorders. Cluster C personality disorders are characterized by anxious, fearful thinking or behavior. Furthermore, personality disorders are often ego-syntonic, meaning that the behaviors and thoughts are consistent with the individual's self-perception and are not seen as problematic by the individual.

Assessing a client with a personality disorder focuses on both verbal and nonverbal assessments. As the nurse conducts follow-up assessments, findings are compared to baseline admission assessments. Assessment includes several components, such as a mental status examination, psychosocial assessment, cultural assessment, spiritual assessment, screening with validated tools, and review of laboratory testing results while also considering lifespan considerations. Review information about performing a mental status examination and psychosocial assessment in the [“Application of the Nursing Process in Mental Health Care”](#) chapter.

## Mental Status Examination

See Table 10.4a for common themes when assessing a client with a personality disorder. Review information about a mental status examination in the “[Assessment](#)” section in Chapter 4.

Table 10.4a Common Findings During A Mental Status Examination Of A Client With A Personality Disorder

Mental Status Examination Component	Common Themes in Personality Disorders
<b>Signs of Distress</b>	<p>Clients with personality disorders may have suicidal ideation, perform self-harm like cutting, or have violence or homicidal ideation.</p> <p><b>*Increased risk for self-injury, suicide, or injury to others must be promptly reported to provider. Do not leave clients alone if statements such as these are being made.</b></p>
<b>Level of Consciousness and Orientation</b>	<p>Clients with personality disorders are typically alert and oriented to person, place, time, and situation. Cognitive functioning is typically intact unless complicated by comorbid conditions or dissociative episodes</p>
<b>Appearance and General Behavior</b>	<p>Behaviors varies by type of personality disorder. Clients may display anger, anxiety, hypersensitivity, emotional lability, or detachment. Behavior may appear dramatic, guarded, rigid, seductive, or odd depending on the subtype. Poor eye contact or unusual mannerisms may be seen in cluster A disorders. In cluster B, the client may present as flamboyant, provocative, or emotionally labile. For example, individuals with borderline personality disorder (BPD) may show intense distress over perceived rejection. In cluster C, the client may present with anxious behaviors.</p>
<b>Speech</b>	<p>Clients with narcissistic or histrionic traits may dominate conversations or speak in an exaggerated manner. Those with paranoid traits may speak with suspicion or defensiveness.</p>
<b>Motor Activity</b>	<p>Clients with BPD or antisocial traits may appear restless or agitated. Minimal movement may be seen in avoidant or schizoid types due to anxiety or withdrawal.</p>
<b>Mood and Affect</b>	<p>Mood may appear anxious, irritable, dysphoric, or constricted depending on the disorder. Mood is often labile or reactive in cluster B disorders (especially in BPD). Clients with BPD often experience feelings of emptiness, intense feelings of abandonment, extreme mood shifts that occur in a matter of hours or days, intense and unstable relationships impulsive behavior such as reckless driving, unsafe sex, substance use, gambling, overspending, or binge eating.<sup>1</sup></p>
<b>Thought and Perception</b>	<p>Thought processes in clients with personality disorders may be rigid or inflexible. Paranoid ideation or transient dissociation may be present under stress. Clients with schizotypal traits may express magical thinking or odd beliefs.</p>
<b>Attitude and Insight</b>	<p>Attitude may be guarded, defensive, manipulative, seductive, or overly compliant based on the disorder. Insight is often limited; clients may externalize blame and minimize their role in conflicts.</p>

<b>Cognitive Abilities and Level of Judgment</b>	Attention, memory, and abstract thinking are typically intact. However, some may show poor judgment or difficulty with impulse control (such as in BPD or antisocial personality disorders). Cognitive distortions are common across all types.
--	---

# Psychosocial Assessment

As previously discussed in the “[Application of the Nursing Process in Mental Health Care](#)” chapter, a psychosocial assessment obtains additional subjective data that detects risks and identifies treatment opportunities and resources.<sup>23</sup>

- Reason for seeking health care (i.e., “chief complaint”)
- Thoughts of self-harm or suicide (both current and historical)
- Cultural assessment
- Spiritual assessment
- Family dynamics
- Current and past medical history
- Current medications
- History of previously diagnosed mental health disorders
- Previous hospitalizations
- Educational background
- Occupational background
- History of exposure to psychological trauma, violence, and domestic

1. Mayo Clinic. (2024). *Borderline personality disorder*. <https://www.mayoclinic.org/diseases-conditions/borderline-personality-disorder/symptoms-causes/syc-20370237>

2. Clinical Practice Guideline for the Treatment of Posttraumatic Stress Disorder. (2017). *What is cognitive behavioral treatment?* American Psychological Association. <https://www.apa.org/ptsd-guideline/patients-and-families/cognitive-behavioral>

3. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

abuse

- Substance use (tobacco, alcohol, recreational drugs, misused prescription drugs)
- Family history of mental illness
- Coping mechanisms
- Functional ability/Activities of daily living

Additional focused questions are used to obtain detailed information used to plan care. The mnemonic PQRSTU can be used to ask questions in an organized fashion. See Table 10.4b for a sample PQRST assessment for clients with a borderline personality disorder and sample responses.

Table 10.4b Sample PQRSTU Questions for Assessing Clients With a Borderline Personality Disorder

PQRSTU	Sample Questions	Sample Client Response
<b>Provocation/ Palliation</b>	“What seems to trigger or worsen your emotional pain or distress? What helps you feel better?”	“It usually starts when I feel like someone is ignoring me or pulling away. I hate being alone—it makes everything worse. Sometimes yelling or cutting makes me feel better, but then I feel ashamed.”
<b>Quality</b>	“Can you describe what your emotional pain feels like?”	“It’s like a storm inside—so intense I can’t control it. One minute I’m fine, the next I’m crying, screaming, or wanting to disappear. It feels like I’m broken or empty.”
<b>Region</b>	“Do you feel any physical symptoms?”	“I feel like I can’t breathe when I get really upset. Sometimes I feel shaky. I’ve had stomachaches or headaches after a fight with my boyfriend.”
<b>Severity</b>	“On a scale of 0 to 10, how intense is your distress when it happens?”	“It’s a 10. I feel totally out of control, like I’ll explode or do something I’ll regret later.”
<b>Timing/ Treatment</b>	“When did these feelings or episodes start? How long do they usually last?”	“I’ve felt like this ever since I was a teenager. The episodes can come on fast, like after a text from my boyfriend, and can last for hours or days.”
<b>Understanding</b>	“How do you explain what you’re going through?”	“I’m very sensitive. I know I can be intense or push people away, but I can’t help it. If people leave me, I don’t know how to handle it. It’s like I feel everything too much.”

## SUICIDE AND SELF INJURY SCREENING

Clients being evaluated or treated for personality disorders may have suicidal ideation. It is important for the nurse to introduce suicide screening in a way that helps the client understand its purpose and normalize questions that might otherwise seem intrusive. The Patient Safety Screener (PSS-3) is an example of a brief screening tool to detect suicide risk in all client presenting to acute care settings.<sup>4</sup>

**Non-suicidal self-injury (NSSI)** refers to intentional self-inflicted destruction of body tissue without suicidal intention and for purposes not socially sanctioned. Common forms of NSSI include behaviors such as cutting, burning, scratching, and self-hitting. It is considered a maladaptive coping strategy without the desire to die. NSSI is a common finding among adolescents and young adults in psychiatric inpatient settings.<sup>5</sup>

Review information about suicide screening, the Patient Safety Screener, and screening for non-suicidal self-injury (NSSI) in the “[Assessment](#)” section of the *Applying the Nursing Process to Mental Health Care* chapter.

## CULTURAL ASSESSMENT

Cultural Formulation Interview (CFI) questions help nurses understand a client’s cultural background and how it influences their experience of mental health symptoms, including personality disorders.<sup>6</sup> Sample CFI questions

4. Suicide Prevention Resource Center. (n.d.). *The patient safety screener: A brief tool to detect suicide risk*. <https://sprc.org/micro-learning/patientsafetyscreener>
5. Cipriano, A., Cella, S., & Cotrufo, P. (2017). Nonsuicidal self-injury: A systematic review. *Frontiers in Psychology*, 8. <https://www.frontiersin.org/articles/10.3389/fpsyg.2017.01946/full>
6. DeSilva, R., Aggarwall, N. K., & Lewis-Fernandez, R. (2015). The DSM-5 cultural formulation interview and the evolution of cultural assessment in psychiatry.



focused specifically on understanding depression within a cultural context include the following:

- Cultural Definition of the Problem
  - “How would you describe the problems you’re having with relationships or emotions?”
  - “What do your family or friend call these difficulties?”
  - “Do others see your behavior or emotions as a problem? If so, how?”
- Cultural Perceptions of Cause, Context, and Support
  - “What do you think causes your emotional ups and downs or relationship conflicts?”
  - “How does your family or cultural group view traits like being very emotional, needing control, being distrustful, or acting out?”
  - “Are there things happening in your life right now that are making these problems worse?”
- Cultural Factors Affecting Coping and Help-Seeking
  - “What have you done to manage these emotions on your own?”
  - “Have you talked to family members, elders, or religious/spiritual leaders about these issues?”
  - “What kind of help do you think would work best for you?”
  - “Are there treatments or approaches you’re uncomfortable with, based on your beliefs or past experiences?”
- Cultural Features of the Nurse–Client Relationship
  - “Are there things about your cultural background that would help me understand you better or make you feel more comfortable?”
  - “Have you had any past experiences with mental health or medical providers that affect how you feel about treatment now?”
  - “Would you prefer working with a provider of a certain background or

*Psychiatric Times*, 32(6). <https://www.psychiatrictimes.com/view/dsm-5-cultural-formulation-interview-and-evolution-cultural-assessment-psychiatry>

gender?”

Findings from the cultural formulation interview are used to individualize a client’s plan of care to their preferences, values, beliefs, and goals.

## SPIRITUAL ASSESSMENT

The FICA Spiritual History Tool is a widely used assessment model for evaluating a client’s spiritual beliefs and how they may influence health, illness, and coping. FICA© is a mnemonic for the domains of Faith, Importance, Community, and Address in Care.<sup>7</sup> The data obtained from a FICA assessment can be helpful in understanding how clients with personality disorders draw on spirituality or religion for support. Spiritual distress is very common for clients experiencing serious illness, and nurses assist clients to adopt healthy coping strategies to deal with these life events. Addressing a client’s spirituality and advocating spiritual care have been shown to improve clients’ health and quality of life.<sup>8,9</sup>

7. GW School of Medicine & Health Sciences. (n.d.). *Clinical FICA tool*.  
<https://smhs.gwu.edu/spirituality-health/program/transforming-practice-health-settings/clinical-fica-tool>
8. Pilger, C., Molzahn, A. E., de Oliveira, M. P., & Kusumota, L. (2016). The relationship of the spiritual and religious dimensions with quality of life and health of patients with chronic kidney disease: An integrative literature review. *Nephrology Nursing Journal: Journal of the American Nephrology Nurses’ Association*, 43(5), 411–426. <https://pubmed.ncbi.nlm.nih.gov/30550069/>
9. Puchalski, C., Jafari, N., Buller, H., Haythorn, T., Jacobs, C., & Ferrell, B. (2020). Interprofessional spiritual care education curriculum: A milestone toward the provision of spiritual care. *Journal of Palliative Medicine*, 23(6), 777–784.  
<https://doi.org/10.1089/jpm.2019.0375>

Table 10.4c summarizes a sample spiritual assessment questions and sample responses from a client experiencing depression.

Table 10.4c Sample FICA Spiritual Assessment Questions for Clients with Personality Disorders

Domain	Sample Assessment Question	Sample Client Response
<b>Faith</b>	“Do you consider yourself spiritual or religious? What gives your life meaning?”	“Lately, I’ve been feeling disconnected from everyone and everything. I pray sometimes, but it feels like no one’s listening.”
<b>Importance</b>	“How important is faith or religious belief in your life?”	“My faith used to help me feel strong, especially during hard times. Now, I feel like I’ve lost that connection. I feel too broken to be helped.”
<b>Community</b>	“Are you part of a spiritual or religious community? Does participation in this community provide support when you’re feeling sad or in a low mood?”	“I haven’t gone to church in months. Some people from church have reached out, but I just feel like being alone when I am home after work.”
<b>Address in Care</b>	“Would you like me (or the health care team) to address spiritual issues during your care? Would you like to speak with a chaplain?”	“Maybe. I don’t want to feel judged for how I feel or for having doubts.”

Feelings of abandonment and spiritual distress are common in individuals experiencing personality disorders and may compound their symptoms. Nurses may recognize cues of spiritual distress and offer to connect the client with a chaplain or spiritual care services. Spiritual goals may be included in the nursing care plan if the client finds them valuable.

## FAMILY DYNAMICS

Family dynamics are included in a psychosocial assessment, especially for children, adolescents, and older adults. **Family dynamics** refers to the patterns of interactions among relatives, their roles and relationships, and the various factors that shape their interactions. Because family members rely on

each other for emotional, physical, and economic support, they are primary sources of relationship security or stress. Family dynamics and the quality of family relationships can have either a positive or negative impact on an individual's health. For example, secure and supportive family relationships can provide love, advice, and care, whereas stressful family relationships can be burdened with arguments, unhealthy relationships, and a lack of support.<sup>10</sup>

Unhealthy family dynamics can cause children to experience trauma and stress as they grow up. This type of exposure, known as adverse childhood experiences (ACEs), is linked to an increased risk of developing mental health disorders.<sup>11</sup> Review information about adverse childhood experiences (ACEs) in the “[Mental Health and Mental Illness](#)” section of Chapter 1.

## Laboratory Testing

There is no specific laboratory test that diagnoses personality disorders. Laboratory or diagnostic tests may be used to rule out other possible causes for the behaviors the client is exhibiting. For example, a thyroid stimulating hormone (TSH) test may be ordered because thyroid disorders can affect mood.

## Life Span Considerations

### CHILDREN AND ADOLESCENTS

Formal diagnosis of a personality disorder is not made before age 18, but traits may be present earlier, such as emerging patterns of emotional instability,

10. Jabbari, B., Schoo, C., & Rouster, A. S. (2023). *Family dynamics*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK560487/>

11. Jabbari, B., Schoo, C., & Rouster, A. S. (2023). *Family dynamics*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK560487/>

conduct issues, or rigid thinking. Children and adolescents may demonstrate increased conflict with authority figures or manipulative behavior, impulsivity, or self-harm.

## OLDER ADULTS

Personality changes in late life may suggest neurocognitive disorder, stroke, or side effects from medications. Personality traits such as paranoia, rigidity, dependency, or social isolation may become more prominent as the individual experiences loss or isolation.

## Diagnosis (Analyze Cues)

Mental health disorders are diagnosed by mental health providers using the diagnostic criteria in the *DSM-5-TR*. Personality disorder diagnoses are typically not made until late adolescence or over the age of 18 because it is important to determine if the symptoms are traits of a developmental stage or pervasive traits of a personality disorder in multiple contexts. Nurses customize nursing diagnoses based on their type of personality disorder, their current signs and symptoms, and the effects on their and their family's functioning.

Nurses create individualized nursing care plans based on the client's response to their mental health disorder(s). Common nursing diagnoses related to the clusters of personality disorders include the following:

- **Cluster A:** *Social Isolation, Disturbed Thought Process, Risk for Loneliness*
- **Cluster B:** *Risk for Suicide, Risk for Self-Directed Violence, Social Isolation, Chronic Low Self-Esteem, Ineffective Coping*
- **Cluster C:** *Anxiety, Risk for Loneliness, Social Isolation*

Common nursing diagnoses for clients diagnosed and hospitalized with personality disorders are further described in Table 10.4d.

Table 10.4d Common Nursing Diagnoses for Clients With Personality Disorders<sup>12,13</sup>

12. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.
13. Ackley, B., Ladwig, G., Makic, M. B., Martinez-Kratz, M., & Zanoliti, M. (2020). *Nursing diagnosis handbook: An evidence-based guide to planning care* (12th ed.). Elsevier.

Nursing Diagnosis	Definition	Selected Defining Characteristics and/or Risk Factors
<b>Risk for Suicide</b>	Susceptible to self-inflicted, life-threatening injury.	<ul style="list-style-type: none"> <li>• Reports desire to die</li> <li>• Statements regarding killing self</li> <li>• Hopelessness</li> <li>• Social isolation</li> </ul>
<b>Risk for Self-Mutilation</b>	Deliberate self-injurious behavior causing tissue damage with the intent of causing nonfatal injury to attain relief of tension.	<ul style="list-style-type: none"> <li>• Cuts or scratches on body</li> <li>• Ingestion or inhalation of harmful substances</li> <li>• Self-inflicted burns</li> </ul>
<b>Risk for Other-Directed Violence</b>	Susceptible to behaviors in which an individual demonstrates they can be physically, emotionally, and/or sexually harmful to others.	<ul style="list-style-type: none"> <li>• History of childhood abuse</li> <li>• History of cruelty to animals</li> <li>• History of witnessing family violence</li> <li>• History of fire-setting</li> </ul>

<b>Ineffective Coping</b>	A pattern of invalid appraisal of stressors, with cognitive and/or behavioral efforts, that fails to manage demands related to well-being.	<ul style="list-style-type: none"> <li>• Destructive behavior toward self or others</li> <li>• Ineffective coping strategies</li> <li>• Ineffective problem-solving skills</li> </ul>
<b>Defensive Coping</b>	Repeated projection of falsely positive self-evaluation based on a self-protective pattern that defends against underlying perceived threats to positive self-regard.	<ul style="list-style-type: none"> <li>• Difficulty maintaining relationships</li> <li>• Hypersensitivity to criticism</li> <li>• Projection of blame</li> <li>• Projection of responsibility</li> </ul>
<b>Social Isolation</b>	Aloneness experienced by the individual and perceived as imposed by others and as a negative or threatening state.	<ul style="list-style-type: none"> <li>• Hostility</li> <li>• Values incongruent with cultural norms</li> <li>• History of rejection</li> </ul>



<b>Ineffective Family Health Management r/t manipulative behavior</b>	Aloneness experienced by the individual and perceived as imposed by others and as a negative or threatening state.	<ul style="list-style-type: none"> <li>• Impaired communication patterns</li> <li>• Disturbed thought processes</li> <li>• Delusional thinking</li> </ul>
<b>Risk for Spiritual Distress as manifested by poor relationships</b>	A state of suffering related to the impaired ability to experience meaning in life through connections with self, others, the world, or a superior being.	<ul style="list-style-type: none"> <li>• Ineffective coping strategies</li> <li>• Perceived insufficient meaning in life</li> <li>• Hopelessness</li> <li>• Social alienation</li> </ul>

## Outcome Identification (Generate Solutions)

Outcomes should address the established nursing diagnoses for each client with prioritization on safety. For example, if the client has a nursing diagnosis of *Risk for Self-Mutilation*, a SMART outcome could be, “The client will refrain from intentional self-inflicted injury.” Read more information about setting SMART outcomes in the “[Application of the Nursing Process in Mental Health Care](#)” chapter.

Examples of other SMART outcomes for clients with personality disorders may include the following<sup>14</sup>:

14. Ackley, B., Ladwig, G., Makic, M. B., Martinez-Kratz, M., & Zanotti, M. (2020).

- The client will remain safe and free of injury.
- The client will seek help when experiencing urges to self-mutilate.
- The client will identify three triggers to self-mutilation after the teaching session.
- The client will describe two preferred healthy coping strategies after the teaching session.

## Planning (Generate Solutions)

Evidence-based interventions are critical and should focus on managing the client's behaviors. Interventions may include establishing clear boundaries, using specific communication styles, and providing reassurance.

Pharmacotherapy may be used to manage specific symptoms, although no medications are specifically approved for personality disorders.

## Promoting Safety

Individuals diagnosed with personality disorder may be suicidal, self-mutilating, impulsive, angry, manipulative, or aggressive. Nurses plan interventions according to the symptoms the client is currently exhibiting with the goal of keeping the client, others, and themselves safe and free of injury. Any threats should be taken seriously. Review interventions for clients diagnosed with *Risk of Suicide* in the "[Application of the Nursing Process in Mental Health Care](#)" chapter.

Clear boundaries and limits should be set and consistently reinforced by the health care team. When behavioral problems emerge, the nurse should calmly review therapeutic goals, limits, and boundaries with the client.<sup>15</sup>

*Nursing diagnosis handbook: An evidence-based guide to planning care* (12th ed.). Elsevier.

15. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

A crisis/safety plan must be developed with the client that includes the following components:

- Identifying thoughts or behaviors that increase the risk of harming self or others
- Identifying people, events, or situations that trigger those thoughts or behaviors
- Implementing coping strategies
- Reaching out to other coping resources

For example, if a client performs superficial self-injurious behavior, the nurse should act based on agency policy while remaining neutral and dressing the client's self-inflicted wounds in a matter-of-fact manner. The client may be asked to write down the sequence of events leading up to the injuries, as well as the consequences, before staff will discuss the event. This cognitive exercise encourages the client to think independently about their triggers and behaviors and facilitates discussion about alternative actions.<sup>16</sup>

Review information regarding developing a safety plan in the “[Establishing Safety](#)” section of the “Foundational Mental Health Concepts” chapter.

## Implementation (Take Action)

As previously discussed in this chapter, there are ten different personality disorders that are categorized into three clusters (A, B, and C) in the DSM-5. Personality disorders within each cluster have similar patterns of behavior. Cluster A personality disorders include paranoid personality disorder, schizoid personality disorder, and schizotypal personality disorder. Cluster A is characterized as the odd, eccentric cluster. Individuals with these types of disorders often experience social awkwardness. Cluster B personality disorders include antisocial, borderline, histrionic, and narcissistic personality

16. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

disorders. Cluster B personality disorders are characterized by dramatic, overly emotional, or unpredictable thinking or behavior. Cluster C personality disorders include avoidant, dependent, and obsessive-compulsive personality disorders. Cluster C personality disorders are characterized by anxious, fearful thinking or behavior.

These disorders are often ego-syntonic, meaning that the behaviors and thoughts are consistent with the individual's self-perception and are not seen as problematic by the individual, despite causing significant issues in their life. This characteristic of personality disorders can make it challenging to implement nursing interventions when the client does not believe a problem exists.

## Nursing Interventions for Personality Disorders Based on Categories of the APNA Implementation Standard

Nursing interventions for clients with personality disorders can be categorized based on the American Psychiatric Nurses Association (APNA) standard for *Implementation* that includes the *Coordination of Care; Health Teaching and Health Promotion; Pharmacological, Biological, and Integrative Therapies; Milieu Therapy; and Therapeutic Relationship and Counseling*. Read more about these subcategories in the “[Application of the Nursing Process in Mental Health Care](#)” chapter. See examples of interventions for each of these categories for clients with depressive disorders in Table 10.4e.

Table 10.4e Examples of Nursing Interventions For Personality Disorders Based on Subcategories of APNA Implementation Standard

Subcategory of the APNA Standard of Implementation	The nurse will ...	Rationale
<b>Coordination of Care</b>	<ul style="list-style-type: none"> <li>– Collaborate with mental health professionals (e.g., therapists, psychiatrists) to ensure continuity of care.</li> <li>– Communicate changes in behavior or mood with the health care provider.</li> <li>– Facilitate referrals to community support resources such as support groups.</li> <li>– Assist with care transitions (e.g., hospital discharge to long-term care or other community-based facilities).</li> </ul>	<p>Personality disorders often require long-term, structured treatment involving multiple providers. Coordinated care ensures consistency and promotes better outcomes.</p>
<b>Health Teaching and Health Promotion</b>	<ul style="list-style-type: none"> <li>-Educate clients on emotional regulation, interpersonal effectiveness, and stress management strategies (e.g., using DBT skills).</li> <li>– Teach clients about triggers, patterns of behavior, and consequences.</li> <li>– Promote self-care routines and adaptive coping techniques.</li> </ul>	<p>Many clients with personality disorders struggle with emotional impulsivity and poor coping skills. Health teaching can enhance insight and self-management, thus reducing crises and potential hospitalizations.</p>

<b>Pharmacological, Biological, and Integrative Therapies</b>	<ul style="list-style-type: none"> <li>– Administer prescribed medications for symptoms (e.g., mood stabilizers for impulsivity, SSRIs for depression/anxiety).</li> <li>– Educate the client on the purposes of the medications-Monitor for side effects and adherence</li> <li>– Encourage integrative therapies (e.g., mindfulness, journaling) to supplement psychotherapy.</li> </ul>	<p>While medications do not cure personality disorders, they can manage co-occurring symptoms. Combined with therapy and self-care strategies, they improve overall stability.</p>
<b>Milieu Therapy</b>	<ul style="list-style-type: none"> <li>– Create a structured and predictable environment to promote safety and reduce chaos.</li> <li>– Model and reinforce appropriate social behaviors in group settings.</li> <li>– Establish clear, consistent boundaries and behavioral expectations.</li> </ul>	<p>Clients with personality disorders often test limits or experience emotional dysregulation. A therapeutic milieu with structure and consistency supports stability and learning.</p>
<b>Therapeutic Relationship and Counseling</b>	<ul style="list-style-type: none"> <li>– Build trust using empathy, consistency, and nonjudgmental communication.</li> <li>– Set and maintain professional boundaries to avoid manipulation or dependency.</li> <li>– Use therapeutic communication to explore core beliefs and relational patterns.</li> <li>– Encourage participation in long-term psychotherapy (e.g., DBT, CBT).</li> </ul>	<p>Clients often have a history of unstable or maladaptive relationships. A consistent, respectful therapeutic relationship helps model healthy interaction and provides a secure base for growth.</p>

## Nursing Interventions for Physiological Signs of Personality Disorders

Although personality disorders are primarily psychiatric in nature, they often contribute to physiological symptoms such as sleep disturbances, somatic

complaints, self-harm behaviors, disordered eating, and fatigue. See common interventions for these conditions in Table 10.4f.

Table 10.4f Nursing Interventions Targeting Physiological Signs of Personality Disorders<sup>17</sup>

17. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

Problem/Intervention	Rationale
<p><b>Nutrition and Eating Behaviors</b></p> <ul style="list-style-type: none"> <li>• <b>Assess for disordered eating patterns (e.g., bingeing or restrictive intake).</b></li> <li>• <b>Collaborate with dietitian for nutritional planning.</b></li> <li>• <b>Monitor intake/output and weight if needed.</b></li> <li>• <b>Support body positivity and emotional awareness around food.</b></li> </ul>	<p>Disordered eating may be associated with efforts to control overwhelming emotions. Balanced nutrition helps stabilize mood and reduce health complications.</p>
<p><b>Sleep</b></p> <ul style="list-style-type: none"> <li>• <b>Assess sleep patterns and contributing factors (e.g., anxiety, racing thoughts, substance use).</b></li> <li>• <b>Encourage regular sleep routines and calming bedtime rituals.</b></li> <li>• <b>Provide education on sleep hygiene (e.g., limiting screen time, reducing caffeine).</b></li> <li>• <b>Consult provider for possible medications.</b></li> </ul>	<p>Many clients with personality disorders experience insomnia due to emotional dysregulation, anxiety, or impulsivity. Improving sleep enhances mood stability and reduces irritability and impulsive behavior.</p>



<p><b>Elimination (Gastrointestinal Discomfort)</b></p> <ul style="list-style-type: none"> <li>• <b>Validate physical symptoms without reinforcing excessive health anxiety.</b></li> <li>• <b>Rule out medical causes in collaboration with the healthcare team.</b></li> <li>• <b>Educate on the mind–body connection and introduce relaxation techniques.</b></li> </ul>	<p>Clients, especially those with borderline or somatization traits, may express emotional distress through physical symptoms like gastrointestinal (GI) complaints. Acknowledging the GI complaint while promoting coping reduces unnecessary interventions and increases insight.</p>
<p><b>Fatigue/Energy Deficit</b></p> <ul style="list-style-type: none"> <li>• <b>Assess energy levels and daily activity patterns.</b></li> <li>• <b>Encourage structured routines with scheduled rest and activity.</b></li> <li>• <b>Promote moderate physical activity to increase energy and reduce tension.</b></li> <li>• <b>Monitor for medication side effects contributing to fatigue.</b></li> </ul>	<p>Emotional volatility and poor sleep can result in persistent fatigue. Routines help regulate biological rhythms, while physical activity improves mood and energy.</p>

<p><b>Self Care Deficits: Self-Harm or Injury</b></p> <ul style="list-style-type: none"> <li>• <b>Monitor for and document signs of cutting, burning, or other non-suicidal self-injury (NSSI).</b></li> <li>• <b>Remove or limit access to sharp objects or means of injury.</b></li> <li>• <b>Engage client in developing a safety plan and alternative coping strategies (e.g., using ice, rubber bands).</b></li> <li>• <b>Refer client to DBT-based interventions.</b></li> </ul>	<p>Self-injury is a maladaptive coping mechanism to relieve emotional pain or dissociation, especially in borderline personality disorder. Safety monitoring and skill-building reduce risk and promote healthier regulation.</p>
--	---

## Communication Tips for Clients with Personality Disorders

Helpful communication techniques for clients with personality disorders and their rationale are described in the following box.

### Communication Tips: Personality Disorders<sup>18,19</sup>

- Establish and maintain clear, consistent boundaries.
  - **Rationale:** Clients with personality disorders

<sup>18</sup>. Ward R. K. (2004). Assessment and management of personality disorders. *American family physician*, 70(8), 1505–1512.

<sup>19</sup>. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

—especially those with borderline or antisocial traits—may test or violate boundaries.

Consistent enforcement of limits fosters safety, trust, and structure.

- Use calm, neutral, and non-reactive language.
  - **Rationale:** Emotional neutrality helps prevent escalation when clients display anger, impulsivity, or mood instability. It also reduces reinforcement of attention-seeking or manipulative behaviors.
- Avoid power struggles or confrontation.
  - **Rationale:** Clients with narcissistic or paranoid traits may become defensive or combative. Use collaborative, non-authoritarian language to de-escalate and preserve the therapeutic alliance. Read more about de-escalating in the following subsection.
- Validate feelings without encouraging distorted behaviors.
  - **Rationale:** Statements like “I can see that you’re really upset” acknowledge emotional pain without reinforcing maladaptive coping (e.g., threats, splitting, self-harm).
- Be direct, honest, and transparent.
  - **Rationale:** Clear and open communication reduces misinterpretation and builds trust, especially with clients who have suspicious or

paranoid tendencies.

- Focus on the present.
  - **Rationale:** Redirect conversations from historical grievances or perceived slights to current goals or feelings. This helps reduce rumination and increases emotional regulation.
- Avoid over-involvement or “rescuing” behaviors.
  - **Rationale:** Maintain professional boundaries to promote client autonomy and reduce dependency.
- Watch for splitting behaviors among staff.
  - **Rationale:** Clients may view staff as all good or all bad, leading to division. Maintain team communication and a unified, consistent approach. Avoid taking sides or reacting emotionally.
- Encourage reflection instead of impulsive reaction.
  - **Rationale:** Help the client slow down and explore what they’re feeling before acting. For example, “Let’s pause for a moment and talk about what just happened.”
- Reinforce positive coping and interpersonal efforts.
  - **Rationale:** Recognize and praise the client’s use of healthy communication, emotional control, or insight to supports behavior change and

build self-esteem.

## De-Escalating

The nurse should implement de-escalation strategies if the client exhibits signs of increasing levels of anxiety or agitation. Strategies include the following:

- Speaking in a calm voice
- Avoiding overreacting
- Implementing active listening
- Expressing support and concern
- Avoiding continuous eye contact
- Asking how you can help
- Reducing stimuli
- Moving slowly
- Remaining patient and not rushing them
- Offering options instead of trying to take control
- Avoiding touching the client without permission
- Verbalizing actions before initiating them
- Providing space so the client doesn't feel trapped
- Avoiding arguing and judgmental comments
- Setting limits early and enforcing them consistently across team members
- Addressing manipulative behaviors therapeutically

If the client continues to escalate, measures must be taken to keep the client and others safe. Review signs of crisis and crisis interventions in the “[Stress, Coping, and Crisis Intervention](#)” chapter. If interventions are not effective in de-escalating a client at risk to themselves or others, seclusion or restraints may be required. Review using seclusion and restraints in the “[Psychosis and](#)

Schizophrenia” chapter. Review ANA guidelines on using restraints in the “Client Rights” section of the “Legal and Ethical Considerations in Mental Health Care” chapter and information on safely implementing restraints in the “Workplace Violence” section of the “Trauma, Abuse, and Violence” chapter.

## Coping Strategies

Teaching self-care and coping strategies is helpful for people diagnosed with personality disorders and their loved ones.<sup>20</sup> Read about stress management and coping strategies in the “Stress, Coping, and Crisis Intervention” chapter.

For clients seeking immediate relief from intense symptoms such as panic or depersonalization, nurses can teach how to stimulate the parasympathetic nervous system. Stimulation of the vagal nerve can result in an immediate, direct relief of intense emotions. This can be accomplished by doing the following<sup>21</sup>:

- Applying ice or ice-cold water to the face
- Performing paced-breathing techniques in which the exhalation phase is at least two to four counts longer than the inhalation phase. For example, advise the client to inhale while counting to four and then exhale while counting to eight.

20. American Psychiatric Association. (n.d.). *What are personality disorders?* <https://www.psychiatry.org/patients-families/personality-disorders/what-are-personality-disorders>

21. Nelson, K. J. (2021). Pharmacotherapy for personality disorders. *UpToDate*. <https://www.uptodate.com>

## Collaborative Interventions

The nurse encourages collaboration with other disciplines and encourage psychotherapy and pharmacotherapy as directed by the healthcare provider. For more information on psychotherapy and pharmacotherapy in regard to treating personality disorders, see the [“Treatment for Personality Disorders”](#) section of this chapter.

## Evaluation (Evaluate Outcomes)

Refer to the SMART outcomes established for each individual client to evaluate the effectiveness of the planned interventions. Modification of the established nursing care plan may be required based on the effectiveness of the interventions.

In general, the nurse can evaluate the effectiveness of interventions in clients with personality disorders with the following questions:

- Does client recognize symptoms are related to their mental health condition?
- Is the client successfully implementing adaptive coping strategies to manage their mental health disorder?
- Is the client following their prescribed treatment plan?
- Is the client adequately performing self-care activities?
- Is the client able to maintain satisfying interpersonal relationships?
- Is the client able to successfully function socially or other important areas of functioning?

## 10.5 Spotlight Application

Kay is a 27-year-old female who is admitted to the mental health unit for feelings of depression and suicidal ideations. She has a history of cutting her legs and arms since she was a teenager. She started psychiatric treatment at age 16 and has been admitted to the psychiatric hospital three times due to overdoses. She admits to a history of promiscuous behavior and occasional marijuana use. She is unmarried but reports having several relationships with male partners. However, she shares that the relationships “never worked out” because the partners didn’t pay enough attention to her or text her as frequently as she expected, which made her angry. She describes feelings she experienced in relationships where one day she felt as if her partner was the “best thing ever,” but the next day she “can’t stand him.” She admits to abusive behaviors toward her partners when she was angry, but states that afterwards she experiences anxiety fearing the loss of the relationship. She acknowledges having trouble getting along with others. She has very few close friends and refers to previous friends as “losers.” She is unemployed with a history of several jobs from which she was terminated because of problems with anger control. There is evidence of scarring and recent cuts on bilateral lower arms. She related these injuries to self-cutting, which she reports “makes me feel better.” She reports attending dialectical behavior therapy (DBT) in the past but hasn’t attended therapy for over a year. She feels therapy helped her to learn how to feel more “in control of her extreme feelings.”

### Critical Thinking Questions:

1. List the symptoms Kay is experiencing that supports the diagnosis of borderline personality disorder (BPD).

*Symptoms of BPD include anger, anxiety, impulsiveness, difficulty controlling emotions, self-injury, suicidal, intense/stormy relationships, and risky behaviors.*

2. Identify possible risk factors contributing to BPD.



*Possible risk factors include environmental factors that may contribute to BPD, including neglect, abuse, and genetics.*

3. Discuss two types of psychotherapy that may be prescribed for clients diagnosed with BPD.

*Cognitive behavioral therapy, dialectical behavioral therapy, interpersonal therapy, mentalization-based therapy, and psychodynamic therapy are types of psychotherapy that may be prescribed for clients.*

4. List at least three nursing interventions that should be included in this client's care plan.

- *Explain policies, expectations, rules, and consequences upon admission.*
- *Search the client's belongings and remove anything that could be used to inflict harm to self or others.*
- *Implement suicide precautions based on a suicide risk screening tool.*
- *Set and document clear boundaries and limits and share them with team members.*
- *Encourage consistent implementation of limits by all team members; do not allow bargaining.*
- *Call out manipulative behavior when it occurs.*
- *If an episode of cutting occurs during hospitalization, remain neutral and dress the client's self-inflicted wounds in a matter-of-fact manner. Ask the client to write down the sequence of events leading up to the injuries before discussing the event. Encourage the client to think independently about her triggers and behaviors and facilitate discussion about alternative actions.*

### Learning Activities

*(Answers to “Learning Activities” can be found in the “Answer Key” at the end of the book. Answers to the interactive activities are provided as immediate feedback.)*

*Please respond to the following questions.*

1. Compare the difference between obsessive-compulsive personality disorder and the anxiety disorder of obsessive-compulsive disorder (OCD).
2. Name that personality disorder. Read each scenario and choose a personality disorder associated with the traits included in the scenario:
  - a. Darla enters the breakroom at her job. She is dressed provocatively with excessive jewelry. She makes eye contact with everyone. She sits down and quickly interjects herself into the conversation. When one of her coworkers shares pictures of her new puppy,

she pulls out her cell phone to show pictures of herself and tells the group how she had a romantic weekend with her handsome, rich boyfriend.

- b. Bob has been summoned to his supervisor's office. Despite his adequate job performance, he thinks to himself, "I knew it! She's out to get me and I am getting fired! She and others walk by my desk several times a day just to check up on me and see what I am doing. And that new employee has been trying to strike conversations with me, but I am not falling for that. I am sure he just wants something from me."
- c. Debbie is distraught after her boyfriend called her and told her their relationship was over. She began yelling on the phone, "I love you and can't live without you." The ex-boyfriend ends the phone call. Debbie begins

calling him over and over;  
however, he doesn't  
answer the phone. She  
then leaves a message  
stating, "I hate you. I never  
want to see you again."  
Overcome with emotion,  
she begins to cut her wrist  
with a bobby pin.

- d. George is a loner and lives alone with his five cats. He has family members who live in the area, but he hasn't seen them in years, and that is okay with him. He has never been married or been in an intimate relationship. He does not have any close friends. He enjoys his job as a night security guard because there is little requirement for social interaction.
- e. Jordan is always trying to please others. She has trouble starting or completing projects because of a lack of self-confidence. She requires much reassurance and advice when making ordinary decisions. She

allows her boyfriend to tell her what to wear, what kind of job to look for, and with whom to associate. She is afraid of him leaving her, so she is very careful not to get him mad at her.

- f. Roger has been arrested several times for domestic abuse and driving while intoxicated. He shows no remorse for any of his actions or the injuries he has caused others. He lies, breaks laws, and has no regard for the feelings of others. As a teenager, he was always in trouble and truant from school. He was incarcerated in the past but found it easy to manipulate the guards into breaking the rules for him.
- g. Deanna has very few friends. She is shy and avoids social interaction, even at work. She is worried that if she did get to know people better, she might say or do something embarrassing, and they

might criticize or reject her. During her performance evaluation with her supervisor, she left the office crying because of minor constructive feedback she received. To prevent these painful experiences, she believes it is best to keep to herself.

- h. Billy tells everyone how important he is to his company. He believes there is no way the company would be successful without him. During work meetings, he monopolizes the conversation and strives to be the center of attention. He often asks others, “Don’t you think I did a great job with that?” Although he does not have any close relationships with coworkers, he does collaborate with a few of them because they are able to help him accomplish his goals.
- i. Bobbi works as an

administrative assistant. She is very organized and spends a lot of time making lists, scheduling, and reviewing details.

Although she has several friends, she often passes on the opportunity to get together because she spends most of her time devoted to work.

Sometimes, despite her efforts, she has difficulty finishing a project because she feels the need to check things “one more time.”

- j. Paul is loner and lacks close friends outside of his immediate family. Most view him as being very unusual, including his odd way of dressing. He wears ill-fitting and bizarre clothing combinations, such as winter boots with shorts. He is very superstitious and believes he is psychic. He believes prime numbers are unlucky and avoids objects (e.g., building floors and house numbers) that are

prime numbers.

3. Check your medication knowledge.  
Select which medications are commonly used to treat severe symptoms of personality disorders.
  - a. What medications are used to treat hallucinations and paranoia?
  - b. What medications are used to treat depressed mood, mood lability, anger, and anxiety?
  - c. What medications are used to treat self-injurious behavior?
4. Compare normal adolescent development with trait similarities of personality disorders.



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=413#h5p-35>



1



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=413#h5p-36>

2



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=413#h5p-37>

3

1. “MH Personality Disorders Glossary Cards” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
2. “MH Personality Disorders Question Set 1” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
3. “MH Personality Disorders Crossword” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

## Case Study

### Nurse Handoff Report:

**Situation:** Tara Mitchell was admitted to the hospital last night for observation after being found outside the back door of a local bar where she works as a waitress, burning her arm with a cigarette. She has a history of non-suicidal self-injurious behavior since she was a teenager, including cutting, burning herself, and pulling out her hair. Eight months ago, she was admitted to the hospital after a suicide attempt in which she jumped out in front of a car after her shift at another local bar. She states that thinking about suicide helps her relax and she feels like she can “escape” from her distracted thoughts that occur throughout the day.

**Background:** Tara was diagnosed with borderline personality disorder following her most recent hospitalization. She has been non-compliant with therapy. Her medical records indicate that when growing up, her father was in the military and was often physically abusive, especially when using alcohol or drugs. He was diagnosed with PTSD after returning from serving in Somalia. Her mother worked two jobs when she was growing up and was not around much. She has two siblings, a brother and a sister, but neither remains in the area and she has not had contact with them for “several years.” Tara dropped out of college with one year of classes left to complete a Graphic Design degree. She has spent the past three years changing jobs frequently, mostly working as a bartender or waitress in local restaurants and bars.

**Assessment:** Since admission, Tara has been quiet and withdrawn in her room. She complains of the “worst pain ever” related to 8 circular burn marks that are present on her left

forearm in the shape of the number 8. She does not make eye contact when communicating, but instead looks outside the door or over the person's shoulder. Her hair is down and her makeup is smeared below her eyes. She is wearing hospital scrubs. During her admission assessment, Tara stated she burned her arm because she was "upset at the bartender because they did not make her drink order first." She said, "They hate me. I don't know why I keep coming back to that job. They are the most incompetent people in the world!"

Tara "zones out" at times during conversation, even in the middle of speaking a sentence. She states, "I don't even know who Tara is. I feel like I'm watching what she does from outside of my body."

Tara explains that she has a long history of frequently changing hobbies, jobs, fashion sense, and relationships based on her current social circle. At times, she thinks her friends, family, and partner are "the best thing that ever happened to me," buying them gifts and sending caring messages frequently. At other times, she completely turns on her relationships, saying, "I can't stand them" and lashes out at them in person, physically, and/or on social media. These actions cause her to feel remorse, and she feels immediate regret and a strong fear that they will leave her. Tara also has a history of sexual promiscuity, even with partners she didn't even know, and doing such acts so that her current partner doesn't know.

When asked about why she burned herself with the cigarette, she explains that she was angry that she had to work while her current boyfriend, Josh, went to a bachelor party for his best friend, Ben. She states, "I was sure that Josh was going to hook up with someone else at the party."

### Reflective Questions:

1. What CUES do you recognize as relevant for providing client-centered care for Tara?
2. What is your hypothesis for Tara's priority nursing problem(s)?
3. What are your first steps in providing nursing care for Tara?

- ▶ Test your clinical judgment with a NCLEX Next Generation-style question: [Chapter 10, Assignment 1](#)<sup>4</sup>



- ▶ Test your clinical judgment with a NCLEX Next Generation-style question: [Chapter 10, Assignment 2](#)<sup>5</sup>



4. "MH Personality Disorders Next Gen Question 1" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

5. "MH Personality Disorders Next Gen Question 2" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

- ▶ Test your clinical judgment with a NCLEX Next Generation-style case study: [Chapter 10, Case Study 1](#)<sup>6</sup>

6. “MH Personality Disorders Next Gen Case Study” by Kellea Ewen is licensed under [CC BY-NC 4.0](#)

## X Glossary

**Body illusion:** A perception that one's own body is significantly different from its actual configuration. For example, a person lying in bed may feel as if they are levitating.

**Cluster A personality disorders:** Include paranoid personality disorder, schizoid personality disorder, and schizotypal personality disorder; characterized by odd, eccentric, and socially awkward behavior.

**Cluster B personality disorders:** Antisocial, borderline, histrionic, and narcissistic personality disorders; characterized by dramatic, overly emotional, or unpredictable thinking or behavior.

**Cluster C personality disorders:** Avoidant, dependent, and obsessive-compulsive personality disorders; characterized by anxious, fearful thinking or behavior.

**Dissociative symptoms:** The experience of detachment or feeling as if one is outside one's body with loss of memory.

**Ego synotic:** Behaviors and thoughts are consistent with the individual's self-perception and are not seen as problematic by the individual, despite causing significant issues in their life

**Entitlement:** Unreasonable expectations of especially favorable treatment.

**Ideas of reference:** False beliefs that coincidental events relate to oneself. For example, a person shopping in a store sees two strangers laughing and believes they are laughing at them, when in reality the other two people do not even notice them.

**Interpersonal therapy:** Therapy which focuses on developing healthy relationships with others.

**Magical thinking:** The idea that one can influence the outcome of specific events by doing something that has no bearing on the circumstances. For example, a person watching a baseball game exhibits magical thinking when

believing that holding the remote control in a certain position caused their favorite player to hit a home run.

**Mentalization-Based Therapy:** A type of psychotherapy designed to help individuals improve their ability to understand and interpret their own and others' mental states—such as thoughts, feelings, and intentions. The goals are to improve self-awareness and understanding of one's own emotions, enhance empathy and understanding of others' perspectives. This subsequently helps individuals develop better interpersonal skills and communication.

**Personality:** A relatively stable pattern of thinking, feeling, and behaving that evolves over a person's lifetime and is unique to each individual. It is influenced by one's experiences, environment (surroundings and life situations), and inherited characteristics.

**Personality disorder:** An enduring pattern of inner experience and behavior that deviates significantly from the expectations of one's culture. Its onset can be traced back to adolescence or early adulthood and is present in a variety of contexts. This pattern of behavior is manifested in two or more of the following areas: cognition/perceptions, affect, interpersonal functioning, and impulse control.

**Personality traits:** Characteristics, whether considered good or bad, that make up one's personality.

**Psychodynamic therapy:** Therapy which promotes self-reflection and self-examination of problematic relationship patterns and unresolved conflicts.

**Psychoeducation:** Education which teaches the client about their medications, psychotherapies, and support groups available in the community

**Splitting:** A pattern of unstable and intense personal relationships characterized by alternating between extremes of idealization and devaluation.









### Learning Objectives

- Identify assessment cues of thought behaviors
- Identify nursing priorities for clients with thought disorders
- Plan outcomes for clients with thought disorders
- Differentiate safety/protective interventions for clients with thought disorders
- Apply evidence-based practice when planning care and interventions for clients with thought disorders
- Analyze treatments for clients with thought disorders
- Apply the nursing process to clients with thought disorders at risk for suicide
- Compare and contrast delirium, psychosis, and schizophrenia

Have you ever cared for a client who was confused, disoriented, had a change in mental status, or was experiencing delirium? These are considered “altered thought processes.” There are several potential medical causes of delirium, such as a urinary tract infection in an elderly client, hypo or hyperglycemia, drug or alcohol intoxication. There are also mental health disorders that can cause altered thought processes, such as schizophrenia. This chapter will discuss psychosis, delirium, and schizophrenia and explain how to care for clients experiencing hallucinations, delusions, and other symptoms of altered thought processes.

## 11.2 Psychosis and Delirium

### Psychosis

Approximately 3 percent of the people in the United States experience psychosis at some point in their lives. About 100,000 adolescents and young adults in the US experience their first episode of psychosis each year.<sup>1</sup>

**Psychosis** is a broad term that refers to a symptom complex characterized by hallucinations, delusions, and disorganized thinking. It can occur in various psychiatric and medical conditions, including schizophrenia, mood disorders, and substance-induced states. Symptoms of psychosis include **delusions** and **hallucinations**, where the client is seeing, hearing, or experiencing other sensations and experiences that aren't real. See Figure 11.1.<sup>2</sup> for artwork titled *Hallucinations*. Other symptoms of psychosis include incoherent or nonsensical speech and behavior that is inappropriate for the situation. A **psychotic episode** is a specific, time-limited occurrence of psychotic symptoms. It can be a single event or part of a recurring pattern, such as in first-episode psychosis. A person experiencing a psychotic episode may also experience depression, anxiety, sleep problems, social withdrawal, lack of motivation, and overall difficulty functioning. When someone experiences this condition, it is referred to as a psychotic episode.<sup>3</sup>

1. National Institute of Mental Health. (n.d.). *What is psychosis?* U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/schizophrenia/raise/what-is-psychosis>
2. "Hallucinations\_MET\_DP808068.jpg" by [Odilon Redon](#) is in the [Public Domain](#)
3. National Institute of Mental Health. (n.d.). *What is psychosis?* U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/schizophrenia/raise/what-is-psychosis>



Figure 11.1 Hallucinations

Psychosis can be a symptom of various mental illnesses, such as schizophrenia, bipolar disorder, severe depression, or severe anxiety, but there are also other potential causes. Sleep deprivation, medical conditions such as hyperglycemia and hyperthyroidism, side effects of some prescription medications, and use of alcohol or other drugs can also cause psychotic symptoms.<sup>4</sup> See Table 11.2a for a side by side comparison of Psychosis and Psychotic Episode

4. National Institute of Mental Health. (n.d.). *What is psychosis?* U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/schizophrenia/raise/what-is-psychosis>

Table 11.2 Comparison of Psychosis and Psychotic Episode<sup>5,6</sup>

Aspect	Psychosis	Psychotic Episode
<b>Definition</b>	A mental condition characterized by a disconnection from reality.	A specific period during which symptoms of psychosis are actively experienced.
<b>Nature</b>	Ongoing or recurring condition; can be chronic or part of another disorder.	Acute, time-limited occurrence of psychotic symptoms.
<b>Symptoms</b>	Delusions, hallucinations, disorganized thinking, impaired insight.	Same symptoms as psychosis, but experienced intensely in a short time frame.
<b>Duration</b>	Can last weeks, months, or be lifelong (depending on cause).	Usually lasts hours to weeks; typically sudden onset.
<b>Cause</b>	May be due to mental illness (e.g., schizophrenia), substance use, trauma, or medical conditions.	Triggered by stress, drug use, sleep deprivation, or illness.
<b>Diagnosis</b>	Used to describe a condition that may be part of disorders like schizophrenia or bipolar disorder.	Not a diagnosis itself; rather, an event within a broader diagnosis.
<b>Treatment Approach</b>	Long-term management with antipsychotics, therapy, and support.	Immediate intervention with medications, hospitalization if needed.
<b>Prognosis</b>	Varies by cause; some recover fully, others may have recurring episodes.	May resolve completely or indicate the beginning of a chronic condition.
<b>Common Misconception</b>	Often confused with violence or multiple personality disorder.	Thought to be a “one-time” event, but may signal a larger issue.

5. Liberman, J. A., & Frist, M. B. (2018). Psychotic disorders. *The New England Journal of Medicine*, 379(3), 270-280 [doi: 10.1056/NEJMr1801490](https://doi.org/10.1056/NEJMr1801490)
6. Fusar-Poli, P., Salazar de Pablo, G., Rajkumar, R. P., López-Díaz, Á., Malhotra, S., Heckers, S., Lawrie, S. M., & Pillmann, F. (2022). Diagnosis, prognosis, and treatment of brief psychotic episodes: A review and research agenda. *Lancet Psychiatry*, 9(1), 72-83. [doi: 10.1016/S2215-0366\(21\)00121-8](https://doi.org/10.1016/S2215-0366(21)00121-8).

## Early Signs of Psychosis

Typically, a person will show changes in their behavior before psychosis develops. Behavioral warning signs for psychosis include the following<sup>7</sup>:

- Drop in grades or worsening job performance
- New trouble thinking clearly or concentrating
- Suspiciousness; paranoid ideas or uneasiness with others
- Withdrawing socially; spending a lot more time alone than usual
- Unusual, bizarre new ideas
- Strange feelings or having no feelings at all
- Decline in self-care or personal hygiene
- Difficulty telling reality from fantasy
- Confused speech or trouble communicating

## Symptoms of a Psychotic Episode

See the following box for signs and symptoms of a psychotic episode according to the *Diagnostic and Statistical Manual of Mental Disorders (DSM-5-TR)* by the American Psychiatric Association.

### **DSM5 Symptoms of a Brief Psychotic Episode<sup>8</sup>**

7. National Institute of Mental Health. (n.d.). *What is psychosis?* U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/schizophrenia/raise/what-is-psychosis>

8. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing

- Delusions
- Hallucinations
- Disorganized speech (e.g., frequent derailment or incoherence)
- Grossly disorganized or catatonic behavior (i.e., a pronounced increase or decrease in the rate and amount of movement; purposeless excessive movement)

A brief psychotic episode is defined by a psychotic event lasting at least one day but less than one month, with an eventual return to previous level of functioning. It does not include a culturally sanctioned response, such as a person's response to the death of a loved one. The disturbance is not better explained by a major depressive disorder or bipolar disorder with psychotic features, another psychotic disorder, or the physiological effects of a substance or a medical condition. The mental health provider may specify if the symptoms are in response to significant stressors (i.e., significant events to anyone experiencing similar circumstances in the individual's culture) or has a peripartum onset (i.e., during pregnancy or within four weeks of delivery).<sup>9</sup>

Review information about delusions, hallucinations, and disorganized speech in the "[Application of the Nursing Process in Mental Health Care](#)" chapter.

## Treatment of Psychosis

Treating psychosis involves a comprehensive, long-term approach tailored to the underlying cause of the condition. For example, if psychosis is caused by a

9. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.



medical condition, side effects of medication, or withdrawal from a substance, it will resolve as these conditions are treated. If psychosis is a symptom of mental illness, treatment typically includes a combination of antipsychotic medicines and psychological therapies. The cornerstone of treatment is antipsychotic medication, which helps reduce or eliminate symptoms like hallucinations and delusions. These medications may be used long-term to prevent relapse, especially in chronic conditions such as schizophrenia. In addition to medication, individuals benefit from psychosocial interventions, including cognitive-behavioral therapy (CBT), which helps manage thought patterns and improve insight. Supportive services like case management, peer support, and vocational rehabilitation can also enhance recovery and quality of life. Family education is often included to promote a supportive environment. Treatment plans are individualized and often involve a multidisciplinary team including psychiatrists, nurses, social workers, and therapists. Read more about the treatment of psychosis associated with schizophrenia in the “[Schizophrenia](#)” section of this chapter.

A psychotic episode is typically treated as a medical emergency, especially if the person is at risk of harming themselves or others. The immediate goal is to stabilize the individual and reduce acute symptoms. Treatment usually begins with antipsychotic medication, often administered in a hospital or closely monitored outpatient setting. Short-term use of sedatives may be considered to manage agitation or insomnia. Once stabilized, care may transition to outpatient follow-up with a psychiatrist or mental health team. Identifying and addressing triggers such as stress, substance use, or underlying medical issues is essential to prevent future episodes. Education and early intervention following a first episode can significantly improve long-term outcomes, and clients are often referred to specialized early psychosis intervention programs when available.

## Delirium

Psychosis caused by medical issues is often referred to as delirium. **Delirium** is a mental state in which the client becomes temporarily confused, disoriented,

and not able to think or remember clearly. It usually starts suddenly and can indicate the onset of a life-threatening medical condition. Delirium resolves as the underlying condition is effectively treated. There are many common causes of delirium, including the following<sup>10 11</sup> :

- Dehydration and electrolyte imbalances
- Dementia
- Hospitalization, especially in intensive care
- Intoxication or withdrawal from alcohol or drugs
- Kidney or liver failure
- Medications, such as sedatives, opioids, anesthesia, antihistamines, anticholinergics, antidepressants, antipsychotics, or anticonvulsants
- Metabolic disorders, such as diabetic ketoacidosis (DKA)
- Serious infections, such as urinary tract infections, pneumonia, and influenza
- Severe pain
- Sleep deprivation

Advanced age makes individuals more vulnerable to delirium, especially when faced with illness or medical stress.<sup>12</sup>

The symptoms of delirium usually start suddenly, over a few hours or a few days, and they often come and go. The most common symptoms are as follows<sup>13</sup> :

10. MedlinePlus (2023). *Delirium*. <https://medlineplus.gov/delirium.html>
11. American Delirium Society. (n.d.). *About delirium*. <https://americandeliriumsociety.org/>
12. Marcantonio, E. R. (2017). Delirium in hospitalized older adults. *The New England Journal of Medicine*, 377(15), 1456-1466. [doi: 10.1056/NEJMc1605501](https://doi.org/10.1056/NEJMc1605501)
13. MedlinePlus (2023). *Delirium*. <https://medlineplus.gov/delirium.html>

- Changes in alertness (usually more alert in the morning, less at night)
- Changing levels of consciousness
- Confusion
- Disorganized thinking or talking in a way that doesn't make sense
- Disrupted sleep patterns or sleepiness
- Emotional changes: anger, agitation, depression, irritability, or overexcitement
- Hallucinations and delusions
- Incontinence
- Memory problems, especially with short-term memory
- Trouble concentrating

Treatment depends on identifying and treating underlying causes. The symptoms of delirium can often be managed with the following interventions<sup>14</sup> :

- Making sure the room is quiet and well-lit
- Having clocks and calendars within view
- Inviting family members to spend time in the room
- Ensuring hearing aids and glasses are worn
- Allowing for uninterrupted sleep when possible
- Getting clients up and out of bed when possible
- Encourage calorie intake and hydration
- Controlling pain with pain relievers (unless the pain medication is causing the psychosis)
- Administering prescribed medications to distressed clients at risk to themselves or to others to calm and settle them, such as haloperidol (However, administer medications with caution because oversedation can worsen delirium.)
- Avoiding the use of restraints

14. MedlinePlus. (2023). *Delirium*. <https://medlineplus.gov/delirium.html>

- ▶ View the [Confusion Assessment Method \(CAM\) PDF](#) commonly used to detect delirium.

- ▶ View the following YouTube video on managing delirium<sup>15</sup>:  
[Managing Delirium Out of Hours.](#)

15. Association for Elderly Medicine Education. (2014, February 6). *Managing delirium out of hours* [Video]. YouTube. All rights reserved.  
<https://youtu.be/1iKe-6lc5b0>

## 11.3 Schizophrenia

There is a spectrum of psychotic disorders, and schizophrenia is one of the disorders on the spectrum. **Schizophrenia** is a serious mental illness that affects how a person thinks, feels, and behaves. It also affects the person's ability to recognize their symptoms as problematic, referred to as a "lack of insight." Continuous signs of the disturbance must be present for at least six months in order for schizophrenia to be diagnosed, and potential medical conditions that could be causing delirium must be ruled out.<sup>1,2</sup>

Schizophrenia is typically diagnosed in the late teen years to the early thirties and tends to emerge earlier in males than females. A diagnosis of schizophrenia often follows the first episode of psychosis when individuals first display symptoms of schizophrenia. Gradual changes in thinking, mood, and social functioning often begin before the first episode of psychosis, usually starting in mid-adolescence. (See "[Early Signs of Psychosis](#)" in the previous section.) Schizophrenia can occur in younger children, but it is rare for it to occur before late adolescence.<sup>3</sup>

1. National Institute of Mental Health. (2024). *Schizophrenia*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/schizophrenia>
2. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.
3. National Institute of Mental Health. (2024). *Schizophrenia*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/schizophrenia>

# Symptoms of Schizophrenia

Symptoms of schizophrenia are classified by three categories, positive, negative, and cognitive:<sup>4</sup>

- **Positive symptoms:** Positive symptoms reflect behaviors or experiences that are added to a person's mental functioning they represent a distortion or excess of normal functions. Positive symptoms include hallucinations, delusions, thought disorders, disorganized speech, and alterations in behaviors. Read more about delusions and hallucinations in the "Thoughts and Perceptions" subsection of the "[Assessment](#)" section of the "Application of the Nursing Process in Mental Health Care" chapter. The most common types of delusions experienced by individuals with schizophrenia are paranoia, persecutory, grandiose, or religious ideas. For example, an individual with persecutory delusions may feel the nursing staff is trying to poison them when they administer medications. People with psychotic symptoms lose a shared sense of reality and experience the world in a distorted way.
- **Negative symptoms:** Negative symptoms refer to the absence or decrease of normal behaviors and emotional responses. They include a loss of motivation, disinterest or lack of enjoyment in daily activities, social withdrawal, difficulty showing emotions, and difficulty functioning normally. Individuals typically experience the following negative symptoms
  - Reduced motivation and difficulty planning, beginning, and sustaining activities (i.e., avolition)
  - Diminished feelings of pleasure in everyday life (i.e., anhedonia)
  - Flat affect (i.e., reduced expression of emotions via facial expression or

4. National Institute of Mental Health. (2024). *Schizophrenia*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/schizophrenia>

voice tone)

- Reduced speaking

- **Cognitive symptoms:** Cognitive symptoms refer to problems in attention, concentration, and memory. For some individuals, the cognitive symptoms of schizophrenia are subtle, but for others, they are more prominent and interfere with activities like following conversations, learning new things, or remembering appointments. Individuals typically experience symptoms such as these<sup>1</sup>:
  - Difficulty processing information to make decisions
  - Problems using information immediately after learning it
  - Trouble focusing or paying attention

The Positive and Negative Syndrome Scale (PANSS) is a widely used tool for measuring the severity of symptoms in clients with schizophrenia. It consists of 30 items, each rated on a scale from 1 (absent) to 7 (extreme), with higher scores indicating more severe symptoms. The scale is divided into three subscales: positive symptoms (7 items), negative symptoms (7 items), and general psychopathology (16 items).

See the following box for signs and symptoms for the diagnosis of schizophrenia according to the *DSM-5-TR*.

### **Symptoms of Schizophrenia<sup>5</sup>**

Schizophrenia is diagnosed when two (or more) of the following characteristics are present for a significant portion of time

5. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

during a one-month period (or less if successfully treated). At least one symptom is delusions, hallucinations, or disorganized speech:

- Delusions
- Hallucinations
- Disorganized speech (i.e., frequent derailment or incoherence)
- Grossly disorganized or catatonic behavior. (**Catatonia** is a state of unresponsiveness.)
- Negative symptoms (i.e., diminished emotional expression or avolition.)

Additionally, for a significant portion of time, the client's level of functioning in one or more areas, such as work, interpersonal relations, or self-care, is significantly below their prior level of functioning. Continuous signs of schizophrenia persist for at least six months (or less if it is successfully treated). Depressive or bipolar disorders with psychotic features must have been previously ruled out, and the disturbance is not attributable to the physiological effects of a substance or other medical condition. The provider may specify if this is the first episode or multiple episodes and if it is an acute episode, in partial remission, or in full remission.<sup>6</sup>

See Figure 11.2<sup>7</sup> for an artwork depiction of the thought disorders associated with schizophrenia.

6. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

7. "[Schizophrenia.png](#)" by William A. Ursprung is in the [Public Domain](#)





Schizophrenia

Figure 11.2 Schizophrenia

## Risk Factors for Schizophrenia

It is believed that several factors contribute to the risk of developing schizophrenia, including genetics, environment, and brain structure and function.<sup>8</sup>

### Genetics

Schizophrenia tends to run in families. Genetic studies strongly suggest that many different genes increase the risk of developing schizophrenia, but that

8. National Institute of Mental Health. (2024). *Schizophrenia*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/schizophrenia>

no single gene causes the disorder by itself. It is not yet possible to use genetic information to predict who will develop schizophrenia.<sup>9</sup>

## Environment

Scientists think that interactions between genetic risk and aspects of an individual's environment play a role in the development of schizophrenia. Environmental factors that may be involved include adverse childhood experiences (ACE) or exposure to viruses or nutritional problems before birth.<sup>10</sup> Other environmental risk factors include urban upbringing, migration, and early life adversities.<sup>11</sup> A systematic review found that cannabis (marijuana) worsens symptoms of psychosis in genetically predisposed individuals and causes more relapses and hospitalizations.<sup>12</sup>

9. National Institute of Mental Health. (2024). *Schizophrenia*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/schizophrenia>
10. National Institute of Mental Health. (2024). *Schizophrenia*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/schizophrenia>
11. Tandon, R., Nasrallah, H., Akbarian, S., Carpenter, W. T., DeLisi, L. E., Gaebel, W., Green, M. F., Gur, R. E., Heckers, S., Kane, J. M., Malaspina, D., Meyer-Lindenberg, A., Murray, R., Owen, M., Smoller, J. W., Yassin, W., & Keshavan, M. (2024). The schizophrenia syndrome, circa 2024: What we know and how that informs its nature. *Schizophrenia Research*. 264, 1-28. [doi: 10.1016/j.schres.2023.11.015](https://doi.org/10.1016/j.schres.2023.11.015).
12. Patel, S., Khan, M. S., & Hamid, P. (2020). The association between cannabis use and schizophrenia: Causative or curative? A systematic review. *Cureus*, 12(7), e9309. <https://doi.org/10.7759/cureus.9309>

# Brain Structure and Function

Scientists think that differences in brain structure, function, and interactions among neurotransmitters may contribute to the development of schizophrenia. For example, differences in the volumes of specific components of the brain, the manner in which regions of the brain are connected and work together, and neurotransmitters, such as dopamine, are found in people with schizophrenia. Differences in brain connections and brain circuits seen in people with schizophrenia may begin developing before birth. Changes to the brain that occur during puberty may trigger psychotic episodes in people who are already vulnerable due to genetics, environmental exposures, or the types of brain differences mentioned previously.<sup>13</sup>



View the following YouTube video on an individual's experience with psychosis<sup>14</sup>: [What is Psychosis?](https://youtu.be/Pgsujx2UQl8)

## Treatment

Early treatment of psychosis increases the chance of a successful remission.<sup>15</sup>

13. National Institute of Mental Health. (2024). *Schizophrenia*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/schizophrenia>

14. Living Well with Schizophrenia. (2019, July 17). *What is psychosis?* [Video]. YouTube. All rights reserved. <https://youtu.be/Pgsujx2UQl8>

15. National Institute of Mental Health. (2024). *Schizophrenia*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/schizophrenia>

Treatments focus on managing symptoms and solving problems related to day-to-day functioning and include antipsychotic medications, psychosocial treatments, family education and support, coordinated specialty care, and assertive community treatment.<sup>16</sup>

## Antipsychotic Medications

Antipsychotic medications reduce the intensity and frequency of psychotic symptoms by inhibiting dopamine receptors. Certain symptoms of psychosis, such as feeling agitated and having hallucinations, resolve within days of starting an antipsychotic medication. Symptoms like delusions usually resolve within a few weeks, but the full effects of the medication may not be seen for up to six weeks.<sup>17</sup>

**First-generation antipsychotics**(also called “typical antipsychotics”) treat positive symptoms of schizophrenia and have several potential adverse effects due to their tight binding to dopamine receptors. Medication is prescribed based on the client’s ability to tolerate the adverse effects. **Second-generation antipsychotics**(also referred to as “atypical antipsychotics”) treat both positive and negative symptoms of schizophrenia. They have fewer adverse effects because they block selective dopamine D2 receptors, as well as serotonin, so they are generally better tolerated than first-generation antipsychotics. Clients respond differently to antipsychotic medications, so it

16. National Institute of Mental Health. (2020). *Schizophrenia*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/schizophrenia>

17. National Institute of Mental Health. (2023). *Mental health medications*. U.S. Department of Health and Human Services. [https://www.nimh.nih.gov/health/topics/mental-health-medications#part\\_2362](https://www.nimh.nih.gov/health/topics/mental-health-medications#part_2362)

may take several trials of different medications to find the one that works best for their symptoms.<sup>18</sup>

See Table 11.3 for a list of common antipsychotic medications. They are usually taken daily in pill or liquid form. Some antipsychotic medications can also be administered as injections twice a month, monthly, every three months, or every six months, which can be more convenient and improve medication adherence.

- ▶ Review information on neuroreceptors affected by antipsychotic medications in the “[Antipsychotics](#)” section of the “Psychotropic Medications” chapter.

Table 11.3 Common Antipsychotic Medications<sup>19, 20, 21</sup>

18. National Institute of Mental Health. (2023). *Mental health medications*. U.S. Department of Health and Human Services. [https://www.nimh.nih.gov/health/topics/mental-health-medications#part\\_2362](https://www.nimh.nih.gov/health/topics/mental-health-medications#part_2362)
19. National Institute of Mental Health. (2023). *Mental health medications*. U.S. Department of Health and Human Services. [https://www.nimh.nih.gov/health/topics/mental-health-medications#part\\_2362](https://www.nimh.nih.gov/health/topics/mental-health-medications#part_2362)
20. Vasan, S., & Padhy, R. K. (2023). *Tardive dyskinesia*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK448207/>
21. Jibson, M. D. (2021). Second-generation antipsychotic medications: Pharmacology, administration, and side effects. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

Medication Class	Mechanism of Action	Adverse Effects
<b>First-Generation (Typical)</b>  Examples:  Chlorpromazine  Haloperidol  Perphenazine  Fluphenazine	Postsynaptic blockade of dopamine receptors in the brain	<ul style="list-style-type: none"> <li>• Extrapyrarnidal side effects (EPS)</li> <li>• Tardive dyskinesia (TD)</li> <li>• Neuroleptic Malignant Syndrome (NMS)</li> </ul>
<b>Second-Generation (Atypical)</b>  Examples:  Risperidone  Olanzapine  Quetiapine  Ziprasidone  Aripiprazole  Paliperidone  Lurasidone  Clozapine	Postsynaptic blockade of dopamine receptors in the brain	<ul style="list-style-type: none"> <li>• Metabolic syndrome</li> <li>• Akathisia</li> <li>• Decreased risk for EPS, TD, and NMS</li> </ul>

## CLOZAPINE

Clients with treatment-resistant schizophrenia may be prescribed clozapine, a specific type of atypical antipsychotic medication. However, people treated with clozapine must undergo routine blood testing to detect a potentially dangerous side effect called **agranulocytosis** (extremely low white blood cell

count). Clozapine also has strong anticholinergic, sedative, cardiac, and hypotensive properties and frequent drug-drug interactions.<sup>22</sup>

## BLACK BOX WARNING

A Black Box Warning states that elderly clients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death.<sup>23</sup>

## ADVERSE EFFECTS

Common side effects of both first- and second-generation antipsychotics include the following<sup>24</sup>:

- Anticholinergic symptoms: dry mouth, constipation, blurred vision, or urinary retention<sup>25</sup>
- Drowsiness
- Dizziness

22. Jibson, M. D. (2021). Second-generation antipsychotic medications: Pharmacology, administration, and side effects. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)
23. United States. Department of Health and Human Services. Office of Inspector General. (2021). *CMS could improve the data it uses to monitor antipsychotic drugs in nursing homes*. <https://collections.nlm.nih.gov/catalog/.nlm:nlmuid-9918384885506676-pdf>
24. National Institute of Mental Health. (2023). *Mental health medications*. U.S. Department of Health and Human Services. [https://www.nimh.nih.gov/health/topics/mental-health-medications#part\\_2362](https://www.nimh.nih.gov/health/topics/mental-health-medications#part_2362)
25. Jibson, M. D. (2021). Second-generation antipsychotic medications: Pharmacology, administration, and side effects. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

- Restlessness
- Weight gain
- Nausea or vomiting
- Low blood pressure

First-generation antipsychotics, also known as neuroleptics or typical antipsychotics, have significant potential to cause extrapyramidal side effects and tardive dyskinesia due to their tight binding to dopamine receptors. The risk for developing these movement disorders is the primary difference between first-generation antipsychotics and second-generation antipsychotics (also known as atypical antipsychotics). In other respects, the two classes of medication have similar side effects and mechanisms of action.<sup>26</sup>

**Extrapyramidal (EPS) side effects** refer to **akathisia** (psychomotor restlessness), rigidity, **bradykinesia** (slowed movement), tremor, and **dystonia** (involuntary contractions of muscles of the extremities, face, neck, abdomen, pelvis, or larynx in either sustained or intermittent patterns that lead to abnormal movements or postures). See Figure 11.3<sup>27</sup> for an image of dystonia.

Acute dystonic reactions affecting the larynx can be a medical emergency requiring intubation and mechanical ventilation. EPS symptoms usually resolve dramatically within 10 to 30 minutes of administration of parenteral anticholinergics such as diphenhydramine and benztropine.<sup>28</sup>

**Tardive dyskinesia (TD)** is a syndrome of movement disorders that can occur

26. Jibson, M. D. (2021). Second-generation antipsychotic medications: Pharmacology, administration, and side effects. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

27. “Dystonia2010.JPG” by [James Heilman, MD](#) is licensed under [CC BY-SA 3.0](#)

28. Lewis, K., & O’Day, C. S. (2023). *Dystonic reactions*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK531466/>



in clients taking first-generation antipsychotics. Hallmark symptoms are smacking and puckering lips, eye blinking, grimacing, and twitching. TD persists for at least one month and can last up to several years despite discontinuation of the medications. Primary treatment of TD includes discontinuation of first-generation antipsychotics and may include the addition of another medication. Second-generation VMAT2 inhibitors such as deutetrabenazine and valbenazine are considered first-line treatment for TD. Clonazepam and ginkgo biloba have also shown good effectiveness for improving symptoms of TD.<sup>29,30</sup>

29. Vasan, S., & Padhy, R. K. (2023). Tardive dyskinesia. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK448207/>
30. Pontone, G. (2020). Treating Tardive Dyskinesia: A Clinical Conundrum and New Approaches. *Drug Induced Disorders: The Clinical Essentials*. Psychopharmacology Institute. <https://psychopharmacologyinstitute.com/section/treating-tardive-dyskinesia-a-clinical-conundrum-and-new-approaches-2557-4810#:~:text=Clonazepam%20probably%20improves%20tard>  
[iv](#)



Figure 11.3 Dystonia




View a supplementary YouTube video on tardive dyskinesia<sup>31</sup>: [Understanding Tardive Dyskinesia](https://youtu.be/gBCiWt-4Cm4).

**Neuroleptic malignant syndrome (NMS)** is a rare but fatal adverse effect that can occur at any time during treatment with antipsychotics. It typically develops over a period of days to weeks and resolves in approximately nine days with treatment. Signs include increased temperature, severe muscular rigidity, confusion, agitation, hyperreflexia, elevation in white blood cell count, elevated creatinine phosphokinase, elevated liver enzymes, myoglobinuria, and acute renal failure. The antipsychotic should be immediately

31. Alliance for Patient Access. (2019, October 14). *Understanding tardive dyskinesia* [Video]. YouTube. All rights reserved. <https://youtu.be/gBCiWt-4Cm4>

discontinued when signs occur. Dantrolene and bromocriptine are typically prescribed for treatment. Nursing interventions include adequate hydration, cooling, and close monitoring of vital signs and serum electrolytes.<sup>32</sup>

 View a supplementary YouTube video<sup>33</sup> on NMS:  
[Neuroleptic Malignant Syndrome in 3 Minutes](https://www.youtube.com/watch?v=s73S6o4wIE0)

Second-generation antipsychotics have a significantly decreased risk of extrapyramidal side effects but are associated with weight gain and the development of metabolic syndrome.<sup>34</sup> **Metabolic syndrome** is a cluster of conditions that occur together, increasing the risk of heart disease, stroke, and type 2 diabetes. Symptoms include increased blood pressure; high blood sugar; excess body fat around the waist (also referred to as having an “apple waistline”); and abnormal cholesterol, triglyceride levels, and high-density lipoprotein (HDL) levels. Weight, glucose levels, and lipid levels should be monitored before treatment is initiated then annually.

32. Chokhawala, K., & Stevens, L. (2023). Antipsychotic medications. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK519503/>

33. TownsendTeaching. 2022, August 28). *Neuroleptic Malignant Syndrome in 3 Minutes* [Video]. YouTube. All rights reserved.  
<https://www.youtube.com/watch?v=s73S6o4wIE0>

34. Chokhawala, K., & Stevens, L. (2023). Antipsychotic medications. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK519503/>



View a supplementary YouTube video<sup>35</sup> on metabolic syndrome: [What is Metabolic Syndrome?](#)



View this PDF comparison: [Adverse Effects of Antipsychotic Medications.](#)

## CLIENT EDUCATION

Clients should be advised to contact their provider if and side effects occur. This includes the development of any involuntary or uncontrollable movements. They should be warned to not suddenly stop taking the medication because abrupt withdrawal can cause dizziness; nausea and vomiting; and uncontrolled movements of the mouth, tongue, or jaw. Clients should be warned to not consume alcohol or other CNS depressants because their ability to operate machinery or drive may be impaired.

## RELAPSE

Some people may experience relapse, meaning their psychosis symptoms come back or get worse. Relapses typically occur when people stop taking their prescribed antipsychotic medication or when they take it sporadically. Some people stop taking prescribed medications because they feel better or they feel that they don't need it anymore, but medication should never be stopped suddenly. After talking with a prescriber, clients can gradually taper

35. Health Link. (2019, December 31). What is metabolic Syndrome? [Video]. YouTube. All rights reserved. [https://youtu.be/fVMvY\\_Lsqzw](https://youtu.be/fVMvY_Lsqzw)

their medications in some situations. However, most people with schizophrenia must stay on an antipsychotic continuously for months, years, or indefinitely for mental wellness.<sup>36</sup>

## Psychosocial Treatments

Cognitive behavioral therapy, behavioral skills training, supported employment, and cognitive remediation interventions are types of psychosocial treatments that can help address the negative and cognitive symptoms of schizophrenia. A combination of these therapies and antipsychotic medication is a common treatment approach for schizophrenia. Psychosocial treatments can help improve an individual's coping skills with the everyday challenges of schizophrenia. Therapies can also help people pursue their life goals, such as attending school, working, or forming relationships. Individuals who participate in regular psychosocial treatments are less likely to relapse or be hospitalized.<sup>37</sup>

## Family Education and Support

Psychosis and schizophrenia can take a heavy toll on a client's family members, significant others, and friends. Educational programs offer instruction about schizophrenia symptoms, treatments, and strategies for assisting their loved one experiencing psychosis and schizophrenia. Increasing their understanding of psychotic symptoms, treatment options, and the course of recovery can lessen their distress, bolster their own coping

36. National Institute of Mental Health. (2023). *Mental health medications*. U.S. Department of Health and Human Services. [https://www.nimh.nih.gov/health/topics/mental-health-medications#part\\_2362](https://www.nimh.nih.gov/health/topics/mental-health-medications#part_2362)

37. National Institute of Mental Health. (2024). *Schizophrenia*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/schizophrenia>

strategies, and empower them to offer effective assistance to their loved one. Family-based services may be provided on an individual basis or through multi-family workshops and support groups.

- ▶ For more information about family-based services in your area, visit the family education and support groups page on the [National Alliance on Mental Illness](#) website.<sup>38</sup>

## Coordinated Specialty Care

**Coordinated specialty care** (CSC) is a general term used to describe recovery-oriented treatment programs for people with first-episode psychosis, an early stage of schizophrenia. A team of health professionals and specialists deliver CSC that includes psychotherapy, medication management, case management, employment and education support, and family education and support. The person with early psychosis and the team work together in a client-centered and family-centered approach to make treatment decisions. Compared to typical care for early psychosis, CSC is more effective at reducing symptoms, improving quality of life, and increasing involvement in work or school.<sup>39</sup>

The goal is to link the individual with a CSC team as soon as possible after

38. National Institute of Mental Health. (2020, May). *Schizophrenia*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/schizophrenia>

39. National Institute of Mental Health. (2024). *Schizophrenia*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/schizophrenia>

psychotic symptoms begin. There are many different programs that are considered CSC in the United States, including (but are not limited to) Comprehensive Community Support (CCS), Community Support Programs (CSP), NAVIGATE, Connection Program, OnTrackNY, Specialized Treatment Early in Psychosis (STEP) program, and Early Assessment and Support Alliance (EASA). Supported Employment/Education (SEE) is an important part of CSC that helps individuals return to work or school because it addresses the client's personal goals. A SEE specialist helps clients develop the skills they need to achieve school and work goals. In addition, the specialist can be a bridge between clients and educators or employers.<sup>40</sup>

Research from the RAISE project (Recovery After an Initial Schizophrenia Episode) has shown that treatments for psychosis work better when they are delivered closer to the time when psychotic symptoms first appear. The goal of the RAISE project is to help decrease the likelihood of future episodes of psychosis, reduce long-term disability, and help people to get their lives back on track so they can pursue their goals.<sup>41</sup>

► Read more about the RAISE project at [RAISE Questions and Answers](#) web page.

With early diagnosis and appropriate treatment, it is possible to recover from psychosis. Many people who receive early treatment never have another psychotic episode. For other people, recovery means the ability to live a

40. National Institute of Mental Health. (n.d.). *What is psychosis?* U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/schizophrenia/raise/what-is-psychosis>

41. National Institute of Mental Health. (n.d.). *Raise researchers.* U.S. Department of Health and Human Services.

fulfilling and productive life, even if psychotic symptoms return at times. However, if untreated, psychotic symptoms can cause disruptions in school and work, strained family relations, and separation from friends. The longer the symptoms go untreated, the greater the risk for developing additional problems. These problems can include abusing substances, having legal trouble, or becoming homeless.<sup>42</sup>

## Assertive Community Treatment

Assertive Community Treatment (ACT) is designed for individuals with schizophrenia who are at risk for repeated hospitalizations or homelessness. Research has demonstrated a prevalence of psychosis as high as 21% among homeless people living on the street.<sup>43</sup> See Figure 11.4<sup>44</sup> for an image of a homeless man. ACT is based on a multidisciplinary team approach, including a medication prescriber, a shared caseload among team members, direct service provision by team members, high frequency of client contact, low client to staff ratios, and outreach to clients in the community. ACT has been shown to reduce hospitalizations and homelessness among individuals with schizophrenia.<sup>45</sup>

- 42. National Institute of Mental Health. (n.d.). *What is psychosis?* U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/schizophrenia/raise/what-is-psychosis>
- 43. Ayano, G., Tesfaw, G., & Shumet, S. (2019). The prevalence of schizophrenia and other psychotic disorders among homeless people: A systematic review and meta-analysis. *BMC Psychiatry*, 19(370). <https://doi.org/10.1186/s12888-019-2361-7>
- 44. “HomelessParis\_7032101.jpg” by Eric Pouhier is licensed under [CC BY-SA 2.5](https://creativecommons.org/licenses/by-sa/2.5/)
- 45. National Institute of Mental Health. (2024). *Schizophrenia*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/schizophrenia>





Figure 11.4 Homelessness

- Read more information about [ACT programs](#) on the Substance Abuse and Mental Health Services Administration (SAMHSA) website.

## 11.4 Applying the Nursing Process to Schizophrenia

The focus of the nursing process for a client with a thought disorder should center on providing client-centered care that addresses the individual's unique needs and experiences. This includes a holistic approach that considers emotional, cognitive, and social factors, while ensuring safety and minimizing risks of self-harm or harm to others. The actions taken by the nurse will differ depending on which phase of the illness the client is in: acute, stabilization, or maintenance.

### Assessment (Recognizing Cues)

Assessment includes several components, such as a mental status examination, psychosocial assessment, cultural assessment, spiritual assessment, screening with validated tools, reviewing laboratory testing results, and including lifespan considerations.

### Mental Status Examination

The nursing assessment adapts to the client's phase of illness, with a focus on acute symptom management in the acute phase, treatment response and stabilization in the stabilization phase, and long-term functional and psychosocial support in the maintenance phase.

During the acute phase, the assessment focuses on identifying and managing positive psychotic symptoms such as hallucinations, delusions, and disorganized speech or behavior. Assessing the severity and impact of these symptoms on the client's functioning, as well as assessing for any immediate safety concerns, including risk of harm to self or others.

In the stabilization phase, the assessment shifts towards monitoring the reduction of acute symptoms and the emergence of any residual symptoms. This phase often involves evaluating the client's response to treatment, adherence to medication, and the presence of any side effects. The focus is also on assessing the patient's cognitive function, mood, and overall ability to engage in daily activities.

During the maintenance phase, the assessment is more comprehensive and long-term, focusing on the management of residual symptoms, particularly negative symptoms such as social withdrawal, apathy, and flat affect. the assessment should include evaluating the client’s support systems and any potential stressors that could trigger a relapse.

Common findings during a mental status examination for a client with schizophrenia who experiencing an acute psychotic episode are described in Table 11.4a. Review information about performing a mental status examination and psychosocial assessment in the “[Application of the Nursing Process in Mental Health Care](#)” chapter. It is also important to assess for suicide risk for clients with psychosis. Review how to assess for suicide risk in the “[Foundational Mental Health Concepts](#)” chapter.

Table 11.4a Common Findings During a Mental Status Examination for Individual With Schizophrenia Experiencing an Acute-Psychotic Episode<sup>1</sup>

1. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

Assessment	Common Findings During Psychotic Episodes  (*Indicates immediately notify health care provider)
Signs of Distress	<b>*May appear internally preoccupied, anxious, fearful, or agitated. Distress may be linked to hallucinations, delusions, or paranoia. Clients may report voices commanding harm or convey fear of being watched or persecuted.</b>
Level of Consciousness and Orientation	<ul style="list-style-type: none"> <li>• Disorientation may occur during acute psychosis or in chronic cases with cognitive decline. Assess for time, place, person, and situation.</li> <li>• Clouded consciousness</li> </ul>
Appearance and General Behavior	<ul style="list-style-type: none"> <li>• Hygiene and grooming may be poor</li> <li>• Clothing may be inappropriate for the weather or mismatched.</li> <li>• Behavior may appear odd, withdrawn, guarded, or disorganized.</li> <li>• May exhibit poor eye contact or social withdrawal.</li> <li>• May demonstrate threatening behavior</li> <li>• May show negativism (a tendency to resist or oppose requests of others)</li> <li>• May have impaired impulse control (which can increase the risk of assault)</li> <li>• May exhibit <b>boundary impairment</b>, an impaired ability to sense where one's influence ends and another person's begins (e.g., the person might walk up to a table and drink out of someone else's glass)</li> </ul>

<b>Speech</b>	<ul style="list-style-type: none"> <li>• May exhibit <b>alogia</b> (reduction or poverty in speech), rapid or pressured speech, or halting speech</li> <li>• May not respond to verbal questions</li> <li>• May not appropriately follow instructions (based on development level)</li> <li>• May demonstrate looseness of associations, clang associations, neologisms, or word salad.</li> <li>• Speech may be tangential or circumstantial.</li> <li>• <b>Neologisms</b> words or phrases created by someone with schizophrenia.</li> </ul>
<b>Motor Activity</b>	<ul style="list-style-type: none"> <li>• May exhibit <b>catatonia</b>, a pronounced increase or decrease in the rate and amount of movement, where excessive movement is purposeless.</li> <li>• May demonstrate psychomotor retardation (pronounced slowness of movement)</li> <li>• May demonstrate psychomotor agitation (running or pacing rapidly in response to internal or external stimuli)</li> <li>• May exhibit <b>echopraxia</b> (mimicking movements of another person)</li> </ul>
<b>Affect and Mood</b>	<ul style="list-style-type: none"> <li>• Affect is often blunted, flat, constricted, labile, or incongruent with stated mood.</li> <li>• Mood may be dysphoric, anxious, or indifferent.</li> <li>• Clients may smile inappropriately or show no emotional response to distressing topics.</li> <li>• Clients may exhibit: <ul style="list-style-type: none"> <li>◦ <b>Anhedonia</b> (reduced ability to experience pleasure)</li> <li>◦ <b>Avolition</b> (reduced motivation or goal-directed behavior)</li> <li>◦ <b>Asociality</b> (decreased desire for social interaction)</li> <li>◦ <b>Apathy</b> (decreased interest in activities that would otherwise be interesting)</li> </ul> </li> </ul>

<b>Thought and Perception</b>	<ul style="list-style-type: none"> <li>• Hallucinations (false sensory perceptions not associated with real external stimuli that can include any of the five senses such as visual, auditory, tactile, gustatory, or olfactory) are a hallmark symptom of schizophrenia.</li> <li>• <b>Paranoia</b> (an irrational fear ranging from being suspicious to thinking someone is trying to kill you) is a common symptom of schizophrenia. <b>*Suicidal or homicidal ideation can result from command hallucinations or defensive actions in response to paranoia</b></li> <li>• Often experience delusions (a fixed, false belief not held by cultural peers and persisting in the face of objective contradictory evidence)</li> <li>• May experience <b>illusions</b>(misperceptions of real stimuli)</li> <li>• Thought process may show disorganization, thought blocking, perseveration, or flight of ideas</li> <li>• May demonstrate <b>loose associations</b> (jumping from one idea to an unrelated idea in the same sentence), <b>clang associations</b>[/pb_glossary, [pb_glossary id="528"]<b>echolalia</b> (pathological repetition of another person's words, or <b>magical thinking</b> (falsely believing that reality can be changed simply by one's thoughts)</li> </ul>
<b>Attitude and Insight</b>	<ul style="list-style-type: none"> <li>• Attitude may be guarded, suspicious, or uncooperative.</li> <li>• Insight is typically impaired; clients often do not recognize their symptoms as part of an illness. They may resist treatment or deny the need for help. <b>Anosognosia</b> is the inability to recognize that one is ill.</li> </ul>

<b>Cognitive Abilities</b>	<ul style="list-style-type: none"> <li>• Memory and attention may be impaired.</li> <li>• Abstract thinking is often poor and concrete thinking is common.</li> <li>• Clients may show difficulty with problem-solving, executive functioning, and reasoning.</li> </ul>
<b>Examiner's Reaction to Client</b>	<ul style="list-style-type: none"> <li>• Frustration</li> <li>• Anxiousness</li> <li>• Countertransference</li> </ul>

When assessing hallucinations, do not imply the perceptions are real. For example, a nurse should ask the client, “What do you hear?” not “What are the voices saying?” It is important to assess for command hallucinations, such as, “Are you hearing a voice that is telling you to do something,” followed by, “Do you believe what you hear is real?” If the answer is “Yes,” the client is at increased risk for acting on the command. Assess when the hallucinations began, their content, and the manner in which the client experiences them (i.e., Are they supportive or distressing? In the background or intrusive?). Ask what makes them worse or better, how the client responds, and what they do to cope with the hallucinations.<sup>2</sup>

When assessing delusions, determine if the client is capable of reality testing (i.e., questioning their thoughts and determining what is real). Ask the client if they believe there is any danger related to the delusion.<sup>3</sup>

2. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

3. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

Countertransference, which refers to the nurse's or health care professional's emotional reactions to a client, can be particularly complex when working with individuals diagnosed with schizophrenia. These reactions can be intense and multifaceted. Understanding and managing countertransference is crucial for effective interventions with this population, as it can significantly impact the nurse-client therapeutic relationship and the client's progress.

## Psychosocial Assessment

As previously discussed in the “[Application of the Nursing Process in Mental Health Care](#)” chapter, a psychosocial assessment obtains additional subjective data that detects risks and identifies treatment opportunities and resources.<sup>4,5</sup>

- Reason for seeking health care (i.e., “chief complaint”)
- Thoughts of self-harm or suicide (both current and historical)
- Cultural assessment
- Spiritual assessment
- Family dynamics
- Current and past medical history
- Current medications
- History of previously diagnosed mental health disorders
- Previous hospitalizations
- Educational background
- Occupational background

4. Clinical Practice Guideline for the Treatment of Posttraumatic Stress Disorder. (2017). *What is cognitive behavioral therapy?* American Psychological Association. <https://www.apa.org/ptsd-guideline/patients-and-families/cognitive-behavioral>

5. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.



- History of exposure to psychological trauma, violence, and domestic abuse
- Substance use (tobacco, alcohol, recreational drugs, misused prescription drugs)
- Family history of mental illness
- Coping mechanisms
- Functional ability/Activities of daily living

After identifying the reason the client is seeking health care, additional focused questions are used to obtain detailed information used to plan care. The mnemonic PQRSTU can be used to ask questions in an organized fashion. See Table 11.4b for a sample PQRST assessment for a client with schizophrenia and sample responses.

Table 11.4b Sample PQRSTU Questions for Assessing Depression

PQRSTU	Sample Questions	Sample Client Response
<b>Provocation/ Palliation</b>	“What tends to trigger or make your symptoms worse? What helps you feel better or more in control?”	“The voices get louder when I’m alone or stressed. Sometimes they calm down if I wear my headphones or talk to someone.”
<b>Quality</b>	“Can you describe what the symptoms feel like or sound like?”	“It feels like people are talking about me behind my back. The voices sound like whispers at first, then they yell. They say I’m being watched.”
<b>Region</b>	“Do these feelings affect your body in any way? Do you feel it in a specific place?”	“I get tense in my chest and shoulders. Sometimes I feel like bugs are crawling under my skin and I try to scratch them off.”
<b>Severity</b>	“On a scale of 0 to 10, how intense or distressing are the symptoms when they are at their worst?”	“It’s a 10 when the voices are yelling. I can’t think straight or focus on anything else. It feels like I’m losing control.”
<b>Timing/ Treatment</b>	“When did the symptoms start? How long do they last when they come on?”	“I started hearing voices when I was in college. They come and go during the day—sometimes for hours. Nighttime is worse.”
<b>Understanding</b>	“What do you think is causing these symptoms? How do you make sense of these feelings?”	“I think the government implanted a chip in the vaccines I received. The voices are part of their surveillance. The doctors tell me I have schizophrenia, but I’m not sure I believe that.”

## SUICIDE AND SELF INJURY SCREENING

Clients with schizophrenia may have hallucinations commanding them to commit suicide or self harm or injure others. The Patient Safety Screener

(PSS-3) is an example of a brief screening tool to detect suicide risk in all client presenting to acute care settings.<sup>6</sup>

**Non-suicidal self-injury (NSSI)** refers to intentional self-inflicted destruction of body tissue without suicidal intention and for purposes not socially sanctioned. Common forms of NSSI include behaviors such as cutting, burning, scratching, and self-hitting. It is considered a maladaptive coping strategy without the desire to die. NSSI is a common finding among adolescents and young adults in psychiatric inpatient settings.<sup>7</sup>

Review information about suicide screening, the Patient Safety Screener, and screening for non-suicidal self-injury (NSSI) in the "[Assessment](#)" section of the Applying the Nursing Process to Mental Health Care" chapter.

## CULTURAL ASSESSMENT

Cultural Formulation Interview (CFI) questions help nurses understand a client's cultural background and how it influences their experience of mental health symptoms, including psychosis.<sup>8</sup> Sample CFI questions focused specifically on understanding depression within a cultural context include the following:

6. Suicide Prevention Resource Center. (n.d.). *The patient safety screener: A brief tool to detect suicide risk*. <https://sprc.org/micro-learning/patientsafetyscreener>
7. Cipriano, A., Cella, S., & Cotrufo, P. (2017). Nonsuicidal self-injury: A systematic review. *Frontiers in Psychology*, 8. <https://www.frontiersin.org/articles/10.3389/fpsyg.2017.01946/full>
8. DeSilva, R., Aggarwall, N. K., & Lewis-Fernandez, R. (2015). The DSM-5 cultural formulation interview and the evolution of cultural assessment in psychiatry. *Psychiatric Times*, 32(6). <https://www.psychiatrictimes.com/view/dsm-5-cultural-formulation-interview-and-evolution-cultural-assessment-psychiatry>

- Cultural Definition of the Problem
  - “How would you describe the problems you are having with your thoughts or experiences?”
  - “Are there any names or terms your family or community use for these experiences?”
  - “How do other people in your culture view experiences like hearing voices or seeing things?”
- Cultural Perceptions of Cause, Context, and Support
  - “What do you think is causing you to hear voices?”
  - “Do you have any cultural or spiritual beliefs that help explain what you’re going through?”
- Cultural Factors Affecting Coping and Help-Seeking
  - “What kinds of things have you done to cope with hearing these voices?”
  - “Are there any traditional remedies, rituals, or religious practices you use to feel better?”
  - “Have you tried to talk to anyone about these experiences, like family members, friends, religious leaders, or traditional healers?”
- Cultural Features of the Nurse–Client Relationship
  - “Is there anything I should know about your background or beliefs that would help me better understand you?”
  - “Do you have any concerns or hesitations you have about seeing a mental health professional?”
  - “What kind of help do you think would work best for you?”

Findings from the cultural formulation interview are used to individualize a client’s plan of care to their preferences, values, beliefs, and goals.

## **SPIRITUAL ASSESSMENT**

The FICA Spiritual History Tool is a widely used assessment model for evaluating a client’s spiritual beliefs and how they may influence health, illness, and coping. FICA© is a mnemonic for the domains of Faith,

Importance, Community, and Address in Care.<sup>9</sup> The data obtained from a FICA assessment can be helpful in understanding how clients with schizophrenia draw on spirituality or religion for support. Addressing a client's spirituality and advocating spiritual care have been shown to improve clients' health and quality of life.<sup>10,11</sup>

Table 11.4c summarizes a sample spiritual assessment questions and sample responses from a client with schizophrenia.

Table 11.4c Sample FICA Spiritual Assessment Questions for Clients with Schizophrenia

9. GW School of Medicine & Health Sciences. (n.d.). *Clinical FICA tool*.  
<https://smhs.gwu.edu/spirituality-health/program/transforming-practice-health-settings/clinical-fica-tool>
10. Pilger, C., Molzahn, A. E., de Oliveira, M. P., & Kusumota, L. (2016). The relationship of the spiritual and religious dimensions with quality of life and health of patients with chronic kidney disease: An integrative literature review. *Nephrology Nursing Journal: Journal of the American Nephrology Nurses' Association*, 43(5), 411–426. <https://pubmed.ncbi.nlm.nih.gov/30550069/>
11. Puchalski, C., Jafari, N., Buller, H., Haythorn, T., Jacobs, C., & Ferrell, B. (2020). Interprofessional spiritual care education curriculum: A milestone toward the provision of spiritual care. *Journal of Palliative Medicine*, 23(6), 777–784.  
<https://doi.org/10.1089/jpm.2019.0375>

Domain	Sample Assessment Question	Sample Client Response
<b>Faith</b>	"Do you consider yourself spiritual or religious? What gives your life meaning?"	"I believe in God. Sometimes I wonder if the voices I hear are from demons?"
<b>Importance</b>	"What importance does your faith or belief have in your life? Has it influenced how you cope with hearing voices?"	When I pray, it helps me feel calmer. But I also get confused when I think God is sending me messages, and it's hard to know what's real."
<b>Community</b>	"Are you part of a spiritual or religious community? Does participation in this community provide support when you're feeling sad or in a low mood?"	"I used to go to church, but I stopped going. I felt like people were judging me for the voices I hear, and I didn't trust them. I want to go back but I don't feel safe there yet."
<b>Address in Care</b>	"How would you like me (or the health care team) to address spiritual issues during your care? Would you like to speak with a chaplain?"	"I'd like someone to help me figure out if what I'm hearing is a symptoms of schizophrenia, messages from God, or voices of demons. Maybe talking to a chaplain would help me figure this out."

Clients with schizophrenia may express religious delusions or spiritual distress. They may express a desire for clarity and connection. Nurses may recognize cues of spiritual distress and offer to connect the client with a chaplain or spiritual care services. Spiritual goals may be included in the nursing care plan if the client finds them valuable.

## FAMILY DYNAMICS

Family dynamics are included in a psychosocial assessment, especially for children, adolescents, and older adults. **Family dynamics** refers to the patterns of interactions among relatives, their roles and relationships, and the various factors that shape their interactions. Because family members rely on each other for emotional, physical, and economic support, they are primary sources of relationship security or stress. Family dynamics and the quality of family relationships can have either a positive or negative impact on an individual's health. For example, secure and supportive family relationships can provide love, advice, and care, whereas stressful family relationships can

be burdened with arguments, unhealthy relationships, and a lack of support.<sup>12</sup> When possible, assess family members and significant others' knowledge of the client's illness and their response. Are they overprotective, frustrated, or anxious? Are they familiar with family support groups, respite, and other community resources?<sup>13</sup>

Unhealthy family dynamics can cause children to experience trauma and stress as they grow up. This type of exposure, known as adverse childhood experiences (ACEs), is linked to an increased risk of developing physical and mental health problems.<sup>14</sup> Review information about adverse childhood experiences (ACEs) in the "[Mental Health and Mental Illness](#)" section of Chapter 1.

## ASSESSMENT OF ACTIVITIES OF DAILY LIVING


Assess the client's ability to perform activities of daily living. Are they getting adequate food, fluid, sleep, and rest? Are they completing daily hygiene tasks and dressing safely for weather conditions? Are they able to control their impulses and make safe decisions?<sup>15</sup> Nursing interventions related to the physiological symptoms of schizophrenia are discussed in the "Implementation" section.

12. Jabbari, B., Schoo, C., & Rouster, A. S. (2023). *Family dynamics*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK560487/>
13. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.
14. Jabbari, B., Schoo, C., & Rouster, A. S. (2023). *Family dynamics*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK560487/>
15. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

## CURRENT MEDICATIONS AND ADVERSE EFFECTS

During the assessment, nurses assess if the client is taking their medications as prescribed, their effectiveness, and if they are experiencing side effects. Are there any barriers to medications or other treatment, such as cost, stigma, or mistrust of health care providers?<sup>16</sup>

Nurses also assess for adverse effects of medications, such as involuntary movements associated with the use of antipsychotic medications (e.g., extrapyramidal side effects or tardive dyskinesia). Clients are routinely assessed for these adverse effects using scales like the Abnormal Involuntary Movement Scale.

View a YouTube video<sup>17</sup> of a nurse performing an  
 assessment using an Abnormal Involuntary Movement  
Scale: [Mental Health AIMS Assessment](https://youtu.be/XuulM7G6T7A)

## Laboratory Testing

The potential laboratory tests and diagnostic procedures for a client with schizophrenia vary depending on whether the client is in the acute phase, stabilization phase, or maintenance phase. During the acute phase, the focus is on identifying any immediate medical issues that could be contributing to the psychotic symptoms and ensuring the client's safety. Ensure the client has had a medical workup for other potential causes of psychosis. For

16. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

17. Dymond Banks. (2021, February 22). Mental Health AIMS Assessment. [Video]. YouTube. All rights reserved. <https://youtu.be/XuulM7G6T7A>



example, dehydration, infection, electrolyte imbalances, abnormal blood glucose level, substance use, or withdrawal from substances can cause psychosis. Concurrent medical disorders are common and should be treated in addition to treating schizophrenia.

In the stabilization phase, the focus shifts to monitoring the client's response to treatment and managing any side effects of medications. During the maintenance phase, the goal is to ensure long-term stability and prevent relapse. If the client is currently taking psychotropic medications, therapeutic drug levels of some types of medications are required. As always, review current information from a medication reference before administering medications.

## Life Span Considerations

Life span considerations influence how the client is assessed, as well as the selection of appropriate nursing interventions. It is important to individualize all interventions to the age and developmental level of the client. Review developmental stages in the “[Application of the Nursing Process in Mental Health Care](#)” chapter.

### CHILDREN AND ADOLESCENTS

Onset of schizophrenia is rare before age 13, but incidence increases in late adolescence and early adulthood. It may be preceded by years of social withdrawal, academic decline, or odd behavior. Children may experience hallucinations (especially auditory), delusions, and disorganized speech/behavior. They may exhibit poor school performance, sleep disturbances, social isolation, or irritability. There is an increased risk of substance use, self-injury, and suicide in adolescents with schizophrenia.

### OLDER ADULTS

New-onset schizophrenia after age 45 is uncommon. Symptoms may be attributed to delirium, dementia, or sensory impairment (e.g., hearing loss).

Polypharmacy and comorbidities complicate diagnosis. In addition to hallucinates, older adults with schizophrenia may include persecutory ideation (e.g., believing a family member is stealing from them or poisoning their food).

## Diagnosis (Analyzing Cues)

Mental health disorders like schizophrenia are diagnosed by mental health providers using the *DSM-5*. Nurses create individualized nursing care plans based on the client's responses to their mental health disorders. See Table 11.4d for a list of common nursing diagnoses and human responses related to schizophrenia.

Table 11.4d Common Nursing Diagnoses Related to Schizophrenia<sup>18 19</sup>

18. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

19. Ackley, B., Ladwig, G., Makic, M. B., Martinez-Kratz, M., & Zanoliti, M. (2020). *Nursing diagnosis handbook: An evidence-based guide to planning care* (12th ed.). Elsevier.

Nursing Diagnosis	Definition	Selected Defining Characteristics
<b>Risk for Suicide</b>	Susceptible to self-inflicted, life-threatening injury.	<ul style="list-style-type: none"> <li>• Reports desire to die</li> <li>• Threats of killing self</li> <li>• Hopelessness</li> <li>• Social isolation</li> </ul>
<b>Ineffective Coping</b>	A pattern of invalid appraisal of stressors, with cognitive and/or behavioral efforts, that fails to manage demands related to well-being.	<ul style="list-style-type: none"> <li>• Alteration in concentration</li> <li>• Alteration in sleep pattern</li> <li>• Inability to meet basic needs</li> <li>• Ineffective coping strategies</li> <li>• Insufficient goal-directed behavior</li> <li>• Risk-taking behavior</li> </ul>

<b>Self-Neglect</b>	A constellation of culturally framed behaviors involving one or more self-care activities in which there is a failure to maintain a socially accepted standard of health and well-being.	<ul style="list-style-type: none"> <li>• Insufficient personal hygiene</li> <li>• Insufficient environmental hygiene</li> <li>• Nonadherence to health activity</li> </ul>
<b>Impaired Communication</b>	Decreased, delayed, or absent ability to receive, process, transmit, and/or use a system of symbols.	<ul style="list-style-type: none"> <li>• Inappropriate verbalizations</li> <li>• Difficulty comprehending communication</li> <li>• Difficulty expressing thoughts verbally</li> </ul>

<b>Imbalanced Nutrition: Less than Body Requirements</b>	<p>Intake of nutrients insufficient to meet metabolic needs.</p>	<ul style="list-style-type: none"> <li>• Food intake less than recommended daily allowance</li> <li>• Insufficient interest in food</li> <li>• Body weight 20% or more below ideal weight range</li> <li>• Read more in the “<a href="#">Nutrition</a>” chapter in <i>Open RN Nursing Fundamentals</i></li> </ul>
<b>Sleep Deprivation</b>	<p>Prolonged periods of time without sustained natural, periodic suspension of relative consciousness that provides rest.</p>	<ul style="list-style-type: none"> <li>• Anxiety</li> <li>• Agitation</li> <li>• Transient paranoia</li> <li>• Intrusive thoughts</li> <li>• Read more in the “<a href="#">Sleep and Rest</a>” chapter of <i>Open RN Nursing Fundamentals</i></li> </ul>

<b>Social Isolation</b>	Aloneness experienced by the individual and perceived as imposed by others and as a negative or threatening state.	<ul style="list-style-type: none"> <li>• Impaired communication patterns</li> <li>• Disturbed thought processes</li> <li>• Delusional thinking</li> </ul>
<b>Hopelessness</b>	Subjective state in which an individual sees limited or no alternatives or personal choices available and is unable to mobilize energy on own behalf.	<ul style="list-style-type: none"> <li>• Long-term stress from mental illness</li> </ul>
<b>Spiritual Distress</b>	A state of suffering related to the impaired ability to experience meaning in life through connections with self, others, the world, or a superior being.	<ul style="list-style-type: none"> <li>• Ineffective coping strategies</li> <li>• Perceived insufficient meaning in life</li> <li>• Hopelessness</li> <li>• Social alienation</li> </ul>

<b>Readiness for Enhanced Hope</b>	A pattern of expectations and desires for mobilizing energy on one's own behalf, which can be strengthened.	<ul style="list-style-type: none"> <li>• Expresses desire to enhance connectedness with others</li> <li>• Expresses desires to enhance sense of meaning in life</li> </ul>
------------------------------------	---	--

## Outcome Identification (Generate Solutions)

Outcomes should be consistent with the recovery model and emphasize hope, resilience, living a full and productive life, and recovery from illness. Expected outcomes are identified based on the client's current phase of their illness: acute, stabilization, or maintenance<sup>20</sup> :

- **Acute:** The overall goal in the acute phase of schizophrenia is client safety and stabilization. An example of an expected outcome is, "The client will consistently be able to label their hallucinations as 'not real' and a symptom of their illness by discharge."<sup>21</sup>
- **Stabilization:** Goals during the stabilization phase focus on understanding the illness and the prescribed treatment plan, as well as controlling and/or coping with symptoms using an optimal medication and psychosocial treatment regimen. Outcomes typically target negative and cognitive symptoms of schizophrenia during this phase because

20. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

21. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

these symptoms respond less well to initial medication treatment than do positive symptoms.<sup>22</sup> An example of an expected outcome during the stabilization phase is, “The client will establish two goal-directed activities by the end of the shift.”

- **Maintenance:** Goals during the maintenance phase focus on maintaining and increasing symptom control and optimal functioning. Factors include treatment adherence, increasing independence, and a satisfactory quality of life.<sup>23</sup> An example of an expected outcome during the maintenance phase is, “The client will identify advantages for taking medications by the end of Week 2.”

## Planning (Generate Solutions)

### Safety

Clients with command hallucinations require close monitoring for suicide, homicide, and other violence risk. Implement interventions to reduce risk of suicide as described in the “[Application of the Nursing Process in Mental Health Care](#)” chapter.

Interpersonal conflict, paranoia, delusions, impaired judgment, limited impulse control, fear, and disagreement with unit rules increase the risk for aggressive behavior.<sup>24</sup> Nursing interventions addressing increased risk for violence to self and others are described in the following box. Read more

22. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

23. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

24. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.



about recognizing signs of crisis and crisis interventions in the “[Stress, Coping, and Crisis Intervention](#)” chapter.

### **Nursing Interventions Addressing Risk for Violence**<sup>25</sup>

- Assess for suicide risk and increase supervision when risk is present. Make rounds at unpredictable intervals and adjust frequency based on risk. Read more about assessing suicide risk in the “[Foundational Mental Health Concepts](#)” chapter and interventions for risk of suicide in the “[Application of the Nursing Process in Mental Health Care](#)” chapter.
- Assess for paranoid thoughts, command hallucinations, impaired impulse control, interpersonal conflict, increasing tension and desperation, and other factors that increase the risk of violence.
- Establish trust and rapport. Engage regularly with the client. Promote communication in a safe manner regarding their concerns that contribute to risk of violence. Engender goodwill and a strong nurse-client relationship.
- Take actions to ensure the client feels safe and secure.
- Teach coping skills to reduce stressors.
- Provide constructive diversion and outlets for physical energy.
- Ensure clients are taking their medications as prescribed. Consider requesting long-acting injectable medications as indicated.

25. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

- If the client targets specific peers or staff, relocate individuals as needed.
- Search client belongings thoroughly on admission and repeat the search whenever circumstances suggest the client may have made or acquired a weapon.
- Use seclusion or restraints when other alternatives have not been successful in keeping the client or others safe. Review ANA guidelines on using restraints in the "[Client Rights](#)" section of the "Legal and Ethical Considerations in Mental Health Care" chapter.

## Acute Phase of Schizophrenia

Hospitalization is indicated during the acute phase of schizophrenia if the client is considered a danger to self (e.g., refuses to eat or is too disorganized to function in the community) or to others (e.g., is behaving in a threatening manner to others).<sup>26</sup> During this phase, planning focuses on selecting interventions that focus on client safety and management of acute symptoms.

During the acute phase of schizophrenia, hospitalization provides safety, structure, and support. As discussed earlier, anosognosia may impair the client's ability to recognize their mental illness. In this case, court-ordered hospitalization may be required.<sup>27</sup> Read more about court-ordered

26. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

27. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

hospitalization in the “[Legal and Ethical Considerations in Mental Health Care](#)” chapter. Nursing interventions focus on providing safety, promoting hygiene and nutrition, improving socialization, encouraging hope and self-esteem, preventing falls, using specific therapeutic techniques, addressing physiological needs, and implementing collaborative interventions.

## Stabilization and Maintenance Phases

During the stabilization and maintenance phases, planning focuses on education, support, and skills training for the client and family. It also addresses how and where these needs can be met within the community. As explained previously in this chapter, relapse prevention efforts are vital. Each relapse can increase residual dysfunction and deterioration and can contribute to despair, hopelessness, and suicide risk. Additionally, recognizing early signs of relapse (e.g., reduced sleep, social withdrawal, and worsening concentration) and implementing intensive treatment are needed to minimize the disruption of the client’s life.<sup>28</sup>

## Implementation (Take Action)

### Acute Phase

#### **PROMOTE HYGIENE AND NUTRITION**

Promote hygiene in clients experiencing psychosis by concisely and explicitly stating expected hygiene tasks. Break tasks into smaller, more manageable tasks and assist when needed. Use visual cues to prompt hygiene tasks, such as putting clean clothes on the bed or clean towels and a toothbrush in the bathroom. Share potential benefits of improved hygiene such as improved

28. Halter, M. (2022). *Vancouver's foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

socialization with others. Reinforce progress in performing hygiene with verbal praise or concrete rewards like additional privileges on the unit.<sup>29</sup>

Clients who are experiencing catatonia require assistance with nutrition, as well as other activities of daily living.

## IMPROVE SOCIALIZATION

Regularly engage with the client. Initially interact briefly about low-anxiety topics like the weather and gradually increase the duration and frequency of interactions as they become more comfortable. Encourage clients to participate in unit activities without pressure, such as “We would like to see you at the morning meeting.” Reinforce the client’s control in their choices, such as, “If you become uncomfortable in the group, you can leave and try again on another day.” Provide positive reinforcement for attempts at socialization, such as, “It was nice to see you in the morning meeting today.”<sup>30</sup>

## ENCOURAGE HOPE AND SELF-ESTEEM

Convey unconditional acceptance, empathy, and support. For example, say, “Sometimes it can feel very discouraging when experiencing a mental health disorder. I am wondering how you are feeling?” If the client cannot identify their feelings, suggest words that may apply, such as, “Sometimes it is hard to say what you are feeling. Do you feel sad, frustrated, or anxious?” Validate the client’s feelings and assure them they are not alone. Help the client identify their positive traits or previous accomplishments. Suggest coping strategies such as journaling and attending a support group. Teach stress management

29. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

30. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

techniques and coping strategies as outlined in the “[Stress, Coping, and Crisis Intervention](#)” chapter.

## PREVENT FALLS

Fall risk may be increased due to orthostatic hypotension, impaired balance, bradykinesia, or other movement disorders. Assess the client’s gait and for orthostatic hypotension. Teach the client to slowly change position from lying to sitting to standing and encourage the use of handrails or seeking assistance when feeling unsteady. Implement additional fall precautions as needed according to agency policy.

- ▶ Read more information about fall precautions in the “[Safety](#)” chapter of *Open RN Nursing Fundamentals*.

## USE THERAPEUTIC TECHNIQUES FOR COGNITIVE IMPAIRMENTS, DELUSIONS, AND HALLUCINATIONS

Recall that clients with schizophrenia may have memory and attention impairments. Repetition with visual and verbal reminders is helpful to promote task completion. Additionally, short but frequent interactions may be less stimulating to the client and better tolerated.<sup>31</sup> Additional techniques for helping clients who are experiencing delusions and hallucinations are described below.

### Helping Clients Who Are Experiencing Delusions

Delusions feel very real to the client and can be frightening. Nurses should

31. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

acknowledge and accept the client's experience and feelings resulting from the delusion while conveying empathy. They can provide reassurance regarding their intentions to help the client feel safer.

Avoid questioning the delusion. Until the client's ability to test reality improves, trying to prove the delusion is incorrect can intensify it and cause the client to view the staff as people who cannot be trusted. Instead, focus on the fear and what would help the client feel safer. For example, if a client states, "The doctor is here. He wants to kill me," the nurse could respond, "Yes, the doctor is here and wants to see you. They talk with all of the clients about their treatment. Would you feel more comfortable if I stayed with you during your meeting with the doctor?" Focusing of events in the present keeps the client focused on reality and helps them distinguish what is real.<sup>32</sup>

If a client is exhibiting paranoia and is highly suspicious, it is helpful to maintain consistent staff assignments. Staff should avoid laughing, whispering, or talking quietly where the client can see these actions but cannot hear what is being said. Staff should ask permission before touching the client, such as before taking their blood pressure.

- ▶ Read additional strategies for working with clients with delusions from the British Columbia Schizophrenia Society: [Steps for Working With Delusions](#).

## Helping Clients Who Are Experiencing Hallucinations

Hallucinations feel very real to the person experiencing them and can be distracting during their interactions with others. Hallucinations can be

32. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

supportive or terrifying, faint or loud, or episodic or constant. For example, listen to simulations of auditory hallucinations in the following box. The nurse should focus on understanding the client's experiences and responses and convey empathy. Command hallucinations, suicidal ideation, or homicidal ideation requires safety measures as previously discussed in the "Provide Safety" subsection.

### Simulations of Auditory Hallucinations

British Columbia Schizophrenia Society created music tracks simulating what auditory hallucinations can feel like to clients. Similar to auditory hallucinations experienced by people living with schizophrenia, when people listen to these songs, they hear voices that can be frightening. Listen to these simulations on YouTube with discretion because some people can find them disturbing:

▶◀ [Track 05: Mark Pelli - Everything \(Songs of schizophrenia mix\)](https://youtu.be/pN-f6AEDNxY) <sup>33</sup>

▶◀ [Track 06: Cassandra Vasik - Sadly mistaken \(Songs of schizophrenia mix\)](https://youtu.be/HCewO3BL1qA) <sup>34</sup>

When working with a client who has a history of hallucinations, watch for hallucination indicators, such as eyes tracking an

33. BC Schizophrenia. (2019, May 6). *Track 05: Mark Pelli – Everything (Songs of schizophrenia mix)* [Video]. YouTube. All rights reserved.

<https://youtu.be/pN-f6AEDNxY>

34. BC Schizophrenia. (2019, May 6). *Track 06: Cassandra Vasik – Sadly mistaken (Songs of schizophrenia mix)* [Video]. YouTube. All rights reserved. <https://youtu.be/HCewO3BL1qA>

unheard speaker, muttering or talking to oneself, appearing distracted, suddenly stopping a conversation as if interrupted, or intently watching a vacant area of the room. Ask about the content of the hallucinations and if they are experiencing command hallucinations. Assess how the client is reacting to the hallucinations, especially if they are exhibiting anxiety, fear, or distress.<sup>35</sup>

Avoid referring to the hallucinations as if they were real to promote reality testing. For example, do not ask, “What are the voices saying to you,” but instead ask, “You look as though you are hearing something. What do you hear?” Do not try to convince the client the hallucinations are not real, but instead offer your perception and convey empathy. For example, “I don’t hear angry voices that you hear, but that must be very frightening for you.” Address any underlying emotion, need, or theme indicated by the hallucination.<sup>36</sup>

Focus on reality-based activities in the “here and now,” such as a conversation or simple project. Promote and guide reality testing. For example, guide the client to look around the room and see if others are frightened; if they are not, encourage them

35. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

36. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.



to consider what they are experiencing are hallucinations. Teach the client to compare their perceptions to trusted others.<sup>37</sup>

See the information in the following box for teaching clients how to manage their hallucinations.

### **Client Education: Teaching Clients How to Manage Hallucinations<sup>38</sup>**

- Manage stress and stimulation.
  - Avoid overly loud or stressful places or activities.
  - Avoid negative or overly critical people and seek out supportive people.
  - Use assertive communication skills so you can tell others “No” if they pressure or upset you.
  - When stressed, focus on your breathing and slow it down. Inhale slowly through your nose as you count from one to four, hold your breath, and then exhale slowly through your mouth.
  - Refer to other stress management and coping strategies in the “[Stress, Coping, and Crisis Intervention](#)” chapter.

<sup>37</sup>. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

<sup>38</sup>. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

- Use other sounds to compete with the hallucinations, such as talking with other people, listening to music or TV, reading aloud, singing, whistling, or humming.
- Determine what is real and unreal by looking at others. Do they seem to be hearing or seeing what you are? Ask trusted others if they are experiencing the same things you are. If the answers to these questions are “No,” then although it feels real, it is not likely real and can be ignored.
- Engage in activities that can take your mind off the hallucinations, such as walking, taking a relaxing bath or shower, or going to a place you find enjoyable where others are present, such as a coffee shop, mall, or library.
- Talk out loud (or silently to yourself if others are nearby) and tell the voices or thoughts to go away. Tell yourself the voices or thoughts are a symptom and not real. Tell yourself that no matter what you hear, you are safe and can ignore what you hear.
- Seek contact with others. Visit a trusted friend or family member. Call a help line or go to a drop-in center. Visit a public place where you feel comfortable.
- Develop a plan with your provider for how to cope with hallucinations. Additional medications may be prescribed to use as needed.

## CATEGORIZING NURSING INTERVENTIONS ACCORDING TO THE APNA STANDARD OF IMPLEMENTATION

Interventions for clients experiencing psychosis previously discussed in this chapter can also be categorized by the standard of *Implementation* by the American Psychiatric Nursing Association (APNA). Read more about this standard in the [“Application of the Nursing Process in Mental Health Care”](#)

chapter. See Table 11.4e for categorization of nursing interventions by this standard.

Table 11.4e Nursing Interventions for Clients with Psychosis Based on the Categories of the APNA Implementation Standard<sup>39</sup>

39. American Psychiatric Association. (2013). *Desk reference to the diagnostic criteria from DSM-5*.

Categories of Interventions Based on the APNA Standard of Implementation	What the nurse will do..	Rationale
Coordination of Care	<ul style="list-style-type: none"> <li>• Maintain safety by implementing safety precautions as needed to prevent self-harm, suicide, or homicide risks.</li> <li>• Ensure consistency of behavioral expectations among all staff on the unit by including expectations in the nursing care plan.</li> <li>• Collaborate with psychiatry, social work, case managers, and family to ensure medication management, housing stability, and follow-up care.</li> <li>• Plan for quality of life, independence, and optimal recovery by referring to local resources and support groups in the community on discharge.</li> </ul>	<p>Schizophrenia often requires long-term, interdisciplinary management due to chronic symptoms and functional impairments. Coordinated care improves adherence and reduces rehospitalization.</p> <p>The client may exhibit high risk for impulsive behaviors that could pose a risk of harm to self/ others. They may experience altered thought processes with poor insight and judgment.</p> <p>Consistent expectations provide a feeling of structure and safety.</p> <p>The nurse coordinates care delivery during inpatient care, and assists in making referrals for optimal recovery after discharge.</p>

<b>Health Teaching</b>	<ul style="list-style-type: none"> <li>• Educate the client and family about schizophrenia, early warning signs of relapse, medication side effects, and self-management strategies. Use simple, concrete language and repetition.</li> <li>• Include teaching topics on stress management, coping strategies, and management of delusions and hallucinations.</li> <li>• Screen for tobacco, alcohol, or drug use. Provide education on interactions between substances and antipsychotics. Offer motivational interviewing or referrals to substance use programs.</li> </ul>	<p>Many clients with schizophrenia have cognitive deficits or poor insight regarding their condition. Psychoeducation enhances adherence, decreases stigma, and supports early intervention. Nurses encourage resilience by promoting adaptive coping strategies.</p> <p>Many individuals with schizophrenia use substances to self-medicate symptoms, worsening outcomes. Integrated care supports recovery and medication efficacy.</p>
------------------------	---	---

<b>Pharmacological, Biological, and Integrative Therapies</b>	<ul style="list-style-type: none"> <li>• Administer antipsychotics as prescribed (monitoring for EPS, metabolic syndrome). Support the use of long-acting injectables (LAIs) if adherence is an issue.</li> <li>• Deliver client education about antipsychotics and expected time frames for improvement.</li> <li>• Open all medications in front of the client.</li> <li>• Observe for and promptly report symptoms of potential adverse effects of first-generation antipsychotics such as tardive dyskinesia (TD) and extrapyramidal side effects (EPS).</li> <li>• Incorporate relaxation or mindfulness practices.</li> </ul>	<p>Medications are the cornerstone of symptom management. Monitoring improves safety, and integrative approaches support overall wellness and self-regulation. Improving the client's understanding of their medications and potential side effects can increase medication compliance.</p> <p>Opening all medications in front of the client may decrease paranoia.</p> <p>Clients experiencing TD or new EPS symptoms should discontinue first-generation antipsychotics and start second-generation antipsychotics per provider order. Medications to treat symptoms may be required.</p>
---	---	--

<b>Milieu Therapy</b>	<ul style="list-style-type: none"> <li>• Maintain a structured, low-stimulation environment with consistent routines. Use redirection when clients become agitated. Ensure safety through observation and early intervention when behaviors escalate. The client may require a private room.</li> <li>• Promote physical exercise to redirect aggressive behavior.</li> <li>• During acute psychosis with agitation, use prescribed medications, seclusion, or restraint to minimize physical harm.</li> <li>• Encourage participation in group therapy addressing social skills, personal grooming, mindfulness, and stress management. Avoid competitive activities or games if the client is agitated.</li> </ul>	<p>A therapeutic milieu reduces stress and promotes safety. Clients with schizophrenia often benefit from clear expectations, boundaries, and minimal environmental confusion. Reducing stimuli may prevent escalation of anxiety and agitation.</p> <p>Physical exercise can decrease tension and provide focus.</p> <p>During acute psychosis, the nurse's priority is to protect the client and others from harm.</p> <p>Group therapy can encourage effective coping skills and socialization. Structured activities provide security and focus. However, avoid competitive activities because they may be too stimulating and can cause escalation of anxiety and agitation.</p>
-----------------------	--	---

<b>Therapeutic Relationship and Counseling</b>	<ul style="list-style-type: none"> <li>• Build trust through consistent, nonjudgmental, and empathic communication. Avoid challenging delusions directly; instead, explore feelings and promote reality orientation gently. Support involvement in therapy (CBT, social skills training).</li> <li>• Use a firm and calm approach with short and concise statements. For example, “John, come with me. Eat this sandwich.”</li> <li>• Identify expectations in simple, concrete terms with consequences. For example, “John, do not yell at or hit Peter. If you cannot control yourself, the seclusion room will help you feel less out of control and prevent harm to yourself and others.”</li> <li>• Acknowledge feelings associated with delusions and hallucinations and convey empathy. Encourage and guide reality testing based on client status.</li> <li>• Redirect excessive energy</li> </ul>	<p>Trust is critical with clients who may be paranoid or withdrawn. A strong therapeutic alliance supports engagement, emotional expression, and adaptive functioning.</p> <p>Clear expectations help the client experience outside controls and understand reasons for medication, seclusion, or restraints if they are not able to control their behaviors.</p> <p>Acknowledging emotion and conveying empathy build trust and a strong nurse-client relationship. Reality testing helps clients manage their delusions and hallucinations.</p> <p>Clients may be impulsive and hyperv verbal and interrupt, blame, ridicule, or manipulate others so setting limits and personal boundaries is vital.</p>
--	--	--



	<p>into appropriate and constructive channels.</p> <ul style="list-style-type: none"><li>• Set limits with personal boundaries.</li></ul>	
--	---	--

**NURSING INTERVENTIONS FOR PHYSIOLOGICAL SIGNS OF SCHIZOPHRENIA**

While schizophrenia is a psychiatric condition, it frequently presents with or leads to physiological health concerns due to self-neglect, poor insight, medication side effects, and lifestyle disruptions. Nursing interventions target common physiological signs of schizophrenia and associated self-care deficits as described in Table 11.4f.

Table 11.4f Nursing Interventions Targeting Physiological Signs of Schizophrenia<sup>40</sup>

40. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

Problem/Intervention	Rationale
<b>Nutrition</b> <ul style="list-style-type: none"> <li>• Monitor weight and regularly.</li> <li>• Encourage healthy eating habits; provide structured meal times.</li> <li>• Educate about antipsychotic-induced weight gain and provide referrals to dietitians if needed.</li> <li>• Involve client in menu planning if possible.</li> </ul>	<p>Second generation antipsychotic medications increase the risk for metabolic syndrome (e.g., weight gain, hyperlipidemia, insulin resistance). Nutritional support reduces health risks and enhances well-being.</p>
<b>Sleep</b> <ul style="list-style-type: none"> <li>• Assess sleep hygiene and patterns.</li> <li>• Encourage routine sleep–wake cycles and a bedtime routine.</li> <li>• Limit caffeine and other stimulants.</li> <li>• Provide a calm, quiet sleep environment.</li> </ul>	<p>Clients often experience sleep disturbances due to psychosis or medication side effects. Adequate sleep supports emotional regulation and reduces relapse risk.</p>
<b>Elimination</b> <ul style="list-style-type: none"> <li>• Monitor for anticholinergic side effects (e.g., constipation, urinary retention).</li> <li>• Encourage fluid and fiber intake.</li> <li>• Provide toileting reminders if needed.</li> <li>• Consult provider for stool softeners or laxatives as indicated.</li> </ul>	<p>Antipsychotic medications can cause constipation and urinary retention, which may go unreported in clients with disorganized thinking. Prevention of constipation avoids additional complications such as fecal impaction.</p>

<p><b>Activity/Mobility</b></p> <ul style="list-style-type: none"> <li>• <b>Encourage daily walking or light exercise.</b></li> <li>• <b>Monitor for extrapyramidal symptoms (EPS), akathisia, or sedation.</b></li> <li>• <b>Engage in structured physical activities or group movement therapy.</b></li> </ul>	<p>Medications may cause motor side effects, and sedentary behavior increases cardiovascular risk. Physical activity improves physical and mental health.</p>
<p><b>Self-Care Deficits</b></p> <ul style="list-style-type: none"> <li>• <b>Assess ability to perform ADLs (e.g., bathing, grooming, dressing).</b></li> <li>• <b>Offer verbal prompts, encouragement, and step-by-step support. When appropriate, give step-by-step reminders, such as “Wash the right side of your face and now your left.”</b></li> <li>• <b>Establish a simple daily hygiene routine.</b></li> <li>• <b>Reinforce successes positively.</b></li> </ul>	<p>Negative symptoms (e.g., avolition, anhedonia) can result in neglect of hygiene, increasing the risk of infection and social isolation. Structured support helps restore dignity and function.</p>

## COMMUNICATION TIPS FOR CLIENTS WITH SCHIZOPHRENIA

Helpful communication techniques for clients with schizophrenia are described in the following box.

### Communication Tips: Schizophrenia

- Use short, simple, and clear sentences.

- **Rationale:** Clients with schizophrenia may have difficulty processing complex information due to thought disorganization or cognitive deficits. Clear, concise language enhances understanding.
- Speak calmly and slowly.
  - **Rationale:** A calm, non-threatening tone reduces anxiety and helps de-escalate agitation. It also models emotional regulation and helps maintain a therapeutic environment.
- Avoid arguing or challenging delusions.
  - **Rationale:** Confronting delusions directly can increase defensiveness and mistrust. Instead, acknowledge the client's feelings without reinforcing the delusion (e.g., "That sounds frightening for you.") Addressing the underlying fear, anxiety, or confusion allows for therapeutic rapport and emotional support, even if the content of their belief is not reality-based.
- Use reality-based statements gently and consistently.
  - **Rationale:** While avoiding direct confrontation, it is still important to reinforce reality (e.g., "I don't see anyone else in the room, but I understand it feels real to you.")
- Limit environmental stimuli during conversations.
  - **Rationale:** Clients may be easily overwhelmed

or distracted. A quiet, low-stimulation setting helps them focus and decreases internal and external confusion.

- Allow extra time for responses.
  - **Rationale:** Thought blocking, slowed cognition, or preoccupation with internal stimuli may delay verbal responses. Patience promotes a respectful, supportive interaction.
- Use the client's name and establish a consistent routine.
  - **Rationale:** Personalizing communication and maintaining predictable interactions increase trust and reduce paranoia or confusion.
- Be consistent and honest in all interactions.
  - **Rationale:** Clients may be suspicious or mistrustful. Consistency in messaging, behavior, and tone builds therapeutic rapport and emotional safety.

## Stabilization Phase

During the stabilization phase of schizophrenia, care is focused on ongoing medication therapy, education, and CBT therapy.<sup>41</sup>

41. Keepers, G. A., Fochtman, L. J., Anzia, J. M., Benjamin, S., Lyness, J. M., Mojtabai, R., Servis, M., Walaszek, A., Buckley, P., Lenzenweger, M. F., Young, A. S., Degenhardt, A., & Hong, S. H. (2020). The American Psychiatric Association

- **Medication Management:** Continue to monitor the effectiveness and side effects of antipsychotic medications. Adjust dosages as needed to minimize side effects and ensure therapeutic efficacy.
- **Psychoeducation:** Educate the client and their family about the illness, treatment options, and the importance of medication adherence.
- **Therapeutic Interventions:** Introduce cognitive-behavioral therapy for psychosis (CBTp) to help the client develop coping strategies and reduce the impact of psychotic symptoms.

## Maintenance Phase

Effective long-term management of individuals with schizophrenia requires a comprehensive, multidisciplinary approach that extends beyond symptom control.

- **Ongoing Monitoring:** Regularly assess the client's mental status, medication adherence, and side effects. Monitor for signs of relapse and intervene early to prevent full-blown episodes.
- **Supportive Services:** Provide supported employment services and assertive community treatment to help the client maintain social and occupational functioning.
- **Lifestyle and Wellness:** Encourage healthy lifestyle choices, including regular exercise, a balanced diet, and smoking cessation. Monitor metabolic parameters, as antipsychotic medications can increase the risk of metabolic syndrome. Ongoing monitoring of the client's functional status, social integration, and quality of life.

## Evaluation (Evaluate Outcomes)

A client's progress is continually assessed using their individualized SMART

practice guideline for the treatment of patients with schizophrenia. *American Journal of Psychiatry*, 177(9), 868-872. [doi: 10.1176/appi.ajp.2020.177901](https://doi.org/10.1176/appi.ajp.2020.177901)

outcomes and current status. Full recovery can take months. By setting small goals, it is easier to identify and recognize progress that may occur in small increments.<sup>42</sup>

## Acute Phase

During the acute phase of schizophrenia treatment, the primary focus is on symptom severity, safety, and the client's immediate response to intervention. Clinicians assess changes in the Positive and Negative Syndrome Scale (PANSS) scores to determine whether there has been a reduction in the frequency and intensity of hallucinations, delusions, or other psychotic features. Monitoring for safety is also crucial—evaluating any shifts in the client's risk of harm to self or others, and observing improvements in their ability to follow safety protocols within the treatment setting. In addition, the client's immediate response to antipsychotic medications is closely observed. Providers track how quickly the symptoms respond to treatment and whether any side effects emerge that could impact further care decisions.<sup>43</sup>

## Stabilization Phase

In the stabilization phase, ongoing symptom monitoring helps ensure the client's progress continues. Updated PANSS scores are compared to both the baseline and acute-phase scores to evaluate trends. Clinicians identify any residual symptoms that persist and outline strategies for their management, such as medication adjustments or therapeutic interventions. Medication adherence becomes a central concern—providers assess whether the client is

42. Halter, M. (2022). *Vancouver's foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

43. Zhang, Y., Liang, W., Gui, J. (2025). Evaluation of the impact of refined nursing care on schizophrenia patients. *Medicine*, 104(3):e40848. [doi:10.1097/MD.00000000000040848](https://doi.org/10.1097/MD.00000000000040848).



consistently following the prescribed regimen and address any side effects that may compromise adherence. This phase is key in solidifying gains made during acute care and building a foundation for long-term stability.<sup>44</sup>


## Maintenance Phase

The maintenance phase focuses on sustaining symptom remission and promoting recovery. Long-term tracking of PANSS scores helps clinicians evaluate symptom stability over time and detect early signs of relapse. Intervention strategies are adjusted accordingly to prevent deterioration. Equally important is evaluating the client's quality of life using standardized tools like the Heinrichs-Carpenter Quality of Life Scale (QLS). Providers assess improvements in the client's ability to engage in meaningful social relationships and maintain occupational or educational roles. Preventing relapse remains a central goal during this phase; clinicians monitor medication adherence rates and identify any new stressors or triggers that could threaten stability. Preventive measures, such as psychoeducation, structured routines, and early intervention strategies, are implemented to support ongoing recovery and well-being.<sup>45</sup>

44. Zhang, Y., Liang, W., Gui, J. (2025). Evaluation of the impact of refined nursing care on schizophrenia patients. *Medicine*, 104(3):e40848. [doi: 10.1097/MD.00000000000040848](https://doi.org/10.1097/MD.00000000000040848).


45. Zhang, Y., Liang, W., Gui, J. (2025). Evaluation of the impact of refined nursing care on schizophrenia patients. *Medicine*, 104(3):e40848. [doi: 10.1097/MD.00000000000040848](https://doi.org/10.1097/MD.00000000000040848).

## 11.5 Spotlight Application

 Elyn Saks, a professor of law, psychology, and psychiatry at USC, shares her experience of living with schizophrenia in a 15-minute TEDGlobal video<sup>1</sup>: “[Elyn Saks: A Tale of Mental Illness– From the Inside.](https://www.ted.com/talks/elyn_saks_seeing_mental_illness)”

### Reflective Questions:

1. What strikes you most about Dr. Sak’s journey with schizophrenia?
2. What are three takeaways points from Dr. Sak’s presentation?

 View the following YouTube video on “[What Living Well With Schizophrenia Means to Me](https://youtu.be/VUhw64iEIQw)”<sup>2</sup>

### Reflective Questions:

1. TEDGlobal. (2012). *A Tale of Mental Illness — From the Inside*. [Video]. TED. All rights reserved. [https://www.ted.com/talks/elyn\\_saks\\_seeing\\_mental\\_illness](https://www.ted.com/talks/elyn_saks_seeing_mental_illness)
2. Living Well with Schizophrenia. (2020, March 7). *What living well with schizophrenia means to me* [Video]. YouTube. All rights reserved. <https://youtu.be/VUhw64iEIQw>

1. How is Lauren's journey with schizophrenia different from Dr. Sak's?
2. What are three things you learned from Lauren's description of her journey with schizophrenia?

## 11.6 Learning Activities



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=536#h5p-38>

1



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=536#h5p-39>

2



*An interactive H5P element has been excluded from this version of the text. You can view it*

1. “MH Psychosis & Schizophrenia Glossary Cards” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
2. “MH Psychosis & Schizophrenia Drag and Drop” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

online here:

<https://wtcs.pressbooks.pub/nursingmhcc/?p=536#h5p-40>

3

## Case Study

**Situation:** Parker Allen, a 28-year-old male, was admitted to a locked behavioral unit in the hospital today for paranoia, psychosis, and violent behavior after he became violent with his landlord. He threw a table at the landlord when she came to his apartment to address a noise complaint by another tenant. She then called the police, who brought him to the emergency room.

**Background:** Parker has a three-year history of schizophrenia. He has a Master's degree in Engineering, and works as a software developer. Parker was diagnosed with schizophrenia shortly after he completed graduate school when he became very isolated and paranoid of those around him. His first psychiatric hospitalization occurred about one year after his diagnosis with schizophrenia when he experienced auditory hallucinations and paranoid delusions. He has tried various medications, but reports he doesn't like the way they make him feel and they make it difficult for him to concentrate on his work. He currently

3. "MH Psychosis & Schizophrenia Question Set 1" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

prescribed medications include olanzapine 10 mg and venlafaxine extended release (XR) 75 mg daily, but he states he recently stopped taking his medications.

**Assessment:** Since arriving in the behavioral health unit, Parker has been withdrawn. He states, “the nurses can see my thoughts,” and explains that is why he has been staying in his room and covering his head with a blanket.

Sometimes Parker answers questions, but does so inappropriately. He tells the nurse, “I’m pretty sure you are all here to kill me, and you’re starting with the hospital food. I know how this is going to go down.” The nursing staff can only speak with Parker in short bits because he either refuses to talk or is distracted easily. He has a disheveled, dirty appearance and has refused to shower since arriving on the unit. His hair is disheveled and his beard is in need of a trim.

The nursing assistant reports that he refused to eat his dinner when he arrived on the unit, but did drink a few sips of water from a sealed water bottle. His vital signs are: Temperature, 37.0°C; Heart rate, 76 beats/min; Respiratory rate, 18 breaths/min; Blood pressure, 128/79 mmHg; and SpO<sub>2</sub>, 95%.

When the nurse enters Parker’s room, Parker is sitting on his bed, wrapped in a blanket with most of it covering his face, facing away from the nurse and looking towards the window. As he is greeted, Parker makes no movements and does not turn to the nurse’s voice. He is mumbling quietly to himself. The nurse walks around to face Parker, and greets him again. He responds quietly, but does not make eye contact and answers with apprehension.

He appears very guarded and suspicious, keeping his arms wrapped around himself with the blanket. Beneath the blanket,

the nurse can see that he is dressed in layers of clothing and has body odor.

The nurse asks Parker if he has having suicidal thoughts. For most questions the nurse asks, Parker repeats the question back to the nurse. He rarely answers the nurse's questions and appears distracted. Parker becomes agitated when asked about hallucinations and states that he is hearing voices that tell him that the food he has been given is poisoned. He reports the voices have been constant since his arrival in the hospital, but at home he could get the voices to go away by listening to music in his headphones.

The nurse notices that Parker makes bizarre statements like, "The bammerhoff got me," and speaks in tangential language. The nurse recalls from previous documentation in the electronic medical record that Parker's landlord told the police that several other neighboring tenants in the apartment building had complained that there was loud, strange-sounding music and noises coming from his apartment into late hours of the night.

Parker does not tolerate the assessment for very long. After 15 minutes, he refuses to answer any more questions, stating, "You better find the cook. They are poisoning all of us." At that time, he lays down in his bed and pulls the covers over his head. The nurse informs Parker that they will return soon with a snack for him and thanks him for his time. He does not respond but continues to mumble in a low speech under the covers in his bed.

**Reflective Questions:**

1. What CUES do you recognize as important for planning Parker's care?
2. What is your hypothesis for Parker based on this information?

3. Write a SMART goal for Parker based on his priority nursing problem at this time.
4. What are your priority nursing interventions for Parker?
5. How will you evaluate if your interventions have been effective?
6. What symptoms require continued monitoring?

- ▶ Test your clinical judgment with a NCLEX Next Generation-style question: [Chapter 11, Assignment 1](#)<sup>4</sup>



- ▶ Test your clinical judgment with a NCLEX Next Generation-style case study: [Chapter 11, Case Study 1](#)<sup>5</sup>



4. "MH Psychosis & Schizophrenia Next Gen Question 1" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

5. "MH Psychosis & Schizophrenia Next Gen Case Study" by Kellea Ewen is licensed under [CC BY-NC 4.0](#)



## XI Glossary

**Agranulocytosis:** Extremely low white blood cell count.

**Akathisia:** Psychomotor restlessness (a feeling of being unable to sit still).

**Alogia:** Reduction or poverty in speech.

**Anhedonia:** The reduced ability to experience pleasure in daily activities.

**Anosognosia:** The inability to recognize that one is ill.

**Apathy:** A decreased interest in activities that would otherwise be interesting.

**Asociality:** A decreased desire for social interaction.

**Avolition:** Reduced motivation or goal-directed behavior.

**Boundary impairment:** An impaired ability to sense where one's influence ends and another person's begins. For example, the person might walk up to a table and drink out of someone else's glass.

**Bradykinesia:** Slowed movement.

**Catatonia:** A pronounced increase or decrease in the rate and amount of movement; excessive movement is purposeless.

**Clang associations:** Stringing words together that rhyme without logical association and do not convey rational meaning. For example, a client exhibiting clang associations may state, "Here she comes with a cat catch a rat match."

**Cognitive symptoms:** A category of symptoms of schizophrenia that refer to problems in attention, concentration, and memory, such as difficulty processing information to make decisions; problems using information immediately after learning it; and trouble focusing or paying attention.

**Command hallucinations:** Auditory hallucinations that command the individual to do something.

**Coordinated specialty care (CSC):** A general term used to describe recovery-oriented treatment programs for people with first-episode psychosis, an early stage of schizophrenia.

**Delusion:** A fixed, false belief not held by cultural peers and persisting in the face of objective contradictory evidence. For example, a client may have the delusion that the CIA is listening to their conversations via satellites.

**Dystonia:** Involuntary contractions of muscles of the extremities, face, neck, abdomen, pelvis, or larynx in either sustained or intermittent patterns that lead to abnormal movements or postures.

**Echolalia:** Pathological repetition of another person's words.

**Echopraxia:** Mimicking the movements of another person.

**Extrapyramidal side effects (EPS):** Adverse effects, such as akathisia, rigidity, bradykinesia, tremor, and acute-dystonic reactions, that can occur from first-generation antipsychotics,

**First-generation antipsychotics:** Also referred to as “typical antipsychotics”; this class of medications has several potential adverse effects due to the blockage of dopamine receptors. Medication is prescribed based on the client's ability to tolerate the adverse effects.

**Flat affect:** A reduced expression of emotions via facial expression or voice tone.

**Hallucinations:** False sensory perceptions not associated with real external stimuli that can include any of the five senses (auditory, visual, gustatory, olfactory and tactile). For example, a client may see spiders climbing on the wall or hear voices telling them to do things. These are referred to as “visual hallucinations” or “auditory hallucinations.”

**Illusions:** Misperceptions of real stimuli. For example, a client may misperceive tree branches blowing in the wind at night to be the arms of monsters trying to grab them.

**Loose associations:** Jumping from one idea to an unrelated idea in the same sentence. For example, the client might state, “I like to dance; my feet are wet.”

**Magical thinking:** Falsely believing that reality can be changed simply by one’s thoughts.

**Metabolic syndrome:** A cluster of conditions that occur together, increasing the risk of heart disease, stroke, and type 2 diabetes. Symptoms include increased blood pressure, high blood sugar, excess body fat around the waist, and abnormal cholesterol or triglyceride levels. Weight, glucose levels, and lipid levels should be monitored regularly.

**Negative symptoms:** A category of symptoms of schizophrenia that includes loss of motivation, diminished feelings of pleasure in everyday life, flat affect, and reduced speaking.

**Neologisms:** Words or phrases created by someone with schizophrenia.

**Neuroleptic malignant syndrome:** A rare but potentially fatal adverse effect that can occur at any time during treatment with antipsychotics. Signs include increased temperature, severe muscular rigidity, confusion, agitation, hyperreflexia, elevation in white blood cell count, elevated creatinine phosphokinase, elevated liver enzymes, myoglobinuria, and acute renal failure. The antipsychotic should be immediately discontinued if these signs occur.

**Paranoia:** An irrational fear that can range from being suspicious to thinking someone is trying to kill you.

**Positive symptoms:** A category of symptoms of schizophrenia that include hallucinations, delusions, thought disorders, disorganized speech, and alterations in behaviors.

**Psychosis:** Conditions where there is loss of contact with reality. Psychosis may be a symptom of a mental illness or other medical conditions. When a person experiences psychosis, their thoughts and perceptions are disturbed,

and the individual has difficulty understanding what is real and what is not real. Symptoms of psychosis include delusions and hallucinations. Other symptoms include incoherent or nonsensical speech and behavior that is inappropriate for the situation.

**Psychotic episode:** An episode of psychosis that can include delusions, hallucinations, disorganized speech, and grossly disorganized or catatonic behavior.<sup>1</sup>

**Schizophrenia:** A severe mental illness characterized by periods of psychosis for at least six months.

**Second-generation antipsychotics:** Also referred to as “atypical antipsychotics”; this class of medication has fewer adverse effects because they block selective dopamine D2 receptors as well as serotonin. For this reason, they are generally better tolerated than first-generation antipsychotics.

**Tardive dyskinesia (TD):** A syndrome of movement disorders that can occur in clients taking first-generation antipsychotics, persisting for at least one month and up to several years despite discontinuation of the medications. The movement disorders include akathisia, dystonia, tics, and other abnormal involuntary movements.

1. American Psychiatric Association. (2013). *Desk reference to the diagnostic criteria from DSM-5*.





### Learning Objectives

- Identify assessment cues of mental health disorders and neurodevelopmental disorders of childhood and adolescence
- Identify nursing priorities for clients with mental health disorders and neurodevelopmental disorders of childhood and adolescence
- Plan outcomes for clients with mental health disorders and neurodevelopmental disorders of childhood and adolescence
- Differentiate safety/protective interventions for clients with mental health disorders and neurodevelopmental disorders of childhood and adolescence
- Apply evidence-based practice when planning care and interventions for clients with mental health disorders and neurodevelopmental disorders of childhood and adolescence
- Analyze treatments for clients with mental health disorders and neurodevelopmental disorders of childhood and adolescence
- Apply the nursing process to clients with mental health disorders and neurodevelopmental disorders of childhood and adolescence at risk for suicide
- Explain conditional confidentiality and mandatory reporting related to caring for children and adolescents

**Mental health disorders among children** are described as “serious changes in the way children typically learn, behave, or handle their emotions, causing

distress and problems getting through the day.”<sup>1</sup> **Neurodevelopmental disorders** are a group of conditions with physical, learning, language, or behavioral impairments that begin during a child’s developmental period, impact their day-to-day functioning, and usually last throughout their lifetime.<sup>2</sup> Attention deficit hyperactivity disorder and autism spectrum disorder are common examples of neurodevelopmental disorders.

One in six children in the United States aged 2 to 8 years (17.4%) has a diagnosed mental health disorder, developmental disorder, or behavioral disorder. Behavior problems are more common among children aged 6 to 11 years.<sup>3</sup>

This chapter will discuss common mental health disorders and neurodevelopmental disorders in children and adolescents and collaborative treatments.

1. Centers for Disease Control and Prevention. (2022). *Data and statistics on children’s mental health*. [https://www.cdc.gov/children-mental-health/data-research/?CDC\\_AAref\\_Val=https://www.cdc.gov/childrensmentalhealth/data.html](https://www.cdc.gov/children-mental-health/data-research/?CDC_AAref_Val=https://www.cdc.gov/childrensmentalhealth/data.html)
2. Ernstmeyer, K., & Christman, E. (Eds.). (2022). *Nursing: Mental health and community concepts*. <https://wtcs.pressbooks.pub/nursingmhcc/>
3. Centers for Disease Control and Prevention. (2022). *Data and statistics on children’s mental health*. [https://www.cdc.gov/children-mental-health/data-research/?CDC\\_AAref\\_Val=https://www.cdc.gov/childrensmentalhealth/data.html](https://www.cdc.gov/children-mental-health/data-research/?CDC_AAref_Val=https://www.cdc.gov/childrensmentalhealth/data.html)



## 12.2 Common Disorders and Disabilities in Children and Adolescents

Mental health disorders, including substance use disorders, are common among children and adolescents in the United States, with nearly 20 percent experiencing a mental health disorder in a given year.<sup>1</sup> The most commonly diagnosed mental health disorders in children aged 13-17 years are attention deficit hyperactivity disorder (ADHD), anxiety problems, behavioral problems, and depression<sup>2</sup>:

- ADHD: 9.8% (approximately 6.0 million)
- Anxiety: 9.4% (approximately 5.8 million)
- Behavioral problems: 8.9% (approximately 5.5 million)
- Depression: 4.4% (approximately 2.7 million)

For adolescents, depression, substance use, and suicide are important concerns. The following statistics demonstrate these concerns in adolescents aged 12-17 years in 2018-2019, prior to the COVID-19 pandemic<sup>3</sup>:

1. Agency for Healthcare Research and Quality. (2025). *Research protocol: Implementation of recommended screening and counseling interventions to prevent mental health disorders in children and adolescents*. <https://effectivehealthcare.ahrq.gov/products/behavioral-health-screening/protocol>
2. Centers for Disease Control and Prevention. (2025). *Data and statistics on children's mental health*. [https://www.cdc.gov/children-mental-health/data-research/?CDC\\_AAref\\_Val=https://www.cdc.gov/childrensmentalhealth/data.html](https://www.cdc.gov/children-mental-health/data-research/?CDC_AAref_Val=https://www.cdc.gov/childrensmentalhealth/data.html)
3. Centers for Disease Control and Prevention. (2025). *Data and statistics on children's mental health*. [https://www.cdc.gov/children-mental-health/data-research/?CDC\\_AAref\\_Val=https://www.cdc.gov/childrensmentalhealth/data.html](https://www.cdc.gov/children-mental-health/data-research/?CDC_AAref_Val=https://www.cdc.gov/childrensmentalhealth/data.html)

- 36.7% had persistent feelings of sadness or hopelessness.
- 18.8% seriously considered attempting suicide; 8.9% attempted suicide.
- 15.1% had a major depressive episode.
- 4.1% had a substance use disorder.
- 1.6% had an alcohol use disorder.

After the COVID-19 pandemic, the Centers for Disease Control and Prevention (CDC) found an 8% increase in persistent feelings of sadness in youth.<sup>4</sup>

These common disorders, as well as developmental disabilities, tics and Tourette syndrome, substance misuse, and gender dysphoria will be discussed in this section. Autism, another type of neurodevelopmental disorder, is discussed in the “[Autism Spectrum Disorder](#)” section.

## Attention Deficit Hyperactivity Disorder

ADHD is a neurodevelopmental disorder characterized by persistent patterns of inattention, hyperactivity, and impulsivity that interfere with functioning or development. It is one of the most common neurodevelopmental disorders of childhood. It is usually first diagnosed in childhood and often lasts into adulthood.<sup>5</sup> Young males are diagnosed more frequently than females. See Figure 12.1<sup>6</sup> for an image depicting a child struggling with symptoms of ADHD in school.

- Centers for Disease Control and Prevention. (2022). *New CDC data illuminate youth mental health threats during the COVID-19 pandemic* [Press Release]. <https://www.cdc.gov/media/releases/2022/p0331-youth-mental-health-covid-19.html>
- Centers for Disease Control and Prevention. (2024). *About attention-deficit/hyperactivity disorder (ADHD)*. [https://www.cdc.gov/adhd/about/?CDC\\_AAref\\_Val=https://www.cdc.gov/ncbddd/adhd/facts.html](https://www.cdc.gov/adhd/about/?CDC_AAref_Val=https://www.cdc.gov/ncbddd/adhd/facts.html)
- “[sad-gad250898b\\_1920](#)” by [Patrick Audet](#) at [Pixabay.com](#) is licensed under [CC0](#)



Figure 12.1 Child Struggling With Symptoms of ADHD in School

## Signs and Symptoms

It is normal for children to exhibit challenging behaviors and have trouble focusing at certain times. However, children with ADHD often have more severe symptoms that cause difficulties at school, at home, or with friends<sup>7</sup>:

- Daydreaming
- Forgetting or losing things
- Squirming or fidgeting
- Talking too much
- Making careless mistakes or taking unnecessary risks
- Difficulty resisting temptation
- Difficulty getting along with others

Diagnosing a child with ADHD is a process requiring several steps by a mental

7. Centers for Disease Control and Prevention. (2024). *About attention-deficit/hyperactivity disorder (ADHD)*. [https://www.cdc.gov/adhd/about/?CDC\\_AAref\\_Val=https://www.cdc.gov/ncbddd/adhd/facts.html](https://www.cdc.gov/adhd/about/?CDC_AAref_Val=https://www.cdc.gov/ncbddd/adhd/facts.html)

health professional. There is no single test to diagnose ADHD, and many other problems such as anxiety, depression, sleep problems, and learning disorders can have similar symptoms as ADHD. The diagnostic process includes a medical exam; hearing and vision tests; and a checklist rating ADHD symptoms completed by parents, teachers, and the child.<sup>8</sup> These symptoms must be present in two or more settings (e.g. home, school) and cause significant impairment in social, academic, or occupational functioning.

## Types of ADHD

There are three subcategories of ADHD, determined by which symptoms are most prominent in the individual<sup>9</sup>:

- **Predominantly Inattentive Presentation:** This involves difficulty organizing or completing tasks, focusing on details, or following instructions and conversations.” The person is easily distracted or forgets details of daily routines. These behaviors are often misunderstood by others, who might label them as ‘laziness’ or a lack of attention.
- **Predominantly Hyperactive-Impulsive Presentation:** This presentation includes constant fidgeting and excessive talking. Sitting still for extended periods, such as during meals or homework, is challenging. Young children may run, jump, or climb constantly. Restlessness and impulsivity are often observed in behaviors such as interrupting others, grabbing objects, speaking at inappropriate moments, or struggling to wait their turn or follow directions. Impulsive tendencies may also

8. Centers for Disease Control and Prevention. (2024). *About attention-deficit/hyperactivity disorder (ADHD)*. [https://www.cdc.gov/adhd/about/?CDC\\_AAref\\_Val=https://www.cdc.gov/ncbddd/adhd/facts.html](https://www.cdc.gov/adhd/about/?CDC_AAref_Val=https://www.cdc.gov/ncbddd/adhd/facts.html)

9. Centers for Disease Control and Prevention. (2024). *About attention-deficit/hyperactivity disorder (ADHD)*. [https://www.cdc.gov/adhd/about/?CDC\\_AAref\\_Val=https://www.cdc.gov/ncbddd/adhd/facts.html](https://www.cdc.gov/adhd/about/?CDC_AAref_Val=https://www.cdc.gov/ncbddd/adhd/facts.html)

increase the likelihood of accidents or injuries compared to peers.

- **Combined Presentation:** This occurs when symptoms from both of the above categories are equally evident.

A person's symptoms of ADHD can change over time, and their predominant presentation may change.

## Causes of ADHD

The cause(s) and risk factors for ADHD are unknown, but current research shows that genetics play an important role. In addition to genetics, other possible risk factors include the following<sup>10</sup> :

- Brain injury
- Exposure to environmental risks (e.g., lead) during pregnancy or at a young age
- Alcohol and tobacco use during pregnancy
- Premature delivery
- Low birth weight

Research does not support popularly held views that ADHD is caused by eating too much sugar, watching too much television, or ineffective parenting. Many of these factors may worsen symptoms, especially in genetically predisposed people, but the evidence is not strong enough to conclude that they are the main causes of ADHD.<sup>11</sup> Recent studies have

10. Centers for Disease Control and Prevention. (2024). *About attention-deficit/hyperactivity disorder (ADHD)*. [https://www.cdc.gov/adhd/about/?CDC\\_AAref\\_Val=https://www.cdc.gov/ncbddd/adhd/facts.html](https://www.cdc.gov/adhd/about/?CDC_AAref_Val=https://www.cdc.gov/ncbddd/adhd/facts.html)

11. Centers for Disease Control and Prevention. (2024). *About attention-deficit/hyperactivity disorder (ADHD)*. [https://www.cdc.gov/adhd/about/?CDC\\_AAref\\_Val=https://www.cdc.gov/ncbddd/adhd/facts.html](https://www.cdc.gov/adhd/about/?CDC_AAref_Val=https://www.cdc.gov/ncbddd/adhd/facts.html)

explored the role of diet in ADHD, yielding mixed findings. While research does not support the widely held belief that excessive sugar intake, excessive screen time, or ineffective parenting directly cause ADHD, certain dietary factors may influence symptom severity. Notably, a “few foods diet” which eliminates potential allergens or hypersensitivity-triggering foods has shown a strong correlation with symptom improvement in some children. This suggests that food allergies or intolerances may play a role in ADHD symptoms for a subset of individuals. However, further research is needed to establish causation.<sup>12</sup>

While the exact cause of attention-deficit/hyperactivity disorder (ADHD) is not fully understood, research suggests that several potential contributing factors may also play a role in its development. Genetic factors appear to be significant, as children with a family history of ADHD or other mental health conditions, such as anxiety or mood disorders, may have an increased risk. Environmental influences—including early exposure to trauma, chronic family conflict, or major life transitions like divorce or relocation—might also contribute to the expression of ADHD symptoms. Additionally, possible differences in brain chemistry and structure have been studied, particularly imbalances in neurotransmitters like dopamine and norepinephrine, and irregularities in brain circuits such as those involving the prefrontal cortex and amygdala. It is important to recognize that these are potential causes and that ADHD is likely the result of a complex interplay of genetic, environmental, and neurological factors rather than any single trigger.<sup>13</sup>

12. Lange, K. W., Lange, K. M., Nakamura, Y., & Reissmann, A. (2023). Nutrition in the management of ADHD: A review of recent research. *Current Nutrition Reports*, 12(3), 383-394. [doi: 10.1007/s13668-023-00487-8](https://doi.org/10.1007/s13668-023-00487-8)
13. Kowalchuk, A., Gonzalez, S. J., & Zoorob, R. J. (2022). Anxiety disorders in children and adolescents. *American Family Physician*, 106(6), 657-664. <https://pubmed.ncbi.nlm.nih.gov/36521463/>



# Treatments for ADHD

The treatment for childhood or adolescent Attention Deficit Hyperactivity Disorder (ADHD) is not curative; it primarily focuses on managing symptoms and improving functional outcomes. The Society for Developmental and Behavioral Pediatrics emphasizes that treatment should focus on areas of functional impairment and include ongoing, scheduled monitoring throughout the client's lifespan.

Management of ADHD typically involves a combination of pharmacologic and non-pharmacologic treatments tailored to the individual needs of the child or adolescent.

## **BEHAVIORAL THERAPY AND PSYCHOTHERAPY**

Behavioral interventions are a cornerstone of ADHD treatment, particularly for younger children. For preschool-aged children (4–5 years old), parent training in behavior management (PTBM) is recommended as the first line of treatment before considering medication, according to the Centers for Disease Control and Prevention. PTBM and behavioral classroom interventions are strongly recommended by the American Academy of Pediatrics and the American Academy of Family Physicians. These strategies are especially effective when combined with pharmacologic treatments. ADHD can significantly affect a child's ability to pay attention, sit still, and maintain relationships with family members and peers due to disruptive behaviors. Psychotherapy can help improve behavior, self-control, and self-esteem.

In addition to behavioral interventions, cognitive-behavioral therapy (CBT) is beneficial, especially for adolescents, helping them develop coping strategies and enhance executive functioning. Educational interventions, such as Individualized Education Programs (IEPs) and 504 plans, provide tailored academic support to help children with ADHD succeed in school.

Complementary and alternative medicine approaches—such as omega-3 fatty acid supplementation, dietary modifications (e.g., elimination of artificial

food colorings), and regular physical activity—have shown some efficacy in symptom management. Finally, neurofeedback and cognitive training aim to improve attention and executive functioning through structured programs, though the evidence supporting these methods remains limited.<sup>14, 15</sup>

Read more in the “[Psychological Therapies and Behavioral Interventions](#)” section of this chapter.

## PARENT EDUCATION ABOUT BEHAVIORAL MANAGEMENT

Nurses can teach parents strategies for managing the behavior of their child or adolescent with ADHD<sup>16</sup>:

- **Create a routine.** Try to follow the same schedule every day, from wake-up time to bedtime.
- **Get organized.** Encourage your child to put school bags, clothing, and toys in the same place every day so that they will be less likely to lose them.
- **Manage distractions.** Limit screentime, limit noise, and provide a clean

14. Chan, E., Fogler, J. M., & Hammerness, P. G. (2016). Treatment of attention-deficit/hyperactivity disorder in adolescents: A systematic review. *JAMA*, 315(18), 1997-2008. [doi: 10.1001/jama.2016.5453](https://doi.org/10.1001/jama.2016.5453).

15. Catalá-López, F., Hutton, B., Núñez-Beltrán, A., Page, M. J., Ridao, M., Macías Saint-Gerons, D., Catalá, M. A., Tabarés-Seisdedos, R., & Moher, D. (2016). The pharmacological and non-pharmacological treatment of attention deficit hyperactivity disorder in children and adolescents: A systematic review with network meta-analyses of randomised trials. *PLoS One*, 12(7):e0180355. [doi: 10.1371/journal.pone.0180355](https://doi.org/10.1371/journal.pone.0180355).

16. Centers for Disease Control and Prevention. (2024). *About attention-deficit/hyperactivity disorder (ADHD)*. [https://www.cdc.gov/adhd/about/?CDC\\_AAref\\_Val=https://www.cdc.gov/ncbddd/adhd/facts.html](https://www.cdc.gov/adhd/about/?CDC_AAref_Val=https://www.cdc.gov/ncbddd/adhd/facts.html)



workspace when your child is doing homework. Some children with ADHD learn better if they are moving or listening to background music.

- **Limit choices.** To help your child avoid feeling overwhelmed or overstimulated, offer only a few options, such as letting them choose between two outfits, meals, or toys.
- **Be clear and specific when you talk with your child.** Let your child know you are listening by describing what you heard them say. Use clear, brief directions when they need to do something.
- **Help your child plan.** Break down complicated tasks into simpler, shorter steps. For long tasks, starting early and taking breaks may help limit stress.
- **Use goals and praise or other rewards.** Use a chart to list goals and track positive behaviors, and then let your child know they have done well by telling them or by rewarding their efforts in other ways. Be sure the goals are realistic because progress towards small steps is important to maintain a child's self-esteem.
- **Discipline effectively.** Instead of scolding, yelling, or spanking, use effective directions, time-outs, or removal of privileges as consequences for inappropriate behavior.
- **Create positive opportunities.** Children with ADHD may find certain situations stressful. Discovering and encouraging what your child does well, whether it's school, sports, art, music, or play, can help create positive experiences.
- **Provide a healthy lifestyle.** Nutritious food, lots of physical activity, and sufficient sleep are important for preventing ADHD symptoms from getting worse.
- **Communicate regularly with teachers.** Clear, regular communication between teachers and parents helps reinforce behavior management strategies at school and at home.

Parents can also be referred to ADHD support groups in the following box.

## Support Groups for ADHD

- ▶ **CHADD**: Information and resources on ADHD, including treatment options, local support groups for clients and parents, and online support communities
- ▶ **Attention Deficit Disorder Association (ADDA)**: Information and resources on ADD for adults living with the disorder, including support groups and workshops
- ▶ **Psychology Today Support Groups**: Support groups near you for ADHD and other conditions

## PHARMACOTHERAPY

Medication may be prescribed to help children aged six and older manage their ADHD symptoms and help them control behaviors that cause difficulties with family, friends, and at school.<sup>17</sup>

Before medications are initiated, a comprehensive medical exam including height, weight, blood pressure, heart rate, and cardiovascular history should be performed. A pretreatment baseline should be established for common side effects such as appetite, sleep, headaches, and abdominal pain. Adolescent clients should also be assessed for substance use.<sup>18</sup>

17. Centers for Disease Control and Prevention. (2024). *About attention-deficit/hyperactivity disorder (ADHD)*. [https://www.cdc.gov/adhd/about/?CDC\\_AAref\\_Val=https://www.cdc.gov/ncbddd/adhd/facts.html](https://www.cdc.gov/adhd/about/?CDC_AAref_Val=https://www.cdc.gov/ncbddd/adhd/facts.html)

18. Krull, K. R. (2022). Attention deficit hyperactivity disorder in children and adolescents: Treatment with medications. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

The choice of medication by the prescriber depends on many factors, such as the following<sup>19</sup> :

- Duration of coverage (e.g., desired coverage for school day plus completion of homework)
- The desire to avoid medication administration at school
- The ability of the child to swallow pills or capsules
- Coexisting emotional or behavioral conditions
- History of substance abuse in the client or a household member (i.e., stimulants with less abuse potential are prescribed)
- Expense
- Preferences of the child and their caregivers

## Stimulants

Stimulants such as methylphenidate (Ritalin), dextroamphetamine (Dexedrine), and dextroamphetamine-amphetamine (Adderall) are considered first-line treatment because of rapid onset of action and a long record of safety and efficacy. Stimulants are available in short-, intermediate-, and long-acting formulations. The exact mechanism of action of stimulants in ADHD is unknown, but they are known to affect the dopaminergic and noradrenergic systems, causing a release of catecholamines. Stimulants have been found to improve caregiver-child interactions, aggressive behavior, and academic productivity.<sup>20</sup>

Stimulants are controlled substances and require a Schedule II prescription.

19. Krull, K. R. (2022). Attention deficit hyperactivity disorder in children and adolescents: Treatment with medications. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

20. Krull, K. R. (2022). Attention deficit hyperactivity disorder in children and adolescents: Treatment with medications. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

CNS stimulants, including methylphenidate and amphetamine-like substances, have a high potential for abuse and dependence. The risk of abuse by the client or their family members should be assessed prior to prescribing stimulants, and signs of abuse and dependence should be evaluated while the client is receiving therapy.<sup>21</sup>

Side Effects

Stimulants may cause minor side effects that resolve when dosage levels are lowered, or a different stimulant is prescribed. The most common side effects include the following<sup>22,23</sup>:

- Difficulty falling asleep or staying asleep
- Loss of appetite and weight loss
- Stomach pain
- Headache

Less common side effects include motor or verbal tics (sudden, repetitive movements or sounds) or personality changes (such as appearing “flat” or without emotion).<sup>24</sup> Sudden death, stroke, and myocardial infarction have

21. National Institutes of Health. (n.d.). *Black box warning*. U.S. National Library of Medicine. <https://www.nlm.nih.gov/>
22. National Institute of Mental Health. (2016). *Mental health medications*. U.S. Department of Health & Human Services. <https://www.nimh.nih.gov/health/topics/mental-health-medications>
23. National Institutes of Health. (n.d.). *Black box warning*. U.S. National Library of Medicine. <https://www.nlm.nih.gov/>
24. National Institute of Mental Health. (2016). *Mental health medications*. U.S. Department of Health & Human Services. <https://www.nimh.nih.gov/health/topics/mental-health-medications>

been reported in adults with CNS-stimulant treatment at recommended doses. Sudden death has been reported in pediatric clients with structural cardiac abnormalities and other serious heart problems taking CNS stimulants at recommended doses. If paradoxical worsening of symptoms or other adverse reactions occur, the provider should be contacted, and the dosage reduced or discontinued. Stimulants are contraindicated in clients using a monoamine oxidase inhibitor (MAOI) or using an MAOI within the preceding 14 days.<sup>25</sup>

Nurses should be aware there is a possibility of diversion or misuse of stimulants by adolescents or their caregivers. Up to 29 percent of school- and college-aged students with stimulant prescriptions have been asked to give, sell, or trade their medication.<sup>26</sup>

## Selective Norepinephrine Reuptake Inhibitors

Selective norepinephrine reuptake inhibitors (SNRIs) such as atomoxetine (Strattera) are an alternative to stimulants for clients who experience side effects with stimulants. They may also be helpful in treating concurrent depressive or anxiety disorders. SNRIs are not controlled substances, so they may be prescribed for adolescents (or their family members) with substance use disorders. The dosage depends on the child's weight, and the duration of action is 10 to 12 hours. Atomoxetine has a Black Box Warning about increased risk of suicidal thinking in children and adolescents.<sup>27</sup>

- 25. National Institutes of Health. (n.d.). *Black box warning*. U.S. National Library of Medicine. <https://www.nlm.nih.gov/>
- 26. Krull, K. R. (2022). Attention deficit hyperactivity disorder in children and adolescents: Treatment with medications. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)
- 27. Krull, K. R. (2022). Attention deficit hyperactivity disorder in children and adolescents: Treatment with medications. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

## Alpha-2 Adrenergic Agonists

Alpha-2 adrenergic agonists such as clonidine are typically used when children respond poorly to stimulants or SNRIs, have unacceptable side effects, or have significant coexisting conditions.

## CLIENT AND PARENT EDUCATION

There are several important client education topics to provide to clients and/or the parents of minor children<sup>28</sup>:

- **Medication information:** Discussing the purpose, potential side effects, and the importance of adherence to prescribed treatments.
- **Controlled Substance Status/High Potential for Abuse and Dependence:** Stimulants are a controlled substance by the FDA that can be abused and lead to dependence. Stimulants should be stored in a safe (preferably locked) place to prevent misuse and should not be shared with anyone. Unused or expired stimulants should be disposed of based on state law and regulations or returned to a medicine take-back program if it is available in the community.
- **Cardiovascular Risks:** Stimulants can increase blood pressure and pulse rate. Potential serious cardiovascular risks include sudden death, cardiomyopathy, myocardial infarction, stroke, and hypertension. Instruct clients to contact a health care provider immediately if they develop symptoms, such as exertional chest pain, dizziness, or passing out.
- **Suppression of Growth:** Stimulants may cause slowing of growth in children and weight loss.
- **Psychiatric Risks:** Stimulants can cause psychosis or manic symptoms, even in clients who have no prior history of these symptoms.
- **Priapism:** Painful or prolonged penile erections can occur; seek immediate medical attention.

28. National Institutes of Health. (n.d.). *Black box warning*. U.S. National Library of Medicine. <https://www.nlm.nih.gov/>

- **Alcohol:** Alcohol should be avoided.

Nurses should reinforce with the client and their family members that the reason for the prescribed medication is to help with self-control and the ability to focus. Possible side effects should be reviewed, and clients and their family members should be reminded it may take one to three months to determine the best pharmacological treatment, dose, and frequency of medication administration. During this time, the child's symptoms and adverse effects will be monitored weekly and the medication dose adjusted accordingly.<sup>29</sup>

## ADHD Into Adulthood

ADHD lasts into adulthood for at least one third of children with ADHD. Treatments for adults can include medication, psychotherapy, or a combination of treatments.<sup>30</sup>

## Anxiety

Some anxiety is a normal part of childhood and occurs at predictable stages of development. For example, from approximately age 8 months through the preschool years, healthy children may show anxiety when separated from their parents or caregivers. Young children also commonly have fears, such as fear of the dark, storms, animals, or strangers.<sup>31</sup> Anxiety is functional and

29. Krull, K. R. (2022). Attention deficit hyperactivity disorder in children and adolescents: Treatment with medications. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

30. Centers for Disease Control and Prevention. (2024). *About attention-deficit/hyperactivity disorder (ADHD)*. [https://www.cdc.gov/adhd/about/?CDC\\_AAref\\_Val=https://www.cdc.gov/ncbddd/adhd/facts.html](https://www.cdc.gov/adhd/about/?CDC_AAref_Val=https://www.cdc.gov/ncbddd/adhd/facts.html)

31. American Academy of Child & Adolescent Psychiatry. (2017). *Anxiety and*

normal human emotion when situational. Consider the fear of dangerous situations such as approaching a rattlesnake or standing on a steep cliff; at crucial times anxiety is important because it provides safety.

Anxiety is also motivational as it drives adolescents to accomplish goals such as passing a test by working hard and studying. **Anxiety disorders** are a group of mental health conditions characterized by excessive fear or worry that interferes with daily activities. Common types include Generalized Anxiety Disorder (GAD), Social Anxiety Disorder, and Panic Disorder. Symptoms can be persistent and may manifest physically and emotionally. Anxiety disorders in children and adolescents are characterized by excessive fear, worry, or avoidance behaviors that are disproportionate to the actual threat and impair daily functioning.

Read more about “[Adverse Childhood Experiences](#)” in the “Trauma, Abuse, and Violence” chapter.

## Symptoms of Anxiety Disorders in Children and Adolescents

Children with anxiety disorders are overly tense or fearful; some may seek lots of reassurance; and their worries may interfere with daily activities. Because anxious children may also be quiet, compliant, and eager to please, their feelings of anxiety can be easily missed. When a child does not outgrow the typical fears and anxieties in childhood or when there are so many fears and worries they interfere with school, home, or play activities, the child may be diagnosed with an anxiety disorder. Examples of symptoms related to

*children.* [https://www.aacap.org/AACAP/Families\\_and\\_Youth/Facts\\_for\\_Families/FFF-Guide/The-Anxious-Child-047.aspx](https://www.aacap.org/AACAP/Families_and_Youth/Facts_for_Families/FFF-Guide/The-Anxious-Child-047.aspx)



different types of anxiety disorders in children and adolescents include the following<sup>32</sup>:

- Being very afraid when away from parents or caregivers (i.e., **separation anxiety**)
- Having extreme fear about a specific thing or situation, such as dogs, insects, or going to the doctor (i.e., phobias)
- Being very afraid of school and other places where there are people (i.e., social anxiety)
- Being very worried about the future and about bad things happening (i.e., general anxiety)
- Having repeated episodes of sudden, unexpected, intense fear associated with symptoms like fast heart rate, trouble breathing, dizziness, GI issues, or shakiness (i.e., panic disorder)

Anxiety can also cause irritability, anger, fatigue, headaches, stomachaches, or trouble sleeping.<sup>33</sup>

## Treatment of Anxiety Disorders in Children and Adolescents

Early treatment of anxiety disorders in children can enhance friendships, social and academic potential, and self-esteem. Interprofessional treatments often include a combination of individual psychotherapy and behavioral therapy, family therapy, medications, and consultations with the child's

32. Centers for Disease Control and Prevention. (2024). *About children's mental health*. <https://www.cdc.gov/children-mental-health/about/index.html>

33. Centers for Disease Control and Prevention. (2024). *About children's mental health*. <https://www.cdc.gov/children-mental-health/about/index.html>

school.<sup>34</sup> The goal of treatment for anxiety disorders in children and adolescents, is to reduce the severity of anxiety symptoms, improve functional impairment, and enhance overall quality of life. Overall, the treatment approach should be individualized, taking into account the severity of symptoms, the specific anxiety disorder, and the child's developmental level. When a child is overly worried or anxious, a nurses' initial assessment should determine if conditions in the child's environment are causing this feeling. For example, is the anxiety resulting from being bullied or from adverse childhood experiences (ACEs)? If so, protective interventions should be put into place. If no realistic threat exists and the anxiety causes significant prolonged life dysfunction, then the child should be referred to a health provider to determine if an anxiety disorder exists.<sup>35</sup>

Read more about psychological and behavioral treatments for children and adolescents in the “[Psychological Therapies and Behavioral Interventions](#)” section.

## PHARMACOTHERAPY

Pharmacotherapy primarily involves the use of selective serotonin reuptake inhibitors (SSRIs), which have been shown to be effective and well-tolerated in children and adolescents with anxiety disorders. The American Academy of Child and Adolescent Psychiatry recommends SSRIs for clients aged 6 to 18 years with social anxiety, generalized anxiety, separation anxiety, or panic disorder. Common SSRIs used include fluoxetine, sertraline, and escitalopram.

34. Centers for Disease Control and Prevention. (2024). *About children's mental health*. <https://www.cdc.gov/children-mental-health/about/index.html>

35. Hilt, R. J., & Nussbaum, A. M. (Eds.). (2016). *DSM-5 pocket guide for child and adolescent mental health*. <https://doi.org/10.1176/appi.books.9781615370511>

In cases of moderate to severe anxiety, a combination of CBT and SSRIs may be more effective than either treatment alone.<sup>36,37</sup>

## CLIENT AND FAMILY EDUCATION

Effective client and family education is essential in the treatment of anxiety disorders in children and adolescents, as it empowers caregivers and young clients with the knowledge, skills, and support needed to manage symptoms, adhere to treatment plans, and promote long-term mental wellness. Key components of an effective education plan include:

- **Understanding Anxiety:** Educating about the anxiety disorders and their symptoms, treatment options, and the importance of early intervention.
- **Coping Strategies:** Providing techniques for managing anxiety, including relaxation exercises and time management.
- **Medication Guidance:** Discussing the role of medications, potential side effects and benefits, and the importance of adherence to treatment plans.
- **Support Resources:** Connecting families with mental health resources, support groups, and counseling services for additional help. The role of CBT. Collaborative decision-making and providing resources such as written materials and reputable online information can support families in managing the condition.

36. Connolly, S. D., Suarez, L., & Sylvester, C. (2011). Assessment and treatment of anxiety disorders in children and adolescents. *Current Psychiatry Reports*, 13(2), 99-110. [doi: 10.1007/s11920-010-0173-z](https://doi.org/10.1007/s11920-010-0173-z).

37. Nicotra, C. M., Strawn, J. R. (2023). Advances in pharmacotherapy for pediatric anxiety disorders. *Child and Adolescent Psychiatric Clinics of North America*, 32(3), 573-587. [doi: 10.1016/j.chc.2023.02.006](https://doi.org/10.1016/j.chc.2023.02.006).

Read additional information about anxiety disorders and associated treatments in the “[Anxiety Disorders](#)” chapter. Post-traumatic stress disorder (PTSD) can develop in children or adolescents who have experienced a shocking, frightening, or dangerous event. It has similar symptoms to severe anxiety. Read more in the “[Post-Traumatic Stress Disorder](#)” section of the “Anxiety Disorders” chapter. There is also an association between obsessive-compulsive disorder (OCD) and children who have been exposed to trauma. Read more in the “[Obsessive-Compulsive Disorder](#)” section of the “Anxiety Disorders” chapter.

## Depression

Every person feels sad occasionally. However, persistent sadness, hopelessness, withdrawal, or loss of interest in previously enjoyed activities may indicate depression. If these symptoms last for two or more weeks, a depressive disorder may be diagnosed. **Depression**, also known as Major Depressive Disorder (MDD), is a mood disorder characterized by persistent feelings of sadness, hopelessness, and a loss of interest or pleasure in activities, and various emotional and physical problems that interfere with daily functioning. It can affect how individuals think, feel, & handle daily activities, & may lead to physical symptoms.<sup>38</sup>

## Symptoms of Depressive Disorders in Children and Adolescents

Examples of behaviors observed in children and adolescents with a depressive disorder are as follows<sup>39</sup>:

- Feeling sad, hopeless, or irritable most of the time

38. Centers for Disease Control and Prevention. (2024). *About children’s mental health*. <https://www.cdc.gov/children-mental-health/about/index.html>

39. American Academy of Child & Adolescent Psychiatry. (2017). *Anxiety and children*. [https://www.aacap.org/AACAP/Families\\_and\\_Youth/Facts\\_for\\_Families/FFF-Guide/The-Anxious-Child-047.aspx](https://www.aacap.org/AACAP/Families_and_Youth/Facts_for_Families/FFF-Guide/The-Anxious-Child-047.aspx)

- Not wanting to do or enjoy fun things
- Showing changes in eating patterns (e.g., eating a lot more or a lot less than usual)
- Showing changes in sleep patterns (e.g., sleeping a lot more or a lot less than normal)
- Showing changes in energy (e.g., being tired and sluggish or tense and restless most of the time)
- Difficulty paying attention
- Feeling worthless, useless, or guilty
- Engaging in self-injury or self-destructive behavior
- Having suicidal thoughts or making a plan for suicide
- Exhibiting physical complaints, such as frequent headaches or stomach aches
- Using alcohol or drugs as a way of trying to feel better

Depression might also cause a child to appear unmotivated or act out, causing others to incorrectly label the child as “lazy” or a “trouble-maker.”<sup>40</sup>

It is important to ask children and adolescents who are withdrawn or sad about self-harm risks. Adolescents may perceive a single disappointment (such as a relationship break-up) as so catastrophic they feel suicidal or begin to hurt themselves.<sup>41</sup>

## Causes of Depression

The cause of depression is not always known. Depression can be hereditary but can also be situational or environmental. Some causes of depression are increased stress, death of a family member or close friend, social media, and

40. Centers for Disease Control and Prevention. (2024). *About children's mental health*. <https://www.cdc.gov/children-mental-health/about/index.html>

41. Hilt, R. J., & Nussbaum, A. M. (Eds.). (2016). *DSM-5 pocket guide for child and adolescent mental health*. <https://doi.org/10.1176/appi.books.9781615370511>

bullying. Having other conditions such as attention problems, learning disorders, anxiety, or conduct disorders create a higher risk for depression.<sup>42</sup> Potential causes can be categorized as the following:

- **Genetic Factors:** A family history of depression or other mood disorders
- **Environmental Stressors:** Traumatic events, significant life changes (e.g., loss, divorce), or chronic stress may contribute.
- **Biological Factors:** Neurochemical imbalances in the brain, hormonal changes, and physical health conditions can also play a role.

## Treatment of Depressive Disorders in Children and Adolescents

First-line treatments for mild to moderate depression in children and adolescents typically include cognitive behavioral therapy (CBT) and interpersonal therapy (IPT). For moderate to severe cases, a combination of psychotherapy and pharmacotherapy is often recommended.

- **Psychotherapy:** CBT is widely used to help young individuals identify and reframe negative thought patterns, build coping skills, and reduce depressive symptoms.
- **Interpersonal Therapy (IPT):** IPT focuses on enhancing interpersonal relationships and social functioning, which are often disrupted by depression.
- **Lifestyle Modifications:** Encouraging regular physical activity, balanced nutrition, and healthy sleep habits can play a supportive role in improving mood and overall well-being.

If depression is suspected, a nurse, school counselor, or other frontline

42. American Academy of Child & Adolescent Psychiatry. (2017). *Anxiety and children*. [https://www.aacap.org/AACAP/Families\\_and\\_Youth/Facts\\_for\\_Families/FFF-Guide/The-Anxious-Child-047.aspx](https://www.aacap.org/AACAP/Families_and_Youth/Facts_for_Families/FFF-Guide/The-Anxious-Child-047.aspx)

professional can collaborate with the child's parents or guardians to refer the child to a qualified mental health provider. A comprehensive assessment should guide treatment planning, which may include individual therapy, family-focused behavioral interventions, and coordination with the child's school to support progress. In more severe cases, antidepressant medications may be prescribed in combination with therapy to achieve better outcomes.

Read more about behavioral treatments in the "[Psychological Therapies and Behavioral Interventions](#)" section.

## Pharmacotherapy

Pharmacologic treatment for depression in children and adolescents is typically reserved for moderate to severe cases and should be guided by a thorough clinical evaluation. The only FDA-approved antidepressants for pediatric use are fluoxetine (Prozac) for children aged 8 and older and escitalopram (Lexapro) for those aged 12 and older. These medications, both selective serotonin reuptake inhibitors (SSRIs), have demonstrated effectiveness but must be prescribed with caution due to the increased risk of suicidal thoughts and behaviors in young clients. Other SSRIs, such as sertraline and citalopram, have shown promising results in clinical studies but are not FDA-approved for this population. Close monitoring, regular follow-up appointments, and collaboration with the child's caregivers are essential when initiating or adjusting medication therapy.<sup>43</sup>

## Client and Family Education

Educating both the client and their family is a vital component of effectively managing depression in children and adolescents. A well-informed support

43. National Institute of Mental Health. (n.d.). *Child and adolescent mental health: Depression*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/child-and-adolescent-mental-health>

system can help recognize early signs, encourage treatment adherence, and promote a more stable and nurturing environment for recovery.

**Understanding Depression:** Families should be educated about what depression is, including its symptoms (such as persistent sadness, irritability, fatigue, changes in appetite or sleep, and loss of interest in activities), how it impacts daily functioning, and its potential effects on academic performance, social relationships, and self-esteem. Education should also include how to recognize warning signs of worsening depression or suicidal ideation, such as increased withdrawal, talk of hopelessness, or sudden behavioral changes. Emphasizing the importance of a supportive home environment and regular follow-up appointments helps ensure continuity of care and early intervention when needed.

**Coping Strategies:** Teaching children and their families healthy coping mechanisms such as mindfulness, deep breathing, journaling, physical activity, and structured routines—can enhance emotional regulation and resilience. Parents and caregivers can model and encourage stress management strategies to support recovery.

**Medication Information:** When pharmacotherapy is part of the treatment plan, it is essential to provide age-appropriate education about the purpose of the medication, potential side effects, and the critical importance of adhering to the prescribed regimen. Families should be encouraged to ask questions and to report any concerns or behavioral changes to the prescribing provider promptly.

**Support Resources:** Families should be connected to community resources, such as school-based mental health programs, peer support groups, and educational websites that provide trustworthy information. Counseling services for the child and family may also be beneficial. Providing written materials or recommending reputable online sources can help reinforce learning and offer continued support outside of clinical settings.<sup>44</sup>

44. National Institute of Mental Health. (n.d.). *Child and adolescent mental*



Read more about depression and pharmacological treatments in the “[Depressive Disorders](#)” chapter.

## Behavior Disorders

Children sometimes argue or act angry or defiant around adults. However, a behavior disorder is diagnosed when disruptive behaviors are uncommon for the child’s age, persist over time, or are severe. Two types of behavior disorders are oppositional defiant disorder and conduct disorder.<sup>45</sup>

### Oppositional Defiant Disorder

All children are oppositional from time to time, particularly when they are feeling tired, hungry, stressed, or upset. They may argue, talk back, disobey, and defy parents, teachers, or other adults. Oppositional behavior is considered a normal part of development for children two to three years of age and early adolescents. However, uncooperative and hostile behavior becomes a serious concern when it is so frequent and consistent that it stands out when compared with other children of the same age or when it significantly affects the child’s social, family, and academic life. According to the DSM-5, these behaviors must be present for at least six months to warrant clinical concern.<sup>46</sup>

*health: Depression. U.S. Department of Health and Human Services.*

<https://www.nimh.nih.gov/health/topics/child-and-adolescent-mental-health>

45. American Academy of Child & Adolescent Psychiatry. (2019). *Oppositional defiant disorder*. [https://www.aacap.org/aacap/families\\_and\\_youth/facts\\_for\\_families/fff-guide/Children-With-Oppositional-Defiant-Disorder-072.aspx](https://www.aacap.org/aacap/families_and_youth/facts_for_families/fff-guide/Children-With-Oppositional-Defiant-Disorder-072.aspx)

46. American Academy of Child & Adolescent Psychiatry. (2019). *Oppositional defiant disorder*. [https://www.aacap.org/aacap/families\\_and\\_youth/](https://www.aacap.org/aacap/families_and_youth/)

When children act out persistently causing serious problems at home, in school, or with peers, they may be diagnosed with **oppositional defiant disorder (ODD)**. Up to 16 percent of all school-age children and adolescents have ODD. ODD usually starts before 8 years of age. Individuals with ODD are more likely to act oppositional or defiant around people they know well, such as family members, a regular care provider, or a teacher.<sup>47</sup>

Examples of ODD behaviors include the following<sup>48</sup>:

- Often being angry or losing one's temper
- Often arguing with adults or refusing to comply with adults' rules or requests
- Often being resentful or spiteful
- Deliberately annoying others or becoming annoyed with others
- Often blaming other people for one's own mistakes or misbehavior

Many children with ODD may have coexisting conditions such as anxiety, post-traumatic stress disorder (PTSD), ADHD, autism, learning disabilities, or substance abuse. Some of these conditions may also be mistaken for ODD. See Figure 12.2<sup>49</sup> for an illustration of conditions that can be mistaken for ODD.

[facts\\_for\\_families/fff-guide/Children-With-Oppositional-Defiant-Disorder-072.aspx](https://www.cdc.gov/children-mental-health/about/about-behavior-or-conduct-problems-in-children.html)

47. Centers for Disease Control and Prevention. (2025). *Behavior or conduct problems in children*. <https://www.cdc.gov/children-mental-health/about/about-behavior-or-conduct-problems-in-children.html>

48. Centers for Disease Control and Prevention. (2025). *Behavior or conduct problems in children*. <https://www.cdc.gov/children-mental-health/about/about-behavior-or-conduct-problems-in-children.html>

49. "Instead of Oppositional Defiant Disorder 1 Wide.png" by MissLunaRose12 is licensed under [CC BY-SA 4.0](https://creativecommons.org/licenses/by-sa/4.0/)

# Conditions and issues that could be mistaken for Oppositional Defiant Disorder



Figure 12.2 Conditions That Can Be Mistaken for Oppositional Defiant Disorder

## Causes of ODD

The etiology of Oppositional Defiant Disorder (ODD) involves a complex interplay of genetic, environmental, and biological factors, with family dynamics playing a particularly significant role in the development and persistence of the disorder. According to the American Academy of Child and Adolescent Psychiatry, children with ODD typically direct their defiant and hostile behaviors toward authority figures, and these behaviors do not include the severe rule-breaking or antisocial acts seen in conduct disorder. Genetic influences, such as a family history of behavioral or mood disorders, may increase a child's vulnerability. Environmental contributors—including inconsistent parenting practices, exposure to trauma, abuse, or neglect—can also play a critical role. Additionally, biological factors, such as differences in brain function and individual temperament, may predispose some children to

exhibit oppositional and defiant behaviors more frequently or intensely than their peers.<sup>50</sup>

## TREATMENT OF ODD

Early intervention is critical to prevent the progression to more severe disorders such as conduct disorder and to mitigate long-term adverse outcomes. Treatment for ODD primarily involves behavioral therapy, focusing on improving parent-child interactions and implementing consistent discipline strategies.

Treatment of ODD includes the following<sup>51</sup>:

- Parent behavioral management training to help parents manage the child's behavior
- Individual psychotherapy to develop more effective anger management skills
- Family psychotherapy to improve communication and mutual understanding
- Cognitive behavioral therapy and other psychotherapies to decrease negativity and enhance effective problem-solving
- Social skills training to increase flexibility and improve frustration tolerance with peers

50. Lin, X., He, T., Heath, M., Chi, P., & Hinshaw, S. (2022). A systematic review of multiple family factors associated with oppositional defiant disorder. *International Journal of Environmental Research and Public Health*, 19(17), 10866. doi: [10.3390/ijerph191710866](https://doi.org/10.3390/ijerph191710866).

51. American Academy of Child & Adolescent Psychiatry. (2019). *Oppositional defiant disorder*. [https://www.aacap.org/aacap/families\\_and\\_youth/facts\\_for\\_families/fff-guide/Children-With-Oppositional-Defiant-Disorder-072.aspx](https://www.aacap.org/aacap/families_and_youth/facts_for_families/fff-guide/Children-With-Oppositional-Defiant-Disorder-072.aspx)

Medications are not recommended as first-line treatment but may be used to address comorbid conditions, which can, in turn, alleviate ODD symptoms. They may be prescribed to control distressing symptoms of ODD, as well as symptoms related to coexisting conditions such as ADHD, anxiety, and mood disorders.<sup>52</sup>

## Parent Education

Parents of children diagnosed with ODD need support and understanding. Nurses can teach parents to help their child with ODD in the following ways<sup>53</sup>:

- **Understanding ODD:** Educating families disorder, symptoms, & impact on functioning.
- **Build on the positives.** Give the child praise and positive reinforcement when they show flexibility or cooperation. Discipline will not work if there are no positive interactions.
- **Take a time-out or break.** If conflict with your child is progressively getting worse instead of better, take a time-out break. Demonstration of taking a break is also good modeling for the child. If the child decides to take a time-out to prevent overreacting, they should receive support for doing so.
- **Prioritize your battles.** Because the child with ODD has trouble avoiding power struggles, prioritize the things you want your child to do. For

52. American Academy of Child & Adolescent Psychiatry. (2019). *Oppositional defiant disorder*. [https://www.aacap.org/aacap/families\\_and\\_youth/facts\\_for\\_families/fff-guide/Children-With-Oppositional-Defiant-Disorder-072.aspx](https://www.aacap.org/aacap/families_and_youth/facts_for_families/fff-guide/Children-With-Oppositional-Defiant-Disorder-072.aspx)

53. American Academy of Child & Adolescent Psychiatry. (2019). *Oppositional defiant disorder*. [https://www.aacap.org/aacap/families\\_and\\_youth/facts\\_for\\_families/fff-guide/Children-With-Oppositional-Defiant-Disorder-072.aspx](https://www.aacap.org/aacap/families_and_youth/facts_for_families/fff-guide/Children-With-Oppositional-Defiant-Disorder-072.aspx)

example, if you give your child a time-out in their room for misbehavior, don't add time to the time-out for arguing. Instead, calmly say, "Your time will start when you go to your room." Listen when they shout, but do not shout back.

- **Set reasonable, age-appropriate limits and consequences.** Set limits and establish consequences that can be enforced consistently. For example, if a child refuses to do homework, a reasonable consequence might be losing screen time for the evening.
- **Maintain other interests.** Manage your stress with healthy life choices such as exercise and relaxation. Maintain personal interests so that managing your child with ODD doesn't take all your time and energy. Use respite care and other breaks as needed.
- **Obtain support.** Collaborate and obtain support from other adults working with your child (e.g., teachers and coaches).

## Conduct Disorder

**Conduct disorder (CD)** is diagnosed when a child shows an ongoing pattern of aggression toward others with serious violations of rules and social norms at home, school, and with peers. These rule violations may involve breaking the law and result in arrest.<sup>54</sup> Adults with antisocial conduct disorder typically show symptoms of CD before age 15.<sup>55</sup>

54. Centers for Disease Control and Prevention. (2025). *Behavior or conduct problems in children*. <https://www.cdc.gov/children-mental-health/about/about-behavior-or-conduct-problems-in-children.html>

55. Mayo Clinic. (2019). *Antisocial personality disorder*. <https://www.mayoclinic.org/diseases-conditions/antisocial-personality-disorder/symptoms-causes/syc-20353928#:~:text=Adults%20with%20antisocial%20personality%20disorder,Destruction%20of%20property>

Examples of CD behaviors are as follows<sup>56</sup> :

- Breaking serious rules, such as running away, staying out all night, or skipping school
- Being aggressive in a way that causes harm, such as bullying, fighting, or being cruel to animals
- Lying, stealing, or purposefully damaging other people's property

Children who exhibit these serious behaviors should receive a comprehensive evaluation and treatment by a mental health professional. Some signs of behavior problems, such as not following rules in school, can be related to learning disorders that require additional assessment and interventions. Without treatment, many youngsters with conduct disorder are likely to have ongoing problems resulting in the inability to adapt to the demands of adulthood.<sup>57</sup>

## TREATMENT FOR CD

Starting treatment early for CD is important. Research indicates the most effective treatment for younger children is behavior therapy training for parents—a therapist helps parents strengthen their relationship with the child and manage behavior effectively. For school-age children and teens, a combination of behavior therapy training that includes the child, the family,

56. Centers for Disease Control and Prevention. (2025). *Behavior or conduct problems in children*. <https://www.cdc.gov/children-mental-health/about/about-behavior-or-conduct-problems-in-children.html>

57. American Academy of Child & Adolescent Psychiatry. (2018). *Conduct disorder*. [https://www.aacap.org/aacap/families\\_and\\_youth/facts\\_for\\_families/fff-guide/conduct-disorder-033.aspx](https://www.aacap.org/aacap/families_and_youth/facts_for_families/fff-guide/conduct-disorder-033.aspx)

and the school is most effective.<sup>58</sup> Read more about treatment in the “[Psychological Therapies and Behavioral Interventions](#)” section.

## Developmental Disabilities

**Developmental disabilities** are a group of conditions with physical, learning, language, or behavioral impairments. These conditions begin during the developmental period, may impact day-to-day functioning, and usually last throughout a person’s lifetime.<sup>59</sup> Research indicates about 17% of children are diagnosed with a developmental disability such as blindness, hearing loss, learning disability, intellectual disability, speech and language impairment, attention deficit hyperactivity disorder (ADHD), or autism spectrum disorder.<sup>60</sup>

Keep in mind that having a developmental disability does not mean the person is not healthy. Being healthy means staying well so one can lead a full, active life.<sup>61</sup>

- 58. American Academy of Child & Adolescent Psychiatry. (2018). *Conduct disorder*. [https://www.aacap.org/aacap/families\\_and\\_youth/facts\\_for\\_families/fff-guide/conduct-disorder-033.aspx](https://www.aacap.org/aacap/families_and_youth/facts_for_families/fff-guide/conduct-disorder-033.aspx)
- 59. Centers for Disease Control and Prevention. (2021). *Increase in developmental disabilities among children in the United States*. <https://www.cdc.gov/ncbddd/developmentaldisabilities/features/increase-in-developmental-disabilities.html>
- 60. Centers for Disease Control and Prevention. (2024). *Developmental disabilities*. <https://www.cdc.gov/environmental-health-tracking/php/data-research/developmental-disabilities.html>
- 61. Centers for Disease Control and Prevention. (2024). *Developmental disabilities*. <https://www.cdc.gov/environmental-health-tracking/php/data-research/developmental-disabilities.html>



Read more information about autism in the “[Autism Spectrum Disorder](#)” section and ADHD in the “Attention Disorder and Hyperactivity Disorder” subsection presented earlier in this section.

## Causes and Risk Factors

Developmental disabilities begin anytime during the developmental period and usually last throughout a person’s lifetime. Most developmental disabilities begin before a baby is born, but some can occur after birth because of injury, infection, or other factors. Many developmental disabilities are thought to be caused by a complex mix of factors including genetics, parental health and behaviors (such as maternal infections or substance use during pregnancy), complications during birth, infections the baby had very early in life, or exposure of the mother or child to high levels of environmental toxins, such as lead. However, there is no known cause for most developmental disabilities.<sup>62</sup>

## Diagnosis of Developmental Disabilities

Developmental disabilities are diagnosed by developmental monitoring and developmental screening through a partnership between parents and health care professionals as a child’s growth and development are monitored.

Every child should receive routine screenings for developmental delays at their well-child visits. During a well-child visit, the provider performs **developmental monitoring** by observing for signs of developmental delays and talking with parents about any concerns they might have about their child’s growth and development. If any problems are noted during

62. Centers for Disease Control and Prevention. (2024). *Developmental disabilities*. <https://www.cdc.gov/environmental-health-tracking/php/data-research/developmental-disabilities.html>

developmental monitoring, **developmental screening** is performed. Standardized tools used during developmental screening are formal questionnaires or checklists based on research that ask questions about a child's development, including language, movement, thinking, behavior, and emotions. Developmental screening can be done by a doctor or nurse but may also be performed by other professionals in health care, early childhood education, community, or school settings to determine if a child is learning as expected or if there are delays.<sup>63</sup>

View an image of developmental milestones in Figure 12.3.<sup>64</sup>

63. Centers for Disease Control and Prevention. (2024). *Developmental disabilities*. <https://www.cdc.gov/environmental-health-tracking/php/data-research/developmental-disabilities.html>

64. "Vroom-Poster\_14x8.5\_FNL-508" by National Center on Birth Defects and Developmental Disabilities, Centers for Disease Control and Prevention is in the [Public Domain](#)



Figure 12.3 Developmental Milestones

- View the [CDC's Developmental Milestones and Milestone Checklists PDF](#).

## Types of Disorders

Based on developmental screenings, children may require further evaluation for language and speech disorders, learning disorders, and intellectual disabilities.

## LANGUAGE AND SPEECH DISORDERS

Some children struggle with understanding language or speaking, and if they do not reach expected developmental milestones, it may be a sign of a

language or speech disorder. Early childhood—especially the first three years—is a critical period when the brain rapidly develops these skills. A rich environment filled with sounds, sights, and regular communication best supports this development.<sup>65</sup>

Language development has many components. Children might have problems with one or more of the following<sup>66</sup> :

- **Receptive Language:** Difficulty understanding what others say due to not hearing the words (hearing loss) or not understanding the meaning of the words.
- **Expressive Language:** Difficulty communicating thoughts using language due to not knowing the words to use, not knowing how to put words together, or not being able to express the words.

Common indicators of language and speech disorders include<sup>67</sup> :

- **Articulation Errors:** Challenges forming specific words or sounds correctly
- **Speech Fluency Issues:** Problems with the smooth flow of words or sentences (e.g., stuttering)
- **Auditory Processing Challenges:** Struggles with interpreting the

65. Centers for Disease Control and Prevention. (2021). *Language and speech disorders in children*. <https://www.cdc.gov/nchs/products/databriefs/db205.htm>

66. Centers for Disease Control and Prevention. (2021). *Language and speech disorders in children*. <https://www.cdc.gov/nchs/products/databriefs/db205.htm>

67. Centers for Disease Control and Prevention. (2021). *Language and speech disorders in children*. <https://www.cdc.gov/nchs/products/databriefs/db205.htm>

meaning of sounds

- **Apraxia of Speech:** Inability to sequence sounds and syllables correctly to form words

Children who struggle with language may become frustrated, which can lead to acting out, appearing helpless, or withdrawing from social interactions. Language or speech disorders may also occur alongside conditions such as ADHD, anxiety, or autism.<sup>68</sup>

If a child's speech or language appears to be delayed, it is important to first determine whether a hearing loss is contributing to the problem. Hearing loss may be subtle—for instance, it might affect only one ear or certain sound frequencies. A doctor may refer the child to a speech-language pathologist, who will evaluate the child's communication skills using spoken tests and conduct a hearing assessment. Based on this evaluation, the specialist can recommend activities to support language development at home, individual or group therapy, or further evaluation by an audiologist or developmental psychologist.<sup>69</sup>

- ▶ For a detailed overview of what is considered typical for each age group, see the [Hearing and Communicative Development Checklist](#). This checklist outlines key milestones from birth to

68. Centers for Disease Control and Prevention. (2021). *Language and speech disorders in children*. <https://www.cdc.gov/nchs/products/databriefs/db205.htm>

69. Centers for Disease Control and Prevention. (2021). *Language and speech disorders in children*. <https://www.cdc.gov/nchs/products/databriefs/db205.htm>

- ▶ five years of age and serves as a guide for identifying potential delays.

## LEARNING DISORDERS

Many children may struggle with learning certain topics or skills in school from time to time. However, when children struggle with learning specific skills over time, it can be a sign of a learning disorder. Having a **learning disorder** means that a child has difficulty in one or more areas of learning, even when their overall intelligence or motivation are not affected.<sup>70</sup>

Some symptoms of learning disorders are as follows<sup>71</sup>:

- Difficulty telling right from left
- Reversing letters, words, or numbers, after the first or second grade
- Difficulties recognizing patterns or sorting items by size or shape
- Difficulty understanding and following instructions or staying organized
- Difficulty remembering what was just said or what was just read
- Lacking coordination when moving around
- Difficulty doing tasks with the hands, like writing, cutting, or drawing
- Difficulty understanding the concept of time

70. Centers for Disease Control and Prevention. (2021). *Learning disorders in children*. <https://www.cdc.gov/ncbddd/childdevelopment/learning-disorder.html>

71. Centers for Disease Control and Prevention. (2021). *Learning disorders in children*. <https://www.cdc.gov/ncbddd/childdevelopment/learning-disorder.html>

Examples of learning disorders include the following<sup>72</sup>:

- **Dyslexia:** Difficulty with reading
- **Dyscalculia:** Difficulty with math
- **Dysgraphia:** Difficulty with writing

Children with learning disorders often experience frustration, which may lead to acting out, withdrawing, or exhibiting low self-esteem. These challenges can coexist with conditions such as ADHD or anxiety, further complicating academic success. Early detection and specialized instruction—such as individualized education programs (IEPs), targeted tutoring, or classroom accommodations—are key to helping these children overcome their difficulties and achieve their full potential. If difficulties persist for at least six months despite additional support, an evaluation by educational and health professionals is recommended.<sup>73</sup>

## INTELLECTUAL DISABILITIES

**Intellectual disability** is defined in the *Diagnostic and Statistical Manual of Mental Disorders (DSM5)* as a disorder with onset during the developmental period that includes both intellectual and adaptive functioning deficits in conceptual, social, and practical domains. The following criteria must be met for an individual to be diagnosed with an intellectual disability<sup>74</sup>:

72. Centers for Disease Control and Prevention. (2021). *Learning disorders in children*. <https://www.cdc.gov/ncbddd/childdevelopment/learning-disorder.html>

73. Centers for Disease Control and Prevention. (2021). *Learning disorders in children*. <https://www.cdc.gov/ncbddd/childdevelopment/learning-disorder.html>

74. American Psychiatric Association. (2013). *Desk reference to the diagnostic criteria from DSM-5*.

- Deficits in intellectual functions, such as reasoning, problem-solving, planning, abstract thinking, judgment, academic learning, and learning from experience that is confirmed by both clinical assessment and individualized, standardized intelligence testing.
- Deficits in adaptive functioning that result in failure to meet developmental and sociocultural standards for personal independence and social responsibility. Without ongoing support, the adaptive deficits limit functioning in one or more activities of daily life, such as communication, social participation, and independent living, across multiple environments such as home, school, work, and the community.
- Onset of intellectual and adaptive deficits occurs during the developmental period.

Levels of intellectual disability vary greatly in children. Children with an intellectual disability might have a hard time communicating their needs, and an intellectual disability can cause them to develop more slowly than other children of the same age. Intellectual disability can be caused by a problem that starts any time before birth to when a child turns 18 years old. It can be caused by injury, disease, or other dysfunction in the brain. For many children, the cause of their intellectual disability is not known.<sup>75</sup>

The more severe the degree of intellectual disability, the earlier the signs can be noticed during developmental monitoring, such as the following<sup>76</sup> :

- Sitting up, crawling, or walking later than other children
- Talking later than other children or having trouble speaking

75. Centers for Disease Control and Prevention. (2021). *Facts about intellectual disabilities in children* [PDF]. [https://www.cdc.gov/ncbddd/actearly/pdf/parents\\_pdfs/intellectualdisability.pdf](https://www.cdc.gov/ncbddd/actearly/pdf/parents_pdfs/intellectualdisability.pdf)

76. Centers for Disease Control and Prevention. (2021). *Facts about intellectual disabilities in children* [PDF]. [https://www.cdc.gov/ncbddd/actearly/pdf/parents\\_pdfs/intellectualdisability.pdf](https://www.cdc.gov/ncbddd/actearly/pdf/parents_pdfs/intellectualdisability.pdf)



- Difficulty remembering things
- Difficulty understanding social rules
- Difficulty seeing the results of their actions
- Difficulty solving problems

The severity of intellectual disability (ID) varies widely and is classified based on the level of support an individual needs in daily life. Both the DSM-5 and the American Association on Intellectual and Developmental Disabilities (AAIDD) describe four levels of severity: mild, moderate, severe, and profound.<sup>77</sup>

Most individuals with ID fall into the mild to moderate range. These individuals typically develop conceptual, social, and daily living skills more slowly than their peers. However, with minimal to moderate support, they can learn practical life skills, communicate effectively, and often live semi-independently. Challenges may include difficulty with reading, managing money, or understanding time, but many can succeed in school and work with proper accommodations.

Severe intellectual disability is characterized by major developmental delays. Individuals in this category may understand some spoken language but have limited verbal communication skills. They often require significant support and supervision in their daily routines and may need help with hygiene, safety, and social interaction. A structured and supportive environment is essential for their well-being.

Profound ID is typically associated with congenital or genetic syndromes. These individuals require extensive support in nearly all areas of life. They have very limited communication abilities and often rely on gestures, sounds, or assistive technology to interact. Around-the-clock care is usually needed, and they benefit greatly from sensory-based therapies and consistent routines.

<sup>77</sup>. American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). <https://doi.org/10.1176/appi.books.9780890425596>

Signs of more severe intellectual disability often appear earlier in childhood. These may include delays in motor milestones (such as sitting, crawling, or walking), limited speech or language development, problems with memory, and difficulty understanding social rules or solving problems. Early identification and intervention are key to helping children with ID reach their full potential.<sup>78</sup>

Intellectual disabilities (ID) can arise from a complex interplay of genetic and environmental factors. Genetic conditions, such as Down syndrome and fragile X syndrome, are well-known contributors. Environmental influences—ranging from prenatal exposures to harmful substances, poor maternal nutrition, and birth complications—also play a significant role. Despite advances in research, the exact cause of an intellectual disability is not always identifiable, making comprehensive assessment essential for effective support planning.

Assessment of ID typically includes standardized tests to measure intellectual functioning, such as the Wechsler Intelligence Scale for Children, as well as evaluations of adaptive behavior, like those conducted with the Vineland Adaptive Behavior Scales. These tools help professionals understand how well an individual manages everyday tasks in comparison to their peers. The results not only confirm a diagnosis but also inform the level and type of support a person may require across various settings.

Treatment for intellectual disabilities centers around three primary areas. First, addressing underlying causes is essential in cases where conditions like metabolic disorders (e.g., phenylketonuria) can be managed through specific interventions such as dietary adjustments. Second, treating comorbid medical or psychiatric conditions is critical. Many individuals with ID also experience issues like epilepsy, attention-deficit/hyperactivity disorder

78. Centers for Disease Control and Prevention. (2022). *Facts about intellectual disability*. <https://www.cdc.gov/ncbddd/developmentaldisabilities/facts-about-intellectual-disability.html>

(ADHD), or anxiety, which may be managed through medications and psychosocial support. Third, enhancing adaptive skills is a major focus. Interventions may include early behavioral therapy, specialized education programs, and life skills training designed to promote independence and improve day-to-day functioning.

Treatment plans for individuals with ID are highly personalized and must adapt over time as needs change. While intellectual disability is typically a lifelong condition, early and targeted interventions have been shown to significantly enhance a person's quality of life, communication abilities, and capacity for independence. Multidisciplinary support and family involvement are also key factors in achieving the best possible outcomes.

Children who are suspected to have an intellectual disability based on developmental screening are referred to a developmental pediatrician or other specialist for treatment.

## Tics and Tourette Syndrome

Tourette syndrome (TS) and other tic disorders affect approximately 1% of school-aged children in the United States.<sup>79</sup> **Tics** are sudden twitches, movements, or sounds that people do repeatedly with the inability to stop their body from doing these actions. There are two types of tics: motor and vocal. **Motor tics** are movements of the body such as blinking, shrugging the shoulders, or jerking an arm. **Vocal tics** are sounds that a person makes with his or her voice such as grunting, humming, clearing the throat, or yelling out a word or phrase. Although the media often portray people with TS as involuntarily shouting out swear words (i.e., coprolalia) or constantly repeating the words of other people (i.e., echolalia), these symptoms are rare.<sup>80</sup>

79. Tourette Association of American. (n.d.). *What is Tourette*. <https://tourette.org/about-tourette/overview/what-is-tourette/#1461071628539-4f1f68e0-cd8d>

80. Centers for Disease Control and Prevention. (2021). *What is Tourette*

The primary symptom of Tourette syndrome (TS) is tics that typically begin when a child is 5 to 10 years old. The first symptoms are often motor tics that occur in the head and neck area. Tics are often worse during times that are stressful or exciting and tend to improve when a person is calm or focused on an activity.<sup>81</sup>

In most cases, tics decrease during adolescence and early adulthood, and sometimes disappear entirely. However, many people with TS experience tics into adulthood and, in some cases, tics can become worse during adulthood.<sup>82</sup> Many individuals with TS also experience ADHD, obsessive-compulsive behaviors, anxiety, learning disabilities, behavioral issues, sleep disturbances, sensory processing issues or social difficulties. These co-occurring conditions may have a greater impact on daily life than the tics themselves. In educational settings, tailored accommodations such as private study areas, modified testing environments, or individualized instruction can be very beneficial.

Tics are typically mild and do not require treatment, but it is essential to educate the individual and others about TS and provide appropriate support across all settings (e.g., school, work, and home). When tics become

*syndrome?* [https://www.cdc.gov/tourette-syndrome/about/?CDC\\_AAref\\_Val=https://www.cdc.gov/ncbddd/tourette/facts.html](https://www.cdc.gov/tourette-syndrome/about/?CDC_AAref_Val=https://www.cdc.gov/ncbddd/tourette/facts.html)

81. Centers for Disease Control and Prevention. (2021). *What is Tourette syndrome?* [https://www.cdc.gov/tourette-syndrome/about/?CDC\\_AAref\\_Val=https://www.cdc.gov/ncbddd/tourette/facts.html](https://www.cdc.gov/tourette-syndrome/about/?CDC_AAref_Val=https://www.cdc.gov/ncbddd/tourette/facts.html)

82. Centers for Disease Control and Prevention. (2021). *What is Tourette syndrome?* [https://www.cdc.gov/tourette-syndrome/about/?CDC\\_AAref\\_Val=https://www.cdc.gov/ncbddd/tourette/facts.html](https://www.cdc.gov/tourette-syndrome/about/?CDC_AAref_Val=https://www.cdc.gov/ncbddd/tourette/facts.html)

problematic or interfere with daily functioning, behavioral treatment or medication may be considered.<sup>83</sup>

 View the following YouTube video of individuals with tics<sup>84</sup>:  
[Tourette Syndrome is...](#)

► To read more about Tourette syndrome and tics, go to [Tourette Association of America](#).

## Substance Use Disorders

Children and adolescents may use alcohol and other substances for many reasons that may include maladaptive coping strategies or for other reasons. Alcohol is the most commonly used substance among young people in the United States. Data from several national surveys document frequent use of alcohol among young people. The 2019 Youth Risk Behavior Survey found these statistics among high school students during the past 30 days<sup>85</sup>:

83. Tourette Association of American. (n.d.). *What is Tourette*. <https://tourette.org/about-tourette/overview/what-is-tourette/#1461071628539-4f1f68e0-cd8d>

84. Tourette Association of American. (2018, January 24). *Tourette Syndrome is...* [Video]. YouTube. All rights reserved. <https://youtu.be/M8clZP-Pl2Y>

85. Centers for Disease Control and Prevention. (2021). *Underage drinking*. [https://www.cdc.gov/alcohol/underage-drinking/?CDC\\_AAref\\_Val=https://www.cdc.gov/alcohol/fact-sheets/underage-drinking.htm](https://www.cdc.gov/alcohol/underage-drinking/?CDC_AAref_Val=https://www.cdc.gov/alcohol/fact-sheets/underage-drinking.htm)

- 22% Drank alcohol
- 16% Rode with a driver who has been drinking alcohol
- 5% Drove after drinking alcohol
- 17% Used electronic vapor products
- 17% Used marijuana
- 4% Smoked cigarettes

**Binge drinking** is defined as a pattern of alcohol consumption that brings the blood alcohol concentration level to 0.08% or more within about two hours. This pattern of drinking usually corresponds to five or more drinks on a single occasion for males or four or more drinks on a single occasion for females within about two hours.

Youth who binge drink alcohol are more likely to experience these issues<sup>86</sup>:

- School problems, such as higher rates of absences or lower grades
- Social problems, such as fighting or lack of participation in youth activities
- Legal problems, such as arrest for driving or physically hurting someone while drunk
- Physical problems, such as hangovers or illnesses
- Unwanted, unplanned, and unprotected sexual activity
- Disruption of normal growth or sexual development
- Physical and sexual violence
- Increased risk of suicide and homicide
- Alcohol-related motor vehicle crashes and other unintentional injuries, such as burns, falls, or drowning
- Memory problems
- Misuse of other substances

86. Centers for Disease Control and Prevention. (2021). *Underage drinking*. [https://www.cdc.gov/alcohol/underage-drinking/?CDC\\_AAref\\_Val=https://www.cdc.gov/alcohol/fact-sheets/underage-drinking.htm](https://www.cdc.gov/alcohol/underage-drinking/?CDC_AAref_Val=https://www.cdc.gov/alcohol/fact-sheets/underage-drinking.htm)

- Changes in brain development that may have lifelong effects
- Alcohol poisoning

**Substance use disorders** occur when the recurrent use of alcohol and/or drugs causes clinically significant impairment, including health problems, disability, and failure to meet major responsibilities at work, school, or home. The 2019 National Survey on Drug Use and Health reports that approximately 20.3 million people aged 12 or older had a substance use disorder in the past year.<sup>87</sup>

The Substance Abuse and Mental Health Services Administration (SAMHSA) has developed a national campaign called “Talk. They Hear You” to help parents and caregivers start talking with their children early about the dangers of alcohol and other drugs.<sup>88</sup> See Figure 12.4<sup>89</sup> for an example of a public service announcement from the “Talk. They Hear You” campaign.

87. Substance Abuse and Mental Health Services Administration. (2021). *Prevention of substance use and mental disorders*. <https://www.samhsa.gov/find-help/prevention>
88. Substance Abuse and Mental Health Services Administration. (2021). *Prevention of substance use and mental disorders*. <https://www.samhsa.gov/find-help/prevention>
89. “TTHY-friends-square-2020” by Substance Abuse and Mental Health Services Administration (SAMHSA) is in the Public Domain.





Figure 12.4 Image from "Talk, They Hear You" Campaign

The vulnerability to substance use disorders is shaped by a complex interplay of factors that include genetics, early experiences, environmental influences, mental health, and family relationships. A family history of substance use can increase risk due to inherited traits and learned behaviors. Initiating substance use at a young age also raises the likelihood of developing a disorder, as early exposure can interfere with brain development and decision-making. Easy access to drugs or alcohol, along with widespread advertising, can lower barriers and increase temptation, especially in communities where substance use is normalized. Mental health challenges, such as anxiety or depression, often contribute to the risk as individuals may use substances to cope with emotional distress. Additionally, family environments characterized by low parental supervision, frequent conflict, or histories of abuse and neglect can further heighten vulnerability by reducing protective supports and increasing emotional instability.



Read more information in the “[Substance Use Disorders](#)” chapter.

## Gender Dysphoria

Distinctions between gender, sexual orientation, and sexual behavior are a critical concept in adolescent health and are greatly influenced by one’s culture. Sex is assigned at birth based on the medical assessment of genitalia. Anatomical characteristics and chromosomes determine whether a person is biologically male or female. Gender identity, gender roles, and gender expression are psychological and cultural constructs referring to various aspects of maleness, femaleness, or other nonbinary designation.<sup>90</sup>

**Gender identity** is an individual’s innate sense of being male, female, androgynous (i.e., of indeterminate sex), nonbinary (i.e., a blend of both genders or don’t identify with either gender), or a preference to reject gender designation. An individual’s gender identity is generally established during early childhood but may evolve across their lifespan. **Gender roles** are social constructs based on masculinity and femininity that embody one’s culture’s expectations, attitudes, behaviors, and personality traits based on one’s biological sex. **Gender expression** refers to how an individual presents one’s gender to the outside world, but it does not necessarily correlate with their gender identity.<sup>91</sup>

An individual’s gender identity is influenced during early childhood by one’s parents and immediate family members defining how a person expresses themselves as members of their gender. As children become adolescents,

90. Forcier, M. (2021). Adolescent sexuality. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

91. Forcier, M. (2021). Adolescent sexuality. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

their influences broaden with peer, media, and community norms of gender and sexuality impacting their individual value systems.<sup>92</sup>

When a person's biological sex differs from their gender identity, they experience feelings of unease about their incongruent maleness or femaleness referred to as **gender dysphoria**. For example, a biologic male with an innate sense of being female may describe himself as “a woman trapped in a man's body.”<sup>93</sup> People who are **transgender** have a gender identity or gender expression that differs from the sex they were assigned at birth. See Figure 12.5<sup>94</sup> for an image of a person displaying a sign at a rally for transgender equality.



Figure 12.5 Transgender Equality Rally

92. Forcier, M. (2021). Adolescent sexuality. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

93. Forcier, M. (2021). Adolescent sexuality. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

94. “[2013\\_Rally\\_for\\_Transgender\\_Equality\\_21175.jpg](#)” by [Ted Eytan](#) is licensed under [CC BY-SA 2.0](#)

According to the *DSM-5*, at least six of the following symptoms must be exhibited for at least six months and associated with distress or impaired functioning in order for a child to be diagnosed with gender dysphoria<sup>95</sup> :

- A strong desire to be of the other gender or an insistence they are another gender
- A strong preference for dressing in clothing stereotypical of the other gender
- A strong preference for playing with toys, games, or activities stereotypical of the other gender
- A strong rejection of the toys, games, and activities stereotypical of their assigned gender
- A strong preference for cross-gender roles when playing
- A strong preference for playmates of the other gender
- A strong dislike of one's sexual anatomy
- A strong desire to have the primary and/or secondary sex characteristics of the other gender

Only a small percentage of children who display gender dysphoria will continue to show these characteristics into adolescence or adulthood. Adolescents with gender dysphoria who dread the appearance of secondary sexual characteristics may seek hormones or surgery to alter their masculinity or femininity.<sup>96</sup>

**Sexual orientation** is different from gender identity and gender expression.

95. American Psychiatric Association. (2013). *Desk reference to the diagnostic criteria from DSM-5*.

96. Forcier, M. (2021). Adolescent sexuality. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

Sexual orientation refers to an individual's pattern of physical, emotional, and romantic arousal (including fantasies, activities, and behaviors) and the gender(s) of persons to whom an individual is physically or sexually attracted. An individual's assessment of their sexual orientation is termed **sexual identity**. Formation of sexual identity may be fluid with experimentation with same-gender sexual contacts as part of adolescent development. Approximately 5 to 10 percent of teens identify as lesbian, gay, or bisexual.<sup>97</sup>

Adolescents and adults who are mature and healthy in their sexuality are able to do the following<sup>98</sup> :

- Take responsibility for one's own behavior
- Practice effective decision-making
- Affirm that human development includes sexual development, which may or may not include reproduction or sexual experience
- Seek further information about sexuality and reproduction as needed and make informed choices about family options and relationships
- Interact with all genders in respectful and appropriate ways
- Affirm one's own gender identity and sexual orientation and respect the gender identities and sexual orientations of others
- Appreciate one's body and enjoy one's sexuality throughout life, expressing one's sexuality in ways that are congruent with one's values
- Express love and intimacy in appropriate ways
- Develop and maintain meaningful relationships, avoiding exploitative or manipulative relationships
- Exhibit skills and communication that enhance personal relationships with family, peers, and romantic partners

Health risks and adverse outcomes can occur more readily among youth who are gender-diverse or from sexual minorities. Risks and adverse outcomes can

97. Forcier, M. (2021). Adolescent sexuality. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

98. Forcier, M. (2021). Adolescent sexuality. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

include child abuse, bullying, sexual harassment, teen dating violence, unprotected sex with risks for sexually transmitted infections and pregnancy, mental health problems (depression, anxiety, suicide, and disordered eating and body image), and substance use.<sup>99</sup>

Nurses should ask all clients, including adolescents, about their gender preferences and provide support if the client indicates a need for help. By promoting sexuality as healthy, respectful, and meaningful in the global context of adolescent development, nurses can encourage a positive model of empowerment for youth exploring their gender and sexual identities.<sup>100</sup>

99. Forcier, M. (2021). Adolescent sexuality. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

100. Forcier, M. (2021). Adolescent sexuality. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

## 12.3 Psychological and Behavioral Therapy

### Treatment of Childhood and Adolescent Mental and Neurodevelopmental Disorders

Mental, emotional, and behavioral disorders in childhood can cause long-term problems. Early treatment of a child's mental health or behavioral problems can reduce problems at home, in school, in forming friendships, and can promote healthy development into adulthood.<sup>1</sup> Treatment for mental health disorders, neurodevelopmental disorders, and behavioral disorders include psychological therapy, parent training in behavioral management, and behavioral interventions for the classroom.

### Psychological Therapies

Depending on the type and severity of the mental health disorder and the developmental age of the child, psychological therapy for children may be used in combination with prescribed medication. Psychological therapy helps a child manage their symptoms so that they can function well at home, in school, and in their community. For example, cognitive-behavioral therapy (CBT) works well for children with disruptive behavior disorders, depression, and anxiety.

Psychological therapy is conducted by trained mental health professionals for children and can be done individually or in groups. Psychological therapy with children can include talking, playing, or other activities to help the child express feelings and thoughts. Therapists may also observe parents and children together and make suggestions for different ways to respond to

1. Centers for Disease Control and Prevention. (2025). *Treating children's mental health with therapy*. [https://www.cdc.gov/children-mental-health/treatment/?CDC\\_AAref\\_Val=https://www.cdc.gov/childrensmentalhealth/parent-behavior-therapy.html](https://www.cdc.gov/children-mental-health/treatment/?CDC_AAref_Val=https://www.cdc.gov/childrensmentalhealth/parent-behavior-therapy.html)

disruptive behaviors.<sup>2</sup> See Table 12.3 for a list of psychological therapies considered effective for common conditions in children and adolescents.<sup>3</sup>

Table 12.3 Psychological Therapies for Various Conditions in Children and Adolescents

2. Centers for Disease Control and Prevention. (2025). *Treating children's mental health with therapy*. [https://www.cdc.gov/children-mental-health/treatment/?CDC\\_AAref\\_Val=https://www.cdc.gov/childrensmentalhealth/parent-behavior-therapy.html](https://www.cdc.gov/children-mental-health/treatment/?CDC_AAref_Val=https://www.cdc.gov/childrensmentalhealth/parent-behavior-therapy.html)
3. Hilt, R. J., & Nussbaum, A. M. (Eds.). (2016). *DSM-5 pocket guide for child and adolescent mental health*. <https://doi.org/10.1176/appi.books.9781615370511>

Therapies	Description	Indications
<b>Cognitive-Behavioral Therapy (CBT)</b>	<p>Teaches clients how to correct cognitive errors in thinking (such as, “nothing ever goes right for me”) and coaches them to try different behaviors (i.e., behavioral activation) to lead to changes in how they feel.</p> <p>Exposure and response prevention (ERP) is a form of CBT for the treatment of obsessive-compulsive disorder (OCD).</p>	<p>Anxiety disorders</p> <p>Depressive disorders</p> <p>Oppositional defiant disorder</p> <p>Eating disorders</p> <p>Substance use disorder</p> <p>Post-traumatic stress disorder</p>
<b>Dialectical Behavior Therapy (DBT)</b>	<p>Specialized version of CBT with skills groups (to teach problem-solving, emotional regulation, distress tolerance, and interpersonal effectiveness skills), as well as individual therapy sessions. Mindfulness and meditation are often encouraged.</p>	<p>Chronic and significant suicide ideation and/or self-harm</p>
<b>Family Therapy</b>	<p>Sessions that focus on the family relationship or interaction patterns that cause dysfunction and assist the family members to amend that pattern (rather than saying the problem resides within an individual).</p>	<p>Eating disorders</p> <p>Conduct disorder</p> <p>Depressive disorders</p> <p>Substance use disorders</p>
<b>Group Therapy</b>	<p>Addresses interaction pattern problems while also providing disorder-specific support within a group of strangers having similar challenges. Peer-based learning can be uniquely effective.</p>	<p>Anxiety disorders</p> <p>Depression</p>



<b>Behavioral Management Training</b>	Also known as parent management training, these programs teach and encourage skillful parent or caregiver responses to challenging child behaviors. Read more in the “Behavioral Interventions” subsection below.	Oppositional defiant disorder  Conduct disorder
<b>Social Skills Training</b>	Variety of group and one-to-one techniques to teach basic behavioral and cognitive skills, reinforce prosocial behaviors, and teach social problem-solving. Group sessions are more beneficial than individual sessions because of peer learning influences.	Oppositional defiant disorder  Attention deficit hyperactivity disorder (ADHD)  Anxiety disorders  Depressive disorders  Autism spectrum disorder
<b>Relaxation Training</b>	Biofeedback, deep breathing, progressive muscle relaxation, and mindfulness are examples of strategies to increase mind-body awareness and enhance the ability to effectively calm emotional reactions. Read more about relaxation training in the “ <a href="#">Stress</a> ” and “ <a href="#">Coping</a> ” sections of the “Stress, Coping, and Crisis Intervention” chapter.	Anxiety disorders  Depressive disorders

<b>Motivational Interviewing (MI)</b>	A specific type of therapeutic communication regarding health behaviors a client needs to change but has significant reluctance. It helps clients state their own reasons for changing behaviors, resolve their own ambivalence, and state actions they can take to change. Read more details in the “ <a href="#">Motivational Interviewing</a> ” section of the “Therapeutic Communication and Nurse-Client Relationship” chapter.	Substance use disorders
---------------------------------------	--	-------------------------

## Behavioral Interventions

Behavioral interventions reward desired behaviors and reduce maladaptive coping behaviors. Most child and adolescent treatment settings used structured programs to motivate and reward age-appropriate behaviors. For example, the point or star system may be used where the child receives points or stars for desired behaviors, and then specific privileges are awarded based on the points or stars earned each day.<sup>4</sup>

## Parent Training in Behavior Management

Parent training in behavior management includes teaching skills and strategies such as positive reinforcement, structure, and consistent discipline to manage their child’s behavior and help them succeed at school, at home, and in relationships.<sup>5</sup>


4. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

5. Centers for Disease Control and Prevention. (2024). *About attention-deficit/*

See Figure 12.6<sup>6</sup> for an illustration of parent training in behavioral management.

### What parents can expect in behavior therapy

Parents typically attend 8-16 sessions with a therapist and learn strategies to help their child. Sessions may involve groups or individual families.



#VitalSigns

**Vital**signs<sup>™</sup>  
www.cdc.gov/vitalsigns/adhd



Figure 12.6 Parent Training in Behavior Management

## BEHAVIORAL MANAGEMENT STRATEGIES

There are several evidence-based strategies that nurses can teach parents and caregivers to help manage behaviors of children and adolescents, such as

*hyperactivity disorder (ADHD).* [https://www.cdc.gov/adhd/about/?CDC\\_AAref\\_Val=https://www.cdc.gov/ncbddd/adhd/facts.html](https://www.cdc.gov/adhd/about/?CDC_AAref_Val=https://www.cdc.gov/ncbddd/adhd/facts.html)

6. "adhd-behavior-therapy-parents-800px.png" by National Center on Birth Defects and Developmental Disabilities is in the Public Domain

Time-Out, Special Time, Functional Analysis of Behavior, Behavioral Activation, Sleep Hygiene, and Responding to Bullying.<sup>7</sup>

## Time-Out

“Time-out” is a strategy for shaping a child’s behavior through selective and temporary removal of the child’s access to desired attention, activities, or other reinforcements following a behavioral transgression. This strategy works for children who experience regular positive praise and attention from their parents or caregivers because they feel motivated to maintain that positive regard. The length of time should be about one minute for each year of age, but adjustments need to be made based on the child’s developmental level. For example, children with developmental delays should have shorter durations.<sup>8</sup>

Tips for caregivers implementing time-outs include the following<sup>9</sup>:

- Set consistent limits to avoid confusion.
- Focus on changing priority misbehaviors rather than everything at once.
- After setting a “time-out,” decline further verbal engagement until a “time-in.”
- Ensure time-outs occur immediately after misbehavior rather than being delayed.
- Follow through if using warnings (e.g., “I’m going to count to three...”).
- State when the time-out is over. Setting a timer can be helpful.

7. Hilt, R. J., & Nussbaum, A. M. (Eds.). (2016). *DSM-5 pocket guide for child and adolescent mental health*. <https://doi.org/10.1176/appi.books.9781615370511>

8. Hilt, R. J., & Nussbaum, A. M. (Eds.). (2016). *DSM-5 pocket guide for child and adolescent mental health*. <https://doi.org/10.1176/appi.books.9781615370511>

9. Hilt, R. J., & Nussbaum, A. M. (Eds.). (2016). *DSM-5 pocket guide for child and adolescent mental health*. <https://doi.org/10.1176/appi.books.9781615370511>

- When the time-out finishes, congratulate the child on regaining personal control and then look for the next positive behavior to praise.
- Give far more positive attention than negative attention.

## Special Time

“Special time” is a strategy for a caregiver and a young child to establish the enjoyment of each other’s company. It is also referred to as “child-directed play” because it emphasizes that caregivers follow the child’s lead. Tips for caregivers implementing special time include the following<sup>10</sup> :

- Commit to setting aside a regular time for “special time.” Daily is best, but two to three times a week consistently also works.
- Select the time of day and label it as “our special time.”
- Choose a time short enough that it can happen reliably, usually 15-30 minutes. Ensure this time happens no matter how good or bad the day’s behaviors were.
- Allow the child to select the activity, which must be something you do not actively dislike or does not involve spending money or completing a chore.
- Follow the child’s lead during play, resisting the urge to tell them what to do.
- End of time; a timer may be helpful. Remind the child when the next special time will be.
- If the child refuses at first, explain you will just sit with them during the “special time.”
- Expect greater success if you set your own special times for yourself, too.

10. Hilt, R. J., & Nussbaum, A. M. (Eds.). (2016). *DSM-5 pocket guide for child and adolescent mental health*. <https://doi.org/10.1176/appi.books.9781615370511>

## Functional Analysis of Behavior

Functional analysis is a strategy for preventing a recurring problematic behavior by first identifying why a behavior keeps recurring and then devising a plan to prevent recurrences. For example, a parent reports their young child “throws temper tantrums every time we go to the store.” As the mental health professional helps the parent analyze the behavior, the parent realizes they have been giving the child candy to halt the tantrums, which actually functions to reward the behavior and encourages it to happen again. If the parent were to stop delivering this unintentional reward, the tantrums would theoretically decrease. Alternatively, the parent may focus on avoiding reexposing the child to a recognized trigger for the behavior, such as the candy aisle.<sup>11</sup>

## Behavioral Activation

Behavioral activation is a strategy to encourage a young person to reengage with other people and do things they find pleasurable. When an individual is sad or anxious, they are less likely to engage in activities they enjoy, and this withdrawal can cause isolation and worsen the mood. The following is a list of tips to provide clients for succeeding with behavioral activation<sup>12</sup>:

- Identify activities that you (not others) find motivating or rewarding. Develop a list with a variety of options because repetitively doing the same thing can become boring.
- Refine the list to things that can be measured as completed rather than vague goals.

11. Hilt, R. J., & Nussbaum, A. M. (Eds.). (2016). *DSM-5 pocket guide for child and adolescent mental health*. <https://doi.org/10.1176/appi.books.9781615370511>

12. Hilt, R. J., & Nussbaum, A. M. (Eds.). (2016). *DSM-5 pocket guide for child and adolescent mental health*. <https://doi.org/10.1176/appi.books.9781615370511>

- Rank the activities in order of those most easily completed to those more difficult to complete.
- Start by selecting something easy to accomplish to get started and then work up the list in increasing level of difficulty.
- Let others know your plans and enlist their help in motivating you.

## Sleep Hygiene

Insomnia is a common problem among children and adolescents. Most sleep problems can be resolved by changing habits and routines that affect sleep, commonly referred to as sleep hygiene. The following is a list of tips for caregivers for improving sleep hygiene<sup>13</sup>:

- Maintain consistent bedtimes and wake times every day of the week.
- Maintain a routine of pre-sleep activities (e.g., brush teeth, read a book).
- Avoid spending non-sleep time in or on one's bed. (i.e., "beds are for sleep").
- Ensure the bedroom is cool and quiet.
- Avoid highly stimulating activities just before bed (e.g., television, video games, social media, or exercise).
- Do not keep video games, televisions, computers, or phones in a child's bedroom. Set a media curfew on technology use.
- Incorporate exercise into the day in order to aid with sleep.
- Avoid caffeine in the afternoons and evenings. Caffeine causes shallow sleep and frequent awakenings.
- Encourage children and adolescents to discuss any worries with the caregiver before bed rather than ruminating on it during sleep time.
- Ensure that children go to bed drowsy but still awake. Falling asleep in places other than a bed forms habits that are difficult to break.
- Use security objects with young children who need a transitional object

13. Hilt, R. J., & Nussbaum, A. M. (Eds.). (2016). *DSM-5 pocket guide for child and adolescent mental health*. <https://doi.org/10.1176/appi.books.9781615370511>

with which to feel safe and secure when their caregiver is not present (e.g., a special blanket or toy)

- When checking on a young child at night, briefly reassure the child you are present and they are OK.
- Avoid afternoon naps for all but very young children because they often interfere with nighttime sleep.
- If a child or adolescent is having sleep difficulties, keep a sleep diary to track sleep time, wake time, activities, and naps to identify patterns.

## Responding to Bullying

Bullying occurs when a person seeks to harm, intimidate, or coerce someone perceived as vulnerable. If there is a sudden change in a child's mood, behavior, sleep, body symptoms, academic performance, or social functioning, there is a possibility they are experiencing bullying. Cyberbullying is a significant public health concern with rates of cyberbullying estimates ranging from 14 to 57%.<sup>14</sup>

The following is a list of tips for assessing and responding to bullying<sup>15</sup>:

- **Detect:** Ask the child: "Sometimes kids get picked on or bullied. Have you ever seen this happen? Has it ever happened to you?" If the child responds "No" but bullying is suspected, suggest the caregivers speak with teachers and/or review the child's social media accounts.
- **Educate:** Let children know that bullying is unacceptable and if they experience bullying, you will help them respond.

14. Zhu, C., Huang, S., Evans, R., & Zhang, W. (2021). Cyberbullying among adolescents and children: A comprehensive review of the global situation, risk factors, and preventive measures. *Frontiers in Public Health*, 9. <https://doi.org/10.3389/fpubh.2021.634909>

15. Hilt, R. J., & Nussbaum, A. M. (Eds.). (2016). *DSM-5 pocket guide for child and adolescent mental health*. <https://doi.org/10.1176/appi.books.9781615370511>



- **Plan:** Plan ways with the child to avoid bullying.
  - Avoid places where bullying happens.
  - Teach them to walk away when bullying occurs and tell a trusted adult who can be quickly accessed.
  - Instruct the child to stay near adults where bullying occurs. Most bullying happens when adults are not around.
  - If a child feels they can confront the bully, teach them to say in a calm, clear voice to stop the behavior and, “Bullying is not OK.”
  - If the child is comfortable with deflating the situation with humor, they can use humor to challenge the bullying.
  - Encourage the child to ask their peers for support and ideas for dealing with the bully.
  - Encourage caregivers to communicate the bullying to the school and other families and jointly devise solutions.

► Read more information about addressing bullying at [www.stopbullying.gov](http://www.stopbullying.gov).

## Behavioral Interventions for the Classroom

Behavioral interventions for the classroom help children and adolescents succeed academically. Behavioral classroom management is a teacher-led approach that encourages a student’s positive behaviors in the classroom through a reward system or a daily report card and discourages their negative behaviors. Organizational training teaches children and adolescents time management, planning skills, and ways to keep school materials organized to optimize student learning and reduce distractions.<sup>16</sup>

16. Centers for Disease Control and Prevention. (2024). *About attention-deficit/*

*hyperactivity disorder (ADHD).* [https://www.cdc.gov/adhd/about/?CDC\\_Aref\\_Val=https://www.cdc.gov/ncbddd/adhd/facts.html](https://www.cdc.gov/adhd/about/?CDC_Aref_Val=https://www.cdc.gov/ncbddd/adhd/facts.html)

## 12.4 Autism Spectrum Disorder

Autism spectrum disorder (ASD) is a neurodevelopmental disorder that affects an individual's communication and behavior. Although autism can be diagnosed at any age, it is classified as a developmental disorder because symptoms generally appear in the first two years of life. About 3% of children in the United States have ASD. ASD is reported in all racial, ethnic, and socioeconomic groups and is three times more common among boys.<sup>1</sup>

ASD is characterized by the following<sup>2</sup>:

- Difficulty with communication and interaction with other people
- Restricted interests and repetitive behaviors
- Symptoms that hurt the person's ability to function properly in school, work, and other areas of life

Autism is known as a “spectrum” disorder because there is wide variation in the type and severity of symptoms experienced by individuals. ASD occurs in all ethnic, racial, and economic groups. Although ASD is a lifelong disorder, treatments and services can improve a person's symptoms and ability to

1. National Institute of Mental Health. (2025). *Autism spectrum disorder (ASD)*. U.S. Department of Health & Human Services. [https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part\\_2547](https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part_2547)
2. National Institute of Mental Health. (2025). *Autism spectrum disorder (ASD)*. U.S. Department of Health & Human Services. [https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part\\_2547](https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part_2547)

function.<sup>3</sup> See Figure 12.7<sup>4</sup> for an image depicting a child coping with the symptoms of autism.



Figure 12.7 Child Coping With Autism Symptoms

## Signs and Symptoms

People with ASD have difficulty with social communication and interaction, experience restricted interests, and exhibit repetitive behaviors. Here are some examples of behaviors in these categories:

### **Social Communication/Interaction Behaviors<sup>5</sup>:**

- Making little or inconsistent eye contact
- Not looking at or listening to people
- Rarely sharing enjoyment of objects or activities by pointing or showing

3. National Institute of Mental Health. (2025). *Autism spectrum disorder (ASD)*. U.S. Department of Health & Human Services. [https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part\\_2547](https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part_2547)

4. “5299266366\_0b6c8ae172\_o” by [hepingting](#) is licensed under [CC BY-SA 2.0](#)

5. National Institute of Mental Health. (2025). *Autism spectrum disorder (ASD)*. U.S. Department of Health & Human Services. [https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part\\_2547](https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part_2547)

things to others

- Being slow to respond (or failing to respond) to someone calling one's name
- Having difficulties with the back-and-forth nature of a conversation
- Talking at length about a favorite subject without noticing that others are not interested or without giving others a chance to respond
- Exhibiting facial expressions, movements, and gestures that do not match what is being said
- Using a tone of voice that may sound flat and robot-like
- Having trouble understanding another person's point of view or being unable to understand other people's actions

### **Restricted Interests or Repetitive Behaviors<sup>6</sup> :**

- Repeating certain behaviors or exhibiting unusual behaviors, such as repeating words or phrases (i.e., **echolalia**)
- Having a lasting intense interest in certain topics, such as numbers, details, or facts
- Having overly focused interests, such as with moving objects or parts of objects
- Getting upset by slight changes in a routine
- Being more or less sensitive than other people to sensory input, such as light, noise, clothing, or temperature
  - This sensitivity can present as physical touch, like a hug, actually being experienced as painful. Items such as seams on pants can be overwhelming and extremely agitating.

6. National Institute of Mental Health. (2025). *Autism spectrum disorder (ASD)*. U.S. Department of Health & Human Services. [https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part\\_2547](https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part_2547)

Although people with ASD experience many challenges, they may also have many strengths, including<sup>7</sup>:

- Being able to learn things in detail and remember information for long periods of time
- Being strong visual and auditory learners
- Excelling in math, science, music, or art

## Causes and Risk Factors

Research suggests that genes can act together with influences from the environment to affect development in ways that can lead to ASD. Risk factors include the following<sup>8</sup>:

- A sibling with ASD
- Older parents at birth
- Genetic conditions such as Down syndrome and Fragile X syndrome
- Very low birth weight
- Gender (ASD is four times more common in boys than in girls)

## Treatments

Current treatments for ASD seek to reduce symptoms that interfere with daily functioning and quality of life. ASD affects each person differently, meaning that people with ASD have unique strengths and challenges and thus

7. National Institute of Mental Health. (2025). *Autism spectrum disorder (ASD)*. U.S. Department of Health & Human Services. [https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part\\_2547](https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part_2547)
8. National Institute of Mental Health. (2025). *Autism spectrum disorder (ASD)*. U.S. Department of Health & Human Services. [https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part\\_2547](https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part_2547)

different treatment needs. Treatment plans typically involve multiple professionals with interventions customized to the individual.<sup>9</sup>

As individuals with ASD exit high school and grow into adulthood, additional services can help improve their health and daily functioning and facilitate social engagement. Support may be provided to complete job training, find employment, and secure housing and transportation.<sup>10</sup>

There are many categories of treatments available, and some treatments involve more than one approach. Treatment categories include the following<sup>11</sup>:

- Behavioral
- Developmental
- Educational
- Social-Relational
- Pharmacological
- Psychological
- Complementary and Alternative

## Behavioral Approaches

Behavioral approaches focus on changing an individual's behaviors by

9. National Institute of Mental Health. (2025). *Autism spectrum disorder (ASD)*. U.S. Department of Health & Human Services. [https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part\\_2547](https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part_2547)
10. National Institute of Mental Health. (2025). *Autism spectrum disorder (ASD)*. U.S. Department of Health & Human Services. [https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part\\_2547](https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part_2547)
11. National Institute of Mental Health. (2025). *Autism spectrum disorder (ASD)*. U.S. Department of Health & Human Services. [https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part\\_2547](https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part_2547)

promoting understanding of what happens before and after the behavior. A notable behavioral treatment for people with ASD is called applied behavior analysis. Applied behavior analysis encourages desired behaviors and discourages undesired behaviors to improve a variety of skills, and progress is tracked and measured.<sup>12</sup>

## Developmental Approaches

Developmental approaches focus on improving specific developmental skills, such as language skills or physical skills. Developmental approaches are often combined with behavioral approaches. The most common developmental therapy for people with ASD is speech and language therapy. Speech and language therapy helps improve the person's understanding and use of speech and language. Some people with ASD communicate verbally. Others with severe symptoms of ASD may communicate through the use of signs, gestures, pictures, or an electronic communication device.<sup>13</sup>

Occupational therapy teaches skills to help the person live as independently as possible. Skills may include dressing, eating, bathing, and relating to other people. Occupational therapy can also include sensory integration therapy to help improve responses to sensory input that may be restrictive or overwhelming.<sup>14</sup>

12. National Institute of Mental Health. (2025). *Autism spectrum disorder (ASD)*. U.S. Department of Health & Human Services. [https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part\\_2547](https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part_2547)
13. National Institute of Mental Health. (2025). *Autism spectrum disorder (ASD)*. U.S. Department of Health & Human Services. [https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part\\_2547](https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part_2547)
14. National Institute of Mental Health. (2025). *Autism spectrum disorder (ASD)*. U.S. Department of Health & Human Services. [https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part\\_2547](https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part_2547)



Physical therapy may be implemented to help improve physical skills, such as fine movements of the fingers or larger movements of the trunk and body.<sup>15</sup>

## Educational Approaches

Educational treatments are provided in a classroom setting. One type of educational approach is the Treatment and Education of Autistic and Related Communication-Handicapped Children (TEACCH) approach. TEACCH is based on the idea that people with autism thrive on consistency and visual learning. It provides teachers with ways to adjust the classroom structure to improve academic and other outcomes. For example, daily routines can be written and placed in clear sight, boundaries can be set around learning stations, and verbal instructions can be complemented with visual instructions or physical demonstrations.<sup>16</sup>

## Social-Relational Approaches

Social-relational treatments focus on improving social skills and building emotional bonds. For example, “social stories” provide simple descriptions of what to expect in a social situation. “Social skills groups” provide opportunities for people with ASD to practice social skills in a structured environment.

## Pharmacological Approaches

There are no medications used to treat ASD, but medications may be used to

15. National Institute of Mental Health. (2025). *Autism spectrum disorder (ASD)*. U.S. Department of Health & Human Services. [https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part\\_2547](https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part_2547)

16. National Institute of Mental Health. (2025). *Autism spectrum disorder (ASD)*. U.S. Department of Health & Human Services. [https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part\\_2547](https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part_2547)

treat symptoms and improve functioning. For example, medication may be used to manage high energy levels, improve focus, or limit self-harming behavior, such as head banging or hand biting. Medication may also be used to treat concurrent psychological and medical conditions, such as anxiety, depression, seizures, or sleep problems.<sup>17</sup>


## Psychological Approaches

Psychological approaches can help people with ASD cope with anxiety, depression, and other mental health issues. For example, cognitive behavioral therapy (CBT) helps individuals focus on the connections between their thoughts, feelings, and behaviors. During CBT a therapist and the individual work together to identify goals and change how the person thinks about a situation to change how they react to the situation.<sup>18</sup>

## Complementary and Alternative Treatments

Some individuals with ASD use special diets, herbal supplements, chiropractic care, animal therapy, art therapy, mindfulness, or relaxation therapies.<sup>19</sup> Treatment is most effective when tailored to the individual, and additional enriching therapies can help individuals with ASD to thrive.

17. National Institute of Mental Health. (2025). *Autism spectrum disorder (ASD)*. U.S. Department of Health & Human Services. [https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part\\_2547](https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part_2547)
18. National Institute of Mental Health. (2025). *Autism spectrum disorder (ASD)*. U.S. Department of Health & Human Services. [https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part\\_2547](https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part_2547)
19. National Institute of Mental Health. (2025). *Autism spectrum disorder (ASD)*. U.S. Department of Health & Human Services. [https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part\\_2547](https://www.nimh.nih.gov/health/statistics/autism-spectrum-disorder-asd#part_2547)

View a supplementary YouTube video<sup>20</sup> providing the  
 perspectives of people with autism: [What's it like to be a person with autism? | In A Different Key | PBS](https://www.youtube.com/watch?v=GoieHK2vR50)

- ▶ To find resources related to caring for children with autism, go to the [Autism Society](#).

20. PBS. (2022, December 9). *What's it like to be a person with autism? | In A Different Key | PBS* [Video]. YouTube. All rights reserved.  
<https://www.youtube.com/watch?v=GoieHK2vR50>

## 12.5 Applying the Nursing Process to Mental Health Disorders in Children and Adolescents

Working with children and adolescents is very different from working with adults. Young people are often reluctant participants who have been brought for care they did not seek on their own. Additionally, their communication skills are limited based on their developmental stage. In addition to gathering information from the child, information must also be obtained from the parent or caregiver.<sup>1</sup>

The first step to successful care is to create a therapeutic nurse-client relationship. A therapeutic alliance can typically be created if the young person feels noticed, heard, and appreciated. It is often helpful to start the conversation with a relatively neutral question like, “Your mom said that you go to \_\_\_\_ school. What is that school like?” School, friends, family, and favorite activities are low-stress conversation starters. For a very young child, a conversation starter could be a simple observation like what they are wearing. For example, “I see you are wearing blue tennis shoes; did you pick those out yourself?”<sup>2</sup>

For young people who seem reluctant to start talking, it may be helpful to describe something you saw that shows you have been paying attention to them. For example, “It looked as though it was hard for you to sit and do nothing while your dad and I were talking. Am I right about that?”<sup>3</sup>

When caring for adolescents, it is helpful to gather data from the parent or

1. Hilt, R. J., & Nussbaum, A. M. (Eds.). (2016). *DSM-5 pocket guide for child and adolescent mental health*. <https://doi.org/10.1176/appi.books.9781615370511>
2. Hilt, R. J., & Nussbaum, A. M. (Eds.). (2016). *DSM-5 pocket guide for child and adolescent mental health*. <https://doi.org/10.1176/appi.books.9781615370511>
3. Hilt, R. J., & Nussbaum, A. M. (Eds.). (2016). *DSM-5 pocket guide for child and adolescent mental health*. <https://doi.org/10.1176/appi.books.9781615370511>

caregiver, and then ask to speak with the adolescent alone. Reinforce that the conversation is “conditionally confidential” and invite the adolescent to sit alone with you to talk. A 1:1 conversation with an adolescent typically creates a better therapeutic alliance with more honest answers obtained.<sup>4</sup>

A more subtle strategy to build a therapeutic-nurse relationship with children and adolescents is to shape how you speak so you are perceived as a responsive problem-solving partner rather than an authority figure who will judge them. Building a therapeutic nurse-client relationship with a young person should lead to learning their true chief complaint because the chief complaint of an adolescent may be different from their parents’ complaints.<sup>5</sup> Furthermore, goal setting and treatment plans will be more effective when the adolescent’s concerns are addressed.

## Privacy, Confidentiality, and Mandatory Reporting

In most regions, minors require parental or guardian consent for treatment. Adolescents may be able to provide informed consent independently for specific services, such as substance use, sexual health). Confidentiality should be balanced with safety and legal obligations.

Confidentiality should be discussed with the adolescent client and their parent/guardian before beginning an assessment or related conversations, and circumstances should be defined for when confidentiality is “conditional” for children and adolescents. State laws determine what information is considered confidential and what requires reporting to law enforcement or Child Protective Services. Examples of what must be reported to law enforcement include child abuse, gunshot or stabbing wounds, sexually transmitted infections, abortions, suicidal ideation, and homicidal ideation.

4. Hilt, R. J., & Nussbaum, A. M. (Eds.). (2016). *DSM-5 pocket guide for child and adolescent mental health*. <https://doi.org/10.1176/appi.books.9781615370511>

5. Hilt, R. J., & Nussbaum, A. M. (Eds.). (2016). *DSM-5 pocket guide for child and adolescent mental health*. <https://doi.org/10.1176/appi.books.9781615370511>

Some state laws make it optional for clinicians to inform parents/guardians if their child is seeking services related to sexual health care, substance abuse, or mental health care. Nurses should be aware of the state laws affecting the confidentiality of child and adolescent care in the state in which they are practicing.<sup>6</sup>

Although it is important for nurses to respect adolescent clients' privacy and confidentiality, it is also important to encourage the adolescent to talk with their parents/guardians about personal issues that affect their health even if they feel uncomfortable doing so. Parent/guardian support can help ensure the adolescent's health needs are met.<sup>7</sup>

Research your state's laws regarding adolescent/child clients' rights.

▶ Here is a link to [Wisconsin Department of Health Services' Client Rights for Minors](#).

## Assessment (Recognizing Cues)

When assessing a child or adolescent's mental health, the nurse incorporates a variety of assessments, such as the following:

- Performing a mental status examination
- Completing a psychosocial assessment including:
  - Reviewing the client's reason for seeking healthcare

6. Middleman, A. B., & Olson, K. A. (2021). Confidentiality in adolescent health care. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

7. Middleman, A. B., & Olson, K. A. (2021). Confidentiality in adolescent health care. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

- Screening the client for suicidal ideation, exposure to trauma or violence, signs of self injury, and substance misuse
- Complete a cultural assessment, spiritual assessment, and assess client family dynamics
- Use of psychotropic medications (drugs that treat psychiatric symptoms) and/or other medications that can cause psychiatric symptoms as side effects
- History of mental disorders, family history of mental illness, and previous hospitalizations
- Educational background, occupational background
- Coping mechanisms and functional ability
- Reviewing specific laboratory results related to the client's use of psychotropic and other medications
- Incorporating life span and developmental considerations

## Mental Status Examination

Specific signs and symptoms of a mental health disorder should be assessed as part of the “health history” component of a comprehensive nursing assessment. While asking questions about specific symptoms and obtaining a health history, the nurse should also be simultaneously performing a mental status examination. The mental status examination includes these items:

- Appearance and General Behavior
- Speech
- Motor Activity
- Affect and Mood
- Thoughts and Perceptions
- Attitude and Insight
- Cognitive Abilities

Review details of a comprehensive mental status examination in the

“Assessment” section of the “Application of the Nursing Process to Mental Health Care” chapter.

## Psychosocial Assessment

A **psychosocial assessment** (also referred to as a health history) is a component of the nursing assessment process that obtains additional subjective data to detect risks and identify treatment opportunities and resources. Agencies have specific forms used for psychosocial assessments/health histories that typically consist of several components<sup>8,9</sup>:

- Reason for seeking health care (i.e., “chief complaint”)
- Thoughts of self-injury or suicide
- Family dynamics
- Cultural assessment
- Spiritual assessment
- Current and past medical history
- Current medications
- History of previously diagnosed mental health disorders
- Previous hospitalizations
- Educational background
- Occupational background
- History of exposure to psychological trauma, violence, and domestic abuse

8. GW School of Medicine & Health Sciences. (n.d.). *Clinical FICA tool*.  
<https://smhs.gwu.edu/spirituality-health/program/transforming-practice-health-settings/clinical-fica-tool>

9. Glasner, J., Baltag, V., & Ambresin, A. E. (2021). Previsit multidomain psychosocial screening tools for adolescents and young adults: A system review [PDF]. *Journal of Adolescent Health*, 68, 449-459.  
[https://www.jahonline.org/article/S1054-139X\(20\)30600-5/pdf](https://www.jahonline.org/article/S1054-139X(20)30600-5/pdf)



- Substance use (tobacco, alcohol, recreational drugs, misused prescription drugs)
- Family history of mental illness
- Coping mechanisms
- Functional ability/Activities of daily living

Review details of a psychosocial assessment in the “[Assessment](#)” section of the “Application of the Nursing Process to Mental Health Care” chapter.

## REASON FOR SEEKING HEALTH CARE

Mental health symptoms in children and adolescents often manifest differently than in adults, often through behavioral changes, school difficulties, or somatic complaints. Parents and caregivers typically bring a child or adolescent in for mental health evaluation due to common concerns, such as the following:

- Poor academic performance
- Developmental delays
- Disruptive or aggressive behavior
- Withdrawn or sad mood
- Irritable or labile mood
- Anxious or avoidant behavior
- Recurrent and excessive physical complaints
- Sleep problems
- Self-harm and suicidality
- Substance abuse
- Disturbed eating

Poor academic performance is a common concern. Keep in mind that assessing a child’s/adolescent’s ability to function in school is like assessing an adult’s ability to function at work. Many factors can affect a young person’s performance at school, such as their ability and effort, the classroom environment, life distractions, or a mental health disorder.

A child's or adolescent's academic ability may be impacted by hearing or visual impairments, learning disorders, or cognitive impairments. The nurse may assist with performing vision or hearing tests or providing a developmental rating scale.

## THOUGHTS OF SUICIDE OR SELF INJURY

Suicide is a leading cause of death in adolescents. Risk may be masked by their behavior or dismissed by caregivers. As discussed in Chapter 1, all clients aged 12 and older presenting for acute care should be screened for suicidal ideation. Universal screening allows for the detection of suicide risk and implementation of early interventions before a person attempts suicide. The ASQ Suicide Risk Screening Tool is commonly used across a variety of health care settings to screen youth for suicidal ideation. ASQ questions include the following<sup>10</sup>:

1. In the past few weeks, have you wished you were dead?
2. In the past few weeks, have you felt that you or your family would be better off if you were dead?
3. In the past week, have you been having thoughts about killing yourself?
4. Have you ever tried to kill yourself?
5. Are you having thoughts of killing yourself right now?

If client answers “No” to all questions 1 through 4, screening is complete, no interventions are necessary, and it is not necessary to ask question #5. However, clinical judgment always overrides a negative screen). If the client answers “Yes” to any of questions 1 through 4, or refuses to answer, they are considered a positive screen and question #5 should be asked to assess acuity. If the client answers “yes” to question 5, they are considered high risk and require a STAT safety/mental health evaluation. They should be kept in

10. National Institute of Mental Health. (n.d.). *Youth ASQ toolkit*.  
<https://www.nimh.nih.gov/research/research-conducted-at-nimh/asq-toolkit-materials/youth-asq-toolkit>

sight at all times while a health care provider is notified. All dangerous objects should be removed from room.<sup>11</sup>

**Non-suicidal self-injury (NSSI)** refers to intentional self-inflicted destruction of body tissue without suicidal intention and for purposes not socially sanctioned. Common forms of NSSI include behaviors such as cutting, burning, scratching, and self-hitting. It is considered a maladaptive coping strategy without the desire to die. NSSI is a common finding among adolescents and young adults in psychiatric inpatient settings.<sup>12</sup>

## FAMILY DYNAMICS

A child's environment – including family dynamics, trauma history, and social factors – can significantly influence their mental health. Nurses assess parent-child relationships, attachment patterns, and exposure to violence, neglect, or instability. Family dynamics refers to the patterns of interactions among family members, their roles and relationships, and the various factors that shape their interactions. Because family members typically rely on each other for emotional, physical, and economic support, they are one of the primary sources of relationship security or stress. Secure and supportive family relationships provide love, advice, and care, whereas stressful family relationships may include frequent arguments, critical feedback, and unreasonable demands.<sup>13</sup>

11. National Institute of Mental Health. (n.d.). *Youth ASQ toolkit*. <https://www.nimh.nih.gov/research/research-conducted-at-nimh/asq-toolkit-materials/youth-asq-toolkit>
12. Cipriano, A., Cella, S., & Cotrufo, P. (2017). Nonsuicidal self-injury: A systematic review. *Frontiers in Psychology*, 8. <https://www.frontiersin.org/articles/10.3389/fpsyg.2017.01946/full>
13. Jabbari, B., Schoo, C., & Rouster, A. S. (2023). *Family dynamics*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK560487>

Interpersonal interactions among family members have lasting impacts and influence the development and well-being of children. Unhealthy family dynamics can cause children to experience trauma and stress as they grow up, known as adverse childhood experiences (ACEs). Conflict between parents and adolescents is associated with adolescent aggression, whereas mutuality (cohesion and warmth) is shown to be a protective factor against aggressive behavior.<sup>14</sup>

- ▶ Review information on adverse childhood experiences (ACEs) in the “[Mental Health and Mental Illness](#)” section of Chapter 1.

Effectively assessing and addressing a client’s family dynamics and its role in a child’s or adolescent’s mental health disorder requires an interprofessional team of health professionals, including nurses, physicians, social workers, and therapists. Nurses are in a unique position to observe interaction patterns, assess family relationships, and attend to family concerns in clinical settings because they are in frequent contact with family members. Collaboration among interprofessional team members promotes family-centered care and provides clients and families with the necessary resources to develop and maintain healthy family dynamics.<sup>15</sup>

Signs of abuse, neglect, or bullying are important for the nurse to observe and report. Read more about abuse and neglect in the “[Trauma, Abuse, and Violence](#)” chapter.

14. Jabbari, B., Schoo, C., & Rouster, A. S. (2023). *Family dynamics*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK560487>

15. Jabbari, B., Schoo, C., & Rouster, A. S. (2023). *Family dynamics*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK560487>

- Read more information about family dynamics in the “[Family Dynamics](#)” chapter of *Open RN Nursing Health Promotion*.

## CULTURAL CONSIDERATIONS

The Cultural Formulation Interview (CFI) is a structured tool in the *DSM-5* used to assess the influence of culture on a client’s experience of distress. See the following box for an adapted version of the CFI tool for children and adolescents.<sup>16</sup>

### Adapted Cultural Formulation Interview for Children and Adolescents<sup>17</sup>

- **Suggested introduction to the child or adolescent:** We have talked about the concerns of your family. Now I would like to know how you are feeling about being \_\_\_\_ years old.
- **Feelings of age appropriateness in different settings:** Do you feel you are like other people your age? In what way? Do you sometimes feel different from other people your age? In what way?

16. Jabbari, B., Schoo, C., & Rouster, A. S. (2023). *Family dynamics*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK560487>

17. Jabbari, B., Schoo, C., & Rouster, A. S. (2023). *Family dynamics*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK560487>

- If they acknowledge sometimes feeling different: Does this feeling of being different happen more at home, at school, at work, and/or some other place? Do you feel your family is different from other families? Does your name have special meaning for you? Is there something special about you that you like or are proud of?
- **Age-related stressors and supports:** What do you like about being at home? At school? With friends? What don't you like at home? At school? With friends? Who is there to support you when you feel you need it? At home? At school? Among your friends?
- **Age-related expectations:** What do your parents or grandparents expect from a person your age in terms of chores, schoolwork, play, or religion? What do your teachers expect from a person your age? What do other people your age expect from a person your age? (If they have siblings, what do your siblings expect from a person your age?)
- **Transition to adulthood (for adolescents):** Are there any important celebrations or events in your community that recognize reaching a certain age or growing up? When is a youth considered ready to become an adult in your family or community? What is good about becoming an adult in your family? In school? In your community? How do you feel about “growing up”? In what ways are your life and responsibilities different from your parents' life and responsibilities?

## LIFE SPAN AND DEVELOPMENTAL CONSIDERATIONS

Children and adolescents are actively experiencing cognitive, emotional, and social development. Young children may lack the vocabulary to describe complex emotions. Nurses use developmentally appropriate language, play therapy, and visual aids to facilitate communication and consider that adolescents may express their feelings through risk-taking or withdrawal. Review theories of development across the life span in the “[Application of the Nursing Process in Mental Health Care](#)” chapter.

Adolescence is a time of exploration regarding gender identity, gender roles, and sexual orientation. As previously discussed, assuring conditional confidentiality is the first step in establishing basic trust and a therapeutic nurse-client relationship with an adolescent client. Most adolescents require privacy to talk candidly about their gender identity and sexuality, so parents/guardians should be asked to leave the examination room at some point during the visit.<sup>18</sup>

## Diagnosis (Analyzing Cues)

Health care professionals use the guidelines in the American Psychiatric Association's *Diagnostic and Statistical Manual (DSM-5)* to diagnose mental health disorders in children.<sup>19</sup>

Nurses customize nursing diagnoses based on the child's or adolescent's response to mental health disorders, their current signs and symptoms, and

18. Middleman, A. B., & Olson, K. A. (2021). Confidentiality in adolescent health care. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

19. American Psychiatric Association. (2013). *Desk reference to the diagnostic criteria from DSM-5*.

the effects on their and their family's functioning. Here are common nursing diagnoses related to childhood and adolescent mental health disorders<sup>20,21,22</sup>.

- Anxiety
- Chronic Low Self-Esteem
- Disabled Family Coping
- Impaired Social Interactions
- Ineffective Impulse Control
- Risk for Delayed Development
- Risk-prone Health Behavior
- Risk for Impaired Parenting
- Risk for Spiritual Distress

## Outcome Identification (Generate Solutions)

Broad goals focus on reducing symptoms of mental health disorders that interfere with the child's or adolescent's daily functioning and quality of life. SMART outcomes stand for outcomes that are specific, measurable, achievable, and realistic with a timeline indicated. They are customized according to each client's diagnoses and needs. Read more about SMART outcomes in the "[Application of the Nursing Process in Mental Health Care](#)" chapter.

For example, a SMART outcome for a child diagnosed with attention deficit hyperactivity disorder is, "The client will demonstrate reduced impulsive

20. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.
21. Ackley, B., Ladwig, G., Makic, M. B., Martinez-Kratz, M., & Zanotti, M. (2020). *Nursing diagnosis handbook: An evidence-based guide to planning care* (12th ed.). Elsevier.
22. Herdman, T. H., & Kamitsuru, S. (Eds.). (2018). *Nursing diagnoses: Definitions and classification, 2018-2020*. Thieme Publishers New York.



behaviors, as reported by parents and their teachers, within two weeks of initiating stimulant medication.”

## Planning (Generate Solutions)

Interventions can be planned by nurses for children and adolescents in a variety of settings, including inpatient, outpatient, day treatment, outreach programs in schools, and home visits. Nurses use collaborative, respectful approaches that involve the youth in goal-setting. They involve parents and caregivers as appropriate, but empower the child’s or adolescent’s voice and participation in their care.<sup>23</sup>

## Safety

Safety receives top priority when planning and implementing interventions for pediatric clients at risk of suicide, and interventions are planned according to their level of risk and developmental level. Review interventions for clients at risk of suicide in the “[Application of the Nursing Process in Mental Health Care](#)” chapter.

It is helpful to encourage older children and adolescents to generate a list of personal warning signs (e.g., thoughts, behaviors, situations, emotional states) that typically precede suicidal thoughts, with as much specificity as possible. These warning signs serve as a cue that it is time for them to use their safety plan. As compared with adults, youths may focus more on general emotional states (e.g., sadness, anger) or external triggers, such as interpersonal situations. Pediatric clients are encouraged to generate a list of coping strategies they can employ to distract themselves from suicidal thoughts and a list of trusted individuals to ask for help during a suicidal crisis. Nurses also work with parents and caregivers to make the client’s

23. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

environment safe and limiting access to means for suicide, such as firearms, medications, knives, and potential household poisons.<sup>24</sup>

## Prevention

A public health approach to children's mental health includes promoting mental health for all children, providing preventative measures for children at risk, and implementing interventions.<sup>25</sup>

It is not known why some children develop disruptive behavior disorders, but children are at greater risk if they are exposed to **adverse childhood experiences (ACEs)**. Toxic stress from ACEs can alter brain development and affect how the body responds to stress. ACEs are linked to chronic health problems, mental illness, and substance abuse. Children with three or more reported ACEs, compared to children with zero reported ACEs, had higher prevalence of one or more mental, emotional, or behavioral disorders (36.3% versus 11.0%).<sup>26</sup> Review information on adverse childhood experiences (ACEs) in the “[Mental Health and Mental Illness](#)” section of Chapter 1.

Preventing ACEs can help children thrive into adulthood by lowering their risk

24. Pettit, J. W., Buitron, V., & Green, K. L. (2018). Assessment and management of suicide risk in children and adolescents. *Cognitive and Behavioral Practice*, 25(4), 460–472. <https://doi.org/10.1016/j.cbpra.2018.04.001>
25. Centers for Disease Control and Prevention. (2021). *Therapy to improve children's mental health*. [https://www.cdc.gov/children-mental-health/treatment/?CDC\\_AAref\\_Val=https://www.cdc.gov/childrensmentalhealth/parent-behavior-therapy.html](https://www.cdc.gov/children-mental-health/treatment/?CDC_AAref_Val=https://www.cdc.gov/childrensmentalhealth/parent-behavior-therapy.html)
26. Centers for Disease Control and Prevention. (2022). *Data and statistics on children's mental health*. [https://www.cdc.gov/children-mental-health/data-research/?CDC\\_AAref\\_Val=https://www.cdc.gov/childrensmentalhealth/data.html](https://www.cdc.gov/children-mental-health/data-research/?CDC_AAref_Val=https://www.cdc.gov/childrensmentalhealth/data.html)

for chronic health problems and substance abuse, improve their education and employment potential, and stop ACEs from being passed from one generation to the next.<sup>27</sup>

Raising awareness about ACEs can help reduce stigma around seeking help for parenting challenges, substance misuse, depression, or suicidal thoughts. Community solutions focus on promoting safe, stable, nurturing relationships and environments where children live, learn, and play. In addition to raising awareness and participating in community solutions, nurses should recognize ACE risk factors and refer clients and their families for effective services and support. See Figure 12.8<sup>28</sup> regarding strategies to prevent ACEs.

27. Centers for Disease Control and Prevention. (2019). *Adverse childhood experiences (ACEs)*. <https://www.cdc.gov/vitalsigns/aces/index.html>

28. This image is derived from “[Preventing Adverse Childhood Experiences \(ACEs\): Leveraging the Best Available Evidence](#)” by [National Center for Injury Prevention and Control, Division of Violence Prevention PDF](#) and is in the [Public Domain](#)



## Preventing ACEs

Strategy	Approach
<b>Strengthen economic supports to families</b>	<ul style="list-style-type: none"><li>• Strengthening household financial security</li><li>• Family-friendly work policies</li></ul>
<b>Promote social norms that protect against violence and adversity</b>	<ul style="list-style-type: none"><li>• Public education campaigns</li><li>• Legislative approaches to reduce corporal punishment</li><li>• Bystander approaches</li><li>• Men and boys as allies in prevention</li></ul>
<b>Ensure a strong start for children</b>	<ul style="list-style-type: none"><li>• Early childhood home visitation</li><li>• High-quality child care</li><li>• Preschool enrichment with family engagement</li></ul>
<b>Teach skills</b>	<ul style="list-style-type: none"><li>• Social-emotional learning</li><li>• Safe dating and healthy relationship skill programs</li><li>• Parenting skills and family relationship approaches</li></ul>
<b>Connect youth to caring adults and activities</b>	<ul style="list-style-type: none"><li>• Mentoring programs</li><li>• After-school programs</li></ul>
<b>Intervene to lessen immediate and long-term harms</b>	<ul style="list-style-type: none"><li>• Enhanced primary care</li><li>• Victim-centered services</li><li>• Treatment to lessen the harms of ACEs</li><li>• Treatment to prevent problem behavior and future involvement in violence</li><li>• Family-centered treatment for substance use disorders</li></ul>

Figure 12.8 Strategies to Prevent ACEs

- Read [Preventing Adverse Childhood Experiences PDF](#) by the CDC with evidence-supporting interventions.

## Implementation (Take Action)

Treatment of mental health disorders in children and adolescents typically requires a combination of psychotherapy and pharmacotherapy. Read about specific multidisciplinary treatments for various disorders in the “[Psychological Therapies and Behavioral Intervention](#),” and “[Autism Spectrum](#)

[Disorder](#)” sections of this chapter. Nurses recognize and capitalize on the client’s and family’s strengths as they develop a nursing care plan and provide education and referral to resources as appropriate. A variety of interventions may be implemented based on the disorder diagnosed and the developmental level of the child or adolescent.

## Behavioral Interventions

Behavioral interventions reward desired behaviors to reduce maladaptive coping behaviors. Most child and adolescent treatment settings use structured programs to motivate and reward age-appropriate behaviors. For example, the point or star system may be used where the child receives points or stars for desired behaviors, and then specific privileges are awarded based on the points or stars earned each day.<sup>29</sup>

Read more about interprofessional behavioral interventions for the child, parents/caregivers, and teachers in the “[Psychological Therapies and Behavioral Interventions](#)” section.

## Play Therapy

Children develop physically, intellectually, emotionally, and socially through play. **Play therapy** encourages children to express feelings such as anxiety, self-doubt, and fear through their natural play. It also allows them to work through painful or traumatic memories. For example, nurses often use dolls or other toys to help children work through feelings of fear prior to a medical

29. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

procedure.<sup>30</sup> See Figure 12.9<sup>31</sup> for an image of children with special needs playing at a school for autism.



Figure 12.9 Children With Special Needs Playing at an Autism School

## Bibliotherapy

**Bibliotherapy** uses books to help children express feelings in a supportive environment, gain insight into feelings and behavior, and learn new ways to cope with difficult situations. When children listen to or read a story, they identify with the characters and experience a catharsis of feelings. Stories and books should be selected based on the child's cognitive and developmental levels that reflect the situations or feelings the child is experiencing and their

30. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

31. "4885680339\_721cd1843e\_k" by [World Bank Photo Collection](#) is licensed under [CC BY-NC-ND 2.0](#)



emotional readiness for the topic.<sup>32</sup> See Figure 12.10<sup>33</sup> for an image of bibliotherapy.



Figure 12.10 Bibliotherapy

## Expressive Arts Therapy

Art provides a nonverbal method of expressing difficult or confusing emotions. Drawing, painting, and sculpting with clay are commonly used art therapies. Children who have experienced trauma often show the traumatic event in their drawing when asked to draw whatever they wish.<sup>34</sup>

32. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

33. "[Bibliotherapy.jpg](#)" by [Shelley Rodrigo](#) is licensed under [CC BY 2.0](#)

34. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

# Journaling

Journaling is an effective technique for older children and teenagers. Journaling is a tangible way to record emotions and begin dialogue with others. A daily journal is also helpful in setting goals and evaluating progress.<sup>35</sup>

# Music Therapy

Music has been recognized for centuries as having healing power. **Music therapy** is an evidence-based approach to improve an individual's physical, psychological, cognitive, behavioral, and social functioning when listening to music, singing, or moving to music.<sup>36</sup> See Figure 12.11<sup>37</sup> for an image of music therapy.

35. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

36. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

37. "[103909417\\_f36b60ceec\\_k](#)" by [midiman](#) is licensed under [CC BY 2.0](#)





Figure 12.11 Music Therapy

## Family Education and Support

Education of family members is a key component for treating child and adolescent mental health disorders. Read more about “Parent Training in Behavior Management” in the “[Psychological Therapies and Behavioral Interventions](#)” section. Nurses can help family members develop specific goals and identify interventions to achieve their family’s goals.

Family members can also be encouraged to attend support groups or group education. Group education can be useful for learning how other families solve problems and build on strengths while developing insight and improved judgment about their own family.<sup>38</sup>

- ▶ Find support groups for many disorders near you at [Psychology Today’s Support Groups](#).

38. Halter, M. (2022). *Varc Carolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

# Disruptive Behavior Management

Nurses can manage a child's or adolescent's disruptive behaviors by implementing many different types of interventions<sup>39</sup> :

- **Behavioral contract:** A verbal or written agreement is made between the client and other parties (e.g., nurses, parents, or teachers) about behaviors, expectations, and needs. The contract is periodically reviewed with positive and negative reinforcement provided.
- **Collaborative and proactive solutions:** The identification of problematic behaviors, their specific triggers, and mutually agreeable solutions.
- **Role playing:** The nurse or client acts out a specific role to practice new behaviors or skills in specific situations.
- **Planned ignoring:** When behaviors are determined to be attention-seeking and not dangerous, they may be ignored. Positive reinforcement is provided for on-task actions.
- **Signals or gestures:** An adult uses a word, gesture, or eye contact to remind the child to use self-control. For example, placing one's finger to one's lips and making eye contact may be used to remind a child to remain quiet during a quiet activity.
- **Physical distance:** It may be helpful to move closer to a child for a calming effect. However, some children may find this agitating and require more space and less physical closeness.
- **Redirection:** The engagement of an individual in an appropriate activity after an undesirable action.
- **Humor:** Appropriate joking can be used as a diversion to help relieve feelings of guilt or fear.
- **Restructuring:** The process of changing an activity to reduce stimulation or frustration.
- **Limit setting:** The process of giving direction, stating an expectation, or

39. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

telling a child what to do or where to go. Caregivers and/or staff should do this firmly, calmly, and consistently without anger, preferably in advance of problem behavior occurring. For example, “I would like you to stop turning the light on and off.”

- **Simple restitution:** The individual is expected to correct the adverse effects of their actions, such as apologizing to the people affected or returning upturned chairs to their proper position.

## Restrictive Interventions

Restrictive interventions are only implemented after attempting less restrictive interventions that did not successfully manage the client's behavior. As a last resort, restrictive interventions, including seclusion or restraints, may be implemented to keep the client or others around them safe.<sup>40</sup>

The Centers for Medicare and Medicaid Services (CMS) and The Joint Commission (TJC) define seclusion as the involuntary confinement of a client alone in a room or area from which the client is physically prevented from leaving. Restraint is defined as any manual method, physical or mechanical device, material, or equipment that immobilizes or reduces the ability of a client to move their arms, legs, body, or head freely. A drug is considered a restraint when it is used as a restriction to manage the client's behavior or restrict the client's freedom of movement and is not a standard treatment or dosage for the client's condition.<sup>41</sup>

40. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

41. Knox, D. K., & Holloman, G. H. (2012). Use and avoidance of seclusion and restraint: Consensus statement of the American Association for Emergency Psychiatry Project BETA seclusion and restraint workgroup. *The Western Journal of Emergency Medicine*, 13(1), 35–40. <https://doi.org/10.5811/westjem.2011.9.6867>

Seclusion is viewed as less restrictive than restraints, but seclusion and restraint are psychologically harmful and can be physically dangerous. Nurses must vigilantly follow agency policy when implementing seclusion or restraints. Members of the health care team who assist with seclusion or restraint of children or adolescents must receive specific training to decrease the risk of injury to the youngster and themselves.<sup>42</sup>

Seclusion and restraints should be discontinued as soon as possible. Clients in seclusion or restraints must be frequently monitored. Hydration, elimination, comfort, and other psychological and physical needs must be monitored regularly and addressed per agency policy.<sup>43</sup>

After the child or adolescent is calm, staff should include the child in a debriefing session and discuss the events leading up to the restrictive interventions to explore ways it may have been prevented.<sup>44</sup>

## Nursing Interventions for Pediatric Clients Based on Categories of the APNA Implementation Standard

Nurses implement interventions related to each subcategory of the *Implementation* Standard of Care for pediatric clients in mental health settings. See Table 12.5a for examples of common nursing interventions for each subcategory.

42. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

43. Knox, D. K., & Holloman, G. H. (2012). Use and avoidance of seclusion and restraint: Consensus statement of the American Association for Emergency Psychiatry Project BETA Seclusion and Restraint Workgroup. *The Western Journal of Emergency Medicine*, 13(1), 35–40. <https://doi.org/10.5811/westjem.2011.9.6867>

44. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

Table 12.5a Examples of Nursing Interventions for Pediatric Mental Health Disorders Based on Subcategories of APNA Implementation Standard

Subcategory of the APNA Standard of Implementation	The nurse will ...	Rationale
<b>Coordination of Care</b>	<ul style="list-style-type: none"> <li>• Collaborate with parents/guardians, school personnel, pediatricians, and therapists to create a comprehensive care plan.</li> <li>• Facilitate communication between home, school, and clinical settings.</li> <li>• Involve child welfare services when safety is a concern.</li> </ul>	Children and adolescents depend on parents, caregivers and school systems for stability. Coordinated care ensures continuity, consistency, and multisystem support.
<b>Health Teaching and Health Promotion</b>	<ul style="list-style-type: none"> <li>• Provide age-appropriate psychoeducation about emotions, coping strategies, and symptom management.</li> <li>• Teach families about diagnoses, developmental expectations, and the impact of trauma or stress.</li> <li>• Promote healthy sleep, nutrition, and physical activity routines.</li> </ul>	Understanding mental health supports empowerment, reduces stigma, and improves engagement in care. Teaching families promotes a supportive home environment.

<b>Pharmacological, Biological, and Integrative Therapies</b>	<ul style="list-style-type: none"> <li>• Administer psychotropic medications as prescribed, with careful monitoring of side effects, growth, and development.</li> <li>• Educate children and families about medications using simple, clear language.</li> <li>• Integrate play, mindfulness, or expressive therapies to support emotional regulation.</li> </ul>	<p>Children may not understand or tolerate medications the same as adults. Combining medication with developmentally appropriate therapies enhances safety and efficacy.</p>
<b>Milieu Therapy</b>	<ul style="list-style-type: none"> <li>• Maintain a structured, supportive, and trauma-informed environment.</li> <li>• Use consistent routines and clearly defined behavioral expectations.</li> <li>• Model and reinforce positive behaviors through praise and positive reinforcement strategies.</li> </ul>	<p>Children need structure and predictability for emotional security. A therapeutic milieu promotes self-regulation and reduces acting-out behaviors.</p>

<b>Therapeutic Relationship and Counseling</b>	<ul style="list-style-type: none"><li>• Build trust through play, nonjudgmental listening, and consistent presence.</li><li>• Use developmentally appropriate communication tools (e.g., drawing, games, storytelling).</li><li>• Support identity development and emotional expression.</li><li>• Engage family in therapy and encourage healthy communication.</li></ul>	A strong therapeutic relationship is the foundation for behavioral change, especially in youth who have experienced trauma or instability. Developmentally-appropriate therapeutic interactions foster connection and insight.
--	--	--

# Communication Tips for Pediatric Clients with Mental Health Disorders

Communicating effectively with pediatric clients must foster feelings of trust, safety, and therapeutic alliance. Table 12.5b provides communication tips when working with pediatric clients.

Table 12.5b Communication Tips for Pediatric Clients<sup>45</sup>

45. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.



Tip	Explanation
<b>Use developmentally appropriate language</b>	Avoid clinical or abstract terms. Use simple, concrete words and metaphors that match the child's age and cognitive level (e.g., "worry bugs" for anxiety). This supports understanding and reduces confusion.
<b>Use play, drawing, or storytelling as communication tools</b>	Many children express emotions and thoughts more effectively through nonverbal activities or play that promote emotional expression while building rapport.
<b>Maintain a calm, warm, and nonjudgmental tone</b>	Children are sensitive to tone and body language. A calm presence helps them feel safe and promotes emotional regulation, especially during distress or behavioral outbursts.
<b>Validate feelings, even if behaviors are inappropriate</b>	Say things like, "It looks like you're feeling really upset right now." Validation shows empathy and teaches emotional literacy without reinforcing negative behavior.
<b>Offer choices when possible</b>	Providing limited, structured choices (e.g., "Would you like to talk here or in the beanbag corner?") fosters autonomy and reduces power struggles.
<b>Be consistent and set clear expectations</b>	Children with mental health challenges often struggle with boundaries. Clear, consistent communication reduces anxiety and helps them feel secure.
<b>Allow time for response and silence</b>	Children, especially those with trauma, anxiety, or neurodevelopmental conditions, may need extra time to process questions. Silence can be therapeutic and allows for reflection.
<b>Use positive reinforcement and praise</b>	Focus on strengths and progress (e.g., "I noticed you tried really hard to use your words instead of yelling."). Positive reinforcement promotes confidence and reinforces desired behavior.
<b>Engage parents and caregivers in communication and modeling</b>	Involve families in treatment goals, behavior support strategies, and shared language for emotional regulation. Consistency across settings supports skill generalization and stability.

## Evaluation (Evaluate Outcomes)


Evaluation focuses on monitoring a child's and adolescent's progress towards meeting their individualized SMART goals and the revision of the nursing care plan as needed. Evaluation criteria are aligned to the specific client diagnosis. Client progress monitoring may include symptom management, behavior

management, academic performance, activities of daily living, and socialization. The nurse also monitors effectiveness of interprofessional treatments, support groups, and community resources for the client's families.

Examples of evaluation questions include:

- “How often are the symptoms occurring?”
- “Have there been any changes in the intensity of the symptoms?”
- “How are the symptoms affecting daily functioning at home, school, and in social settings?”
- “Are there any improvements in academic performance or social interactions?”
- “Have there been any changes in mood or behavior?”
- “Are there any new behaviors or worsening of previous behaviors?”
- “What changes have parents or teachers observed in the child’s behavior or emotional state?”
- “How are the symptoms of coexisting conditions being managed?”

## 12.6 Spotlight Application

 Dr. Temple Grandin was one of the first individuals to document the insights gained from her personal experience of autism. View the following YouTube video<sup>1</sup> of Dr. Temple Grandin describing her personal experience as, “Everything in my mind works like a search engine set for the image function”: [Temple Grandin on Her Search Engine | Blank on Blank](https://youtu.be/Ifsh6sojAvg).

### Reflective Questions

1. What symptoms of autism does Dr. Temple Grandin illustrate?
2. What strengths does Dr. Temple Grandin illustrate and what has she achieved?

1. Blank on Blank. (2016, March 1). *Temple Grandin on her search engine | Blank on Blank*. [Video]. YouTube. All rights reserved. <https://youtu.be/Ifsh6sojAvg>

## 12.7 Learning Activities



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=640#h5p-41>

1



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=640#h5p-42>

2



*An interactive H5P element has been excluded from this version of the text. You can view it*

1. “MH Childhood and Adolescence Disorders Glossary Cards” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
2. “MH Childhood and Adolescence Disorders Fill in Blank” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

online here:

<https://wtcs.pressbooks.pub/nursingmhcc/?p=640#h5p-44>

3

- ▶ Test your clinical judgment with a NCLEX Next Generation-style question: [Chapter 12, Assignment 1](#)<sup>4</sup>



- ▶ Test your clinical judgment with a NCLEX Next Generation-style case study: [Chapter 12, Case Study 1](#)<sup>5</sup>



3. "MH Childhood and Adolescence Disorders Question Set 1" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
4. "MH Childhood and Adolescence Disorders Next Gen Question 1" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
5. "MH Childhood and Adolescence Disorders Next Gen Case Study" by Kellea Ewen is licensed under [CC BY-NC 4.0](#)

## XII Glossary

**Adverse childhood experiences (ACEs):** Potentially traumatic events that occur in childhood, such as violence, abuse, and growing up in a family with mental health or substance use problems.

**Anxiety disorders:** A group of mental health conditions characterized by excessive fear or worry that interferes with daily activities.

**Bibliotherapy:** A behavioral intervention that uses books to help children express feelings in a supportive environment, gain insight into feelings and behavior, and learn new ways to cope with difficult situations.

**Conduct disorder (CD):** A behavioral disorder diagnosed when a child shows an ongoing pattern of aggression toward others with serious violations of rules and social norms at home, school, and with peers.

**Depression:** A mood disorder characterized by persistent feelings of sadness, hopelessness, and a loss of interest or pleasure in activities, and various emotional and physical problems that interfere with daily functioning.

**Developmental disabilities:** A group of conditions with physical, learning, language, or behavioral impairments.

**Developmental monitoring:** Routine screenings for developmental delays during well-child visits based on observations of the child and discussion with parents.

**Developmental screening:** Formal questionnaires or checklists based on research that ask questions about a child's development, including language, movement, thinking, behavior, and emotions.

**Dyscalculia:** A learning disorder with difficulty with math.

**Dysgraphia:** A learning disorder with difficulty with writing.

**Dyslexia:** A learning disorder with difficulty with reading.

**Echolalia:** Pathological repetition of another person's words.

**Expressive language disorder:** Difficulty communicating thoughts using language due to not knowing the words to use, not knowing how to put words together, or not being able to express the words.

**Intellectual disability:** A person's ability to learn at an expected level and function in daily life is limited.

**Learning disorder:** Difficulty in one or more areas of learning, even when a child's overall intelligence or motivation are not affected.

**Mental health disorders among children:** Serious changes in the way children typically learn, behave, or handle their emotions, causing distress and problems getting through the day.

**Music therapy:** A behavioral intervention to improve an individual's physical, psychological, cognitive, behavioral, and social functioning by listening to music, singing, or moving to music.

**Oppositional defiant disorder (ODD):** A behavioral disorder diagnosed when children act out persistently, causing serious problems at home, in school, or with peers.

**Play therapy:** A behavioral intervention that encourages children to express feelings such as anxiety, self-doubt, and fear through their natural play. It also allows them to work through painful or traumatic memories.

**Receptive language disorder:** Difficulty understanding what others say due to not hearing the words (hearing loss) or not understanding the meaning of the words.

**Restraint:** Any manual method, physical or mechanical device, material, or equipment that immobilizes or reduces the ability of a client to move their arms, legs, body, or head freely.

**Seclusion:** The involuntary confinement of a client alone in a room or area from which the client is physically prevented from leaving.

**Separation anxiety:** A condition of children being very afraid when away from parents or caregivers.

**Tics:** Sudden twitches, movements, or sounds that people do repeatedly with the inability to stop their body from doing these actions.







### Learning Objectives

- Identify assessment cues of eating disorder behaviors
- Identify nursing priorities for clients with eating disorders
- Plan outcomes for clients with eating disorders
- Differentiate safety/protective interventions for clients with eating disorders
- Apply evidence-based practice when planning care and interventions for clients with eating disorders
- Analyze treatments for clients with eating disorders
- Apply the nursing process to clients with eating disorders at risk for suicide

Many individuals experience concerns about their weight at some point in their lives. However, eating disorders involve a distorted perception of weight and body image, leading to an unhealthy relationship with food. This psychological struggle can result in behaviors such as extreme food restriction, compulsive exercise, binge eating, and purging. Eating disorders can affect people of any gender, age, race, or socioeconomic background and often have a profound impact on their loved ones. Without proper identification and treatment, the physiological consequences can be life-threatening. This chapter explores the various types of eating disorders, common assessment findings, treatment approaches, and nursing interventions.

## 13.2 Basic Concepts

Eating disorders are severe mental health conditions characterized by significant disturbances in eating behaviors and related thoughts and emotions. These disorders can have profound impacts on physical, psychological, and social functioning. The most common types of eating disorders include anorexia nervosa, bulimia nervosa, and binge eating disorder.

The way we talk about eating disorders with others matters.<sup>1</sup> See Figure 13.1<sup>2</sup> with facts nurses can use to help shape the conversation around eating disorders.

1. National Institute of Mental Health. (2024). *Eating disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/eating-disorders>
2. “[2020\\_eatingdisorderinfographics\\_final.jpg](#)” by U.S. Department of Health and Human Services, National Institutes of Health, National Institute of Mental Health is in the [Public Domain](#)



Figure 13.1 Facts About Eating Disorders

## Common Eating Disorders

Common eating disorders include anorexia nervosa, bulimia nervosa, and binge eating disorder.<sup>3</sup> Individuals may have more than one type of disorder.

### Anorexia Nervosa

**Anorexia nervosa** is a condition where people avoid food, severely restrict food, or eat very small quantities of only certain foods. They have an intense

3. National Institute of Mental Health. (2024). *Eating disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/eating-disorders>

fear of gaining weight or becoming fat, even if they are significantly underweight. People with anorexia nervosa often have distorted body image and view themselves as overweight.<sup>4</sup> See Figure 13.2<sup>5</sup> for an illustration of how a person with anorexia nervosa may perceive themselves.




Figure 13.2 Body Perception of a Person With Anorexia Nervosa

There are two subtypes of anorexia nervosa<sup>6,7</sup>:

- **Restricting Type:** Individuals severely limit the amount and type of food

4. National Institute of Mental Health. (2024). *Eating disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/eating-disorders>
5. “Bulimia-Anorexia-Nervosa-Skimmed-Delusional-4049661.png” by unknown author is licensed under CC0.
6. National Institute of Mental Health. (2024). *Eating disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/eating-disorders>
7. American Psychiatric Association. (2013). *Desk reference to the diagnostic criteria from DSM-5*.

they consume and may engage in excessive exercise.

- **Binge-Eating/Purging Type:** Individuals experience episodes of binge eating and/or purging.
  - **Binge eating** involves consuming large amounts of food in a short period with a sense of loss of control.
  - **Purging** refers to self-induced vomiting or the misuse of laxatives, diuretics, or enemas to eliminate consumed food.

Anorexia nervosa can be fatal due to medical complications associated with starvation. It has an extremely high mortality (death) rate compared with other mental health disorders. Young people between the ages of 15 and 24 with anorexia have 10 times the risk of dying compared to their same-aged peers. Males represent 25% of individuals with anorexia nervosa and are at a higher risk of dying because they are often diagnosed later than females.<sup>8,9</sup>

Signs and symptoms of anorexia nervosa include the following<sup>10</sup>:

- Severely restricted eating
- Extreme thinness (also referred to as **emaciation**)
- A relentless pursuit of thinness and unwillingness to maintain a healthy weight
- Intense fear of gaining weight

8. National Institute of Mental Health. (2024). *Eating disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/eating-disorders>

9. National Eating Disorders Association. (n.d.). *Statistics and research on eating disorders*. <https://www.nationaleatingdisorders.org/statistics-research-eating-disorders>

10. National Institute of Mental Health. (2024). *Eating disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/eating-disorders>

- Distorted body image where a person's self-esteem is heavily influenced by their perception of their of body weight and shape
- Denial of the seriousness of low body weight

Other signs may develop over time, including these issues<sup>11</sup>:

- Thinning of the bones (i.e., osteopenia or osteoporosis)
- Anemia
- Muscle wasting and weakness
- Brittle hair and nails
- Dry and yellowish skin
- Growth of fine hair all over the body (**lanugo**)
- Severe constipation
- Low blood pressure
- Slowed breathing and pulse
- Drop in internal body temperature, causing the person to feel cold all the time
- Lethargy, sluggishness, or feeling tired all the time
- Infertility
- Damage to the structure and function of the heart
- Brain damage
- Multiorgan failure

See the following box for criteria used to diagnose anorexia nervosa according to the *Diagnostic and Statistical Manual of Disorders (DSM-5-TR)*.

11. National Institute of Mental Health. (2024). *Eating disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/eating-disorders>



## DSM-5 Criteria for Anorexia Nervosa<sup>12</sup>

- Restriction of energy intake relative to requirements, leading to significantly low body weight in the context of age, sex, developmental trajectory, and physical health. Significantly low weight is defined as weight less than minimally normal, or in children and adolescents, less than that minimally expected.
- Intense fear of gaining weight or becoming fat or persistent behavior that interferes with weight gain, even though already at a significantly low weight.
- Disturbance in the way in which one's body weight or shape is experienced, undue influence of body weight or shape on self-evaluation, or persistent lack of recognition of the seriousness of the current low body weight.

The type is specified as follows:

- **Restricting type:** Weight loss is accomplished primarily through dieting, fasting, or excessive exercise. During the last three months, the individual has not engaged in recurrent episodes of binge eating or purging behavior.
- **Binge eating/purging type:** During the last three months, the individual has engaged in recurrent episodes of binge eating or purging behavior (i.e., self-induced vomiting or the misuse of laxatives, diuretics, or enemas).

12. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

# Bulimia Nervosa

**Bulimia nervosa** is a condition where people have recurrent and frequent episodes of binge eating (i.e., eating unusually large amounts of food in a short amount of time while also feeling a lack of control over these episodes). Binge eating is followed by behaviors used to eliminate the excess food such as forced vomiting, excessive use of laxatives or diuretics, fasting, excessive exercise, or a combination of these behaviors. People with bulimia nervosa may be slightly underweight, normal weight, or overweight.<sup>13</sup> Clients with combination Anorexia Nervosa-Bulimia Nervosa have a significantly low body weight, whereas clients with Bulimia Nervosa typically have a normal or above-normal body weight. Unlike anorexia nervosa, bulimia nervosa can be harder to detect, as individuals may fall anywhere along the weight spectrum.

Signs and symptoms of bulimia nervosa include the following<sup>14</sup>:

- Chronically inflamed and sore throat
- Swollen salivary glands in the neck and jaw area
- Worn tooth enamel and increasingly sensitive and decaying teeth as a result of exposure to stomach acid
- Acid reflux disorder and other gastrointestinal problems
- Intestinal distress and irritation from laxative abuse
- Severe dehydration from purging of fluids
- Disruptions in sodium, calcium, potassium, and other minerals, increasing the risk of dysrhythmias and cardiac arrest

13. National Institute of Mental Health. (2024). *Eating disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/eating-disorders>

14. National Institute of Mental Health. (2024). *Eating disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/eating-disorders>

See the following box for *DSM-5-TR* criteria for diagnosing bulimia nervosa.

### **DSM-5-TR Criteria for Bulimia Nervosa<sup>15</sup>**

- Recurrent episodes of binge eating. A binge eating episode is characterized by both of the following:
  - Eating in a discrete period of time (e.g., within any two-hour period) an amount of food that is definitely larger than what most individuals would eat in a similar period of time under similar circumstances.
  - A sense of lack of control overeating during the episode (i.e., feeling that one cannot stop eating or control what or how much one is eating).
- Recurrent inappropriate compensatory behavior to prevent weight gain, such as self-induced vomiting; misuse of laxatives, diuretics, or other medications; fasting; or excessive exercise.
- The binge eating and inappropriate compensatory behaviors both occur, on average, at least once a week for three months.
- Self-evaluation is unduly influenced by body shape and weight.
- The disturbance does not occur exclusively during episodes of anorexia nervosa.

15. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

# Binge Eating Disorder

Binge eating disorder is the most common eating disorder in the United States.<sup>16</sup> **Binge eating disorder** is a condition where people lose control over their eating and have recurring episodes of eating unusually large amounts of food. However, unlike bulimia nervosa, periods of binge eating are not followed by purging, excessive exercise, or fasting. As a result, people with binge eating disorder often are overweight or obese.<sup>17</sup>

These are the signs and symptoms of a binge eating disorder<sup>18</sup>:

- Eating unusually large amounts of food in a specific amount of time, such as a two-hour period
- Eating even when feeling full or not hungry
- Eating at a fast pace
- Eating until uncomfortably full
- Eating alone or in secret to avoid embarrassment
- Feeling distressed, ashamed, or guilty about eating
- Frequent dieting, possibly without weight loss

See the *DSM-5-TR* criteria for diagnosing binge eating disorder in the following box.

16. National Institute of Mental Health. (2024). *Eating disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/eating-disorders>

17. National Institute of Mental Health. (2024). *Eating disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/eating-disorders>

18. National Institute of Mental Health. (2024). *Eating disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/eating-disorders>

## DSM-5-TR Criteria for Binge Eating Disorder<sup>19</sup>

- Recurrent episodes of binge eating. An episode of binge eating includes eating within any 2-hour period) an amount of food that is definitely larger than what most people would eat in a similar amount of time under similar circumstances, and a sense of lack of control over eating during the episode.
- The binge eating episodes are associated with three (or more) of the following:
  - Eating much more rapidly than normal.
  - Eating until feeling uncomfortably full.
  - Eating large amounts of food when not feeling physically hungry.
  - Eating alone because of being embarrassed by how much one is eating.
  - Feeling disgusted with oneself, depressed, or very guilty afterward.
- Marked distress regarding binge eating is present.
- The binge eating occurs, on average, at least once a week for three months.
- The binge eating is not associated with the recurrent use of inappropriate compensatory behavior, such as in bulimia nervosa, and does not occur exclusively during the course

<sup>19</sup>. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

of bulimia nervosa or anorexia nervosa.

## Other Eating Disorders

**Pica** is another type of eating disorder in which an individual repeatedly eats things that are not considered food and have no nutritional value, such as paper, dirt, soap, hair, glue, or chalk. Individuals with pica do not usually have an aversion to food, and ingested items vary with age. The behavior is inappropriate to the developmental level of the individual and is not part of a culturally supported practice. A person diagnosed with pica is at risk for potential intestinal blockages or toxic effects of substances consumed (such as lead in paint chips). Treatment for pica involves testing for nutritional deficiencies and addressing them if needed. Behavioral interventions used to treat pica may include redirecting the individual from the nonfood items and rewarding them for setting aside or avoiding nonfood items.<sup>20</sup>

While pica is most commonly seen in children, it also occurs in individuals with intellectual disabilities and, in some cases, during pregnancy.

Diagnosis, according to the DSM-5-TR is based on the following criteria<sup>21</sup>:

- Persistent eating of non-nutritive, nonfood substances for at least one month

20. American Psychiatric Association. (2023). *What are eating disorders?* <https://www.psychiatry.org/patients-families/eating-disorders/what-are-eating-disorders>

21. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

- The eating of these substances is inappropriate for the individual's developmental level
- The eating behavior is not part of culturally supported or socially normative practice

Health risks associated with pica depend on the type of substance ingested:

- **Geophagia** (Clay Eating): Can lead to constipation, hypokalemia, nutritional deficiencies, and exposure to infectious agents like parasites. Clay contaminated with lead increases the risk of lead poisoning.
- **Pagophagia** (Ice Eating): Associated with iron deficiency, particularly during pregnancy, can lead to tooth decay and sensitivity.
- **Amylophagia** (Starch Eating): Also linked to iron deficiency, and in excessive amounts, may contribute to high blood sugar and obesity due to its carbohydrate content.
- **Toxic Substances:** Some nonfood items contain harmful contaminants such as lead, mercury, arsenic, and fluoride, which can cause poisoning, neurological damage, or organ dysfunction.

Pica during pregnancy can have severe consequences for both the mother and fetus. Lead exposure from pica has been linked to long-term neurological disabilities in children, and maternal geophagy has been associated with childhood motor function delays.

The cause of pica remains unknown. However, several factors may contribute to its development, including:

- Stress
- Cultural influences
- Learned behavior
- Low socioeconomic status
- Underlying mental health disorders
- Nutritional deficiencies
- Child neglect
- Pregnancy

- Epilepsy
- Familial psychopathology

Treatment and Management for pica involves testing and addressing any underlying nutritional deficiencies, particularly iron and zinc. Behavioral interventions, such as redirecting individuals from nonfood items and reinforcing appropriate eating behaviors, are commonly used. In cases where pica coexists with other mental health conditions, treatment may involve therapy or medications to manage underlying psychiatric symptoms<sup>22</sup>.

**Avoidant restrictive food intake disorder (ARFID)** is a condition where individuals limit the amount or type of food eaten. Unlike anorexia nervosa, people with ARFID do not have a distorted body image or extreme fear of gaining weight. ARFID is most common in middle childhood and usually has an earlier onset than other eating disorders. Many children go through phases of picky eating, but a child with ARFID does not eat enough calories to grow and develop properly, and an adult with ARFID does not eat enough calories to maintain basic body function.<sup>23</sup>

Signs and symptoms of ARFID include the following<sup>24</sup>:

- Dramatic restriction of types or amount of food eaten
- Lack of appetite or interest in food

22. Al Nasser, Y., Muco, E., & Alsaad, A. J. (2025). *Pica*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK532242/>

23. National Institute of Mental Health. (2024). *Eating disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/eating-disorders>

24. National Institute of Mental Health. (2024). *Eating disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/eating-disorders>



- Dramatic weight loss
- Upset stomach, abdominal pain, or other gastrointestinal issues with no other known cause
- Limited range of preferred foods that becomes even more limited (i.e., “picky eating” that gets progressively worse)

ARFID does not include food restriction related to lack of availability of food; dieting; cultural practices, such as religious fasting; or developmentally normal behaviors, such as toddlers who are picky eaters. Food avoidance or restriction commonly develops in infancy or early childhood and may continue in adulthood, but it can start at any age. Regardless of the age of the person affected, ARFID can impact families, causing increased stress at mealtimes and in other social eating situations. Treatment for ARFID involves an individualized plan and may involve several specialists, including a mental health professional and a registered dietitian.

## Body Mass Index

**Body mass index (BMI)** is a person’s weight in kilograms divided by the square of height in meters. BMI is an easy screening method to determine if an individual’s weight is classified as underweight, healthy weight, overweight, or obese. However, it is important to remember that BMI is a screening method and does not take into account muscle mass, bone density, overall body composition, racial, and sex differences.<sup>25</sup> See Table 13.2a for adult weight status according to BMI ranges.

Table 13.2a Adult Weight Status by BMI<sup>26</sup>

25. Nordqvist, C. (2022). *Why BMI is inaccurate and misleading*. Medical News Today. <https://www.medicalnewstoday.com/articles/265215>

26. Centers for Disease Control and Prevention. (2024). *Adult BMI calculator*. <https://www.cdc.gov/bmi/adult-calculator/>

BMI	Weight Status
Below 18.5	Underweight
18.5 – 24.9	Healthy Weight
25.0 – 29.9	Overweight
30.0 or higher	Obese

For children and teens, the interpretation of BMI depends upon their age and sex. After BMI is calculated for children and teens, it is expressed as a percentile obtained from either a percentile calculator or growth chart. See categories of weight status based on BMI percentiles in Table 13.2b. Links to calculators and growth charts are provided in the following box.

Table 13.2b Child or Adolescent Weight Status by BMI Percentile<sup>27</sup>

BMI Percentile	Weight Status
Less than 5th Percentile	Underweight
5th Percentile to Less Than 85th Percentile	Healthy Weight
85th Percentile to Less Than 95th Percentile	Overweight
95th Percentile or Higher	Obese

**Calculating BMI**

- ▶ The CDC’s Adult [BMI Calculator](https://www.cdc.gov/bmi/child-teen-calculator/) conveniently calculates a

27. Centers for Disease Control and Prevention. (2024). *Child and teen BMI calculator*. <https://www.cdc.gov/bmi/child-teen-calculator/>

- ▶ person's BMI based on their height and weight. The following formulas can also be used to calculate a person's BMI<sup>28</sup>:
  - $\text{weight (kg)} / [\text{height (m)}]^2$  or
  - $\text{weight (lb)} / [\text{height (in)}]^2 \times 703$
- ▶ The CDC's [BMI Percentile Calculator for Children and Adolescents](#) conveniently calculates a child's or adolescent's BMI percentile based on their height, weight, gender, and age. Otherwise, gender growth charts are used to determine BMI percentiles and then their weight status is determined using the appropriate table below:
  - [CDC extended BMI-for-age growth charts](#)

## Risk Factors

The exact cause of eating disorders is not fully understood, but research suggests that a combination of genetic, biological, behavioral, psychological, and cultural factors can increase a person's risk. Eating disorders are hereditary. Researchers are working to identify DNA variations that are linked to the increased risk of developing eating disorders. Brain imaging studies are also providing a better understanding of eating disorders. For example, researchers have found differences in patterns of brain activity in women with eating disorders in comparison to healthy women. This kind of research can help guide the development of new means of diagnosis and treatment of eating disorders.<sup>29</sup>

28. Nordqvist, C. (2022). *Why BMI is inaccurate and misleading*. Medical News Today. <https://www.medicalnewstoday.com/articles/265215>

29. National Institute of Mental Health. (2024). *Eating disorders*. U.S. Department

Although genetics increases the risk for developing eating disorders, the individual's environment plays a significant role. Onset can occur at nearly any point in the life span, but eating disorders commonly start in adolescence and young adulthood when there can be exceptional pressure from peers, social media, advertisements, etc., to diet or lose weight. For those genetically vulnerable to eating disorders, initial weight loss may reinforce a reward feedback mechanism and establish a maladaptive eating behavior pattern. Physiological and sensorial changes result in alterations in hunger and satiety, gastrointestinal motility, and decision-making around food and eating.

As you consider the genetic, environmental, and social influences of eating disorders, recognize that a combination of these factors can affect the severity and presenting characteristics for each client. Additionally, although eating disorders are commonly thought of as affecting young women, it can affect people of all genders, ages, races/ethnicities, body weights, and socioeconomic statuses. See Figure 13.3<sup>30</sup> for an illustration of its diverse impact. Some individuals are mildly affected throughout their lives but then triggered by a significant physical or emotional life event that manifests clinical worsening.

of Health and Human Services. <https://www.nimh.nih.gov/health/topics/eating-disorders>

30. [“eating-disorders-everyone-sm.jpg”](#) by [U.S. Department of Health and Human Services, National Institutes of Health, National Institute of Mental Health](#) is in the [Public Domain](#)



Figure 13.3 Eating Disorders Affect Anyone

## Cultural Considerations

Cultural beliefs impact self-concept and satisfaction with body size. Anorexia nervosa is associated with cultures that value thinness. When caring for a client with an eating disorder, cultural considerations need to be addressed to ensure effective and culturally sensitive care. For example, Western cultures often emphasize thinness, which can contribute to body dissatisfaction and eating disorders. Furthermore, studies indicate that social media significantly influences these beliefs. For example, one study found that participants with higher use of social media had significantly greater odds of having eating concerns.<sup>31</sup> See Figure 13.4<sup>32</sup> for an image of an extremely underweight fashion model in a culture that values thinness.

<sup>31</sup>. Sidani, J. E., Shensa, A., Hoffman, B., Hanmer, J., & Primack, B. A. (2016). The association between social media use and eating concerns among us young adults. *Journal of the Academy of Nutrition and Dietetics*, 116(8), 1465-1472. <https://doi.org/10.1016/j.jand.2016.03.021>

<sup>32</sup>. “7991065935\_8f05b38f46\_k” by Farrukh is licensed under [CC BY-NC 2.0](#).



Figure 13.4 Underweight model

Black and Hispanic teenagers are more likely to suffer from bulimia nervosa. Additional considerations are the influences of some sports cultures on athletes where weight is a consideration, such as in wrestling, gymnastics, figure skating, and body building.<sup>33</sup> Nurses should be culturally competent and aware of the specific cultural context of their clients. This includes understanding cultural norms, values, and potential barriers to treatment.

View a supplementary YouTube video<sup>34</sup> about eating disorders: [Eating Disorders: Psychiatric Mental Health for Nursing Students | @LevelUpRN](https://www.youtube.com/watch?v=IFBXsrSvIOI)

33. National Eating Disorders Association. (n.d.). *Statistics and research on eating disorders*. <https://www.nationaleatingdisorders.org/statistics-research-eating-disorders>

34. Level Up RN. (2023, September 11). *Eating disorders: psychiatric mental health for nursing students | @LevelUpRN* [Video]. Youtube. All rights reserved. <https://www.youtube.com/watch?v=IFBXsrSvIOI>

## 13.3 Treatment for Eating Disorders

Early treatment is important for individuals with eating disorders because of increased risk for suicide, self-injury behaviors, and medical complications. People with eating disorders may also have other mental health disorders (such as depression or anxiety) or problems with substance use. There are a variety of treatments that have been shown to be effective in treating eating disorders. Generally, treatment is more effective before the disorder becomes chronic, but even people with long-standing eating disorders can recover.<sup>1</sup> It is important to note that treatment can be difficult for some individuals to obtain due to limited options and scarce resources in many rural areas.

The choice of treatment depends on the specific type of eating disorder, the severity of the condition, and the individual's needs. Treatment plans may include one or more of the following:

- Individual, group, and/or family psychotherapy
- Medications
- Nutritional counseling
- Medical care and monitoring

### Psychotherapy

Cognitive behavioral therapy (CBT) is used to reduce or eliminate binge eating and purging behaviors. Therapy focuses on modifying dysfunctional

1. National Institute of Mental Health. (2024). *Eating disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/eating-disorders>

thoughts and behaviors related to eating, body image, and weight.<sup>2</sup> Limited provider availability may require individuals to complete therapy virtually.

Read more about CBT in the “[Treatments for Depression](#)” section of the “Depressive Disorders chapter.

**Family-based therapy** a type of psychotherapy in which parents take an active role in managing their child’s eating behaviors and weight restoration. This approach involves parents carefully monitoring all of the client’s eating and activities to prevent excessive exercise and purging. The goal is to promote healthy eating behaviors and weight restoration, with parents providing structure and support to reduce parental distress and anxiety. As treatment progresses, autonomy in feeding is gradually shifted back to the adolescent, and the focus shifts to improving family communication and independence. This therapy has been found to be very effective in helping adolescents gain weight and improve eating habits and moods.<sup>3</sup>

**Interpersonal Psychotherapy (IPT):** IPT is another effective treatment, focusing on improving interpersonal functioning and reducing eating disorder symptoms.

## Medications

Evidence also suggests that medications such as antidepressants, antipsychotics, or mood stabilizers may also be helpful for treating eating

2. National Institute of Mental Health. (2024). *Eating disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/eating-disorders>
3. National Institute of Mental Health. (2024). *Eating disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/eating-disorders>



disorders and other co-occurring mental illnesses such as anxiety or depression.<sup>4</sup> Fluoxetine, a selective serotonin reuptake inhibitor (SSRI), is FDA-approved for the treatment of bulimia nervosa (BN) and has demonstrated efficacy in reducing binge-purge behaviors. Lisdexamfetamine, approved for binge eating disorder (BED), effectively decreases the frequency of binge eating episodes. Although not FDA-approved for anorexia nervosa (AN), olanzapine has shown potential benefits by promoting weight gain and alleviating anxiety symptoms in individuals with AN.<sup>5</sup>

- ▶ Read more about antidepressants in the “[Treatments for Depression](#)” section of the “Depressive Disorders” chapter.
- ▶ Read more about antipsychotics in the “[Schizophrenia](#)” section of the “Psychosis and Schizophrenia” chapter.

## Nutritional Rehabilitation & Counseling

Nutritional rehabilitation is essential for all eating disorders, nutritional rehabilitation aims to restore healthy eating patterns and address malnutrition. Registered dietitian nutritionists play a crucial role in this aspect of treatment. Nutritional counseling by a dietician with specialized training is necessary for individuals with eating disorders. The counseling should

4. National Institute of Mental Health. (2024). *Eating disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/eating-disorders>
5. Muratore, A. F., & Attia, E. (2022). Psychopharmacologic management of eating disorders. *Current Psychiatry Reports*, 24(7), 345-351. doi: [10.1007/s11920-022-01340-5](https://doi.org/10.1007/s11920-022-01340-5).

incorporate education about nutritional needs, as well as planning and monitoring healthy food choices.<sup>6</sup>

## Medical Care and Monitoring

Medical treatments for eating disorders can be delivered in a variety of settings and are guided by the severity and chronicity of the illness, medical stability, insurance coverage, and available community resources. The following characteristics apply to the setting selected for an individual's treatment<sup>7</sup>:

- **Outpatient Treatment:** This is suitable for clients who are medically stable and can maintain their daily activities. It includes regular visits to a healthcare team comprising a physician, therapist, and dietitian.
- **Intensive Outpatient:** The client is medically and psychiatrically stable and does not need daily medical monitoring. Symptoms are under sufficient control for the individual to be able to function in normal social, educational, or vocational situations and continue to make progress in recovery.
- **Partial Hospitalization:**
  - The client is medically stable. The eating disorder impairs functioning but is without immediate risk. The client requires daily assessment of physiologic and mental status.
  - The client is psychiatrically stable but is unable to function in normal social, educational, or vocational situations. They engage in daily binge eating, purging, fasting, restricted food intake, or other pathogenic weight control techniques.

6. National Eating Disorders Association. (n.d.). *Treatment*.  
<https://www.nationaleatingdisorders.org/treatment>

7. National Eating Disorders Association. (n.d.). *Treatment*.  
<https://www.nationaleatingdisorders.org/treatment>

- **Residential:** The client is medically stable and requires no intensive medical intervention. They are psychiatrically impaired and unable to respond to partial hospital or outpatient treatment.
- **Inpatient:**
  - The client is medically unstable as determined by:
    - Unstable or depressed vital signs
    - Laboratory findings presenting acute health risk
    - Complications due to coexisting medical problems such as diabetes
  - The client is psychiatrically unstable as determined by:
    - Rapidly worsening symptoms
    - Suicidal ideation with a plan and unable to contract for safety

## 13.4 Applying the Nursing Process to Eating Disorders

People with eating disorders may appear healthy even when they are very ill. Additionally, individuals with anorexia nervosa often do not view their behavior as a problem. They are typically only seen in health care settings due to concerned family or friends who encourage them to seek treatment. Conversely, individuals with bulimia nervosa or binge eating disorder may feel shame and sensitivity to the perceptions of others regarding their illness. Therefore, it is vital for the nurse to build a therapeutic nurse-client relationship with clients with eating disorders and empathize with possible feelings of low self-esteem and lack of control over eating.<sup>1</sup>

This section will apply the nursing process to anorexia and bulimia nervosa.

### Assessment (Recognizing Cues)

When assessing an individual with a potential or diagnosed eating disorder, it is vital to obtain their perception of the problem while assessing for signs and symptoms. Care planning that does not address their perspective will not be effective. As previously mentioned, clients with anorexia nervosa often do not perceive their behaviors as a problem, so specialized therapeutic techniques are required. Review signs and symptoms associated with various eating disorders in the “[Basic Concepts](#)” section.

A complete nursing assessment includes mental status examination, psychosocial assessment, and screening for risk of suicide or self-harm. Nutritional patterns, fluid intake, and daily exercise should also be assessed. If the client has a bingeing or purging pattern, the amount of food eaten and/or the frequency of these behaviors should be assessed.

1. Halter, M. (2022). *Varc Carolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

## Mental Status Examination

See Table 13.4a for common findings when assessing a client with an eating disorder. (See expected findings for these components of a mental status examination in the “[Assessment](#)” section in Chapter 4.) Critical findings that require immediate notification of the provider are bolded with an asterisk.

Table 13.4a Common Findings During Mental Status Examinations for Clients With Eating Disorders<sup>2</sup>

2. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

Mental Status Examination Component	Common Findings in Eating Disorders (*Indicates immediately notify provider)
<b>Signs of Distress</b>	May appear anxious, preoccupied with food, weight, or body image; may exhibit shame, denial, or resistance when discussing eating behaviors. Clients with bulimia may appear more outwardly distressed or dysregulated; those with anorexia may appear emotionally blunted.
<b>Level of Consciousness and Orientation</b>	Typically alert and oriented unless severely malnourished, in which case confusion, poor concentration, or delayed responses may occur.
<b>Appearance and General Behavior</b>	May appear underweight (anorexia), within normal weight range (bulimia), or overweight (binge eating disorder). Common findings: baggy clothing to hide body shape, pale or dry skin, lanugo (fine body hair), brittle hair/nails, signs of vomiting (e.g., Russell's sign, dental erosion). Behavior may be avoidant or perfectionistic.
<b>Speech</b>	Generally normal in rate and volume, but may be guarded or evasive about food-related topics. Some clients may exhibit pressured speech if anxious; others may appear overly controlled.
<b>Motor Activity</b>	May be restless or hyperactive (common in anorexia), or sluggish due to malnutrition. Repetitive or ritualistic movements (e.g., fidgeting, checking body parts) may be observed.
<b>Mood and Affect</b>	Mood is often anxious, irritable, or depressed. Affect may be blunted or restricted, especially in anorexia. Clients may express low self-worth or describe feelings of shame, guilt, or emptiness.

<b>Thought and Perception</b>	<p>Thought content often centers on body image distortion, weight obsession, and fear of gaining weight. Clients may express rigid thinking patterns or perfectionism.</p> <p><b>*Suicidal ideation may be present in severe cases or with coexisting depression.</b></p>
<b>Attitude and Insight</b>	<p>Often demonstrate poor insight into the severity of the disorder. Clients may deny being underweight or minimize behaviors such as purging or restricting. May resist treatment, particularly if weight restoration is a goal.</p>
<b>Cognitive Abilities and Level of Judgment</b>	<p>Memory and attention typically intact unless severely malnourished. Clients may exhibit obsessive thoughts, poor concentration, and rigid cognitive patterns. Judgment and decision-making may be impaired by body image distortions.</p>

## Psychosocial Assessment

As previously discussed in the “[Application of the Nursing Process in Mental Health Care](#)” chapter, a psychosocial assessment obtains additional subjective data that detects risks and identifies treatment opportunities and resources.<sup>34</sup>

- Reason for seeking health care (i.e., “chief complaint”)
- Thoughts of self-harm or suicide (both current and historical)

3. Clinical Practice Guideline for the Treatment of Posttraumatic Stress Disorder. (2017). *What is cognitive behavioral treatment?* American Psychological Association. <https://www.apa.org/ptsd-guideline/patients-and-families/cognitive-behavioral>
4. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

- Cultural assessment
- Spiritual assessment
- Family dynamics
- Current and past medical history
- Current medications
- History of previously diagnosed mental health disorders
- Previous hospitalizations
- Educational background
- Occupational background
- History of exposure to psychological trauma, violence, and domestic abuse
- Substance use (tobacco, alcohol, recreational drugs, misused prescription drugs)
- Family history of mental illness
- Coping mechanisms
- Functional ability/Activities of daily living

After identifying the reason the client is seeking health care, additional focused questions are used to obtain detailed information used to plan care. The mnemonic PQRSTU can be used to ask questions in an organized fashion. See Table 13.4b for a sample PQRST assessment for eating disorders and sample responses by a client.

Table 13.4b Sample PQRSTU Questions for Assessing Eating Disorders



PQRSTU	Sample Questions	Sample Client Response
<b>Provocation/ Palliation</b>	<p>“What seems to trigger your eating behaviors or concerns about food or your body?”</p> <p>“What helps you cope or feel better?”</p>	<p>“It usually starts when I feel stressed or out of control. Looking at myself in the mirror makes it worse. Skipping meals helps me feel like I’m in control again.”</p>
<b>Quality</b>	<p>“Can you describe the thoughts or feelings you have about food or your body?”</p>	<p>“I constantly think about calories and weight. Even when I eat something small, I feel guilty and disgusting. It feels like I’m never thin enough.”</p>
<b>Region</b>	<p>“Do you feel any physical effects from your eating behaviors—like pain, fatigue, or other sensations?”</p>	<p>“I’m always cold and tired. Sometimes I get dizzy when I stand up too fast. My stomach hurts, but I don’t really feel hungry anymore.”</p>
<b>Severity</b>	<p>“How much does this affect your daily life?”</p>	<p>I think about food and my body all day. It’s hard to focus on school or even talk to my friends.”</p>
<b>Timing/ Treatment</b>	<p>“When did these behaviors or thoughts start? How often do they happen?”</p>	<p>“It started around two years ago, but got worse this past year. I am very careful about what I eat most days, but sometimes I binge and then purge at night.”</p>
<b>Understanding</b>	<p>“What do you believe is causing this? How do you understand what you’re going through?”</p>	<p>“When my life feels overwhelming, I focus on food and weight because it’s the one thing I can manage successfully.”</p>

## SUICIDE AND SELF INJURY SCREENING

Clients being evaluated or treated for eating disorders often have suicidal

ideation, especially if they are also experiencing a depressive disorder. The Patient Safety Screener (PSS-3) is an example of a brief screening tool to detect suicide risk in all client presenting to acute care settings.<sup>5</sup>

**Non-suicidal self-injury (NSSI)** refers to intentional self-inflicted destruction of body tissue without suicidal intention and for purposes not socially sanctioned. Common forms of NSSI include behaviors such as cutting, burning, scratching, and self-hitting. It is considered a maladaptive coping strategy without the desire to die. NSSI is a common finding among adolescents and young adults in psychiatric inpatient settings.<sup>6</sup>

Review information about suicide screening, the Patient Safety Screener, and screening for non-suicidal self-injury (NSSI) in the “[Assessment](#)” section of the Applying the Nursing Process to Mental Health Care” chapter.

## CULTURAL ASSESSMENT

Cultural Formulation Interview (CFI) questions help nurses understand a client’s cultural background and how it influences their experience of eating disorders.<sup>7</sup> Sample CFI questions focused specifically on understanding eating disorders within a cultural context include the following:

5. Suicide Prevention Resource Center. (n.d.). *The patient safety screener: A brief tool to detect suicide risk*. <https://sprc.org/micro-learning/patientsafetyscreener>
6. Cipriano, A., Cella, S., & Cotrufo, P. (2017). Nonsuicidal self-injury: A systematic review. *Frontiers in Psychology*, 8. <https://www.frontiersin.org/articles/10.3389/fpsyg.2017.01946/full>
7. DeSilva, R., Aggarwall, N. K., & Lewis-Fernandez, R. (2015). The DSM-5 cultural formulation interview and the evolution of cultural assessment in psychiatry. *Psychiatric Times*, 32(6). <https://www.psychiatrictimes.com/view/dsm-5-cultural-formulation-interview-and-evolution-cultural-assessment-psychiatry>

- Cultural Definition of the Problem
  - “How would you describe the difficulties you’re having with eating, food, or your body?”
  - “Are there any words or used in your community or family to describe what you’re going through?”
- Cultural Perceptions of Cause, Context, and Support
  - “What do you think is causing these problems with eating or your body image?”
  - “Are there cultural or family expectations around food, weight, or appearance that you think affect you?”
  - “How do your family and friends view your eating habits? Do they express concerns about your weight?”
  - “Do you feel any pressure from your friends to look or eat a certain way?”
- Cultural Factors Affecting Coping and Help-Seeking
  - “What kinds of things have you done to cope with these feelings so far?”
  - “Are there any traditional remedies, rituals, or religious practices you use to feel better?”
  - “Have you tried to talk to anyone about these feelings, like family members, friends, religious leaders, or traditional healers?”
  - “Do people in your community seek help for issues with food or body image? If so, from whom?”
- Cultural Features of the Nurse–Client Relationship
  - ““Is there anything I should know about your background or beliefs that would help me better understand you?”
  - “Do you have any concerns or hesitations you have about seeing a mental health professional?”
  - “What kind of help do you think would work best for you?”

Findings from the cultural formulation interview are used to individualize a client’s plan of care to their preferences, values, beliefs, and goals.

## Objective Assessment

Common objective assessment findings for individuals with anorexia nervosa and bulimia nervosa are compared in Table 13.4c. Note that clients with binge eating disorder may have obesity and gastrointestinal symptoms but do not typically have other associated abnormal assessment findings.

Table 13.4c Comparison of Assessment Findings in Anorexia Nervosa and Bulimia Nervosa<sup>8</sup>

8. Halter, M. (2022). *Varc Carolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

<b>Anorexia Nervosa</b>	<b>Bulimia Nervosa</b>
Low weight	Normal to slightly low weight
Muscle weakening (from starvation and electrolyte imbalance)	Muscle weakening (from electrolyte imbalance)
Peripheral edema (from hypoalbuminemia)	Peripheral edema (from rebound fluids if diuretics are used)
Cardiovascular abnormalities (hypotension, bradycardia, heart failure from starvation, and dehydration)	Cardiovascular abnormalities (cardiomyopathy and cardiac dysrhythmias from electrolyte imbalances)
Abnormal lab results (hypokalemia and anemia from starvation)	Electrolyte imbalances (hypokalemia and hyponatremia from diuretics, laxatives, or vomiting)
<p>Other signs:</p> <p>Amenorrhea (lack of menstruation)</p> <p>Lanugo (growth of fine hair all over the body)</p> <p>Cold extremities</p> <p>Constipation</p> <p>Impaired renal function</p> <p>Decreased bone density</p> <p>Wearing many layers of clothing due to feeling cold and also to hide body frame.</p>	<p>Other signs:</p> <p>Tooth erosion or dental caries (from vomiting reflux over enamel)</p> <p>Parotid swelling (due to increased serum amylase levels)</p> <p>Calluses or scars on hand (from self-induced vomiting)</p> <p>Seizures (purging via self-induced vomiting lowers seizure threshold)</p>

## Diagnostic and Lab Work

Laboratory and diagnostic testing are typically performed to rule out thyroid imbalances and to evaluate for potential physiological complications resulting

from starvation, dehydration, and electrolyte imbalances. Laboratory testing may include the following<sup>9</sup>:

- Complete blood count
- Electrolyte levels
- Glucose level
- Thyroid function tests
- Erythrocyte sedimentation rate (ESR)
- Creatine phosphokinase (CPK)

Diagnostic testing may include these tests:

- Electrocardiogram (ECG)
- Dual energy X-ray absorptiometry (DEXA) to measure bone density

## Lifespan Considerations

Life span considerations influence how the client is assessed, as well as the selection of appropriate nursing interventions. It is important to individualize all interventions to the age and developmental level of the client. Review developmental stages in the “[Application of the Nursing Process in Mental Health Care](#)” chapter.

## CHILDREN AND ADOLESCENTS

Eating disorders (EDs) are often underdiagnosed and many adolescents go untreated, do not recover, or only reach partial recovery. Denial of symptoms is common. Higher rates of EDs are being seen in younger children, boys, and in minority groups. EDs are also being increasingly recognized in clients with obesity. Family influence must be considered. Individual treatment including

9. Halter, M. (2022). *Varc Carolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

cognitive behavioral therapy and family-based treatment focused on weight restoration and reducing blame can be effective.<sup>10</sup>

## OLDER ADULTS

Older adults may present with unintentional weight loss, disinterest in food, or somatic complaints. Their focus may be on control rather than weight or body image. Symptoms may be misattributed to aging, other medical conditions, dementia, or side effects of medications. Perceived stigma may prevent older adults from discussing eating habits, body image concerns, or feelings. Grief, isolation, depression, and anxiety may increase risk for eating disorders.<sup>11</sup>

## Diagnosis (Analyzing Cues)

Common nursing diagnoses for individuals diagnosed with anorexia nervosa or bulimia nervosa include these diagnoses<sup>12</sup>:

- Imbalanced Nutrition: Less Than Body Requirements
- Imbalanced Nutrition: More Than Body Requirements
- Risk for Electrolyte Imbalance
- Risk for Imbalanced Fluid Volume
- Impaired Body Image

10. Campbell, K., & Peebles, R. (2014). Eating disorders in children and adolescents: State of the art review. *Pediatrics*, 134(3), 582–592. <https://doi.org/10.1542/peds.2014-0194>

11. Lapid, M. I., Prom, M. C., Burton, M. C., et al. (2010). *Eating disorders in the elderly*. *International Psychogeriatrics*, 22(4), 523–536, <https://doi.org/10.1017/S1041610210000104>

12. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

- Ineffective Coping
- Interrupted Family Processes
- Chronic Low Self-Esteem
- Powerlessness
- Risk for Spiritual Distress

## Outcome Identification (Generate Solutions)

Outcomes are individualized to client's situation and diagnosis and symptoms and should address the acute nursing diagnosis with prioritization on safety. These are the typical overall treatment goals for individuals with eating disorders<sup>13</sup>:

- Restoring adequate nutrition
- Bringing weight to a healthy level
- Reducing excessive exercise
- Stopping binge-purge and binge eating behaviors

SMART expected outcomes are individualized for each client based on their established nursing diagnoses and current status. (SMART is an acronym for Specific, Measurable, Attainable/Actionable, Relevant, and Timely.) Examples of SMART outcomes for an individual hospitalized with anorexia nervosa who is experiencing electrolyte imbalances are:

- *The client will maintain a normal sinus heart rhythm with a regular rate during their hospitalization.*
- *"The client will achieve and maintain normal electrolyte levels (potassium, sodium, magnesium, and phosphorus) within the reference range, as measured by daily blood tests, for at least 7 consecutive days before discharge, through adherence to the prescribed nutritional plan*

13. National Institute of Mental Health. (2024). *Eating disorders*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/eating-disorders>



*and medical interventions.”*

## Planning (Generate Solutions)

### Safety

Safety receives top priority when planning and implementing interventions for clients with eating disorders who are at risk of suicide. If the client is exhibiting risk for suicide, a safety plan should be immediately implemented. Review nursing care for clients with risk for suicide in the “[Application of the Nursing Process in Mental Health Care](#)” chapter.

Planning other interventions depends on the acuity of the client’s clinical status and their established expected outcomes. As previously discussed, clients are hospitalized for stabilization. Common criteria for hospitalization include extreme electrolyte imbalance, weight below 75% of healthy body weight, arrhythmias, hypotension, temperature less than 98 degrees Fahrenheit, or risk for suicide.<sup>14</sup> After a client is medically stable, the treatment plan includes a combination of psychotherapy, medications, and nutritional counseling. Review the “[Treatment for Eating Disorders](#)” section for more details.

## Implementation (Take Action)

### Inpatient Care

Severely malnourished clients may require therapeutic enteral nutrition. Any client with negligible food intake for more than five days is at risk of developing a potentially fatal complication called **refeeding syndrome**. The hallmark feature of refeeding syndrome is hypophosphatemia but may also involve serious sodium and fluid imbalances; changes in glucose, protein, and

14. Halter, M. (2022). *Varc Carolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

fat metabolism; thiamine deficiency; hypokalemia; and hypomagnesaemia. To avoid this syndrome, a thorough nutritional assessment must be performed followed by the slow reintroduction of nutrients and fluids according to evidence-based guidelines.<sup>15</sup>

After resolving acute symptoms, clients with anorexia nervosa begin a weight restoration program for incremental weight gain with a treatment goal set for 90% of ideal body weight. Specially trained dietitians assist in developing daily meal plans and caloric intake, and clients are generally weighed two or three times a week to gauge progress.<sup>16</sup>

Nurses should be aware that clients with bulimia nervosa typically establish a therapeutic nurse-client relationship more quickly than clients with anorexia nervosa. As previously discussed in this chapter, clients with anorexia nervosa often do not view their condition as a disorder and value their obsessive-compulsive behaviors with eating as a way to feel safe and secure and avoid negative feelings. Conversely, clients with bulimia nervosa view their behaviors as problematic and desire help.<sup>17</sup>

## Outpatient Care

Outpatient partial hospitalization is an option for clients who have been medically stabilized. In this setting, clients are in a clinical setting during the day and then go home to practice skills in the afternoon. Outpatient

15. Mehanna, H. M., Moledina, J., & Travis, J. (2008). Refeeding syndrome: What it is, and how to prevent and treat it. *BMJ*, 336, 1495–1498. <https://doi.org/10.1136/bmj.a301>
16. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.
17. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

treatment continues if the client maintains a contracted weight, vital signs are within a normal range, and there is an absence of disordered eating behaviors.<sup>18</sup>

A significant part of the recovery process includes rebuilding relationships with family. Family members or significant others often feel frustrated, powerless, and hopeless. Family members may feel overwhelmed and exhausted at the level of care and monitoring required to support the client with an eating disorder into recovery because the strategies they previously attempted, such as forcing the client to eat or begging the client to eat, were not successful. The nurse helps with this recovery process by providing education to the client and their loved ones about the illness, treatment, and meal planning. Adaptive coping skills to address disordered thoughts should be reinforced.<sup>19</sup> Review information about coping strategies in the “[Stress, Coping, and Crisis Intervention](#)” chapter.

## Nursing Interventions for Eating Disorders Based on Categories of the APNA Implementation Standard

Nurses individualize interventions based on the client’s current clinical status and their phase of treatment. Interventions can be categorized based on the American Psychiatric Nursing Association (APNA) standard for *Implementation* that includes the *Coordination of Care; Health Teaching and Health Promotion; Pharmacological, Biological, and Integrative Therapies; Milieu Therapy; and Therapeutic Relationship and Counseling*. (Review information about these subcategories in the “[Application of the Nursing](#)

18. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

19. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

Process in Mental Health Care” chapter.) Read nursing interventions for clients with eating disorders categorized by APNA categories in Table 13.4d.

Table 13.4d Examples of Nursing Interventions for Clients with Eating Disorders by APNA Subcategories<sup>20, 21</sup>

20. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.
21. American Nurses Association, American Psychiatric Nurses Association, and International Society of Psychiatric-Mental Health Nurses. (2014). *Psychiatric-Mental Health Nursing: Scope and Standards of Practice* (2nd ed.).

Subcategory of the APNA Standard of Implementation	The nurse will ...	Rationale
<b>Coordination of Care</b>	<ul style="list-style-type: none"> <li>• Communicate client trends with interprofessional team members, such as risk for suicide and target weight. A target weight and daily caloric intake are set in collaboration with the dietitian and the provider.</li> <li>• Collaborate with interdisciplinary team (physician, dietitian, therapist, social worker) to create a personalized care plan. Include family members when appropriate.</li> <li>• Refer to community resources and outpatient treatment.</li> </ul>	<p>Eating disorders require multidisciplinary management to address medical, psychological, and nutritional needs. Involving the family supports continuity and adherence.</p>

<b>Health Teaching and Health Promotion</b>	<ul style="list-style-type: none"> <li>• Provide education about the physical and psychological effects of eating disorders.</li> <li>• Teach about nutrition, body image, and coping strategies using age-appropriate language. Address myths about weight, food, and health.-Promote health by teaching adaptive coping strategies such as journaling. Support basic skills such as learning how to create meal plans, shopping at the grocery store, and navigating family or social eating situations.</li> </ul>	<p>Education increases insight, empowers recovery, and reduces maladaptive beliefs. Promoting healthy lifestyle choices supports long-term change. Nurses encourage resilience by promoting healthy coping strategies, communication, and problem-solving skills.</p>
---	---	---

<b>Pharmacological, Biological, and Integrative Therapies</b>	<ul style="list-style-type: none"> <li>• Deliver client education about antidepressants or other prescribed medications with expected time frames for improvement.</li> <li>• Monitor for side effects and signs of malnutrition (e.g., bradycardia, electrolyte imbalance).</li> <li>• Incorporate mindfulness, journaling, or guided imagery to reduce anxiety.</li> </ul>	<p>Medications may help with comorbid anxiety or depression. Integrative therapies enhance self-regulation and reduce relapse risk. Medical monitoring ensures safety during nutritional rehabilitation.</p>
---	--	--

<b>Milieu Therapy</b>	<ul style="list-style-type: none"> <li>• Create a structured, nonjudgmental environment with consistent meal and weigh-in routines.</li> <li>• Establish therapeutic norms for group therapy and communal eating.</li> <li>• Monitor for food hoarding, purging, or excessive exercise.-Provide a pleasant, calm atmosphere at mealtimes. Emphasize the social nature of eating. Encourage conversations during mealtimes that do not involve the topics of eating or exercise.-Observe clients during meals to prevent hiding or throwing away food and at least one hour after eating to prevent purging.-Encourage the client to make their own menu choices as they</li> </ul>	<p>A therapeutic milieu provides predictability and safety, reduces secrecy, and supports behavioral stabilization. It also models healthy social interactions. The milieu of an eating disorder specialty unit is purposefully organized to assist the client in establishing healthy eating patterns and normalization of eating. The highly structured environment provides precise mealtimes, adherence to the meal plan, close observation of bathroom trips, and monitoring potential access to laxatives or diuretics. Mealtimes can cause episodes of high anxiety. The client should feel accepted and safe from judgmental evaluations in the milieu with a focus on eating behaviors and underlying feelings of anxiety, dysphoria, low self-esteem, and a lack of control.</p>
-----------------------	--	--



	approach their goal weight.	
--	-----------------------------	--

<b>Therapeutic Relationship and Counseling</b>	<ul style="list-style-type: none"> <li>• Provide 1:1 therapeutic communication to encourage the client to develop adaptive coping strategies, use stress management techniques, develop supportive relationships, and seek spiritual resources.</li> <li>• Acknowledge the emotional and physical difficulty the client is experiencing.</li> <li>• Use motivational interviewing and contract with the client to increase their ownership of treatment goals.</li> <li>• Weigh the client daily in their underwear for the first week and then three times a week. Do not allow oral intake before the morning weigh-in. It is permissible for the client to not view the scale during the</li> </ul>	<p>Effective therapeutic techniques for clients with depression can promote hope and positive self-esteem. Clients with eating disorders often struggle with control and perfectionism; a supportive approach fosters motivation for recovery.</p> <p>The first priority is to establish a therapeutic relationship. A client's feelings of fatigue can be used to engage cooperation in the treatment plan.</p> <p>Motivational interviewing is a collaborative, goal-oriented style of communication. It is designed to strengthen personal motivation and commitment to specific goals by eliciting and exploring the person's reasons for change within an atmosphere of acceptance and compassion.<sup>22</sup></p> <p>Accurate weight taking and monitoring are vital. The client may try to control and sabotage the weight monitoring. The client is typically expected to gain 0.5 pound on a specific schedule. However, weight gain of more than five pounds in one week can cause pulmonary edema. The particulars of how clients should be weighed (i.e., open vs. blind weighed) is a point of debate in the field. Because viewing the scale can cause anxiety, blind weighing is typically used during the acute stage of treatment, whereas open weighing may be suitable at later stages of recovery.<sup>23</sup></p>
--	--	--

22. Miller, W. R., & Rollnick, S. (2013). *Motivational Interviewing: Helping people to change* (3rd ed.). Guilford Press.
23. Foreich, F. V., Ratcliffe, S. E., & Vartanian, L. R. (2020). Blind versus open weighing from an eating disorder patient perspective. *Journal of Eating Disorders* 8, 39. <https://doi.org/10.1186/s40337-020-00316-1>

	<p>weigh-in.</p> <ul style="list-style-type: none"> <li>• Administer liquid supplements as prescribed.</li> <li>• Be empathetic with the client's struggle to give up control of their eating and weight as they are expected to regain weight. Encourage the clients to verbalize or use a journal to record their feelings surrounding eating disorder behaviors. Confront irrational thoughts and beliefs to promote healthy eating behaviors.</li> <li>• Monitor physical activity and individualize the client's plans for exercise.</li> <li>• Focus on the client's strengths, including their work on normalizing weight and eating behaviors. Reinforce the knowledge and skills gained from</li> </ul>	<p>Oral or enteral supplements may be prescribed based on the client's status. However, be alert for refeeding syndrome in severely malnourished clients.</p> <p>External control is required initially to promote good nutrition and a healthy weight. Cognitive and behavioral changes will occur gradually.</p> <p>The client often experiences a strong drive to exercise. Nurses can assist in planning a reasonable amount of exercise.</p> <p>Acknowledge milestones and encourage other sources of gratification other than eating.</p>
--	--	---

	individual, family, and group therapy sessions.	
--	---	--

## Nursing Interventions for Physiological Signs of Eating Disorders

Nursing interventions also target common physiological signs of eating disorders as summarized in Table 13.4e.

Table 13.4e Nursing Interventions Targeting Physiological Signs of Eating Disorders<sup>24</sup>

24. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

Problem/Intervention	Rationale
<p><b>Nutrition</b></p> <ul style="list-style-type: none"> <li>• Monitor weight, caloric intake, and food behavior during meals.</li> <li>• Collaborate with dietitian to develop a structured, individualized meal plan.</li> <li>• Supervise meals and post-meal periods to prevent purging.</li> <li>• Monitor vital signs and signs of dehydration (dry mucosa, hypotension, tachycardia).</li> <li>• Assess lab values (K<sup>+</sup>, Na<sup>+</sup>, Mg<sup>++</sup>, phosphate) frequently.</li> <li>• Monitor for bradycardia, hypotension, orthostatic changes, and arrhythmias.</li> <li>• Encourage oral fluid intake and restrict excessive water consumption (often used for weight manipulation).</li> </ul>	<p>Clients with eating disorders often experience malnutrition, electrolyte imbalances, and risk of refeeding syndrome. Structured support ensures gradual, safe nutritional rehabilitation. Electrolyte disturbances (especially hypokalemia) are life-threatening and common with vomiting, laxative use, and restriction. Malnutrition, purging, and starvation can lead to life-threatening cardiac changes, especially in anorexia. Early detection is critical to prevent sudden cardiac arrest. Close monitoring prevents cardiac and renal complications.</p>
<p><b>Sleep/Fatigue</b></p> <ul style="list-style-type: none"> <li>• Assess sleep patterns and disturbances.</li> <li>• Encourage consistent sleep-wake cycles and limit daytime naps. Provide periods of rest after activities, if needed.</li> <li>• Teach about relaxation techniques.</li> </ul>	<p>Anxiety, hyperactivity, and malnutrition often disrupt sleep. Adequate rest supports mood stabilization, cognition, and healing.</p>

<b>Elimination</b> <ul style="list-style-type: none"> <li>• <b>Assess for constipation, diarrhea, or laxative abuse.</b></li> <li>• <b>Encourage dietary fiber and hydration.</b></li> <li>• <b>Educate on normal bowel patterns.</b></li> </ul>	<p>Disordered eating and purging can cause gastrointestinal dysfunction. Laxative abuse and food avoidance alter elimination patterns and must be managed to prevent long-term complications.</p>
<b>Activity and Exercise</b> <ul style="list-style-type: none"> <li>• <b>Assess for compulsive or secretive exercise behaviors.</b></li> <li>• <b>Set limits on physical activity during medical refeeding or low BMI states.</b></li> <li>• <b>Provide alternative coping methods for emotional distress.</b></li> </ul>	<p>Excessive exercise is a common compensatory behavior in eating disorders and may worsen malnutrition and cardiac risk. Structured limits ensure safety during stabilization.</p>
<b>Self-Care Deficits</b> <ul style="list-style-type: none"> <li>• <b>Assess ability to perform ADLs (e.g., bathing, grooming, dressing).</b></li> <li>• <b>Reinforce positive self-care behaviors.</b></li> </ul>	<p>Severe malnutrition and depressive symptoms may lead to poor hygiene and low motivation. Encouraging self-care promotes dignity and recovery of function.</p>

## Communication Tips for Clients with Eating Disorders

Helpful communication techniques for building therapeutic rapport, fostering trust, and supporting recovery in individuals with eating disorders are described in the following box.

## Communication Tips for Clients with Eating Disorders<sup>25</sup>

- Use nonjudgmental, empathetic language.
  - **Rationale:** Clients with eating disorders often experience intense shame, guilt, or secrecy. An empathetic tone reduces defensiveness and encourages honesty and openness in discussing symptoms.
- Avoid focusing on weight or appearance.
  - **Rationale:** Commenting on weight, even positively, can reinforce disordered thinking. Instead, focus on behaviors, emotions, and health (e.g., “I noticed you seemed more energized today.”).
- Validate emotional experiences, not just behaviors.
  - **Rationale:** Acknowledge feelings behind behaviors (e.g., “It sounds like you were feeling overwhelmed when that happened.”). This builds trust and shifts focus from control to coping.
- Be consistent and set clear, supportive boundaries.
  - **Rationale:** Clients may test boundaries due to anxiety or control struggles. Consistent

25. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.



expectations and compassionate firmness help them feel safe and contained.

- Normalize discussion of food and body image struggles.
  - **Rationale:** Statements like “Many people in recovery have similar fears” reduce isolation and stigma, encouraging clients to share more freely.
- Collaborate in goal setting.
  - **Rationale:** Involve clients in creating achievable recovery goals. This enhances autonomy and reduces resistance, especially in those who fear loss of control.
- Frame meal plans and interventions as health-supportive, not punitive. Use open-ended questions (such as, “What makes meals difficult for you?”) instead of coercion.
  - **Rationale:** Clients with eating disorders often experience intense fear, shame, and anxiety around food, eating, and body image. Their relationship with food is not merely about nutrition but is often deeply tied to control, self-worth, identity, or coping with distress. When meal plans and interventions are presented in a rigid or authoritative manner, clients may perceive them as punitive or threatening, which can lead to increased resistance or noncompliance.

## Referral to Resources

Nurses refer clients and their loved ones to community resources. Examples of community resources are described in the following box.

### **Examples of Resources for Individuals with Eating Disorders**

- ▶ [National Eating Disorders Association \(NEDA\)](#): Support, resources, and treatment options
- ▶ [Eating Disorders Resource Group](#): Resources including treatment apps
- ▶ [ANAD](#): Eating disorder peer support groups

## Evaluation (Evaluate Outcomes)

Evaluation is a continuous process of reviewing a client's progress towards their individualized goals and SMART outcomes. Interventions are continually evaluated and modified based on their success in meeting these short-term goals.

Recovery is a long and difficult process. It is common for individuals to relapse. Attention should be paid to identifying triggers and warning signs of disordered eating before a relapse occurs. Ongoing support is critical to long term recovery. Potential long term medical consequences of eating disorders are neurological disease, cardiac damage, blood pressure complications and early onset osteoporosis.

General questions to include when evaluating a client's plan of care include the following:

- Has the patient's weight stabilized or moved closer to a healthy range?
- Are vital signs (e.g., heart rate, blood pressure, temperature) within normal limits?

- Have laboratory values (e.g., electrolytes, liver function) improved or normalized?
- Has the patient's menstrual cycle returned (if applicable)?
- Are physical symptoms (e.g., dizziness, fatigue, hair loss) improving?
- Is the patient adhering to a structured meal plan?
- Has the frequency of restrictive eating, binge eating, or purging behaviors decreased?
- Is the patient complying with treatment recommendations (e.g., attending therapy, following medical advice)?
- Are harmful behaviors (e.g., self-induced vomiting, laxative misuse, excessive exercise) being reduced or eliminated?
- Are the patient's relationships with family, friends, or peers improving?
- Is the patient participating in social activities, hobbies, or responsibilities they had previously avoided?
- Has the patient's overall quality of life and ability to function in daily life improved?
- Has the patient's preoccupation with food, weight, or body image decreased?
- Are there improvements in mood, anxiety, or emotional stability?
- Does the patient demonstrate increased insight into their illness and motivation for recovery?
- Is the patient developing healthier coping mechanisms to manage stress and emotions?

## 13.5 Spotlight Application

Consider these real-life stories of people who have struggled with eating disorders.

1. Princess Diana raised public awareness about eating disorders after sharing her experience with bulimia nervosa in recordings featured in the documentary *Diana: In Her Own Words*. After the publication, there was a sudden spike in the number of reported cases of bulimia, and many people came forward to receive the treatment they needed. Because of Diana's openness and honesty about her struggles with her eating disorder, countless people were empowered to be open and admit their own similar struggles.

▶ Read more about Princess Diana's experience in a NEDA blog titled [20 Years Later: How Princess Diana's Legacy Continues to Help People With Eating Disorders](https://staging.nationaleatingdisorders.org/princess-diana-legacy-people-with-eating-disorders/).<sup>1</sup>

2. Mike Majama, a major league baseball player, experienced an eating disorder that impacted his sports career.

1. Sundquist, A. (n.d.). *20 years later: How Princess Diana's legacy continues to help people with eating disorders* [Blog]. National Eating Disorders Association. <https://staging.nationaleatingdisorders.org/princess-diana-legacy-people-with-eating-disorders/>

- ▶ Read a *Good Morning America* [interview](#) with Mike Majama.<sup>2</sup>

3. Karla Mosley, an actress on the television show *The Bold and the Beautiful*.

- ▶ Read an [NPR interview](#) with Karla Mosley.<sup>3</sup>

4. Veterans with eating disorders.

- ▶ Read a Connecticut Health Team [story](#) about veterans with eating disorders.<sup>4</sup>

2. Thorbecke, C. (2018). *MLB star opens up about his eating disorder struggles and overcoming his 'demons.'* ABC News Internet Ventures. <https://www.goodmorningamerica.com/wellness/story/mlb-star-opens-eating-disorder-struggles-overcoming-demons-54056732>

3. Meraji, S. M. (2019). *When it comes to race, eating disorders don't discriminate.* NPR. <https://www.npr.org/sections/health-shots/2019/03/03/699410379/when-it-comes-to-race-eating-disorders-dont-discriminate>

4. McCarthy, P. (2021). *Eating disorders among veterans attributed to trauma, military weight requirements.* Connecticut Health I-Team.

## Reflective Questions

1. Compare and contrast the experiences of these individuals.
2. What strikes you most about these real-life stories?

<http://c-hit.org/2021/11/03/eating-disorders-among-veterans-attributed-to-trauma-military-weight-requirements/>

## 13.6 Learning Activities

(Answers to “Learning Activities” can be found in the “Answer Key” at the end of the book. Answers to the interactive activities are provided as immediate feedback.)

### Case Study:

Tiffany is a 24-year-old nursing student who works weekends as a certified nursing assistant. Her sister Sara brought Tiffany to a clinic appointment with a nurse practitioner because of their family’s concerns about Tiffany’s weight and eating habits. During the intake process, Tiffany states, “I don’t think there is a problem; I just don’t want to be fat.” Tiffany is 5’7” and weighs 101 pounds in her street clothes, with a calculated BMI of 15.8. When asked to perform a 24-hour diet recall, Tiffany states, “I drink coffee for breakfast, eat a low-fat yogurt for lunch, and then eat a head of iceberg lettuce for dinner most days.” She cannot recall the last time she had a menstrual cycle. Tiffany states, “I used to run five miles every day, but lately I have been too tired to run.” She denies any purging episodes. Sara states, “Tiffany does not sit down and eat meals with the rest of us. We have tried everything and cannot get her to eat. My mom has tried talking to her, begging her, and even yelling at her. If Tiffany does eat something besides lettuce, she either goes in the bathroom and vomits or runs ten miles. We are very worried about her and don’t know what to do.”

### Questions:

1. What symptoms of an eating disorder is Tiffany demonstrating?
2. What other assessments does the nurse plan for Tiffany?
3. What findings does the nurse anticipate as a result of a comprehensive assessment?
4. What laboratory tests will likely be ordered during this visit?
5. What is the goal of treatment for Tiffany?
6. What type of psychotherapy would be helpful for Tiffany?
7. What conditions would cause Tiffany to be hospitalized?
8. Tiffany is hospitalized. Create a brief nursing care plan for Tiffany including

a nursing diagnosis, SMART goal, and 3-5 nursing interventions.



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=672#h5p-45>

1



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=672#h5p-46>

2

1. "MH Eating Disorders Glossary Cards" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
2. "MH Eating Disorders Question Set 1" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)



- ▶ Test your clinical judgment with an NCLEX Next Generation-style question: [Chapter 13, Assignment 1](#)<sup>3</sup>



- ▶ Test your clinical judgment with an NCLEX Next Generation-style case study: [Chapter 13, Case Study 1](#)<sup>4</sup>



3. "MH Eating Disorders Next Gen Question by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

4. "MH Eating Disorders Next Gen Case Study by Kellea Ewen is licensed under [CC BY-NC 4.0](#)

## XIII Glossary

**Amylophagia:** Starch eating.

**Anorexia nervosa:** A condition where people avoid food, severely restrict food, or eat very small quantities of only certain foods.

**Avoidant restrictive food intake disorder (ARFID):** A condition where individuals limit the amount or type of food eaten.

**Binge eating disorder:** A condition where people lose control over their eating and have recurring episodes of eating unusually large amounts of food. Unlike bulimia nervosa, periods of binge eating are not followed by purging, excessive exercise, or fasting.

**Binge eating episode:** An episode characterized by both eating in a discrete period of time (e.g., within any two-hour period) an amount of food that is definitely larger than what most individuals would eat and a sense of a lack of control over eating during the episode; followed by purging behaviors.

**Binge-Eating/Purging Type:** Individuals experience episodes of binge eating and/or purging.

**Body mass index (BMI):** A person's weight in kilograms divided by the square of height in meters.

**Bulimia nervosa:** A condition where people have recurrent and frequent episodes of eating unusually large amounts of food and feeling a lack of control over these episodes.

**Emaciation:** A condition of extreme thinness.

**Family-based therapy:** A type of psychotherapy in which parents take an active role in managing their child's eating behaviors and weight restoration.

**Geophagia:** Clay eating.

**Lanugo:** Growth of fine hair all over the body.

**Pagophagia:** Ice eating.

**Pica:** An eating disorder in which an individual repeatedly eats things that are not considered food with no nutritional value, such as paper, dirt, soap, hair, glue, or chalk.

**Purging:** Behavior in eating disorders used to compensate for overeating, such as forced vomiting, excessive use of laxatives, or diuretics.

**Refeeding syndrome:** A potentially fatal complication for any client with negligible food intake for more than five days that involves hypophosphatemia; serious sodium and fluid imbalances; changes in glucose, protein, and fat metabolism; thiamine deficiency; hypokalemia; and hypomagnesaemia.

**Restricting Type:** Individuals severely limit the amount and type of food they consume and may engage in excessive exercise.







### Learning Objectives

- Identify assessment cues of substance use behaviors
- Identify nursing priorities for clients with substance use disorders
- Plan outcomes for clients with substance use disorders
- Differentiate safety/protective interventions for clients with substance use disorders
- Apply evidence-based practice when planning care and interventions for clients with substance use disorders
- Analyze treatments for clients with substance use disorders
- Apply the nursing process to clients with substance use disorders at risk for suicide
- Assess clients for signs of intoxication, dependency, and withdrawal symptoms and intervene appropriately
- Apply the nursing process to clients experiencing withdrawal or toxicity from substances
- Describe care for clients with nonsubstance-related dependencies (e.g., gambling, sexual addiction)
- Describe the neurobiology and risk factors for substance use disorders
- Describe protective factors and prevention programs for substance use disorders

Misuse of alcohol, drugs, and prescribed medications is estimated to cost the United States more than \$400 billion in health care expenses, law enforcement and criminal justice costs (due to drug-related crimes), lost

workplace productivity, and losses from motor vehicle crashes.<sup>1</sup> The Centers for Disease Control and Prevention (CDC) National Center for Health Statistics estimates 107,622 drug overdose deaths and 52,000 alcohol-induced deaths occurred in the United States in 2021.<sup>2</sup>

- ▶ View a visualization of recent drug overdose death rates nationally and by state at the CDC National Center for Health Statistics' Vital Statistics Rapid Release web page: [Provisional Drug Overdose Death Counts](#).

Substance use disorders significantly impact individuals, families, communities, and our society. According to the 2020 National Survey on Drug Use and Health (NSDUH), 40.3 million people in the United States aged 12 or older (14.5 percent) have a substance use disorder (SUD).<sup>3</sup>

This chapter will provide an overview of many topics related to substance use. It begins by reviewing signs of intoxication of various psychoactive substances and treatment for overdose and withdrawal symptoms. The neurobiology of substance use disorders and risk factors are explored, and then evidence-based treatment and prevention interventions are discussed based on the

1. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
2. National Center for Health Statistics. (2022). *U.S. overdose deaths in 2021 increased half as much in 2020 – but are still up 15%* [Press release]. [https://www.cdc.gov/nchs/pressroom/nchs\\_press\\_releases/2022/202205.htm](https://www.cdc.gov/nchs/pressroom/nchs_press_releases/2022/202205.htm)
3. National Survey on Drug Use and Health. (n.d.). <https://nsduhweb.rti.org/respweb/homepage.cfm>



Surgeon General's Report on Alcohol, Drugs, and Health PDF. Finally, the nursing process is applied to a hospitalized client receiving treatment for alcohol withdrawal.

# 14.2 Substances: Use, Intoxication, and Overdose

A **substance** is defined as a psychoactive compound with the potential to cause health and social problems, including substance use disorder. Substances can be divided into four major categories: alcohol, illicit drugs (including nonmedical use of prescription drugs), over-the-counter drugs, and other substances. See examples of substances known to have a significant public health impact in Table 14.2a. **Substance use** refers to the use of any of the psychoactive substances listed in Table 14.2a.

Table 14.2a Categories and Examples of Substances<sup>1</sup>

Substance Category	Examples
Alcohol	Beer, malt liquor, wine, and distilled spirits
Illicit drugs (including prescription drugs used nonmedically)	<ul style="list-style-type: none"><li>• Opioids, including heroin</li><li>• Cannabis</li><li>• Sedatives, hypnotics, and anxiolytics</li><li>• Hallucinogens</li><li>• Stimulants, including methamphetamine-like substances, cocaine, and crack</li><li>• Dextromethorphan and other cold medications</li></ul>
Over-the-counter drugs (used nonmedically)	Dextromethorphan, pseudoephedrine, and other cold medications
Other substances	Inhalants such as spray paint, gasoline, and cleaning solvents; Delta-8 THC

1. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

Millions of Americans misuse substances. See Figure 14.1<sup>2</sup> regarding the number of people aged 12 and older who reported using various substances in a one-month period of time.




Figure 14.1 Number of People Aged 12 and Older Who Reported Using Substances in the Past Month in 2020. Used under Fair Use.

2. This image is a derivative of “Past Month General Substance Use and Nicotine Vaping: Among People Aged 12 and Older; 2020” table by [Substance Abuse and Mental Health Services Administration](#). (2021). Key substance use and mental health indicators in the United States: Results from the 2020 National Survey on Drug Use and Health (HHS Publication No. PEP21-07-01-003, NSDUH Series H-56). Rockville, MD: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration. Used under Fair Use. Retrieved from <https://www.samhsa.gov/data/report/2020-nsduh-annual-national-report>

# Controlled Substances

The Controlled Substances Act is a federal law that places all **controlled substances** (i.e., substances regulated by the U.S. Drug Enforcement Agency) into one of five categories called schedules. This placement is based on the substance’s medical use, its potential for abuse or dependency, and related safety issues. For example, Schedule I drugs have a high potential for abuse and potentially cause severe psychological and/or physical dependence, whereas Schedule V drugs represent the least potential for abuse.<sup>3</sup>

**Dependence** means that when a person suddenly stops using a drug, their body goes through **withdrawal**, a group of physical and mental symptoms that can range from mild to life-threatening. See examples of controlled substances categorized by schedule in Table 14.2b.

Table 14.2b Examples of Substances by Schedule<sup>4</sup>

- 3. United States Drug Enforcement Administration. (n.d.). *The Controlled Substance Act. U.S. Department of Justice.* <https://www.dea.gov/drug-information/csa>
- 4. United States Drug Enforcement Administration. (n.d.). *The Controlled Substance Act. U.S. Department of Justice.* <https://www.dea.gov/drug-information/csa>

Schedule	Definition	Examples
<b>Schedule I</b>	No currently accepted medical use and a high potential for abuse.	Heroin, LSD, MDMA (Ecstasy), and cannabis (marijuana)
<b>Schedule II</b>	High potential for abuse, with use potentially leading to severe psychological or physical dependence. These drugs are also considered dangerous.	Hydrocodone, cocaine, methamphetamine, methadone, hydromorphone, meperidine, oxycodone, fentanyl, amphetamine/dextroamphetamine salts (Adderall), methylphenidate (Ritalin), and phencyclidine (PCP)
<b>Schedule III</b>	Moderate to low potential for physical and psychological dependence. Abuse potential is less than Schedule I and Schedule II drugs but more than Schedule IV.	Acetaminophen with codeine, ketamine, anabolic steroids, and testosterone
<b>Schedule IV</b>	Low potential for abuse and low risk of dependence.	Alprazolam (Xanax), diazepam (Valium), lorazepam (Ativan), zolpidem (Ambien), and tramadol (Ultram)
<b>Schedule V</b>	Lower potential for abuse than Schedule IV and consist of preparations containing limited quantities of certain narcotics. Generally used for antidiarrheal, antitussive, and analgesic purposes.	Cough medications with codeine, diphenoxyate/atropine (Lomotil), and pregabalin (Lyrica)

► See the [DEA alphabetized list of controlled substances PDF](#).

## Substance Misuse

**Substance misuse** is defined as the use of alcohol or drugs in a manner, situation, amount, or frequency that could cause harm to the user or to those

around them.<sup>5</sup> Substance misuse can involve a wide range of substances, including alcohol, illicit drugs, prescription medications, over-the-counter drugs, and inhalants. Misuse can be of low severity and temporary, but it can increase the risk for serious and costly consequences such as motor vehicle crashes, overdose death, suicide, various types of cancer, heart, liver, and pancreatic/liver diseases, HIV/AIDS, and unintended pregnancies. Substance use during pregnancy can cause complications for the baby such as fetal alcohol spectrum disorders (FASDs) or neonatal abstinence syndrome (NAS). Substance misuse is also associated with intimate partner violence, child abuse, and neglect.<sup>6</sup>

Substance abuse, an older diagnostic term, referred to unsafe substance use (e.g., drunk or drugged driving), use that caused legal problems, or use that continued despite failure to meet work and family responsibilities. However, the term “substance abuse” is now avoided by professionals because it does not accurately account for the neurobiological knowledge we now have about addictive disorders. Instead, the term “substance use disorder” is preferred and is further discussed in the “[Substance Use Disorder](#)” subsection of this chapter.<sup>7</sup>

5. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
6. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
7. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

# Intoxication and Overdose

**Intoxication** refers to a disturbance in behavior or mental function during or after the consumption of a substance. **Overdose** is the biological response of the human body when too much of a substance is ingested. Signs of intoxication and overdose for categories of psychoactive substances are described in the following subsections.

Anyone can call a regional poison control center at 1-800-222-1222 for consultation regarding toxic ingestion of substances and overdoses. Poison control centers are available at all times, every day of the year. Some hospitals also have toxicologists available for bedside consultation for overdoses.

## Alcohol Use and Intoxication

Based on the 2015-2020 Dietary Guidelines for Americans, a **standard drink** is defined as 14 grams (0.6 ounces) of pure alcohol. Examples of a standard drink are one 12-ounce beer, 8 – 9 ounces of malt liquor, 5 ounces of wine, or 1.5 ounces of distilled spirits. See Figure 14.2<sup>8</sup> for images of standard drinks.

8. "[NIH\\_standard\\_drink\\_comparison.jpg](#)" by [National Institutes of Health](#) is in the [Public Domain](#)




Figure 14.2 Standard Drink

The 2020 National Survey on Drug Use and Health reports that 50 percent (139.7 million) of Americans aged 12 or older use alcohol, 22.2 percent are binge drinkers, and 6.4 percent are heavy alcohol users.<sup>9</sup> **Heavy drinking** is defined as a female consuming 8 or more drinks per week or a male consuming 15 or more standard drinks per week, or either gender binge drinking on 5 or more days in the past 30 days. **Binge drinking** is defined as consuming several standard drinks on one occasion in the past 30 days; for men, this refers to drinking five or more standard alcoholic drinks on one occasion, and for women this refers to drinking four or more standard drinks on one occasion.<sup>10</sup>

**Alcohol intoxication** refers to problematic behavioral or psychological changes (e.g., inappropriate sexual or aggressive behavior, mood lability, or impaired judgment) that develop during or shortly thereafter alcohol

9. National Survey on Drug Use and Health. (n.d.). <https://nsduhweb.rti.org/respweb/homepage.cfm>

10. National Survey on Drug Use and Health. (n.d.). <https://nsduhweb.rti.org/respweb/homepage.cfm>



ingestion. Signs and symptoms of alcohol intoxication include a range of behavioral, neurological, and systemic effects. Some examples include<sup>11</sup>:

- Slurred speech
- Incoordination
- Unsteady gait
- Nystagmus
- Impairment in attention or memory
- Nausea/vomiting

## ALCOHOL OVERDOSE

An alcohol overdose occurs when there is so much alcohol in the bloodstream that areas of the brain controlling autonomic nervous system functions (e.g., breathing, heart rate, and temperature control) begin to shut down. Alcohol intoxication while also taking opioids or sedative-hypnotics (such as benzodiazepines or sleep medications) increases the risk of an overdose. Alcohol overdose can cause permanent brain damage or death.<sup>12</sup> Anyone who consumes too much alcohol too quickly is in danger of an alcohol overdose, especially for individuals who engage in binge drinking. As blood alcohol concentration (BAC) increases, so does the risk of harm. When BAC reaches high levels, blackouts (gaps in memory), loss of consciousness (passing out), and death can occur. BAC can continue to rise even when a person stops drinking or is unconscious because alcohol in the stomach and intestine

11. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

12. National Institute on Alcohol Abuse and Alcoholism, & National Institutes of Health. (2024). *Understanding the dangers of alcohol overdose*. <https://www.niaaa.nih.gov/publications/brochures-and-fact-sheets/understanding-dangers-of-alcohol-overdose>

continues to enter the bloodstream and circulate throughout the body. See Figure 14.3<sup>13</sup> for the impairments related to rising BAC.

13. “NIAAA\_BAC\_Increases\_Graphic.jpg” by [The National Institute on Alcohol Abuse and Alcoholism](#) is in the [Public Domain](#). Access for free at <https://www.niaaa.nih.gov/publications/brochures-and-fact-sheets/understanding-dangers-of-alcohol-overdose>




Figure 14.3 Impairments of Rising Blood Alcohol Concentrations

It is dangerous to assume that a person who drank an excessive amount of alcohol will “sleep it off.” One potential danger of alcohol overdose is choking on one’s vomit and dying from lack of oxygen because high levels of alcohol intake hinder the gag reflex, resulting in the inability to protect the airway. Asphyxiation can occur due to an obstructed airway or from aspiration of gastric contents into the lungs. For this reason, do not leave a person alone

who has passed out due to alcohol misuse. Keep them in a partially upright position or roll them onto one side with an ear toward the ground to prevent choking if they begin vomiting.<sup>14</sup> Critical signs and symptoms of an alcohol overdose include the following<sup>15</sup> :

- Mental confusion or stupor
- Difficulty remaining conscious or inability to wake up
- Vomiting
- Seizures
- Slow respiratory rate (fewer than 8 breaths per minute)
- Irregular breathing (10 seconds or more between breaths)
- Slow heart rate
- Clammy skin
- Dulled responses, such as no gag reflex (which prevents choking)
- Extremely low body temperature
- Bluish skin color or paleness

## MEDICAL TREATMENT OF ACUTE ALCOHOL INTOXICATION

Acute alcohol intoxication can cause hypotension and tachycardia as a result of peripheral vasodilation or fluid loss. Treatment begins with the evaluation of the client's blood alcohol level (BAC). The primary focus of care involves client stabilization and supportive care. This includes monitoring vital signs, ensuring airway protection, and preventing complications such as

14. National Institute on Alcohol Abuse and Alcoholism, & National Institutes of Health. (2024). *Understanding the dangers of alcohol overdose*.

<https://www.niaaa.nih.gov/publications/brochures-and-fact-sheets/understanding-dangers-of-alcohol-overdose>

15. National Institute on Alcohol Abuse and Alcoholism, & National Institutes of Health. (2024). *Understanding the dangers of alcohol overdose*.

<https://www.niaaa.nih.gov/publications/brochures-and-fact-sheets/understanding-dangers-of-alcohol-overdose>

hypoglycemia, hypotension, and electrolyte imbalances. It is important to know if other drugs like opioids, benzodiazepines, or street drugs have been ingested because this increases the risk of overdose, and other treatments (such as naloxone) may be required. For clients with moderate to severe intoxication, routine lab work includes serum glucose and electrolytes to assess for hypoglycemia, hypokalemia, hypomagnesemia, hypocalcemia, hypophosphatemia, and hyperlactatemia. If hypoglycemia is present, a dextrose intravenous infusion is provided.<sup>16</sup> Severely intoxicated clients may receive intravenous thiamine, along with dextrose, to prevent Wernicke's encephalopathy. Wernicke's encephalopathy is an acute, life-threatening neurological condition characterized by nystagmus, ataxia, and confusion caused by thiamine (B1) deficiency associated with alcohol use disorder. Thiamine is required for cerebral energy utilization.<sup>17</sup> If untreated, Wernicke's encephalopathy can progress to Korsakoff's syndrome, a chronic, irreversible memory disorder resulting from thiamine deficiency.<sup>18</sup>

Some clients with acute alcohol intoxication can become agitated, violent, and uncooperative. Chemical sedation with administration of benzodiazepines may be required to prevent the client from harming themselves or others. However, benzodiazepines must be used with caution because they worsen the respiratory depression caused by alcohol. Approximately one percent of clients with acute alcohol intoxication

16. Cowan, E., & Su, M. K. (2024). Ethanol intoxication in adults. In *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

17. Vasan, S., & Kumar, A. (2023). *Wernicke encephalopathy*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK470344/>

18. National Institute of Neurological Disorders and Stroke. (n.d.). *Wernicke-Korsakoff syndrome*. [https://www.ninds.nih.gov/health-information/disorders/wernicke-korsakoff-syndrome#:~:text=Korsakoff%20syndrome%20\(also%20called%20Korsakoff's,t he%20brain%20involved%20with%20memory.](https://www.ninds.nih.gov/health-information/disorders/wernicke-korsakoff-syndrome#:~:text=Korsakoff%20syndrome%20(also%20called%20Korsakoff's,t he%20brain%20involved%20with%20memory.)

require critical care. Risk factors for admission to the intensive care unit (ICU) include abnormal vital signs (hypotension, tachycardia, fever, or hypothermia), hypoxia, hypoglycemia, and the need for parenteral sedation. If the client has inadequate respirations or airway maintenance, intubation and mechanical ventilation are required. Activated charcoal and gastric lavage are generally not helpful because of the rapid rate of absorption of alcohol from the gastrointestinal tract. Some acutely intoxicated clients experience head traumas from injuries sustained while intoxicated. If the client's mental status does not improve as their BAC level decreases, a CT scan of the head may be obtained.<sup>19</sup>

## MEDICAL TREATMENT OF ALCOHOL OVERDOSE

In addition to the supportive care measures used for alcohol intoxication, treatment for alcohol overdose may require more aggressive interventions. This includes advanced airway management, mechanical ventilation if necessary, and intensive monitoring in an ICU setting.<sup>20</sup>

## Opioid Use and Intoxication

In 2020, 9.5 million (3.4%) of Americans aged 12 and older reported using opioids in the past year. Among this population, 9.3 million people misused prescription pain relievers, and 902,000 people used heroin.<sup>21</sup> Opioids are substances that act on opioid receptors in the central nervous system. Medically, they are used for relief of moderate to severe pain and anesthesia.

19. Cowan, E., & Su, M. K. (2024). Ethanol intoxication in adults. In *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

20. Cowan, E., & Su, M. K. (2024). Ethanol intoxication in adults. In *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

21. National Survey on Drug Use and Health. (n.d.) <https://nsduhweb.rti.org/respweb/homepage.cfm>

When misused, opioids cause a person to feel relaxed and euphoric (i.e., experience an intense feeling of happiness). Opioid prescription medications include Schedule II medications such as morphine, oxycodone, hydrocodone, fentanyl, and hydromorphone. Heroin, an illegal street drug, is also an opioid, but it is classified as a Schedule I drug.<sup>22</sup> Injected opioid misuse is a risk factor for contracting HIV, hepatitis B, hepatitis C, and bacterial endocarditis.<sup>23</sup>

**Opioid intoxication** a condition where a person is affected by opioid drugs to the point of being impaired and potentially ill. It causes problematic behavioral or psychological changes such as initial euphoria followed by apathy, dysphoria, psychomotor retardation or agitation, and impaired judgment. These are some signs of opioid intoxication<sup>24</sup>:

- Pupillary constriction (or dilation from severe overdose) and one of the following:
  - Drowsiness or coma
  - Slurred speech
  - Impairment in attention or memory

## OPIOID OVERDOSE

From 1999 to 2019, nearly 500,000 people died from an overdose involving prescription or illicit opioids. This rise in opioid overdose deaths can be

22. A.D.A.M. Medical Encyclopedia [Internet]. (2022). *Opioid intoxication*.  
<https://medlineplus.gov/ency/article/000948.htm>

23. Substance Abuse and Mental Health Services Administration. (2024). *Learn about substances*. U.S. Department of Health & Human Services.  
<https://www.samhsa.gov/find-help/atod>

24. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

outlined in three distinct waves. See Figure 14.4<sup>25</sup> for an illustration of these three waves of opioid overdose. The first wave of overdose deaths began with the increased prescription rate of opioids in the 1990s. The second wave began in 2010 with rapid increases in overdose deaths involving heroin. The third wave began in 2013 with significant increases in overdose deaths involving synthetic opioids, particularly illicitly manufactured fentanyl.<sup>26</sup>

There are two types of fentanyl: pharmaceutical fentanyl prescribed for severe pain, and illicitly manufactured fentanyl. Most recent cases of fentanyl-related overdose are associated with illicitly manufactured fentanyl that is added to other street drugs that make them more powerful, more addictive, and more dangerous.<sup>27</sup>

25. “3-waves-2019-medium.PNG” by [The Centers for Disease Control and Prevention](#) is in the [Public Domain](#). Access for free at <https://www.cdc.gov/drugoverdose/epidemic/index.html>
26. Centers for Disease Control and Prevention. (2021). *Understanding the epidemic*. U.S. Department of Health & Human Services. <https://www.cdc.gov/drugoverdose/epidemic/index.html>
27. Centers for Disease Control and Prevention. (n.d.). *The facts about fentanyl* [PDF Handout]. [https://www.cdc.gov/stopoverdose/fentanyl/pdf/fentanyl\\_fact\\_sheet\\_508c.pdf](https://www.cdc.gov/stopoverdose/fentanyl/pdf/fentanyl_fact_sheet_508c.pdf)



## Three Waves of the Rise in Opioid Overdose Deaths




Figure 14.4 Three Waves of Opioid Overdose

Carfentanil is a new factor in opioid overdose rates. Carfentanil is a synthetic opioid used to tranquilize large mammals like elephants. It is approximately 10,000 times more potent than morphine and 100 times more potent than fentanyl. A miniscule amount of powder containing two milligrams of carfentanil can be lethal. Carfentanil can be accidentally absorbed through the skin or inhaled by unsuspecting parties. If carfentanil or another fentanyl-related drug is suspected to be present on an individual, first responders and health care professionals must carefully follow safety protocols to avoid accidental lethal exposure.<sup>28</sup> Signs of opioid overdose include the following:

- Unconsciousness or inability to awaken
- Pinpoint pupils
- Slow, shallow breathing; breathing difficulty manifested by choking sounds or a gurgling/snoring noise from a person who cannot be

28. Drug Enforcement Administration. (n.d.). *Officer safety alert*. U.S. Department of Justice. <https://www.justice.gov/usao-edky/file/898991/download>

awakened; or respiratory arrest

- Fingernails or lips turning blue or purple

▶ Read a U.S. Department of Justice Officer Safety Alert PDF:  
[Carfentanil: A Dangerous New Factor in the U.S. Opioid Crisis.](#)

## TREATING OPIOID OVERDOSE

Naloxone reverses the effects of an opioid overdose. A single-step nasal spray delivery of naloxone is the easiest and most successful route of administration for members of the community and first responders. Naloxone intramuscular injection is also available.<sup>29</sup>

Five basic steps are recommended for nurses, first responders, health professionals, and other bystanders to rapidly recognize and treat opioid overdose to prevent death.<sup>30</sup>

1. **Evaluate Signs of Overdose & Obtain Emergency Assistance:** If an opioid overdose is suspected, try to stimulate the person by calling their name or vigorously grinding one's knuckles into their sternum. If the person does

29. Eggleston, W., Podolak, C., Sullivan, R. W., Pacelli, L., Keenan, M., & Wojcik, S. (2018). A randomized usability assessment of simulated naloxone administration by community members. *Addiction*, 113(12), 2300-2304.  
<https://doi.org/10.1111/add.14416>

30. Substance Abuse and Mental Health Services Administration. (2018). *SAMHSA opioid overdose prevention toolkit: Five essential steps for first responders* [PDF Manual]. U.S. Department of Health & Human Services.  
<https://store.samhsa.gov/sites/default/files/d7/priv/five-essential-steps-for-first-responders.pdf>

not respond, call 911 or obtain emergency assistance.

## 2. **Provide Rescue Breathing, Chest Compressions, and Oxygen as Needed**

- Provide rescue breathing if the person is not breathing on their own. A barrier device is recommended to reduce the risk of disease transmission. Rescue breathing for adults involves the following steps:
  - Be sure the person's airway is clear.
  - Place one hand on the person's chin, tilt the head back, and pinch the nose closed.
  - Place your mouth over the person's mouth to make a seal and give two slow breaths.
  - Watch for the person's chest (but not the stomach) to rise.
  - Follow up with one breath every five seconds.
- If the individual becomes pulseless, provide cardiopulmonary resuscitation (CPR).
- Administer oxygen as needed.

## 3. **Administer the First Dose of Naloxone**

- Naloxone should be administered to anyone suspected of an opioid overdose when there is evidence of significant respiratory depression or CNS depression.
- Research has shown that women, older adults, and those without obvious signs of opioid use disorder are undertreated with naloxone and, as a result, have a higher death rate. Therefore, naloxone should be considered for women and the elderly who are found unresponsive.
- Naloxone can be used in life-threatening opioid overdose circumstances in pregnant women.
- Naloxone can be given intranasally, intramuscularly, subcutaneously, or intravenously. The nasal spray is a prefilled device that requires no assembly and delivers a single dose into one nostril. An auto-injector is injected into the outer thigh to deliver naloxone intramuscularly or subcutaneously.

- All naloxone products are effective in reversing opioid overdose, including fentanyl-involved opioid overdoses, although overdoses involving potent or large quantities of opioids may require additional doses of naloxone.
- Withdrawal triggered by naloxone can feel unpleasant; some people may awaken confused, agitated, or aggressive. Provide safety, reassurance, and explain what is happening.

#### **4. Administer a Second Dose of Naloxone If the Person Does Not Respond:**

- If the person overdosing does not respond within 2 to 3 minutes after administering a dose of naloxone, administer a second dose of naloxone.
- People who have taken long-acting or potent opioids (like fentanyl) may require additional intravenous bolus doses or an infusion of naloxone.
- The duration of effect of naloxone depends on dose, route of administration, and overdose symptoms. It is shorter than the effects of some opioids, so a second dose may be required.

#### **5. Monitor the Person's Response:**

- Most people respond to naloxone by returning to spontaneous breathing within 2 to 3 minutes. Continue resuscitation while waiting for the naloxone to take effect.
- The goal of naloxone therapy is to restore adequate spontaneous breathing but not necessarily achieve complete arousal.
- The individual should be monitored for recurrence of signs and symptoms of opioid toxicity for at least four hours from the last dose of naloxone. People who have overdosed on long-acting opioids like fentanyl require prolonged monitoring.
- Because naloxone has a relatively short duration of effect, overdose symptoms may return. Therefore, it is essential to get the person to an emergency department or other source of medical care as quickly as possible, even if the person revives after the initial dose of naloxone

and seems to feel better.

## PREVENTING ILLICIT DRUG OVERDOSE

Nearly 85% of overdose deaths involve illicitly manufactured fentanyl, heroin, cocaine, or methamphetamine. Potential opportunities to link people to care or to implement life-saving actions have been identified for more than 3 in 5 people who died from drug overdose. Circumstances that represent a potential touchpoint for linkage to care are as follows<sup>31</sup>:

- **Bystander present:** Nearly 40% of opioid and stimulant overdose deaths occurred while a bystander was present.
- **Recent release from an institution:** Among the people who died from overdoses involving opioids, about 10% had recently been released from an institution (such as jail/prison, inpatient rehabilitation facilities, or psychiatric hospitals). Risk increased for this population because they were unaware of decreased tolerance to the drug due to abstinence from it while in the institution.
- **Previous overdose:** Among the people who died from overdoses involving opioids, about 10% had a previous overdose.
- **Mental health diagnosis:** Among all the people who died of a drug overdose, 25% had a documented mental health diagnosis.
- **Substance use disorder treatment:** Among the people who died from opioid overdose, nearly 20% had previously been treated for substance use disorder.

Opioid overdoses can be prevented by helping individuals struggling with opioid use disorder find the right treatment and recovery services, as well as

31. Centers for Disease Control and Prevention. (2020). *Overdose deaths and the involvement of illicit drugs*. U.S. Department of Health & Human Services. <https://www.cdc.gov/drugoverdose/featured-topics/VS-overdose-deaths-illicit-drugs.html>

providing public education about administering naloxone. See Figure 14.5.<sup>32</sup> for an image related to these strategies.<sup>33</sup> Read more about treatment and recovery in the “[Treatment of Substance Use Disorders](#)” section.



Figure 14.5 Preventing Overdose Deaths

32. “prevent-overdose-deaths-71k-large.jpg” by [Centers for Disease Control and Prevention, National Center for Injury Prevention and Control](#) is in the [Public Domain](#). Access for free at <https://www.cdc.gov/drugoverdose/resources/graphics/overdose.html>
33. Centers for Disease Control and Prevention. (2020). *Overdose deaths and the involvement of illicit drugs*. U.S. Department of Health & Human Services. <https://www.cdc.gov/drugoverdose/featured-topics/VS-overdose-deaths-illicit-drugs.html>

# Cannabis (Marijuana) Use and Intoxication

Approximately 48.2 million (17.5 percent) of Americans aged 12 or older used cannabis (marijuana) in the past year, and 14.2 million people (5.1%) have a cannabis use disorder.<sup>34</sup> Changes in marijuana policies across states have legalized marijuana for recreational and/or medicinal uses (e.g., pain control, increased appetite for individuals undergoing chemotherapy, intractable seizures, etc.). Although many states now permit dispensing marijuana for medicinal purposes, the U.S. Food and Drug Administration has not approved “medical marijuana.” Therefore, it is important for nurses to educate people about both the adverse health effects and the potential therapeutic benefits linked to marijuana.<sup>35</sup>

The main psychoactive chemical in marijuana is Delta-9-tetrahydrocannabinol (THC). THC alters the functioning of the hippocampus and other areas of the brain that enable a person to form new memories and shift their attentional focus. As a result, marijuana causes impaired thinking and interferes with a person’s ability to learn and perform complicated tasks. THC also disrupts functioning of the cerebellum and basal ganglia that regulate balance, posture, coordination, and reaction time. For this reason, people who have used marijuana may not drive safely and may have problems playing sports or engaging in other physical activities.<sup>36</sup>

When marijuana is smoked, THC and other chemicals pass from the lungs into the bloodstream and are rapidly carried to the brain. The person begins

34. National Institute on Drug Abuse. (2020). *Marijuana research report*.  
<https://nida.nih.gov/publications/research-reports/marijuana/letter-director>

35. National Institute on Drug Abuse. (2020). *Marijuana research report*.  
<https://nida.nih.gov/publications/research-reports/marijuana/letter-director>

36. National Institute on Drug Abuse. (2020). *Marijuana research report*.  
<https://nida.nih.gov/publications/research-reports/marijuana/letter-director>



to experience effects such as euphoria and sense of relaxation almost immediately. Other common effects include heightened sensory perception (e.g., brighter colors), laughter, altered perception of time, and increased appetite. Other people experience anxiety, fear, distrust, or panic, especially if they take too much, the marijuana has high potency, or the person is inexperienced in using cannabis. People who have taken large or highly potent doses of marijuana may experience acute psychosis, including hallucinations, delusions, and a loss of the sense of personal identity.<sup>37</sup>

If marijuana is consumed in foods or beverages, the effects are delayed for 30 minutes to 1 hour, because the drug must first pass through the digestive system. Eating or drinking marijuana delivers significantly less THC into the bloodstream. Because of the delayed effects, people may inadvertently consume more THC than they intend to.<sup>38</sup> Although detectable amounts of THC may remain in the body for days or even weeks after use, the noticeable effects of smoked marijuana generally last from 1 to 3 hours, and marijuana consumed in food or drink may last for many hours.<sup>39</sup>

Delta-8 THC products are manufactured from hemp-derived cannabidiol (CBD) and have psychoactive and intoxicating effects similar to Delta-9 THC. Some Delta-8 THC products are labeled as “hemp products,” which can mislead consumers who associate “hemp” with being non-psychoactive. Delta-8 THC is available for purchase online and in stores but has not been approved by the U.S. Food and Drug Administration for safe use in any context. It should be kept out of reach of children and pets. Some

37. National Institute on Drug Abuse. (2020). *Marijuana research report*.  
<https://nida.nih.gov/publications/research-reports/marijuana/letter-director>

38. National Institute on Drug Abuse. (2020). *Marijuana research report*.  
<https://nida.nih.gov/publications/research-reports/marijuana/letter-director>

39. National Institute on Drug Abuse. (2020). *Marijuana research report*.  
<https://nida.nih.gov/publications/research-reports/marijuana/letter-director>



manufacturers use unsafe chemicals to make Delta-8 THC through a chemical synthesis process that can contaminate the end product. As a result, there has been a recent increase in reports of adverse events with 8% of cases requiring admission to a critical care unit.<sup>40</sup>

THC affects brain systems that are still maturing through young adulthood, so regular use by teens may have negative and long-lasting effects on their cognitive development. Marijuana smoking is associated with large airway inflammation, increased airway resistance, lung hyperinflation, and chronic bronchitis. Vaping products that contain THC are associated with serious lung disease and death. Also, contrary to popular belief, marijuana can be addictive. THC stimulates neurons in the brain's reward system to release higher levels of dopamine and encourages the brain to repeat the rewarding behavior.<sup>41</sup>

**Cannabis intoxication** is defined as problematic behavior or psychological changes (e.g., impaired motor coordination, euphoria, anxiety, sensation of slowed time, impaired judgment, social withdrawal) that developed during or shortly after cannabis use. Signs and symptoms of cannabis intoxication include a range of acute psychological and physiological effects, which include some of the following<sup>42</sup>:

- Conjunctival redness

40. U.S. Food & Drug Administration. (2022). *5 things to know about Delta-8 tetrahydrocannabinol – Delta-8 THC*. <https://www.fda.gov/consumers/consumer-updates/5-things-know-about-delta-8-tetrahydrocannabinol-delta-8-thc>

41. National Institute on Drug Abuse. (2020). *Marijuana research report*. <https://nida.nih.gov/publications/research-reports/marijuana/letter-director>

42. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

- Increased appetite
- Dry mouth
- Tachycardia

## CANNABIS OVERDOSE

According to the CDC, a fatal overdose caused solely by marijuana is unlikely. However, effects from marijuana can lead to unintentional injury, such as a motor vehicle crash, fall, or poisoning. Overconsumption of marijuana can occur when using marijuana-infused products like edibles and beverages because it can take up to two hours to feel the effects from the drug.<sup>43</sup> Additionally, marijuana purchased as a street drug may be laced with other substances like synthetic fentanyl that can cause overdose.

The potential medicinal properties of marijuana and THC have been the subject of research and heated debate for decades. THC itself has proven medical benefits in particular formulations. For example, the U.S. Food and Drug Administration (FDA) has approved the use of THC-based oral medications dronabinol (Marinol) and nabilone (Cesamet) for the treatment of nausea in clients undergoing cancer chemotherapy and to stimulate appetite in clients with wasting syndrome due to AIDS.<sup>44</sup> Marijuana is also used by individuals with certain illnesses such as multiple sclerosis for the management of spasticity, tics, convulsions, and dyskinesia.<sup>45</sup> It can also be

43. Centers for Disease Control and Prevention. (2025). *Cannabis frequently asked questions*. U.S. Department of Health & Human Services.  
[https://www.cdc.gov/cannabis/faq/?CDC\\_AAref\\_Val=https://www.cdc.gov/marijuana/faqs.htm](https://www.cdc.gov/cannabis/faq/?CDC_AAref_Val=https://www.cdc.gov/marijuana/faqs.htm)

44. National Institute on Drug Abuse. (2020). *Marijuana research report*.  
<https://nida.nih.gov/publications/research-reports/marijuana/letter-director>

45. Shepard, S. (2021). *Can medical marijuana help your MS?* WebMD.  
<https://www.webmd.com/multiple-sclerosis/multiple-sclerosis-medical->

used for symptom management in other diseases such as epilepsy, Alzheimer's Disease, Crohn's Disease, PTSD, and glaucoma.

## MEDICAL TREATMENT FOR CANNABIS INTOXICATION

Treatment of cannabis intoxication is generally supportive. Most cases are mild and self-limited, requiring no formal medical intervention. Clients should be placed in a quiet environment and offered reassurance. Severe agitation or anxiety can be managed with benzodiazepines. If psychotic symptoms are present, second-generation antipsychotic agents may be used, with dosages adjusted based on the severity of the symptoms. In cases involving severe mood or psychotic symptoms, inpatient treatment may be warranted. For children who ingest cannabis, more severe symptoms such as coma, convulsions, or cardiopulmonary instability may develop, necessitating more intensive medical care.<sup>46</sup>

## Sedative, Hypnotic, and Anxiolytic Use and Intoxication

Examples of medications in the sedative, hypnotic, or anxiolytic class include benzodiazepines, such as alprazolam (Xanax), diazepam (Valium), lorazepam (Ativan), chlordiazepoxide (Librium), and clonazepam (Klonopin). An example of a hypnotic is zolpidem (Ambien). Although these are prescription medications, they are commonly misused.

Chronic use of benzodiazepines causes changes in gamma-aminobutyric acid (GABA) receptors, resulting in decreased GABA activity and the

[marijuana#:~:text=Surveys%20show%20that%20many%20people%20with%20MS%20already,that%20attacks%20your%20brain%2C%20spinal%20cord%2C%20and%20nerves.](#)

46. Gorelick, D.A. (2023). Cannabis-related disorders and toxic effects. *New England Journal of Medicine*, 389(24), 2267-2275. doi: 10.1056/NEJMr2212152

development of tolerance. When benzodiazepines are no longer present, or suddenly present at lower doses, withdrawal occurs.

Sedatives, hypnotics, and anxiolytic intoxication cause behavioral or psychological changes similar to alcohol intoxication, such as inappropriate sexual or aggressive behavior, mood lability, and impaired judgment. Symptoms of intoxication are as follows<sup>47</sup>:

- Slurred speech
- Incoordination
- Unsteady gait
- Nystagmus
- Impaired cognition (attention and memory)
- Stupor or coma

Benzodiazepines are not detected by standard urine tests for drugs of abuse. However, specific benzodiazepine urine tests identify the metabolites of some benzodiazepines.<sup>48</sup>

## **SEDATIVE, HYPNOTIC, AND ANXIOLYTIC OVERDOSE**

Benzodiazepines cause CNS depression and are commonly involved in drug overdose. They are often co-ingested with other drugs, such as alcohol or opioids that cause stupor, coma, and respiratory depression. When treating benzodiazepine overdose, end tidal CO<sub>2</sub> (i.e., capnography) is used to monitor clients at risk for hypoventilation. Endotracheal intubation and mechanical

47. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

48. Greller, H., & Gupta, A. (2024). Benzodiazepine poisoning. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

ventilation may be required.<sup>49,50</sup> Flumazenil is an antidote to reverse benzodiazepine-induced sedation following general anesthesia and procedural sedation. However, it is used cautiously for benzodiazepine overdose because it is associated with withdrawal seizures in individuals who have developed a tolerance to benzodiazepines.

Medical Treatment for sedative, hypnotic, and anxiolytic overdose include supportive care and pharmacotherapy. Supportive care involves securing the airway, providing respiratory support, and ensuring adequate circulation. This may involve bag-mask ventilation and, if necessary, endotracheal intubation and mechanical ventilation. Medication intervention includes the use of Flumazenil is a benzodiazepine antagonist that can be used to reverse the effects of benzodiazepines. The American Heart Association (AHA) guidelines recommend flumazenil in specific scenarios, such as pediatric exploratory ingestions and iatrogenic overdoses during procedural sedation, where high-risk conditions like chronic benzodiazepine dependence and coingestion of other dangerous substances can be reliably excluded. The initial dose for adults is 0.2 mg IV over 15 seconds, which can be repeated every 60 seconds up to a total of 1 mg.<sup>51</sup>

49. Kang, M., Galuska, M. A., & Ghassemzadeh, S. (2023). *Benzodiazepine toxicity*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK482238/>
50. Greller, H., & Gupta, A. (2024). Benzodiazepine poisoning. *UpToDate*. [www.uptodate.com](https://www.uptodate.com)
51. Lavonas, E. J., Akpunonu, P. D., Arens, A. M., Babu, K. M., Cao, D., Hoffman, R. S., Hoyte, C. O., Mazer-Amirshahi, M. E., Stolbach, A., St-Onge, M., Thompson, T. M., Wang, G. S., Hoover, A. V., & Drennan, I. R. (2023). 2023 American Heart Association focused update on the management of patients with cardiac arrest or life-threatening toxicity due to poisoning: An update to the American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation*, 148(16), e149-e184. [doi:10.1161/CIR.0000000000001161](https://doi.org/10.1161/CIR.0000000000001161).

# Hallucinogen Use and Intoxication

In 2020, 7.1 million (2.6 percent) people in America aged 12 or older used hallucinogens.<sup>52</sup> Hallucinogens are a diverse group of drugs that alter a person's awareness of their surroundings, as well as their own thoughts and feelings. They are commonly split into two categories: classic hallucinogens (such as LSD and peyote) and dissociative drugs (such as phencyclidine and ketamine). Dextromethorphan is an over-the-counter cough suppressant misused for its hallucinogenic and dissociative properties at high doses. It is important to note that ketamine is being utilized in monitored environments for pediatric pain control in emergency departments, acute and chronic pain management, depression and other mental health disorders.<sup>53</sup>

Hallucinogens cause hallucinations (sensations and images that seem real though they are not), and dissociative drugs can cause users to feel out of control or disconnected from their bodies and environments. Historically, some cultures have used hallucinogens like peyote as a part of religious or healing rituals. Users of hallucinogens and dissociative drugs have increased risk for serious harm because of altered perceptions and moods. As a result, users might do things they would never do when not under the influence of a hallucinogen, like jump off a roof or act on suicidal thoughts..<sup>54</sup>

Phencyclidine (PCP) is an example of a hallucinogen. It is an illegal street drug that usually comes as a white powder that can be inhaled through the nose,

52. National Survey on Drug Use and Health. (n.d.). <https://nsduhweb.rti.org/respweb/homepage.cfm>

53. Gao, M., Rejaei, D., & Liu, H. (2016). Ketamine use in current clinical practice. *Acta Pharmacologica Sinica*, 37(7), 865-72. [doi: 10.1038/aps.2016.5](https://doi.org/10.1038/aps.2016.5).

54. National Institute on Drug Abuse, & National Institutes of Health. (2023). *Psychedelics and dissociative drugs*. U.S. Department of Health & Human Services. <https://nida.nih.gov/research-topics/psychedelic-dissociative-drugs>

injected into a vein, smoked, or swallowed.<sup>55</sup> PCP intoxication causes problematic behavioral changes (e.g., belligerence, assaultiveness, impulsiveness, unpredictability, psychomotor agitation, impaired judgment) that occur during or shortly thereafter use.<sup>56</sup> Because of these symptoms, PCP is associated with violent and aggressive behavior including self-injury and violent criminal offenses (such as assaults, intimate partner violence, and homicide).<sup>57</sup> Within one hour of ingestion, two or more of the following signs or symptoms occur<sup>58</sup>:

- Vertical or horizontal **nystagmus** (an involuntary eye movement that causes the eye to rapidly move up and down or from side to side)
- Hypertension
- Tachycardia
- Numbness or diminished responsiveness to pain
- Ataxia (impaired balance or coordination)
- Dysarthria
- Muscle rigidity
- Seizures or coma

55. A.D.A.M. Medical Encyclopedia [Internet]. Atlanta (GA): A.D.A.M., Inc.; c1997-2022. Substance use – phencyclidine (PCP); [updated 2022, February 18]. [https://medlineplus.gov/ency/patientinstructions/000797.htm#:~:text=Phencyclidine%20\(PCP\)%20is%20an%20illegal,into%20a%20vein%20\(shooting%20up\)](https://medlineplus.gov/ency/patientinstructions/000797.htm#:~:text=Phencyclidine%20(PCP)%20is%20an%20illegal,into%20a%20vein%20(shooting%20up))

56. American Psychiatric Association. (2013). *Desk reference to the diagnostic criteria from DSM-5*.

57. Crane, C. A., Easton, C. J., & Devine, S. (2013). The association between phencyclidine use and partner violence: An initial examination. *Journal of Addictive Diseases*, 32(2), 150–157. <https://doi.org/10.1080/10550887.2013.797279>.

58. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

- Hyperacusis (sensitivity to noise)

## MEDICAL TREATMENT FOR HALLUCINOGEN INTOXICATION

Medical treatment for hallucinogen intoxication focuses primarily on supportive care and reassurance, as most cases are self-limiting. A calm, low-stimulation environment is essential to reduce sensory overload, anxiety, and disorientation. When significant agitation or anxiety is present, benzodiazepines such as lorazepam are typically the first-line treatment. In situations where a client exhibits severe psychomotor agitation or poses a risk to themselves or others, physical restraints may be necessary to ensure safety, followed by chemical sedation with intravenous benzodiazepines. If psychotic symptoms such as paranoia, delusions, or hallucinations are persistent and distressing, short-term use of antipsychotics may be indicated. In rare but severe cases, such as those involving hyperthermia, seizures, or cardiovascular instability, intensive care is required for close monitoring and life-saving interventions.<sup>59</sup> Review ANA guidelines on using restraints in the “[Client Rights](#)” section of the “Legal and Ethical Considerations in Mental Health Care” chapter and information on safely implementing restraints in the “[Workplace Violence](#)” section of the “Trauma, Abuse, and Violence” chapter.

## HALLUCINOGEN OVERDOSE

Overdose can occur with some dissociative drugs like PCP. High doses of PCP can cause seizures, coma, and death, especially if taken with depressants such as alcohol or benzodiazepines.<sup>60</sup>

59. Heard, K., & Hoppe, J. (2024). Phencyclidine(PCP) intoxication in adults. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

60. National Institute on Drug Abuse, & National Institutes of Health. (2021). *Hallucinogens drug facts*. U.S. Department of Health & Human Services. <https://nida.nih.gov/publications/drugfacts/hallucinogens>



Dextromethorphan is an over-the-counter medication that is misused due to its hallucinogenic effects and can also cause overdose. Nonmedical use of dextromethorphan results in approximately 6,000 emergency department visits annually in the United States, often with co-ingestion of alcohol. Signs of toxic doses include neurobehavioral changes (e.g., hallucinations, inappropriate laughing, psychosis with dissociative features, agitation, and coma); tachycardia; dilated pupils; diaphoresis; and a “zombie-like” ataxic gait. Because acetaminophen is commonly present in cough and cold medications, toxic doses can cause severe delayed hepatotoxicity, hepatic failure, and death; serum acetaminophen levels should be obtained in all clients presenting with toxic levels of dextromethorphan.<sup>61</sup>

## MEDICAL TREATMENT FOR HALLUCINOGEN OVERDOSE

The medical treatment for a hallucinogen overdose involves a combination of supportive care, symptomatic management, and pharmacologic interventions. For severe cases involving hyperthermia, seizures, or hypertensive crises, more intensive medical interventions may be necessary, including cooling measures and anticonvulsants.<sup>62</sup>

61. Rosenbaum, C., & Boyer, E. (2023). Dextromethorphan misuse and poisoning: Clinical features and diagnosis. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)
62. Lavonas, E. J., Akpunonu, P. D., Arens, A. M., Babu, K. M., Cao, D., Hoffman, R. S., Hoyte, C. O., Mazer-Amirshahi, M. E., Stolbach, A., St-Onge, M., Thompson, T. M., Wang, G. S., Hoover, A. V., & Drennan, I. R. (2023). 2023 American Heart Association focused update on the management of patients with cardiac arrest or life-threatening toxicity due to poisoning: An update to the American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation*, 148(16), e149-e184. [doi:10.1161/CIR.0000000000001161](https://doi.org/10.1161/CIR.0000000000001161).

## Stimulant Use and Intoxication

Stimulants include amphetamine-type substances, cocaine, and crack. Stimulants cause the release of dopamine in the brain and are highly addictive because the flood of dopamine in the brain's reward circuit strongly reinforces drug-taking behaviors. With continued drug use, the reward circuit adapts and becomes less sensitive to the drug. As a result, people take stronger and more frequent doses in an attempt to feel the same high and to obtain relief from withdrawal symptoms. Because the high from stimulants starts and fades quickly, people often take repeated doses in a form of bingeing, often giving up food and sleep while continuing to take the drug every few hours for several days. Both the use and withdrawal from amphetamines can cause psychosis with symptoms of hallucinations and paranoia.<sup>63</sup>

Approximately 2 million Americans used methamphetamine in the past year.<sup>64</sup> Methamphetamine comes in many forms and can be ingested by smoking, swallowing a pill, snorting, or injecting the powder that has been dissolved in water or alcohol. Methamphetamine can be easily made in small clandestine laboratories with relatively inexpensive over-the-counter ingredients such as pseudoephedrine, a common ingredient in cold medications. To curb this illegal production, federal law requires pharmacies take steps to limit sales and obtain photo identification from purchasers. Methamphetamine production also involves a number of other very dangerous chemicals. Toxic effects from these chemicals can remain in the environment long after the lab has been shut down, causing a wide range of

63. National Institute on Drug Abuse, & National Institutes of Health. (2024). *Methamphetamine*. <https://nida.nih.gov/research-topics/methamphetamine.gov/publications/drugfacts/methamphetamine>

64. National Institute on Drug Abuse, & National Institutes of Health. (2024). *Methamphetamine*. <https://nida.nih.gov/research-topics/methamphetamine.gov/publications/drugfacts/methamphetamine>

health problems for people living in the area. These chemicals can also result in deadly lab explosions and house fires.<sup>65</sup>

Long-term use of methamphetamine has many negative consequences, including extreme weight loss, severe dental problems, intense itching leading to skin sores from scratching, involuntary movements (dyskinesia), anxiety, memory loss, and violent behavior.<sup>66</sup>

Cocaine is another powerfully addictive stimulant drug made from the leaves of the coca plant native to South America. It is estimated that 5.5 million people aged 12 or older have used cocaine, including about 778,000 users of crack.<sup>67</sup> Users may snort cocaine powder through the nose, rub it into their gums, or dissolve the powder and inject it into the bloodstream. Cocaine that has been processed to make a rock crystal is called “crack.” The crystal is heated (making crackling sounds) to produce vapors that are inhaled into the lungs. In the short-term, cocaine use can result in increased blood pressure, restlessness, and irritability. In the long-term, severe medical complications of cocaine use include heart attacks and seizures.<sup>68</sup>

65. National Institute on Drug Abuse, & National Institutes of Health. (2024). *Methamphetamine*. <https://nida.nih.gov/research-topics/methamphetamine.gov/publications/drugfacts/methamphetamine>

66. National Institute on Drug Abuse, & National Institutes of Health. (2024). *Methamphetamine*. <https://nida.nih.gov/research-topics/methamphetamine.gov/publications/drugfacts/methamphetamine>

67. National Institute on Drug Abuse, & National Institutes of Health. (2024). *Cocaine*. <https://nida.nih.gov/research-topics/cocainegov/research-topics/methamphetamine>

68. National Institute on Drug Abuse, & National Institutes of Health. (2024). *Cocaine*. <https://nida.nih.gov/research-topics/cocainegov/research-topics/methamphetamine>

Stimulant intoxication causes problematic behavioral or psychological changes such as euphoria or blunted affect; changes in sociability; hypervigilance; interpersonal sensitivity; anxiety, tension, or anger; and impaired judgment. These are some symptoms of stimulant intoxication<sup>69</sup>:

- Tachycardia or bradycardia
- Pupillary dilation
- Elevated or decreased blood pressure
- Perspiration or chills
- Nausea or vomiting
- Weight loss
- Psychomotor agitation or retardation
- Muscular weakness, respiratory depression, chest pain or cardiac arrhythmias
- Confusion, seizures, dyskinesia (involuntary, erratic, writhing movements of the face, arms, legs, or trunk), dystonia (involuntary muscle contractions that result in slow repetitive movements), or coma

## MEDICAL TREATMENT FOR STIMULANT INTOXICATION

It is important for nurses to be aware that individuals with acute stimulant intoxication may, without provocation, develop severe agitation with extreme violence and place themselves, family members, medical staff, and other clients at risk of major injury. Control of agitation and hyperthermia (body temperature over 41 degrees Celsius) receive top priority for treatment with the following interventions<sup>70</sup>:

- Intravenous benzodiazepines are administered immediately for chemical

69. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

70. Boyer, E. W., & Hernon, C. (2024). Methamphetamine: Acute intoxication. *UpToDate*. [www.uptodate.com](https://www.uptodate.com)

sedation of severely agitated individuals.

- Physical restraints should be avoided because clients who physically struggle against restraints undergo isometric muscle contractions that are associated with lactic acidosis, hyperthermia, sudden cardiac collapse, and death.
- Airway management with endotracheal intubation and mechanical ventilation may be required.
- Aggressive cooling is achieved through sedation, fluid resuscitation, external cooling blankets, or evaporative cooling techniques. Antipyretics are not used because the increased body temperature is caused by muscular activity, not an alteration in the hypothalamic temperature set point.

## MEDICAL TREATMENT FOR STIMULANT OVERDOSE

Medical treatment for stimulant overdose requires rapid, comprehensive interventions due to the risk of life-threatening complications such as seizures, arrhythmias, and hyperthermia. Immediate goals include sedation, cardiovascular stabilization, airway protection, and cooling. Intravenous benzodiazepines remain the first-line treatment to manage agitation and prevent seizures. In cases of severe overdose, intensive care support may be necessary, including endotracheal intubation, mechanical ventilation, and advanced cardiac life support. Continuous cardiac monitoring is critical due to the high risk of arrhythmias and myocardial infarction. Hyperthermia is treated aggressively with external cooling methods, intravenous fluids, and sedation; antipyretics are ineffective and not indicated. Rhabdomyolysis, acute kidney injury, and multi-organ failure may develop, requiring supportive care such as IV hydration or renal replacement therapy. Early recognition and escalation to intensive care are essential to improve outcomes.<sup>71</sup>

71. Boyer, E. W., & Hernon, C. (2024). Methamphetamine: Acute intoxication. *UpToDate*. [www.uptodate.com](https://www.uptodate.com)

- ▶ Read [DrugFacts](#) by the National Institute on Drug Abuse for more information about substances, intoxication, and overdose.

## Inhalant Use and Intoxication

In 2020, 2.4 million (0.9 percent) of people aged 12 or older in America used inhalants. Unlike other illicit drugs, the percentage of inhalant use was highest among adolescents aged 12 to 17.<sup>72</sup> Inhalants are various products easily bought or found in the home, such as spray paints, markers, glue, gasoline, and cleaning fluids. People who use inhalants breathe in the fumes through their nose or mouth, usually by sniffing, snorting, bagging, or huffing. Although the high that inhalants produce usually lasts just a few minutes, people often try to make it last by continuing to inhale again and again over several hours.<sup>73</sup>

Prescription medications can also be misused as inhalants. For example, amyl nitrate is a prescription medication administered via inhalation to relieve chest pain. However, it is misused by individuals to cause a high. It is referred to by the street drug name as “poppers.”

Inhalant intoxication causes problematic behavioral or psychological changes

72. National Institute on Drug Abuse, & National Institutes of Health. (2024). *Inhalants*. <https://nida.nih.gov/research-topics/inhalants>

73. National Institute on Drug Abuse, & National Institutes of Health. (2024). *Inhalants*. <https://nida.nih.gov/research-topics/inhalants>

such as belligerence, assaultiveness, apathy, and impaired judgment. Inhalant intoxication includes these symptoms<sup>74</sup>:

- Dizziness
- Nystagmus
- Incoordination
- Slurred speech
- Unsteady gait
- Lethargy
- Depressed reflexes
- Psychomotor retardation
- Tremor
- Generalized muscle weakness
- Blurred or double vision
- Stupor or coma
- Euphoria

Long-term effects of inhalant use may include liver and kidney damage, hearing loss, bone marrow damage, loss of coordination and limb spasms (from nerve damage), delayed behavioral development (from brain problems), and brain damage (from cut-off oxygen flow to the brain).<sup>75</sup>

Acute intoxication with inhalants can cause life-threatening seizures and coma. Many solvents and aerosol sprays are highly concentrated with many other active ingredients; sniffing these products can cause the heart to stop

74. American Psychiatric Association. (2022). *Desk reference to the diagnostic criteria from DSM-5-TR* (5th ed.). American Psychiatric Association Publishing.

75. National Institute on Drug Abuse, & National Institutes of Health. (2024). *Inhalants*. <https://nida.nih.gov/research-topics/inhalants>

within minutes. This condition, known as sudden sniffing death, can happen to otherwise healthy young people the first time they use an inhalant.<sup>76</sup>

## MEDICAL TREATMENT FOR INHALANT INTOXICATION

Treatment consists of maintaining cardiorespiratory function with 100 percent oxygen administration by rebreather mask and implementation of Pediatric Advanced Life Support (PALS) protocols that may require endotracheal intubation and mechanical ventilation.<sup>77</sup>

## MEDICAL TREATMENT FOR INHALANT OVERDOSE

Treatment of inhalant overdose focuses on immediate stabilization and supportive care. Clients should be removed from the exposure source and placed in a well-ventilated area or given supplemental oxygen to address hypoxia. Continuous monitoring of vital signs, cardiac rhythm, and respiratory status is essential due to the risk of sudden sniffing death syndrome, which results from cardiac arrhythmias. In cases of respiratory depression or loss of consciousness, airway management and mechanical ventilation may be necessary. Seizures, if present, are treated with intravenous benzodiazepines. Because some inhalants can cause chemical pneumonitis, metabolic acidosis, or hepatic and renal injury, further care may include fluid resuscitation, correction of acid-base imbalances, and monitoring for organ dysfunction. Admission to intensive care may be required for clients with severe symptoms or multi-organ involvement.

76. National Institute on Drug Abuse, & National Institutes of Health. (2024). *Inhalants*. <https://nida.nih.gov/research-topics/inhalants>

77. Boyer, E. W., & Hernon, C. (2024). Methamphetamine: Acute intoxication. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)



## 14.3 Withdrawal Management/Detoxification

Nurses working in hospital settings or emergency departments commonly provide care for clients receiving withdrawal treatment for alcohol, opioids, or other substances. Furthermore, clients frequently underreport alcohol and substance use, so nurses must be aware of signs of withdrawal in clients receiving medical care for other issues and notify the health care provider.<sup>1</sup>

**Withdrawal management**, also called detoxification, includes interventions aimed at managing the physical and emotional symptoms that occur after a person suddenly stops using a psychoactive substance. Withdrawal symptoms vary in intensity and duration based on the substance(s) used, the duration and amount of use, and the overall health of the individual. Some substances, such as opioids, sedatives, and tranquilizers, produce significant physical withdrawal effects, especially if they have been used in combination with heavy alcohol use. Rapid or unmanaged cessation from these substances can result in a longer than expected course of withdrawal with seizures and other health complications.

Assessment and treatment of withdrawal symptoms will be further discussed in the following sections.

### Alcohol Withdrawal

The prevalence of alcohol use disorder is estimated to be as high as 32 percent among hospitalized clients. Approximately half of clients with alcohol use disorder experience alcohol withdrawal when they reduce or stop drinking, with as many as 20 percent experiencing serious manifestations such as hallucinations, seizures, and delirium tremens.<sup>2</sup> Severe alcohol

1. Sellers, E. M. (n.d.). *CIWA-AR for alcohol withdrawal*. MDCalc. <https://www.mdcalc.com/ciwa-ar-alcohol-withdrawal#why-use>

2. Pace, C. (2025). Alcohol withdrawal: Epidemiology, clinical manifestations, course, assessment, and diagnosis. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

withdrawal is considered a medical emergency that is best managed in an intensive care unit.

Symptoms of early or mild alcohol withdrawal include anxiety, minor agitation, restlessness, insomnia, tremor, diaphoresis, palpitations, headache, and alcohol craving. Clients often experience loss of appetite, nausea, vomiting, and diarrhea. Their risk for falls often increases when they try to go unassisted to the bathroom with these gastrointestinal symptoms. Physical signs include sinus tachycardia, systolic hypertension, hyperactive reflexes, and tremor. Without treatment, symptoms of mild alcohol withdrawal generally begin within 6 to 24 hours after the last drink and resolve within one to two days.<sup>3</sup>

Some clients develop moderate to severe withdrawal symptoms that can last up to six days, such as withdrawal hallucinations, seizures, or delirium tremens:

- Hallucinations typically occur within 12 to 24 hours after the last drink. They are typically visual and commonly involve seeing insects or animals in the room, although auditory and tactile phenomena may also occur..<sup>4</sup>
- Alcohol withdrawal-related seizures can occur within 6 to 48 hours after the last drink. Risk factors for seizures include concurrent withdrawal from benzodiazepines or other sedative-hypnotic drugs..<sup>5</sup>
- **Delirium tremens (DTs)** is a rapid-onset, fluctuating disturbance of attention and cognition that is sometimes associated with hallucinations.

3. Pace, C. (2025). Alcohol withdrawal: Epidemiology, clinical manifestations, course, assessment, and diagnosis. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

4. Pace, C. (2025). Alcohol withdrawal: Epidemiology, clinical manifestations, course, assessment, and diagnosis. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

5. Pace, C. (2025). Alcohol withdrawal: Epidemiology, clinical manifestations, course, assessment, and diagnosis. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

In its most severe manifestation, DTs are accompanied by agitation and signs of extreme autonomic hyperactivity, including fever, severe tachycardia, hypertension, and drenching sweats. DTs typically begin between 72 and 96 hours after the client's last drink. Mortality rates from withdrawal delirium have been historically as high as 20 percent, but with appropriate medical management, the mortality rate is between 1 and 4 percent. Death is attributed to cardiovascular complications, hyperthermia, aspiration, and severe fluid and electrolyte disorders..<sup>6</sup>

## Clinical Institute Withdrawal Assessment for Alcohol Scale (CIWA-Ar)

The Clinical Institute Withdrawal Assessment for Alcohol Scale (CIWA-Ar) is the most widely used scale to determine the need for medically supervised withdrawal management. It is used in a wide variety of settings, including outpatient, emergency, psychiatric, general medical-surgical, and intensive care units when there is a clinical concern regarding a client's alcohol withdrawal.

The CIWA-Ar scale is typically utilized in association with a protocol containing medications to guide symptom-triggered treatment. Clients with an alcohol use disorder who have a CIWA-Ar score of less than 10 do not typically require medical management.<sup>7</sup> Treatment score may vary depending upon agency protocol.

There are ten questions on the CIWA-Ar related to nausea/vomiting, tremor, paroxysmal sweats, anxiety, agitation, tactile disturbances, auditory disturbances, visual disturbances, headache, and level of orientation. The

6. Pace, C. (2025). Alcohol withdrawal: Epidemiology, clinical manifestations, course, assessment, and diagnosis. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

7. Pace, C. (2025). Alcohol withdrawal: Epidemiology, clinical manifestations, course, assessment, and diagnosis. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

questions are rated from 0 to 7, except for orientation that is rated from 0 to 4. View the full CIWA-Ar scale in the following box.

▶ View the [CIWA-Ar](#) on the MDCalc medical reference website.

View a supplementary YouTube video<sup>8</sup> of a physician  
▶ performing a CIWA assessments on a simulated client: [A  
CIWA-Ar assessment](#)

## Treatment of Alcohol Withdrawal

Benzodiazepines are used to treat the psychomotor agitation associated with alcohol withdrawal and prevent progression from minor symptoms to severe symptoms of seizures, hallucinations, or delirium tremens. Diazepam (Valium), lorazepam (Ativan), and chlordiazepoxide (Librium) are used most frequently to treat or prevent alcohol withdrawal symptoms.<sup>9</sup> Precedex (dexmedetomidine), a selective alpha-2 adrenergic agonist, is increasingly being utilized as an adjunctive treatment for alcohol withdrawal, particularly in severe cases such as alcohol withdrawal delirium or when traditional

8. META:PHI Ontario. (2023, November 22). *A CIWA-Ar assessment* [Video]. YouTube. All rights reserved. [https://www.youtube.com/watch?v=Go\\_CJS7dhVI](https://www.youtube.com/watch?v=Go_CJS7dhVI)

9. Hoffman, R. & Weinhouse, G. (2025). Management of moderate and severe alcohol withdrawal symptoms. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

therapies like benzodiazepines are insufficient. Its sedative and anxiolytic properties help manage agitation and autonomic hyperactivity without causing significant respiratory depression, making it a valuable option in intensive care settings. By reducing sympathetic outflow, Precedex can help stabilize vital signs, improve comfort, and decrease the need for high doses of other sedatives. However, its use should be carefully monitored due to potential side effects such as bradycardia and hypotension.<sup>10</sup>

- ▶ Review information about benzodiazepines in the “[Antianxiety Medications](#)” section of the “Psychotropic Medications” chapter.

Anticonvulsants may be used concurrently or instead of benzodiazepines. Anticonvulsants decrease the probability of withdrawal seizures.

- ▶ Review information about anticonvulsants in the “[Treatments for Bipolar Disorders](#)” section of the “Bipolar Disorders” chapter.

Chronic alcohol use is associated with depletion of thiamine and magnesium. Clients receiving alcohol withdrawal treatment typically receive intravenous thiamine, along with dextrose, to prevent Wernicke’s encephalopathy. Wernicke’s encephalopathy is an acute, life-threatening neurological condition characterized by nystagmus, ataxia, and confusion caused by

10. Yavarovich, E. R., Bintvihok, M., & McCarty, J. C. (2019). Association between dexmedetomidine use for the treatment of alcohol withdrawal syndrome and intensive care unit length of stay. *Journal of Intensive Care*, 7(49)  
<https://doi.org/10.1186/s40560-019-0405-1>

thiamine (B1) deficiency associated with alcohol use disorder.<sup>11</sup> If untreated, Wernicke's encephalopathy can progress to Korsakoff's syndrome, a chronic, irreversible memory disorder resulting from thiamine deficiency.<sup>12</sup> Treatment of other electrolyte deficiencies may be included during alcohol withdrawal treatment and medications may be given via the oral route if tolerated by the client.

## Opioid Withdrawal

Medically supervised opioid withdrawal, also known as detoxification, involves administering medication to reduce the severity of withdrawal symptoms that occur when an opioid-dependent client stops using opioids. However, supervised withdrawal alone does not generally result in sustained abstinence from opioids, nor does it address reasons the client became dependent on opioids.<sup>13</sup>

Clients may undergo detoxification for several reasons<sup>14</sup>:

- Initiating the process to “get clean and stay clean” from opioids. Some clients may follow up with inpatient or outpatient treatment after

11. Vasan, S., & Kumar, A. (2023). *Wernicke encephalopathy*. StatPearls [Internet]. <https://www.ncbi.nlm.nih.gov/books/NBK470344/>

12. National Institute of Neurological Disorders and Stroke. (n.d). *Wernicke-Korsakoff syndrome*. [https://www.ninds.nih.gov/health-information/disorders/wernicke-korsakoff-syndrome#:~:text=Korsakoff%20syndrome%20\(also%20called%20Korsakoff's,t he%20brain%20involved%20with%20memory.](https://www.ninds.nih.gov/health-information/disorders/wernicke-korsakoff-syndrome#:~:text=Korsakoff%20syndrome%20(also%20called%20Korsakoff's,t he%20brain%20involved%20with%20memory.)

13. Sevarino, K. A. (2022). Medically supervised opioid withdrawal during treatment for addiction. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

14. Sevarino, K. A. (2022). Medically supervised opioid withdrawal during treatment for addiction. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

completing the detoxification process.

- Treating withdrawal symptoms when a client dependent on opioids or heroin becomes hospitalized and lacks access to the misused substance.
- Beginning the first step in treating opioid use disorder and transitioning to medication-assisted treatment like methadone or suboxone treatment.
- Establishing an abstinent state without withdrawal symptoms required for the client's setting or status (e.g., incarceration, probation, or a drug-free residential program).

## Clinical Opiate Withdrawal Scale

The Clinical Opiate Withdrawal Scale (COWS) is used in both inpatient and outpatient settings for the monitoring of withdrawal symptoms during opioid detoxification. It can be serially administered to track changes in the severity of withdrawal symptoms over time or in response to treatment.<sup>15</sup>

Symptoms of opioid withdrawal include drug craving, anxiety, restlessness, gastrointestinal distress, diaphoresis, and tachycardia. COWS rates the severity of 11 signs and symptoms of opioid withdrawal on a scale from 0 to 5, as described in the following box.

▶ View the [COWS](#) on the MedCalc medical reference website.

## Treatment of Opioid Withdrawal

A calm, quiet environment with supportive and reassuring staff is instrumental for helping clients overcome most symptoms of acute opioid

15. Wesson, D. R. (n.d.). *COWS score for opiate withdrawal*. MDCalc.  
<https://www.mdcalc.com/cows-score-opiate-withdrawal#why-use>

withdrawal and can decrease the need for pharmacologic interventions. Clients who have associated diarrhea, vomiting, or sweating should be monitored for dehydration and have fluid levels maintained with oral and/or intravenous fluids.<sup>16</sup>

Medications commonly used to treat opioid withdrawal symptoms include opioid agonists such as buprenorphine and naloxone, or methadone or alpha-2 adrenergic agonists such as clonidine. Other medications may be prescribed to treat specific symptoms.<sup>17</sup>

## **BUPRENORPHINE**

Buprenorphine is an effective treatment for opioid withdrawal symptoms. A disadvantage of buprenorphine is it can worsen opioid withdrawal symptoms if not administered carefully. To avoid this situation, the client must be in a state of mild to moderate withdrawal before receiving their first dose of buprenorphine (i.e., have a COWS score greater than 10). The first dose of buprenorphine is typically 2 to 4 mg sublingually.<sup>18</sup>

Buprenorphine can cause respiratory depression. Common side effects include sedation, headache, nausea, constipation, and insomnia.<sup>19</sup>

16. Sevarino, K. A. (2023). Medically supervised opioid withdrawal during treatment for addiction. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

17. Sevarino, K. A. (2023). Medically supervised opioid withdrawal during treatment for addiction. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

18. Sevarino, K. A. (2023). Medically supervised opioid withdrawal during treatment for addiction. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

19. Sevarino, K. A. (2023). Medically supervised opioid withdrawal during treatment for addiction. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)



## BUPRENORPHINE/NALOXONE

The combination medication buprenorphine/naloxone (Suboxone) is commonly used for detoxification, as well as maintenance of abstinence from opioids. It may be used in outpatient settings as an opioid-blocker.<sup>20</sup>

## METHADONE

If buprenorphine is not available, methadone may be prescribed. Methadone is a long-acting, synthetic opioid that reduces opioid craving and withdrawal symptoms by blocking the effect of opioids. It is typically prescribed in one of two ways<sup>21</sup>:

- Substitution Therapy: Methadone is prescribed to replace the use of an opioid and then is gradually tapered to prevent severe withdrawal symptoms.
- Maintenance Therapy: It is prescribed long-term as one component of a comprehensive medication-assisted treatment plan for opioid use disorder. With counseling and other behavioral therapies, methadone helps individuals achieve and sustain recovery and lead active and meaningful lives.

In contrast to buprenorphine, methadone does not induce withdrawal symptoms when administered to a client with opioid in their system because it has an additive effect on opioids that are already present.

20. Sevarino, K. A. (2023). Medically supervised opioid withdrawal during treatment for addiction. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

21. Substance Abuse and Mental Health Services Administration (SAMHSA). (2022). *Methadone*. <https://www.samhsa.gov/medication-assisted-treatment/medications-counseling-related-conditions/methadone>

## ALPHA-2 ADRENERGIC AGONISTS

Alpha-2 adrenergic agonists, such as clonidine, decrease many symptoms of opioid withdrawal and effectively relieve the autonomic symptoms of sweating, diarrhea, intestinal cramps, nausea, anxiety, and irritability. They are least effective for symptoms of myalgias, restlessness, insomnia, and craving.<sup>22</sup>

Clonidine can be taken orally or administered via a clonidine patch and changed weekly. Relief from withdrawal symptoms typically occurs within 30 minutes after a dose. However, common side effects of hypotension and sedation limit the use of these drugs. Contraindications to Alpha-2 adrenergic agonists include hypotension, renal insufficiency, cardiac instability, pregnancy, and psychosis. Tricyclic antidepressants should be stopped three weeks prior to use.<sup>23</sup>

## SYMPTOM-SPECIFIC MEDICATIONS

Various medications are prescribed to provide targeted relief for symptoms of opioid withdrawal<sup>24</sup>:

- Anxiety, irritability, restlessness: Diphenhydramine, hydroxyzine, lorazepam, and clonazepam
- Abdominal cramping: Dicyclomine
- Diarrhea: Bismuth and loperamide
- Nausea/vomiting: Ondansetron, prochlorperazine, and promethazine
- Insomnia: Trazodone, doxepin, mirtazapine, quetiapine, and zolpidem

22. Sevarino, K. A. (2023). Medically supervised opioid withdrawal during treatment for addiction. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

23. Sevarino, K. A. (2023). Medically supervised opioid withdrawal during treatment for addiction. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

24. Sevarino, K. A. (2023). Medically supervised opioid withdrawal during treatment for addiction. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

- Muscle aches, joint pain, and headache: Ibuprofen, acetaminophen, ketorolac, and naproxen
- Muscle spasms and restless legs: Cyclobenzaprine, baclofen, diazepam, and methocarbamol

Warm baths, rehydration, and gentle stretching are also helpful for relieving muscle aches and cramps. Use of benzodiazepines and zolpidem is not recommended for clients receiving methadone or buprenorphine therapy unless they are under close medical supervision due to the risk of oversedation.<sup>25</sup>

## Benzodiazepine Withdrawal

Rapid recognition and treatment of benzodiazepine withdrawal is critical because it can be life threatening, especially in individuals who have been taking high doses or have used the medication for an extended period. The severity of withdrawal symptoms can range from mild to life threatening, depending on the duration of use, dosage, type of benzodiazepine, and presence of comorbid conditions. Initial symptoms may include tremors, anxiety, restlessness, and general malaise, often resembling rebound anxiety. As withdrawal progresses, more severe features can develop, including perceptual disturbances such as visual or auditory hallucinations, intense agitation, and depersonalization. In some cases, clients may experience psychosis with paranoid delusions or disorganized thinking.<sup>26</sup>

Of particular concern is the risk of seizures, which may occur within one to five days of abrupt cessation or rapid tapering, especially in those with high dose or long term use. Seizures can be generalized and are potentially life threatening. Autonomic instability may present as tachycardia, hypertension,

25. Sevarino, K. A. (2023). Medically supervised opioid withdrawal during treatment for addiction. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

26. Kang, A. (2024). Benzodiazepine withdrawal. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

hyperthermia, and diaphoresis, further complicating the clinical picture and increasing the risk of morbidity and mortality if not promptly managed.

## TREATMENT OF BENZODIAZEPINE WITHDRAWAL

Withdrawal from benzodiazepines is managed carefully due to the risk of severe symptoms, including anxiety, insomnia, irritability, tremors, perceptual disturbances, and in some cases, seizures or psychosis. Treatment typically involves substituting the short-acting benzodiazepine the individual has been using with a long-acting one, such as diazepam or clonazepam, which provides more stable blood levels and reduces the intensity of withdrawal symptoms. The medication is then gradually tapered over time, with the goal of minimizing withdrawal effects while avoiding excessive sedation or other complications. The tapering process can last several weeks to several months, depending on the dose, duration of use, and individual response. A slow and individualized taper is especially important for those who have been taking high doses or have used benzodiazepines over an extended period. Clinicians often recommend reducing the dose by no more than 10–25% every 1–2 weeks, adjusting based on the client's symptoms and tolerability.<sup>27</sup>

## Withdrawal Management and Stabilization

Withdrawal management is highly effective in preventing immediate and serious medical consequences associated with discontinuing substance use, but by itself, it is not an effective treatment for any substance use disorder. It is considered stabilization, meaning the client is assisted through a period of acute detoxification and withdrawal to be medically stable and substance-free. Stabilization often prepares the individual for treatment. It is considered a first step toward recovery, similar to the acute management of a diabetic coma as a first step toward managing the underlying illness of diabetes.

27. Kang, A. (2024). Benzodiazepine withdrawal. *UpToDate*. [www.uptodate.com](https://www.uptodate.com)

Similarly, acute stabilization and withdrawal management are most effective when followed by evidence-based treatments and recovery services.<sup>28</sup>

Unfortunately, many individuals who receive withdrawal management do not become engaged in treatment. Studies have found that half to three quarters of individuals with substance use disorders who receive withdrawal management services do not enter treatment. One of the most serious consequences when individuals do not begin continuing care after withdrawal management is overdose. Because withdrawal management reduces acquired tolerance, those who attempt to reuse their former substance in the same amount or frequency may overdose, especially those with opioid use disorders.<sup>29</sup>

The remaining sections of this chapter will discuss substance abuse disorders and treatments, as well as prevention strategies.

28. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
29. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

## 14.4 Substance-Related and Other Addictive Disorders

### Substance-Related Disorders

Prolonged, repeated misuse of substances can produce changes to the brain that can lead to a substance use disorder. According to the *Diagnostic and Statistical Manual of Mental Disorders (DSM-5)*, **substance use disorder (SUD)** is an illness caused by repeated misuse of substances such as alcohol, caffeine, cannabis, hallucinogens, inhalants, opioids, sedatives, hypnotics, anxiolytics, stimulants (amphetamines, cocaine, and other stimulants), and tobacco. All of these substances taken in excess have a common effect of directly activating the brain reward system and producing such an intense activation of the reward system that normal life activities may be neglected. Nonsubstance related disorders such as gambling disorder activate the same reward system in the brain.<sup>1</sup>

Substance use disorders are diagnosed based on cognitive, behavioral, and psychological symptoms. See the *DSM-5* diagnostic criteria used for SUD in the following box. SUD can range from mild to severe and from temporary to chronic.<sup>2</sup>

#### ***DSM-5 Criteria for Substance Use Disorder***

1. American Psychiatric Association. (2013). *Desk reference to the diagnostic criteria from DSM-5*.
2. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

SUD diagnosis requires the presence of two or more of the following criteria in a 12-month period<sup>3</sup>:

- The substance is often taken in larger amounts or over a longer period than intended.
- There is a persistent desire or unsuccessful efforts to cut down or control substance use.
- A great deal of time is spent in activities necessary to obtain the substance, use the substance, or recover from its effects.
- There is a craving, or a strong desire or urge, to use the substance.
- There is recurrent substance use resulting in a failure to fulfill major role obligations at work, school, or home.
- There is continued substance use despite having persistent or recurrent social or interpersonal problems caused by or exacerbated by the effects of the substance.
- Important social, occupational, or recreational activities are given up or reduced because of substance use.
- There is recurrent substance use in situations in which it is physically hazardous.
- Substance use is continued despite knowledge of having a persistent or recurrent physical or psychological problem that is likely to have been caused or exacerbated by the substance.
- Tolerance develops to the substance, as defined by:
  - A need for markedly increased amounts of the

3. American Psychiatric Association. (2013). *Desk reference to the diagnostic criteria from DSM-5*.

substance to achieve intoxication or the desired effect.

- There is a markedly diminished effect with continued use of the same amount of the substance.
- Withdrawal symptoms occur when substance use is cut back or stopped following a period of prolonged use.

The disorder is classified as mild, moderate, or severe. Individuals exhibiting two or three symptoms are considered to have a “mild” disorder, four or five symptoms constitute a “moderate” disorder, and six or more symptoms are considered a “severe” substance use disorder.

Millions of Americans are diagnosed with SUD. See Figure 14.6<sup>4</sup> for a graphic of the number of people aged 12 and older with a substance use disorder in 2020.

4. This image is a derivative of “People Aged 12 or Older with a Past Year Substance Use Disorder (SUD); 2020” table by [Substance Abuse and Mental Health Services Administration](#). (2021). Key substance use and mental health indicators in the United States: Results from the 2020 National Survey on Drug Use and Health (HHS Publication No. PEP21-07-01-003, NSDUH Series H-56). Rockville, MD: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration. Used under Fair Use. Retrieved from <https://www.samhsa.gov/data/report/2020-nsduh-annual-national-report>






Figure 14.6 Number of People Aged 12 and Older With a Substance Use Disorder in 2020. Used under Fair Use.

SUD often develops gradually over time due to repeated misuse of a substance, causing changes in brain areas that control reward, stress, and executive functions like decision-making and self-control. Multiple factors influence whether a person will develop a substance use disorder such as the substance itself; the genetic vulnerability of the user; and the amount, frequency, and duration of the misuse.

Severe substance use disorders are commonly referred to as addictions.

**Addiction** is associated with compulsive or uncontrolled use of one or more substances. Addiction is a chronic illness that has the potential for both relapse and recovery. **Relapse** refers to the return to substance use after a significant period of abstinence. **Recovery** is a process of change through which individuals improve their health and wellness, live a self-directed life, and strive to reach their full potential. Although abstinence from substance misuse is a primary feature of a recovery lifestyle, it is not the only healthy feature.<sup>5</sup> The chronic nature of addiction means that some individuals may

5. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

relapse after an attempt at abstinence, which can be a normal part of the recovery process. Relapse does not mean treatment failure. Relapse rates for substance use are similar to rates for adherence to therapies for other chronic medical illnesses. There are a variety of medications that can be prescribed to assist with relapse prevention.<sup>6</sup>

Individuals with severe substance use disorders can overcome their disorder with effective treatment and regain health and social function, referred to as **remission**. When positive changes and values become part of a voluntarily adopted lifestyle, this is referred to as “being in recovery.”<sup>7</sup> Among the 29.2 million adults in 2020 who have ever had a substance use problem, 72.5 percent considered themselves to be in recovery.<sup>8</sup>

Many individuals seeking care in health care settings, such as primary care, obstetrics and gynecology, emergency departments, and medical-surgical units, have undiagnosed substance use disorders. Recognition and early treatment of substance use disorders can improve their health outcomes and reduce overall health care costs.<sup>9</sup>

6. National Institute on Drug Abuse. (2022). Drugs, brains, and behavior: The science of addiction treatment and recovery. <https://nida.nih.gov/publications/drugs-brains-behavior-science-addiction/treatment-recovery>
7. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
8. Substance Abuse and Mental Health Services Administration. (2021). *Key substance use and mental health indicators in the United States: Results from the 2020 national survey on drug use and health*. Center for Behavioral Health Statistics and Quality, Substance Abuse, & Mental Health Services Administration. <https://www.samhsa.gov/data/>
9. Substance Abuse and Mental Health Services Administration, & Office of the

# Substance Use Disorder in Nurses

Health care professionals are not immune to developing SUD. SUD is a chronic illness that can affect anyone regardless of age, occupation, economic circumstances, ethnic background, or gender. The National Council of State Boards of Nursing (NCSBN) created a brochure called *A Nurse's Guide to Substance Use Disorder in Nursing*. This brochure states that many nurses with substance use disorder (SUD) are unidentified, untreated, and may continue to practice when their impairment may endanger the lives of their clients. Because of the potential safety hazards to clients, it is a nurse's legal and ethical responsibility to report a colleague's suspected SUD to their manager or supervisor. It can be hard to differentiate between the subtle signs of SUD and stress-related behaviors, but three significant signs include behavioral changes, physical signs, and drug diversion.<sup>10</sup>

Behavioral changes include decreased job performance, absences from the unit for extended periods, frequent trips to the bathroom, arriving late or leaving early, and making an excessive number of mistakes including medication errors.<sup>11</sup>

Physical signs include subtle changes in appearance that may escalate over time; increasing isolation from colleagues; inappropriate verbal or emotional responses; and diminished alertness, confusion, or memory lapses. Signs of

Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

10. NCSBSN. (2025). *A nurse's guide to substance use disorder in nursing* [Brochure]. <https://www.ncsbn.org/nursing-regulation/practice/substance-use-disorder/substance-use-in-nursing.page>
11. NCSBSN. (2025). *A nurse's guide to substance use disorder in nursing* [Brochure]. <https://www.ncsbn.org/nursing-regulation/practice/substance-use-disorder/substance-use-in-nursing.page>

diversion include frequent discrepancies in opioid counts, unusual amounts of opioid wastage, numerous corrections of medication records, frequent reports of ineffective pain relief from clients, offers to medicate coworkers' clients for pain, and altered verbal or phone medication orders.<sup>12</sup>

**Drug diversion** occurs when medication is redirected from its intended destination for personal use, sale, or distribution to others. It includes drug theft, use, or tampering (adulteration or substitution). Drug diversion is a felony that can result in a nurse's criminal prosecution and loss of license.<sup>13</sup>

The earlier that a nurse is diagnosed with SUD and treatment is initiated, the sooner that client safety is protected and the better the chances for the nurse to recover and return to work. In most states, a nurse diagnosed with a SUD enters a nondisciplinary program designed by the Board of Nursing for treatment and recovery services. When a colleague treated for an SUD returns to work, nurses should create a supportive environment that encourages their continued recovery.<sup>14</sup>

► View the NCSBN PDF pamphlet [A Nurse's Guide to Substance Use Disorder in Nursing](https://www.ncsbn.org/nursing-regulation/practice/substance-use-disorder/substance-use-in-nursing.page).

12. NCSBSN. (2025). *A nurse's guide to substance use disorder in nursing* [Brochure]. <https://www.ncsbn.org/nursing-regulation/practice/substance-use-disorder/substance-use-in-nursing.page>
13. Nyhus, J. (2021). Drug diversion in healthcare. *American Nurse*. <https://www.myamericannurse.com/drug-diversion-in-healthcare/>
14. NCSBSN. (2025). *A nurse's guide to substance use disorder in nursing* [Brochure]. <https://www.ncsbn.org/nursing-regulation/practice/substance-use-disorder/substance-use-in-nursing.page>

## Nonsubstance-Related Disorders

**Nonsubstance-related disorders** are excessive behaviors related to gambling, viewing pornography, compulsive sexual activity, Internet gaming, overeating, shopping, overexercising, and overusing technologies. These behaviors are thought to stimulate the same addiction centers of the brain as addictive substances. However, gambling disorder is the only nonsubstance use disorder with diagnostic criteria listed in the *DSM-5*. See the *DSM-5* criteria for the diagnosis of a gambling disorder in the following box.

### ***DSM-5* Criteria for Gambling Disorder<sup>15</sup>**

Gambling disorder is defined as persistent and recurrent problematic gambling behavior leading to clinically significant impairment or distress, as indicated by four or more of the following criteria in a 12-month period. Additionally, the gambling behavior is not better explained by a manic episode.

- Needs to gamble with increasing amounts of money to achieve the desired excitement
- Is restless or irritable when attempting to cut down or stop gambling
- Has made repeated unsuccessful efforts to control, cut back, or stop gambling
- Is often preoccupied with gambling (e.g., persistent thoughts of reliving past gambling experiences, planning the next venture, or thinking of ways to get money with which to gamble)

15. American Psychiatric Association. (2013). *Desk reference to the diagnostic criteria from DSM-5*.

- Often gambles when feeling distressed (e.g., helpless, guilty, anxious, or depressed)
- After losing money gambling, often returns another day to get even (otherwise known as “chasing one’s losses”)
- Lies to conceal the extent of involvement with gambling
- Has jeopardized or lost a significant relationship, job, or educational or career opportunity because of gambling
- Relies on others to provide money to relieve desperate financial situation caused by gambling

## 14.5 Neurobiology of Substance Use Disorders

Severe substance use disorders, also called addictions, were once considered a moral failing or character flaw but are now known to be chronic illnesses. Scientific evidence shows that addiction to substances is a chronic brain disease that has potential for relapse and recovery.<sup>1</sup>

All addictive substances have powerful effects on the brain. These effects account for the euphoric or intensely pleasurable feelings that people experience during their initial use of alcohol or other substances. These feelings motivate people to use those substances again and again, despite the risks for significant harm. As individuals continue to misuse alcohol or other substances, progressive changes, called **neuroadaptations**, occur in the structure and function of the brain. These neuroadaptations drive the transition from controlled, occasional substance use to chronic misuse that can be difficult to control and can endure long after an individual stops using the substances. These changes can lead to the need for increased amounts of substances to achieve the same effect, referred to as tolerance. They may produce continued, periodic cravings for the substance that can lead to relapse. More than 60 percent of people treated for a substance use disorder experience relapse within the first year after they are discharged from treatment, and a person can remain at increased risk of relapse for many years.<sup>2</sup>

1. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
2. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

For many people, initial substance use involves an element of impulsivity (i.e., acting without foresight or regard for the consequences). For example, an adolescent may impulsively take a first drink, smoke a cigarette, experiment with marijuana, or succumb to peer pressure and try a party drug like Ecstasy. If the experience is pleasurable, this feeling positively reinforces the substance use, making the person more likely to take the substance again. Another person may take a substance to relieve negative feelings such as stress, anxiety, or depression. In this case, the temporary relief the substance brings from the negative feelings reinforces substance use, increasing the likelihood that the person will use again.<sup>3</sup>

Many other environmental and social stimuli can reinforce a behavior. For example, peer approval positively reinforces substance use for some people. Likewise, if drinking or using drugs with others provides a feeling of relief from social isolation, substance use is reinforced.<sup>4</sup>

Eventually, in the absence of the substance, a person may experience negative emotions such as stress, anxiety, or depression or the individual may feel physically ill. This is called withdrawal, which often leads the person to use the substance again to relieve the withdrawal symptoms. As use becomes an ingrained behavior, impulsivity shifts to compulsivity, and the primary drivers of repeated substance use shift from positive reinforcement (feeling pleasure) to negative reinforcement (feeling relief) as the person seeks to stop the negative feelings and physical illness that accompany withdrawal. Eventually,

3. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
4. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>



the person begins taking the substance not to get “high,” but rather to escape the “low” feelings from withdrawal. Compulsive substance seeking is a key characteristic of addiction, as well as loss of control over use. Compulsivity helps to explain why many people with addiction experience relapses after abstinence.<sup>5</sup>

Three regions of the brain are the key components in the development and persistence of substance use disorders: the basal ganglia, the extended amygdala, and the prefrontal cortex:

- The basal ganglia controls the rewarding, pleasurable effects of substance use and is responsible for the formation of habitual substance taking. Two subregions of the basal ganglia are particularly important in substance use disorders:
  - The nucleus accumbens, involved in motivation and the experience of reward.
  - The dorsal striatum, involved in forming habits and other routine behaviors.
- The extended amygdala is involved in the stress response and the feelings of unease, anxiety, and irritability that typically accompany substance withdrawal.
- The prefrontal cortex is involved in executive function (e.g., the ability to organize thoughts and activities, prioritize tasks, manage time, and make decisions), including exerting control over substance use.

These changes in the brain persist long after substance use stops and are associated with a high incidence of relapse with substance use disorders.

5. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

## Risk Factors

Many factors influence the development of substance use disorders, including developmental, environmental, social, and genetic factors, and co-occurring mental health disorders. Other conditions called protective factors protect people from developing a substance use disorder or addiction. Examples of protective factors include positive self-image, self-control, and social competence. The relative influence of these factors varies across individuals and the life span.

Whether an individual ever uses alcohol or another substance and whether that initial use progresses to a substance use disorder of any severity depends on a number of factors including the following<sup>6</sup>:

- A person's genetic makeup and biological factors
- The age when substance use begins
- Psychological factors related to a person's unique history and personality
- Environmental factors, such as the availability of drugs, family and peer dynamics, financial resources, cultural norms, exposure to stress, and access to social support

## Early Life Experiences

The experiences a person has early in childhood and in adolescence can set the stage for substance use and sometimes escalate to substance use disorder. Early life stressors (referred to as adverse childhood experiences) include physical, emotional, and sexual abuse; neglect; household instability

6. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

(such as parental substance use and conflict, mental illness, or incarceration of household members); and poverty.<sup>7</sup>

- ▶ Review information on adverse childhood experiences in the “[Mental Health and Mental Illness](#)” section of Chapter 1.

Adolescence is a critical vulnerable period for substance misuse and the development of substance use disorders because a characteristic of this developmental period is risk taking and experimentation. For some young people, this includes trying alcohol, marijuana, or other drugs. Additionally, the brain undergoes significant changes during this life stage, making it particularly vulnerable to substance exposure. For example, the frontal cortex, a region in the front part of the brain that includes the prefrontal cortex, does not fully develop until the early to mid-20s. Research shows that heavy drinking and drug use during adolescence affects development of this critical area of the brain.<sup>8</sup>

Approximately 74 percent of 18- to 30-year-olds admitted to treatment programs began using substances at the age of 17 or younger. Individuals who start using substances during adolescence often experience more chronic and intensive use, and they are at greater risk of developing a

7. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
8. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

substance use disorder compared with those who begin use at an older age. In other words, the earlier the exposure, the greater the risk.<sup>9</sup>

## Genetic and Molecular Factors

Genetic factors are thought to account for 40 to 70 percent of individual differences in risk for addiction. Although multiple genes are likely involved in the development of addiction, only a few specific gene variants have been identified that either predispose to or protect against addiction. Some of these variants have been associated with the metabolism of alcohol and nicotine, while others involve receptors and other proteins associated with key neurotransmitters and molecules involved in all parts of the addiction cycle. Genes involved in strengthening the connections between neurons and in forming drug memories have also been associated with addiction risk. Like other chronic health conditions, substance use disorders are influenced by the complex interplay between a person's genes and environment.<sup>10</sup>

## Concurrent Mental Health Disorders

In 2020, 17 million adults (6.7%) had both a substance use disorder (SUD) and any mental health illness (AMI) as illustrated in Figure 14.7.<sup>11</sup> The relationship

9. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
10. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
11. Image is a derivative of "Past Year Substance Abuse Disorder (SUD) and Any Mental Illness (AMI): Among Adults Aged 18 or Older; 2020" table by [Substance Abuse and Mental Health Services Administration](#). (2021). Key

between SUDs and mental disorders is known to be bidirectional, meaning the presence of a mental health disorder may contribute to the development or exacerbation of an SUD, or an SUD may contribute to the development or exacerbation of a mental health disorder. The combined presence of SUDs and mental health disorders results in greater functional impairment; worse treatment outcomes; higher morbidity and mortality; increased treatment costs; and higher risk for homelessness, incarceration, and suicide.




Figure 14.7 Concurrent Mental Illness (AMI) and Substance Use Disorder (SUD) in Adults in 2020. Used under Fair Use.

The reasons why substance use disorders and mental health disorders often occur together are not clear, but there are three possible explanations. One reason may be because certain substances may temporarily mask the

substance use and mental health indicators in the United States: Results from the 2020 National Survey on Drug Use and Health (HHS Publication No. PEP21-07-01-003, NSDUH Series H-56). Rockville, MD: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration. Used under Fair Use. Retrieved from <https://www.samhsa.gov/data/report/2020-nsduh-annual-national-report> .

symptoms of mental health disorders (such as anxiety or depression). A second reason may be that certain substances trigger a mental health disorder that otherwise would not have developed. For example, research suggests that alcohol use increases risk for PTSD by altering the brain's ability to recover from traumatic experiences. A third possible reason is that both substance use disorders and mental health disorders are caused by overlapping factors, such as particular genes, neurobiology, or exposure to traumatic or stressful life experiences.<sup>12</sup>

Mental health disorders and substance use disorders have overlapping symptoms, making diagnosis and treatment planning challenging. For example, people who use methamphetamine for a long period of time may experience paranoia, hallucinations, and delusions that can be mistaken for symptoms of schizophrenia.<sup>13</sup>

## Gender

Some groups of people are more vulnerable to substance misuse and substance use disorders. For example, biological males tend to drink more than biological females and are at higher risk for alcohol use disorder. However, biological females who use cocaine, opioids, or alcohol may progress from initial use to a substance use disorder faster than males. Compared with biological males, biological females also exhibit greater

12. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
13. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

withdrawal symptoms from some drugs such as nicotine and have higher levels of the stress hormone cortisol.<sup>14</sup>

## Race and Ethnicity

Neurobiological factors contributing to differential rates of substance use disorders across racial and ethnic groups have been researched. A study using functional magnetic resonance imaging (fMRI) found that African American smokers showed greater activation of the prefrontal cortex upon exposure to smoking-related cues than did White smokers, an effect that may partly contribute to the lower smoking-cessation success rates among African Americans.<sup>15</sup>

Alcohol research on racial and ethnic groups has shown that approximately 36 percent of East Asians carry a gene variant that alters the rate of alcohol metabolism, causing a buildup of acetaldehyde, a toxic by-product that produces symptoms such as flushing, nausea, and rapid heartbeat. Although these effects may protect some individuals of East Asian descent from alcohol use disorder, those who drink despite the effects are at increased risk for esophageal and head and neck cancers.<sup>16</sup>

14. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
15. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
16. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

# Individual, Family, and Community Level Risk Factors

An individual's risk factors for developing SUD include the following<sup>17</sup>:

- **Early initiation of substance use:** Engaging in alcohol or drug use at a young age
- **Early and persistent problem behavior:** Emotional distress, aggressiveness, and “difficult” temperaments in adolescents
- **Rebelliousness:** High tolerance for deviance and rebellious activities
- **Favorable attitudes toward substance use:** Positive feelings towards alcohol or drug use; low perception of risk
- **Peer substance use:** Friends and peers who engage in alcohol or drug use
- **Genetic predictors:** Genetic susceptibility to alcohol or drug use
- **Academic failure beginning in late elementary school:** Poor grades in school
- **Lack of commitment to school:** When a young person no longer considers the role of being a student as meaningful and rewarding or lacks investment or commitment to school

17. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>



## 14.6 Addiction Cycle

### Addiction Cycle

The addiction process involves a three-stage cycle that becomes more severe as a person continues to misuse substances, causing neuroadaptations in brain function that reduce a person's ability to control their substance use. Each stage is associated with one of the brain regions previously described (i.e., basal ganglia, extended amygdala, and prefrontal cortex). See Figure 14.8<sup>1</sup> for an image of the brain regions associated with the three stages of addiction. This three-stage model provides a useful way to understand the symptoms of addiction, the ways it can be prevented and treated, and the steps for recovery.<sup>2</sup>

These are the three stages of addiction:

- **Binge/Intoxication:** The stage at which an individual consumes an intoxicating substance and experiences its rewarding or pleasurable effects.
- **Withdrawal/Negative Affect:** The stage at which an individual experiences a negative emotional state in the absence of the substance.

1. Substance Abuse and Mental Health Services Administration (US); Office of the Surgeon General (US). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health* [Internet]. Washington (DC): US Department of Health and Human Services; 2016 Nov. Figure 2.3, The Three Stages of the Addiction Cycle and the Brain Regions Associated with Them. Used under Fair Use. Available from <https://www.ncbi.nlm.nih.gov/books/NBK424849/figure/ch2.f3/>
2. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

- **Preoccupation/Anticipation:** The stage at which one seeks substances again after a period of abstinence.




Figure 14.8 Brain Regions Associated With the Three Stages of Addiction. Used under Fair Use.

## Binge/Intoxication: Basal Ganglia

All addictive substances produce feelings of pleasure. These “rewarding effects” positively reinforce their use and increase the likelihood of repeated use. The rewarding effects of substances involve activity in the nucleus accumbens, including activation of the brain’s dopamine and opioid signaling system. Studies show that antagonists, or inhibitors, of dopamine and opioid

receptors can block drug and alcohol-seeking behaviors.<sup>3</sup> See Figure 14.9<sup>4</sup> for an illustration of rewarding actions of addictive substances on the nucleus accumbens.

3. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
4. Substance Abuse and Mental Health Services Administration (US); Office of the Surgeon General (US). *Facing Addiction in America: The Surgeon General's Report on Alcohol, Drugs, and Health* [Internet]. Washington (DC): US Department of Health and Human Services; 2016 Nov. Figure 2.5, Actions of Addictive Substances on the Brain. Used under Fair Use. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK424849/figure/ch2.f5/>




Figure 14.9 Rewarding Actions of Addictive Substances on the Nucleus Accumbens (NAc). Used under Fair Use.

Dopamine neurons (DA) are activated, either directly or indirectly, by all addictive substances, but particularly by stimulants and nicotine.<sup>5</sup>

The brain's opioid system includes naturally occurring opioid molecules (i.e., endorphins) and three types of opioid receptors (i.e., mu, delta, and kappa) and plays a key role in mediating the rewarding effects of opioids and alcohol.

5. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

Activation of the opioid system stimulates the nucleus accumbens (NAc) directly or indirectly through the GABA and dopamine (DA) system.<sup>6</sup>

Tetrahydrocannabinol (THC), the primary psychoactive component of marijuana, targets GABA neurons that stimulate the dopamine (DA) system, as well as directly stimulating the nucleus accumbens (NAc).<sup>7</sup>

Activation of the brain's reward system by alcohol and drugs not only generates the pleasurable feelings associated with those substances, but also ultimately triggers changes in the way a person responds to stimuli associated with the use of those substances. A person learns to associate the stimuli present while using a substance, including people, places, drug paraphernalia, and even internal states, such as mood, with the substance's rewarding effects. Over time, these stimuli can activate the dopamine system on their own and trigger powerful urges to take the substance. These "cravings" can persist even after the rewarding effects of the substance have diminished. As a result, exposure to people, places, or things previously associated with substance use can serve as triggers or cues that promote substance seeking and taking, even in people who are in recovery.<sup>8</sup>

6. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
7. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
8. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

## Withdrawal/Negative Affect Stage: Amygdala

The withdrawal/negative affect stage of addiction follows the binge/intoxication stage, and, in turn, sets up future rounds of binge/intoxication. During this stage, a person who has been using alcohol or drugs experiences withdrawal symptoms, including negative emotions and symptoms of physical illness when they stop taking the substance. Symptoms of withdrawal may occur with all addictive substances, although they vary in intensity and duration depending on both the type of substance and the severity of use. The negative feelings associated with withdrawal are thought to come from two sources: diminished activation in the reward circuitry of the basal ganglia and activation of the brain's stress systems in the extended amygdala.<sup>9</sup>

When used long-term, all addictive substances cause dysfunction in the brain's dopamine reward system. For example, brain imaging studies in individuals with addictions show long-lasting decreases in dopamine receptors even after stopping substance abuse. See Figure 14.10<sup>10</sup> for an image of brain imaging studies comparing an individual with cocaine addiction to someone without addiction. Decreases in the activity of the dopamine system have also been observed during withdrawal from stimulants, opioids, nicotine, and alcohol. Other studies also show that when an addicted person

9. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
10. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. Used under Fair Use. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>



is given a stimulant, it causes a smaller release of dopamine than when the same dose is given to a person who is not addicted.<sup>11</sup>




Figure 14.10 Decreased Dopamine Receptors in an Individual With a Cocaine Addiction. Used under Fair Use.

These findings suggest that people addicted to substances experience an overall reduction in the sensitivity of the brain's reward system (especially in the brain circuits involving dopamine), both to addictive substances and also to natural reinforcers, such as food and sex. This impairment explains why individuals who develop a substance abuse disorder often do not derive the same level of satisfaction or pleasure from once-pleasurable activities and may neglect eating. This loss of reward sensitivity may also account for the

11. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

compulsive escalation of substance use as addicted individuals attempt to regain the pleasurable feelings the reward system once provided.<sup>12</sup>

At the same time, a second process occurs during the withdrawal stage related to the activation of stress neurotransmitters in the extended amygdala. These stress neurotransmitters include corticotropin releasing factor (CRF), norepinephrine, and dynorphin. These neurotransmitters play a key role in the negative feelings associated with withdrawal and in stress-triggered substance use. When researchers used antagonists to block activation of addicted individuals' stress neurotransmitter systems, it reduced their substance intake in response to withdrawal and stress.<sup>13</sup>

## Preoccupation/Anticipation Stage: Prefrontal Cortex

The preoccupation/anticipation stage of the addiction cycle is the stage in which a person may begin to seek to use substances again after a period of abstinence. In this stage, an addicted person becomes preoccupied with using substances again, referred to as craving. This stage of addiction involves the brain's prefrontal cortex, the region that controls executive function (e.g., the ability to organize thoughts and activities; prioritize tasks; manage time; make decisions; and regulate one's own actions, emotions, and impulses). Executive function is essential for a person to make appropriate choices about whether or not to use a substance and to override strong urges to use it, especially when experiencing triggers related to the substance (e.g., being at

12. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
13. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>



a party where alcohol is served) or stressful experiences. People with alcohol, cocaine, or opioid use disorders show impairments in executive function, including disruption of decision-making and behavioral inhibition.<sup>14</sup>

14. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

## 14.7 Treatment of Substance Use Disorders

Research has shown that substance use disorders are similar in course, management, and outcome to other chronic illnesses, such as hypertension, diabetes, and asthma. It is possible to adopt the same type of chronic care management approach to the treatment of substance use disorders as is used to manage chronic physical illnesses. Remission of substance use disorders and even full recovery can be achieved if evidence-based care is provided for adequate periods of time by properly trained health care professionals and augmented with supportive monitoring, recovery support services, and social services.<sup>1</sup>

There are a spectrum of effective strategies and services available to identify, treat, and manage substance use disorders. Research shows that the most effective way to help someone with a substance misuse problem who is at risk for developing a substance use disorder is to intervene early, before the condition can progress. Screening for substance misuse is increasingly being provided in general health care settings so that emerging problems can be detected and early intervention provided. The addition of services to address substance use problems and disorders in mainstream health care has extended the continuum of care and includes a range of effective, evidence-based medications; behavioral therapies; and supportive services. However, a number of barriers have limited the widespread adoption of these services, including common myths about addiction, lack of resources, insufficient training, and workforce shortages. This is particularly true for the treatment of individuals with co-occurring substance use and mental health disorders.<sup>2</sup>

1. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
2. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's*

There are several common myths about addiction that make it harder for people with substance use disorders to seek treatment to get well. Review these myths in the following box.

### Common Myths About Addiction<sup>3</sup>

- Using drugs or alcohol is a choice, so if someone gets addicted, it's their fault.
  - **Reality:** Addiction is a consequence of many contributing factors, including genetics, neurobiology, adverse childhood effects (ACEs), trauma, and other influences.
- If someone just uses willpower, they should be able to stop using the substance.
  - **Reality:** For people who are genetically vulnerable to addiction, substance use can cause profound changes in the brain that hijack the reward pathway of the brain. Addictive substances flood the brain with neurotransmitters that signal pleasure. These changes create intense impulses to continue using the substances despite negative consequences of doing so.

*report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

3. Face It Together. (n.d.) *Common myths about addiction*. <https://www.wefaceittogether.org/learn/common-myths>

- Addiction affects certain types of people.
  - **Reality:** Addiction can affect anyone, no matter one's age, income, ethnicity, religion, family, or profession. Nationally, about one in eight people ages 12 and older are impacted.
- If someone has a stable job and family life, they can't be suffering from addiction.
  - **Reality:** Anyone is vulnerable to addiction. Many people hide the severity of their illness or don't get help because of stigma or shame.
- People have to become seriously ill before they can get well.
  - **Reality:** The longer a person waits to get help, the more changes happen to the brain, which can have deadly consequences like overdose. Studies show that people forced into treatment have an equal chance of successful recovery as people who initiate treatment on their own.
- Going to rehab will fix the problem.
  - **Reality:** Addiction is a chronic disease, similar to hypertension or diabetes, that can be controlled but not cured. Treatment is the first step towards wellness, but it is just the beginning. Staying well requires a lifelong commitment to managing the chronic disease.
- If someone relapses, they can never get better.

- **Reality:** Relapse is no more likely with addiction than other chronic illnesses like diabetes. Getting well involves changing deeply embedded behaviors that are significantly rewarded in the brain. Behavioral change takes time and effort, and setbacks can occur. A relapse can signal that the treatment approach or other supports need to change or that other treatment methods are needed. There is hope that people who experience a relapse will return to recovery.

Substance use disorder treatment is designed to help individuals stop or reduce harmful substance misuse, improve their health and social function, and manage their risk for relapse. For example, mild substance use disorders can be identified quickly in many medical settings and often respond to brief motivational interventions and/or supportive monitoring, referred to as guided self-change. In contrast, severe and chronic substance use disorders often require specialty substance use disorder treatment and continued post-treatment support to achieve full remission and recovery. To address the spectrum of problems associated with substance use disorders, a continuum of care is planned and implemented based on an individual's needs, including early intervention, treatment, and recovery support services.<sup>4</sup>

4. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

## Early Intervention

Early intervention services can be provided in a variety of settings (e.g., school clinics, primary care offices, mental health clinics) to people who have substance misuse problems or mild substance use disorders. These services are usually provided when an individual presents for another medical condition or social service need and is not necessarily seeking treatment for a substance use disorder. The goals of early intervention are to reduce the harms associated with substance misuse, reduce risk behaviors before they lead to injury, improve health and social function, and prevent progression from misuse to a substance use disorder. Early intervention consists of providing education about risks of substance use, safe levels of alcohol and medication use, and strategies to quit substance use. Referral to treatment services is provided as needed.<sup>5</sup>

## Populations Who Should Receive Early Intervention

Early intervention should be provided to children, adolescents, and adults who show signs of substance misuse or a mild substance use disorder. One group in need of early intervention is people who binge drink, particularly those aged 12 to 17, who are at higher risk for future substance use disorders because of their young age. Available research shows that brief, early interventions given by a respected care provider (such as a nurse, nurse educator, or physician) in the context of routine medical care can educate and motivate many individuals who are misusing substances to understand and acknowledge their risky behavior and reduce their substance use.<sup>6</sup>

5. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
6. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's*

# Components of Early Intervention

One structured approach to delivering early intervention to people showing signs of substance misuse and/or early signs of a substance use disorder is through screening and brief intervention (SBI). Research has shown that several methods of SBI are effective in decreasing substance misuse for a variety of populations in a variety of health care settings.

## SCREENING

Regardless of the substance, the first step of early intervention is using a screening tool to identify behaviors that put the individual at risk for harm or for developing a substance use disorder. Positive screening results should be followed by brief educational sessions tailored to the specific problems and interests of the individual. It should be delivered in a nonjudgmental manner, emphasizing both the importance of reducing substance use and the individual's ability to accomplish this goal. Follow-up evaluation should assess whether the screening and the brief intervention were effective in reducing the substance misuse or if formal treatment is required.<sup>7</sup>

Ideally, substance misuse screening should occur for all individuals who present to health care settings, including primary care, urgent care, mental health care, and emergency departments. Several validated screening instruments have been developed to help nonspecialty providers identify individuals who may have, or be at risk for, a substance use disorder. An example of an evidence-based screening tool is the Alcohol Use Disorders

*report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

7. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

Identification Test (AUDIT).<sup>8</sup> Based on a World Health Organization study, the AUDIT has become the world's most widely used alcohol screening instrument and is available in approximately 40 languages. It is a self-administered questionnaire consisting of ten questions pertaining to an individual's alcohol use. Read additional details about the AUDIT tool in the following box.

### **AUDIT Screening Tool**

- ▶ Review the full AUDIT tool: [Check Your Drinking: An Interactive self-test.](#)

The screening tool includes ten questions with answers ranging from 0 (Never) to 4 (4 or more times a week):

- How often do you have a drink containing alcohol?
- How many standard drinks containing alcohol do you have on a typical day when drinking?
- How often do you have six or more drinks on one occasion?
- During the past year, how often have you found that you were not able to stop drinking once you had started?
- During the past year, how often have you failed to do what was normally expected of you because of drinking?
- During the past year, how often have you needed a drink in the morning to get yourself going after a heavy drinking session?
- During the past year, how often have you had a feeling of

8. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>



guilt or remorse after drinking?

- During the past year, how often have you been unable to remember what happened the night before because you had been drinking?
- Have you or someone else been injured as a result of your drinking?
- Has a relative, friend, doctor, or other health worker been concerned about your drinking or suggested you cut down?

## BRIEF INTERVENTIONS

**Brief interventions** is a term used to describe quick therapeutic techniques used to initiate change in individuals with unhealthy behaviors. For example, motivational interviewing (MI) is a client-centered therapeutic technique that can be used to address a person's ambivalence to change their use of substances. It uses a conversational approach to help the client discover their interest in changing their behavior. The nurse asks the client to express their desire for change and any ambivalence they might have and then begins to work with the client on making a plan to change their behavior and a commitment to the change process. Individuals who receive MI are more likely to adhere to a treatment plan and subsequently achieve better outcomes.<sup>9</sup>

9. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

## REFERRAL

When an individual's substance use problem meets criteria for a substance use disorder or when brief interventions do not produce change, referral to specialized treatment should occur. This is called Screening, Brief Intervention, and Referral to Treatment (SBIRT). A referral for assessment and development of a clinical treatment plan is created with the client and tailored to meet their needs. Effective referral processes should incorporate strategies to motivate the client to accept the referral and assistance in navigating barriers for treatment.<sup>10</sup>

Compare the effectiveness of communication with a client being treated in the emergency department for injuries sustained when driving under the influence of alcohol in these YouTube videos:

▶ Video 1<sup>11</sup>: [Anti-SBIRT \(Doctor A\)](#)

▶ Video 2 (Using SBIRT)<sup>12</sup>: [Using SBIRT Effectively \(Doctor B\)](#)

10. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

11. SBIRTInsitute. (2011, March 2). *Anti-SBIRT (Doctor A)*. [Video]. YouTube. All rights reserved. <https://youtu.be/ZGETDcFcAbI>

12. SBIRTInsitute. (2011, March 2). *Using SBIRT Effectively (Doctor B)*. [Video]. YouTube. All rights reserved. <https://youtu.be/uL8QyJF2wVw>

# Treatment

Evidence-based treatment interventions include medications, behavioral therapies, and recovery services. Treatment can occur in a variety of settings, but treatment for severe substance use disorders has traditionally been provided in specialty substance use disorder treatment programs. The National Institute on Drug Abuse (NIDA) outlines the following evidence-based principles for effective treatment of adults and adolescents with substance use disorders<sup>13 14</sup>:

- Substance use disorders are complex but treatable diseases that affect brain function and behavior. Psychoactive substances alter the brain's structure and function, resulting in changes that persist long after substance use has ceased. This may explain why individuals with substance use disorder are at risk for relapse even after long periods of abstinence and despite the potentially devastating consequences of their behaviors.
- No single treatment is appropriate for everyone. Treatment varies depending on the type of drug and the characteristics of the individuals. Matching treatment settings, interventions, and services to an individual's particular problems and needs is critical to their ultimate success in returning to productive functioning in the family, workplace, and society.
- Treatment must be readily available. Because individuals may be uncertain about entering treatment, taking advantage of available services the moment people are ready for treatment is critical. Potential clients can be lost if treatment is not immediately available or readily

13. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

14. National Institute on Drug Abuse. (2025). *Treatment*. <https://nida.nih.gov/research-topics/treatment>

accessible.

- Effective treatment attends to multiple needs of the individual, not just their substance abuse. To be effective, treatment must address the individual's substance abuse, as well as associated medical, psychological, social, vocational, and legal problems. It is also important that treatment is tailored to the individual's age, gender, ethnicity, and culture.
- Remaining in treatment for an adequate period of time is critical. The appropriate duration for an individual depends on the type and degree of the client's problems and needs. Research indicates that most addicted individuals need at least three months in treatment to significantly reduce or stop their substance use and that the best outcomes occur with longer durations of treatment. Recovery from substance use disorder is a long-term process and frequently requires multiple episodes of treatment. As with other chronic illnesses, relapses can occur and should signal a need for treatment to be reinstated or adjusted. Because individuals often leave treatment prematurely, programs should include strategies to engage and keep clients in treatment.
- Behavioral therapies, including individual, family, or group counseling, are the most common types of treatment. Behavioral therapies vary in focus and may address a client's motivation to change, provide incentives for abstinence, build skills to resist drug use, replace substance-using activities with constructive and rewarding activities, improve problem-solving skills, and facilitate interpersonal relationships. Additionally, participation in group therapy and other peer support programs during and following treatment can help maintain abstinence.
- An individual's treatment plan must be continually evaluated and modified as needed to ensure it meets their changing needs. A client may require varying combinations of services and treatment components during the course of treatment and recovery. In addition to counseling or psychotherapy, clients may require medication, medical services, family therapy, parenting instruction, vocational rehabilitation, and/or social and legal services. For many clients, a recovery-oriented systems approach provides the best results, with the treatment intensity varying according to a person's changing needs.

- Many individuals with substance use disorders also have other mental health disorders. Treatment should address all conditions using appropriate medications.
- Treatment does not need to be voluntary to be effective. Sanctions or enticements from family, employment settings, and/or the criminal justice system can significantly increase treatment entry, retention rates, and the ultimate success of drug treatment interventions. Substance use during treatment must be monitored continuously because relapses during treatment do occur. Knowing their substance use is being monitored can be a powerful incentive for individuals to withstand urges to use substances. Monitoring also provides an early indication of a relapse, signaling an adjustment is needed in the individual's treatment plan to better meet their needs.
- Treatment programs should test clients for the presence of HIV/AIDS, hepatitis B and C, tuberculosis, and other infectious diseases, as well as provide targeted risk-reduction counseling. Many substance misuse-related behaviors put people at risk of infectious diseases. Targeted counseling reduces infectious disease. Counseling can also help those who are already infected to manage their illness. Moreover, engaging in substance abuse treatment can facilitate adherence to other medical treatments.

## Medications

Medications are an important element of treatment for many clients, especially when combined with counseling and other behavioral therapies. For example, methadone, buprenorphine, and naltrexone are effective in helping individuals addicted to heroin or other opioids stabilize their lives and reduce their substance misuse. Acamprosate, disulfiram, and naltrexone are medications approved for treating alcohol dependence. For persons addicted to nicotine, a nicotine replacement product (available as patches, gum, lozenges, or nasal spray) or an oral medication (such as bupropion or varenicline) can be an effective component of treatment when part of a

comprehensive behavioral treatment program.<sup>15</sup> See Table 14.7 for a list of medications approved by the FDA to treat alcohol and opioid use disorders.

Table 14.7 Pharmacotherapy to Treat Alcohol and Opioid Use Disorders<sup>16</sup>

15. National Institute on Drug Abuse. (2025). *Treatment*. <https://nida.nih.gov/research-topics/treatment>
16. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

Medication	Use	DEA Schedule	Application
<b>Buprenorphine-naloxone</b>	Opioid use disorder	CIII	Used for detoxification or maintenance of abstinence.
<b>Methadone</b>	Opioid use disorder	CII	Used for withdrawal and long-term maintenance of abstinence of opioid addiction. Dispersed only at opioid treatment centers certified by SAMHSA and approved by state authority.
<b>Naltrexone</b>	Opioid use disorder and alcohol use disorder	Not scheduled under the Controlled Substances Act	Block opioid receptors, reduce cravings, and diminish rewarding effects of opioids and alcohol. Extended-release injections are recommended to prevent relapse.
<b>Acamprosate</b>	Alcohol use disorder	Not scheduled under the Controlled Substances Act	Used for maintenance of alcohol abstinence.
<b>Disulfiram</b>	Alcohol use disorder	Not scheduled under the Controlled Substances Act	Causes severe physical reactions when alcohol is ingested, such as nausea, flushing, and heart palpitations. The knowledge that the reaction will occur acts as a deterrent to drinking alcohol.

Medically assisted detoxification is only the first stage of treatment and by itself does little to change long-term substance misuse. Although medically assisted detoxification can safely manage the acute physical symptoms of withdrawal, detoxification alone is rarely sufficient to help individuals achieve long-term abstinence of substances. Clients should be encouraged to continue long-term treatment and recovery services following detoxification.<sup>17</sup>

17. National Institute on Drug Abuse. (2025). *Treatment*. <https://nida.nih.gov/research-topics/treatment>

# Behavioral Therapies

In addition to medications, effective treatment of SUD includes behavioral therapies to help clients recognize the impact of substance misuse on their interpersonal relationships and ability to function in a healthy, safe, and productive manner. Behavioral therapies also teach and motivate clients to change their behaviors as a way to control their substance use disorders.<sup>18</sup> Evidence-based behavioral therapies include cognitive-behavioral therapy, dialectical behavior therapy, family therapy, contingency management, community reinforcement approach, motivational enhancement therapy, matrix model, and twelve-step facilitation. These therapies are further described in the following subsections.

## COGNITIVE-BEHAVIORAL THERAPY

The theoretical foundation for cognitive-behavioral therapy (CBT) is that substance use disorders develop, in part, as a result of maladaptive behavior patterns and dysfunctional thoughts. As a result, CBT treatments involve techniques to modify such behaviors and improve coping skills by emphasizing the identification and modification of dysfunctional thinking. CBT is a short-term approach, usually involving 12 to 24 weekly individual sessions. These sessions typically explore the positive and negative consequences of substance use with self-monitoring as a mechanism to recognize cravings and other situations that may lead the individual to relapse. They also help the individual develop healthy coping strategies.

## DIALECTICAL BEHAVIOR THERAPY

Dialectical behavior therapy (DBT) is an evidence-based therapy that teaches

18. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>



a skill called mindfulness. Multiple research studies have noted that mindfulness is potentially useful in helping people diagnosed with SUD gain mastery over substance cravings.<sup>19</sup>

- ▶ Review information about cognitive-behavioral therapy and dialectical behavior therapy in the “[Treatments for Depression](#)” section of the “Depressive Disorders” chapter.

## FAMILY THERAPY

Family behavior therapy (FBT) is a therapeutic approach used for both adolescents and adults that addresses not only substance use but also other issues the family may be experiencing, such as mental health disorders and family conflict. FBT includes up to 20 treatment sessions that focus on developing skills and setting behavioral goals. Basic necessities are reviewed and inventoried with the client, and the family pursues resolution strategies and addresses activities of daily living, including violence prevention and HIV/AIDS prevention.<sup>20</sup>

19. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
20. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

## CONTINGENCY MANAGEMENT

Behavior change involves learning new behaviors and changing old behaviors. Positive rewards or incentives for changing behavior can aid this process. Contingency management involves giving tangible rewards to individuals to support positive behavior change and has been found to effectively treat substance use disorders. In this therapy clients receive a voucher with monetary value that can be exchanged for food items, healthy recreational options (e.g., movies), or other sought-after goods or services when they exhibit desired behavior such as drug-free urine tests or participation in treatment activities. Clinical studies comparing voucher-based reinforcement to traditional treatment regimens have found that voucher-based reinforcement is associated with longer treatment engagement, longer periods of abstinence, and greater improvements in personal function. These positive findings, initially demonstrated with individuals with cocaine use disorders, have been reproduced in individuals with alcohol, opioid, and methamphetamine use disorders.<sup>21</sup>

## COMMUNITY REINFORCEMENT APPROACH

Community reinforcement approach (CRA) plus vouchers is an intensive 24-week outpatient program that uses incentives and reinforcers to reward individuals who reduce their substance use. Individuals are required to attend one to two counseling sessions each week that emphasize improving relations, acquiring skills to minimize substance use, and reconstructing social activities and networks to support recovery. Individuals receiving this treatment are eligible to receive vouchers with monetary value if they provide drug-free urine tests several times per week. Research has demonstrated that

21. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

CRA plus vouchers promotes treatment engagement and facilitates abstinence.<sup>22</sup>

## MOTIVATIONAL ENHANCEMENT THERAPY

Motivational enhancement therapy (MET) uses motivational interviewing techniques to help individuals resolve any uncertainties they have about stopping their substance use. MET works by promoting empathy, developing client awareness of the discrepancy between their goals and their unhealthy behavior, avoiding argument and confrontation, addressing resistance, and supporting self-efficacy to encourage motivation and change. The therapist supports the client in executing the behaviors necessary for change and monitors progress toward client-expressed goals.<sup>23</sup>

## MATRIX MODEL

The matrix model is a structured, multi-component behavioral treatment that consists of evidence-based practices, including relapse prevention, family therapy, group therapy, drug education, and self-help, delivered in a sequential and clinically coordinated manner. The model consists of 16 weeks of group sessions held three times per week, which combine CBT, family education, social support, individual counseling, and urine drug testing.<sup>24</sup>

22. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

23. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

24. Substance Abuse and Mental Health Services Administration, & Office of the

## TWELVE-STEP FACILITATION

Twelve-step facilitation (TSF), an individual therapy typically delivered in 12 weekly sessions, is designed to prepare individuals to understand, accept, and become engaged in Alcoholics Anonymous (AA), Narcotics Anonymous (NA), or similar 12-step programs.<sup>25</sup> Twelve-step programs are further discussed under the “Recovery Support Services” section below.

TSF focuses on three key ideas:

- Acceptance: Realizing that their substance use is part of a disorder, that life has become unmanageable because of alcohol or drugs, that willpower alone will not overcome the problem, and that abstinence is the best alternative
- Surrender: Giving oneself to a higher power, accepting the fellowship and support structure of other recovering individuals, and following the recovery activities laid out by a 12-step program
- Active involvement in a 12-step program

## Recovery Support Services

In addition to medications and behavioral therapies, effective treatment of SUD includes recovery support services (RSS). Recovery support services provided by substance use disorder treatment programs and community organizations provide support to individuals receiving treatment for SUD, as well as ongoing support after treatment. These supportive services are

Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

25. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

typically delivered by trained case managers, recovery coaches, and/or peers. Specific RSS include assistance in navigating systems of care, removing barriers to recovery, staying engaged in the recovery process, and providing a social context for individuals to engage in community living without substance use. RSS can be effective in promoting healthy lifestyle techniques to increase resilience skills, reduce the risk of relapse, and help achieve and maintain recovery. Individuals who participate in RSS typically have better long-term recovery outcomes.<sup>26</sup>

Recovery goes beyond abstinence and the remission of substance use disorder to include a positive change in the whole person. There are many paths to recovery. People choose their individual pathway based on their cultural values, socioeconomic status, psychological and behavioral needs, and the nature of their substance use disorder.<sup>27</sup>

A study of over 9,000 individuals with previous substance use disorders asked how they defined recovery. These three themes emerged<sup>28</sup>:

- **Abstinence:** 86 percent viewed abstinence as part of their recovery, but

26. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
27. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
28. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

the remainder did not think abstinence was important for their recovery. However, abstinence was considered “essential” by those affiliated with 12-step mutual aid groups.

- **Personal growth:** “Being honest with myself” was endorsed as part of recovery by 98 percent of participants. Other almost universally endorsed elements included “handling negative feelings without using alcohol or drugs” and “being able to enjoy life without alcohol or drugs.” Almost all study participants viewed their recovery as a process of growth and development, and about two thirds saw it as having a spiritual dimension.
- **Service to others:** Engaging in service to others was another prominent component of how study participants defined recovery. This is perhaps because during periods of heavy substance misuse, individuals may damage interpersonal relationships, which they later regret and attempt to resolve during recovery. Service to others has evidence of helping individuals maintain their own recovery.<sup>29</sup>

Recovery-Oriented Systems of Care (ROSC) embrace the idea that severe substance use disorders are most effectively addressed through a chronic care management model that includes long-term, outpatient care, recovery housing, and recovery coaching and management checkups. ROSC are designed to be easy to navigate for people seeking help, transparent in their operations, and responsive to the cultural diversity of the communities they serve. ROSC often use long-term recovery management protocols, such as recovery management checkups and telephone case monitoring.<sup>30</sup>

29. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

30. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's*

Even when remission is achieved after one or two years, it can take four to five more years before an individual's risk of relapse drops below 15 percent (the level of risk that people in the general population have of developing a substance use disorder in their lifetime). As a result, similar to other chronic illnesses, a person with a serious substance use disorder often requires ongoing monitoring and management to maintain remission and to provide early reintervention should relapse occur. Recovery support services (RSS) refer to the collection of community services that can provide emotional and practical support for continuing remission, as well as daily structure and rewarding alternatives to substance use. Recovery supports include services such as mutual aid groups, recovery coaches, recovery housing, community care, and education-based recovery support.

## MUTUAL AID GROUPS

Mutual aid groups, such as 12-step groups, are well-known recovery supports, and they share a number of features. Members share their substance use problem and value learning from each other's experiences as they focus on personal-change goals. The groups are voluntary associations that charge no fees and are member-led.

### Alcoholics Anonymous

Alcoholics Anonymous (AA) has been in existence since 1935. Its philosophy, approach, and format have been adapted by groups focusing on recovery from other substances, such as Narcotic Anonymous, Cocaine Anonymous, Marijuana Anonymous, and Crystal Meth Anonymous. AA and derivative programs share two major components: social fellowship and a 12-step program of action that was formulated based on members' experiences of recovery from severe alcohol use disorders. These 12 steps are ordered in a

*report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

logical progression, beginning with accepting that one cannot control one's substance use, followed by abstaining from substances permanently, and transforming one's spiritual outlook, character, and relationships with other people.<sup>31</sup>

Research studying 12-step mutual aid groups, specifically those focused on alcohol, has shown that participation in the groups promotes an individual's recovery by strengthening recovery-supportive social networks; increasing members' abilities to cope with risky social contexts and negative emotions; augmenting motivation to recover; reducing depression, craving, and impulsivity; and enhancing psychological and spiritual well-being.<sup>32</sup>

► Find a local Alcoholics Anonymous group near you: [Find A.A. Near You](#)

## Al-Anon and Alateen

Friends and family members often suffer when a loved one has a substance use disorder. This can include worrying about their loved one experiencing accidents, injuries, legal consequences, diseases, or death or experiencing verbal or physical abuse. Mutual aid groups provide emotional support to

31. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
32. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>



concerned significant others to help them systematically and strategically cope with the problems related to the substance misuse of their loved one.<sup>33</sup>

Al-Anon is a mutual aid group for family members dealing with substance misuse in a loved one. Like AA, Al-Anon is based on a 12-step philosophy and provides support whether or not members' loved ones seek help or achieve remission or recovery. More than 80 percent of Al-Anon members are women. The principal goal of Al-Anon is to foster emotional stability and "loving detachment" from the loved one rather than coaching members to "get their loved one into treatment or recovery." Al-Anon includes Alateen that focuses on the specific needs of adolescents affected by a parent's or other family member's substance use. Research studies regarding the effectiveness of Al-Anon show that participating family members experience reduced depression, anger, and relationship unhappiness at rates comparable to those of individuals receiving psychological therapies.<sup>34</sup>

► Find a local Al-Anon group near you: [Al-Anon Meetings](#)

## RECOVERY COACHES

Voluntary and paid recovery coaches help individuals discharging from

33. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
34. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

treatment to connect to community services while addressing any barriers or problems that may hinder the recovery process. A recovery coach's responsibilities may include providing strategies to maintain abstinence, connecting people to recovery housing and social services, and helping people develop personal skills that maintain recovery.<sup>35</sup>

## RECOVERY HOUSING

Recovery-supportive houses provide both a substance-free environment and mutual support from fellow recovering residents. Many residents stay in recovery housing during or after outpatient treatment, with self-determined residency lasting for several months to years. Residents often informally share resources and give advice based on their experience in accessing health care, finding employment, managing legal problems, and interacting with the social service system.<sup>36</sup>

## COMMUNITY CARE

Recovery community centers may host mutual aid group meetings; offer recovery coaching, education, and social events; and provide access to other resources such as housing, education, and employment. Some recovery community centers encourage community members to engage in advocacy

35. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

36. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

to combat negative public attitudes, educate the community, and improve supports for recovery in the community.<sup>37</sup>

## EDUCATION-BASED RECOVERY SUPPORT

Education-based recovery support services are designed to help individuals in early substance use disorder recovery achieve their educational goals while also focusing on the areas of their social, emotional, spiritual, and physical well-being needed to help sustain recovery. High school and college environments can be difficult for students in recovery because of high levels of substance misuse among other students, peer pressure to engage in substance use, and widespread availability of alcohol and drugs. High school and collegiate recovery support programs provide recovery-supportive environments and peer engagement with other students in recovery.<sup>38</sup>

## Planning Individualized Treatment

After an individual is assessed and diagnosed with substance use disorder (SUD) by a trained professional based on criteria in the *Diagnostic and Statistical Manual of Mental Disorders (DSM-5-TR)*, a collaborative, personalized treatment plan is designed with the client to meet their specific needs. The treatment plan and goals should be person-centered and include strength-based approaches that draw upon an individual's strengths and resources to keep them engaged in care. Individualized treatment plans

37. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
38. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

should consider age, gender identity, race and ethnicity, language, health literacy, religion/spirituality, sexual orientation, culture, trauma history, and co-occurring physical and mental health problems. This increases the likelihood of successful treatment engagement and retention. Throughout treatment, individuals should be periodically reassessed to determine response to treatment and to make any needed adjustments to the treatment plan.<sup>39</sup>

Nurses can improve engagement and retention in treatment programs by building a strong therapeutic alliance with the client, effectively using evidence-based motivational strategies, acknowledging the client's individualized barriers, and creating a positive environment. Referring individuals to recovery support services, such as child care, housing, and transportation, can also improve retention in treatment.<sup>40</sup>

## Treatment Settings and Continuum of Care

The treatment of severe substance use disorder is typically delivered in freestanding programs in various settings (e.g., hospital, residential, or outpatient settings) that vary in the frequency of care delivery (e.g., daily sessions to monthly visits), range of treatment components offered, and planned duration of care. As clients progress in treatment and begin to meet the goals of their individualized treatment plan, they transfer from clinical

39. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

40. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

management in residential or intensive outpatient programs to less clinically intensive outpatient programs that promote client self-management.<sup>41</sup>

A typical progression for someone who has a severe substance use disorder might start with 3 to 7 days in a medically managed withdrawal program, followed by a 1- to 3-month period of intensive rehabilitative care in a residential treatment program, followed by intensive outpatient program (2 to 5 days per week for a few months) and later in a traditional outpatient program that meets 1 to 2 times per month. For many clients whose current living situations are not conducive to recovery, outpatient services should be provided in conjunction with recovery-supportive housing. In general, clients with serious substance use disorders are recommended to stay engaged for at least one year in the treatment process, which may involve participating in three to four different programs or services at reduced levels of intensity, all of which are ideally designed to help the client prepare for continued self-management after treatment ends.<sup>42</sup>

The levels of the treatment continuum include the following:

- **Medically monitored and managed inpatient care:** An intensive service delivered in an acute, inpatient hospital setting. These programs are typically necessary for individuals who require withdrawal management, primary medical and nursing care, and for those with co-occurring mental and physical health conditions. Treatment is available 24 hours a

41. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

42. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

day and usually provided by an interdisciplinary team of health care professionals who can address serious mental and physical health needs.

- **Residential services:** Organized services, also in a 24-hour setting but outside of a hospital. These programs typically provide support, structure, and an array of evidence-based clinical services. Such programs are appropriate for physically and emotionally stabilized individuals who may not have a living situation that supports recovery, may have a history of relapse, or have co-occurring physical and/or mental health illnesses.
- **Partial hospitalization and intensive outpatient services:** Services range from counseling and education to clinically intensive programming. Partial hospitalization programs are used as a step-down treatment option after completing residential treatment and are usually available 6 to 8 hours a day during the work week. These services are considered to be approximately as intensive but less restrictive than residential programs and are appropriate for clients living in an environment that supports recovery but who need structure to avoid relapse.
- **Outpatient services:** Group and individual behavioral interventions and medications when appropriate. These components of care can be offered during the day, before or after work or school, or in the evenings and weekends. Typically, outpatient programs are appropriate as the initial level of care for individuals with a mild to moderate substance use disorder or as continuing care after completing more intensive treatment. Outpatient programs are also suitable for individuals with concurrent mental health conditions.

## Helping Individuals in Need of Treatment

Despite the fact that substance use disorders are widespread, only a small percentage of people receive treatment because of the barriers previously discussed. Results from the 2020 National Survey of Drug Use and Health (NSDUH) indicate that among people aged 12 or older in 2020, 14.9 percent (or 41.1 million people) needed substance use treatment in the past year (defined if they had a SUD diagnosed in the past year or if they received substance use

treatment at a specialty facility in the past year). There are many common reasons people do not seek treatment<sup>43</sup>:

- Not ready to stop using (40.7 percent)
- Do not have health care coverage/could not afford (30.6 percent)
- Perceived negative effect on their job (16.4 percent) or cause neighbors/community to have a negative opinion (8.3 percent)
- Do not know where to go for treatment (12.6 percent) or no program has the type of treatment desired (11.0 percent)
- Do not have transportation, the programs are too far away, or hours are inconvenient (11.8 percent)

Nurses can use motivational interviewing strategies to explore clients' reasons for not seeking treatment and address their perceived barriers to treatment.

## Strategies to Reduce Harm

Strategies to reduce the harm associated with substance use have been developed to engage people in treatment, as well as address the needs of individuals who are not yet ready to participate in treatment. Harm reduction programs provide public health services to prevent and reduce substance use-related risks among those actively using substances. Strategies include outreach and education programs, needle/syringe exchange programs, overdose prevention education, and improving public access to naloxone to reverse potentially lethal opioid overdose. These strategies are designed to reduce negative consequences for people with substance use disorders and those around them, such as overdose and the transmission of HIV and other

43. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

infectious diseases. They also encourage individuals to seek treatment to reduce, manage, and stop their substance use.<sup>44</sup>

44. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>




## 14.8 Prevention of Substance Use Disorders

The misuse of alcohol and drugs and substance use disorders has a significant impact on public health in the United States. The Centers for Disease Control and Prevention (CDC) reports over 80,000 drug overdose deaths occurred in the United States in 2024, and overdose deaths in urban and rural populations have shown a significant upward trend over the past 20 years.<sup>1</sup> See Figure 14.11<sup>2</sup> for a graphic of overdose rates in urban and rural areas.

1. National Center for Health Statistics. (2025). *Provisional drug overdose death counts*. <https://www.cdc.gov/nchs/nvss/vsrr/drug-overdose-data.htm>
2. “db403-fig1.png” from by Hedegaard, H., & Spencer, M. R. is in the [Public Domain](#). Access the full report at <https://www.cdc.gov/nchs/products/databriefs/db403.htm#fig1>.

Figure 1. Age-adjusted rates of drug overdose deaths, by urban and rural residence: United States, 1999–2019



<sup>1</sup>Significant increasing trend from 1999 to 2017, with different rates of change over time; stable trend from 2017 through 2019,  $p < 0.05$ .

<sup>2</sup>Significant increasing trend from 1999 through 2019, with different rates of change over time,  $p < 0.05$ .

NOTES: Drug overdose deaths were identified using *International Classification of Diseases, 10th Revision* underlying cause-of-death codes X40–X44, X60–X64, X85, and Y10–Y14. Age-adjusted death rates were calculated using the direct method and the 2000 U.S. standard population. Decedent's county of residence was classified as urban or rural based on the 2013 NCHS Urban–Rural Classification Scheme for Counties. Access data table for Figure 1 at: <https://www.cdc.gov/nchs/data/databriefs/db403-tables-508.pdf#1>.

SOURCE: National Center for Health Statistics, National Vital Statistics System, Mortality.

Figure 14.11 Drug Overdose Rates in Urban and Rural Areas

Substance misuse is associated with a wide range of health and social problems, including heart disease, stroke, high blood pressure, various cancers (e.g., liver, lung, and colon cancer), mental health disorders, neonatal abstinence syndrome, driving under the influence (DUI) injuries and fatalities, incarcerations, sexual assaults and rapes, unintended pregnancies, sexually transmitted infections, and bloodborne pathogens like hepatitis and human immunodeficiency virus (HIV).<sup>3</sup> Given the impact of substance misuse on

3. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

public health, it is critical to implement preventative interventions to stop substance misuse from starting, as well as to identify and intervene early with individuals who have already begun to misuse substances.

## Promoting Protective Factors

Experiencing a family member's substance use disorder is considered an adverse childhood event (ACE) that can impact a child's risk for developing behavioral problems, substance misuse, and chronic illness. Targeted prevention programs implemented at the family, school, and individual levels can complement broader population-level policies by promoting protective factors for children and adolescents.

- ▶ Read more about ACEs in the "[Adverse Childhood Experiences](#)" section of the "Trauma, Abuse, and Violence" chapter.

Protective factors help prevent substance use disorders from developing despite the presence of risk factors. Protective factors exist at the individual, family, school, and community levels.

Examples of interventions to promote protective factors at the individual level include the following<sup>4</sup>:

- **Social, emotional, and behavioral competence:** Promoting interpersonal skills that help youth integrate feelings, thoughts, and actions to achieve specific social and interpersonal goals.
- **Self-efficacy:** Enhancing an individual's belief that they can modify,

4. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

control, or abstain from substance use.

- **Spirituality:** Supporting beliefs in a higher being or involvement in spiritual practices or religious activities.
- **Resiliency:** Encouraging an individual's capacity for adapting to change and coping with stressful events in healthy and flexible ways.

Interventions to promote protective factors at the family, school, and community levels are as follows<sup>5</sup>:

- **Opportunities for positive social involvement:** Creating developmentally appropriate opportunities to be meaningfully involved with the family, school, or community.
- **Recognition for positive behavior:** Providing community and family recognition of individuals' efforts and accomplishments to encourage positive future behaviors.
- **Bonding:** Promoting attachment, commitment, and positive communication with family members, schools, and communities.
- **Marriage or committed relationships:** Encouraging committed relationships with people who do not misuse alcohol or drugs.
- **Healthy beliefs and standards for behavior:** Establishing family, school, and community norms that communicate clear and consistent expectations about not misusing alcohol or drugs.

## Prevention Interventions

The Institute of Medicine describes three categories of prevention interventions: universal, selective, and indicated. Universal interventions are aimed at all members of a given population, selective interventions are aimed at a subgroup determined to be at high-risk for substance use, and indicated

5. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

interventions are targeted for individuals who are already misusing substances but have not developed a substance use disorder. Examples of evidence-based prevention interventions for each category are provided in the following subsections.

## Universal Interventions

Universal interventions include policies that affect the entire population, such as the setting the minimum legal drinking age or reducing the availability of substances in a community. For example, laws targeting alcohol-impaired driving, such as license revocation for impaired driving and 0.08 legal blood alcohol (BAC) limits have helped cut alcohol-related traffic deaths per 100,000 in half since the early 1980s.<sup>6</sup>

Several family-focused, universal prevention interventions show substantial preventive effects on substance use. Two examples are the Strengthening Families Program and the I Hear What You're Saying program<sup>7</sup>:

- **Strengthening Families Program (SFP):** A widely used seven-session, family-focused program that enhances parenting skills, such as nurturing, setting limits, and communicating, as well as promoting adolescents' skills in refusing substances. Across multiple studies conducted in rural United States communities, SFP showed reductions in tobacco, alcohol, and drug use up to nine years after the intervention (i.e., to age 21)

6. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

7. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

compared with youth who were not assigned to the SFP. SFP also shows reductions in prescription drug misuse up to 13 years after the intervention (i.e., to age 25). Strong African American Families, a cultural adaptation of SFP, shows reductions in early initiation and rate of alcohol use for Black or African American rural youth.

- **I Hear What You're Saying:** An Internet-based program that involves nine 45-minute sessions to improve communication, establish family rules, and manage conflict. Specifically focused on mothers and daughters, follow-up results showed lower rates of substance use in an ethnically diverse sample.<sup>8</sup>

## Selective Interventions

Selective interventions are delivered to particular communities, families, or children who, due to exposure to risk factors, are at increased risk of substance misuse problems. Target audiences may include families living in poverty, the children of substance-misusing parents, or children who have difficulties with social skills. Selective interventions deliver specialized prevention services to individuals with the goal of reducing identified risk factors, increasing protective factors, or both. Examples of selective intervention programs are the Nurse-Family Partnership, Familias Unidas, and Brief Alcohol Screening and Intervention for College Students<sup>9</sup>:

8. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
9. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

- **Nurse-Family Partnership:** A program focused on children younger than age 5 has shown significant reductions in the use of alcohol in the teen years compared with those who did not receive the intervention. Trained nurses provide an intensive home visitation intervention for at-risk, first-time mothers during pregnancy. This intervention provides ongoing education and support to improve pregnancy outcomes, infant health, and development of parenting skills.<sup>10</sup>
- **Familias Unidas:** A family-based intervention for Hispanic or Latino youth that includes both multi-parent groups (eight weekly 2-hour sessions) and four to ten 1-hour individual family visits. It has been shown to lower substance use or delay the start of substance use among adolescents.<sup>11</sup>
- **Brief Alcohol Screening and Intervention for College Students (BASICS):** A brief motivational intervention designed to help students reduce alcohol misuse and negative consequences of their drinking. It consists of two 1-hour interviews with a brief online assessment after the first session. The first interview gathers information about alcohol consumption patterns and personal beliefs about alcohol while providing instructions for self-monitoring drinking between sessions. The second interview uses data from the online assessment to develop personalized, normative feedback that reviews negative consequences and risk factors, clarifies perceived risks and benefits of drinking, and provides options for reducing alcohol use and its consequences. Follow-up studies of students who used BASICS have shown reductions in the quantity of drinking in

10. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

11. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

the general college population, as well as fraternity members.

## Indicated Prevention Interventions

Indicated prevention interventions are directed to those who are already involved in substance misuse but who have not yet developed a substance use disorder. An example of an indicated prevention intervention is Coping Power, a 16-month program for children in Grades 5 and 6. The program uses skills-based training to increase social competence, self-regulation, and positive parental involvement. Results include reduced substance use, delinquency, and aggressive behaviors.<sup>12</sup>

12. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>



# 14.9 Applying the Nursing Process to Substance Use Disorders

## Assessment (Recognizing Cues)

### Mental Status Examination

See Table 14.9a for common findings when assessing a client with a substance use disorder. (See expected findings for these components of a mental status examination in the “[Assessment](#)” section in Chapter 4.) Critical findings that require immediate notification of the provider are bolded with an asterisk. This table helps identify clinical features that may present during intoxication, withdrawal, or long-term substance use and supports a comprehensive nursing assessment.

Table 14.9a Common Findings During Mental Status Examinations for Clients With Substance Use Disorders<sup>1</sup>

1. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

Mental Status Examination Component	Common Findings in Substance Use Disorders (*Indicates immediately notify provider)
<b>Signs of Distress</b>	May present with anxiety, agitation, physical discomfort, or withdrawal symptoms (e.g., tremors, sweating, nausea). May appear drowsy, euphoric, or dysregulated depending on the substance used and timing.
<b>Level of Consciousness and Orientation</b>	Varies with substance type and use pattern. May be fully alert or display fluctuating levels of consciousness, confusion, or disorientation—especially during withdrawal or overdose of alcohol, opioids, and benzodiazepines.
<b>Appearance and General Behavior</b>	May appear disheveled or neglected, especially with chronic use of a substance. Signs of intoxication or withdrawal may include dilated or constricted pupils, slurred speech, or slowed/increased movements. Behavior may be guarded, restless, or impulsive. Track marks may be present for those using IV substances.
<b>Speech</b>	Speech may be slurred, rapid, pressured, or incoherent, depending on substance being used.
<b>Motor Activity</b>	May display tremors, psychomotor agitation (e.g., pacing, jitteriness), retardation (e.g., sedation), or repetitive movements. Falls, unsteadiness, or hyperactivity may also occur.
<b>Mood and Affect</b>	Mood may fluctuate based on substance used and timing. Mood may be euphoric, irritable, anxious, depressed, or flat. Clients may appear emotionally labile or detached. Intoxication may mask the client's true emotional state and withdrawal may increase distress.

<b>Thought and Perception</b>	Thought content may include denial, guilt, paranoia, or preoccupation with obtaining substances. In severe intoxication or withdrawal, hallucinations, delusions, or psychosis may occur (e.g., alcohol-induced hallucinosis).
<b>Attitude and Insight</b>	Insight is often impaired—clients may minimize or deny the extent of use or its consequences. Attitude may range from cooperative to guarded, defensive, or evasive depending on trust and withdrawal status.
<b>Cognitive Abilities and Level of Judgment</b>	Impaired attention, memory deficits, or poor concentration are common. Long-term use may cause cognitive slowing (e.g., in alcohol or benzodiazepine use). Executive function and judgment may be compromised.

## Psychosocial Assessment

A comprehensive psychosocial assessment includes the following components:

- Reason for seeking health care (i.e., “chief complaint”)
- Thoughts of suicide or self injury
- Cultural assessment
- Spiritual assessment
- Family dynamics
- Current and past medical history
- Current medications
- History of previously diagnosed mental health disorders
- Previous hospitalizations
- Educational background
- Occupational background
- History of exposure to psychological trauma, violence, and domestic abuse

- Substance use (tobacco, alcohol, recreational drugs, misused prescription drugs)
- Family history of mental illness
- Coping mechanisms
- Functional ability/Activities of daily living

After identifying the reason the client is seeking health care, additional focused questions are used to obtain detailed information. The mnemonic PQRSTU can be used to ask questions in an organized fashion. See Table 14.9b for a sample PQRST assessment for a client experiencing an alcohol use disorder.

Table 14.9b Sample PQRSTU Questions for Assessing Alcohol Use Disorder

PQRSTU	Sample Questions
<b>Provocation/ Palliation</b>	“What usually leads you to drink?” “What helps you cut back or stop, even temporarily?”
<b>Quality</b>	“How would you describe your drinking experience—what does it do for you?”
<b>Region</b>	“Do you notice any physical effects when you drink or stop drinking?”
<b>Severity</b>	“How much has drinking affected your health, work, or relationships?”
<b>Timing/ Treatment</b>	“When did you start drinking regularly?” “How has your drinking changed over time?”
<b>Understanding</b>	“Why do you think this problem developed?”  “What do you believe needs to happen to stop this problem?”

## SUICIDE AND SELF INJURY SCREENING

Clients being evaluated or treated for substance use disorders may have suicidal ideation. It is important for the nurse to introduce suicide screening in a way that helps the client understand its purpose and normalize questions that might otherwise seem intrusive. The Patient Safety Screener (PSS-3) is an

example of a brief screening tool to detect suicide risk in all client presenting to acute care settings.<sup>2</sup>

**Non-suicidal self-injury (NSSI)** refers to intentional self-inflicted destruction of body tissue without suicidal intention and for purposes not socially sanctioned. Common forms of NSSI include behaviors such as cutting, burning, scratching, and self-hitting. It is considered a maladaptive coping strategy without the desire to die. NSSI is a common finding among adolescents and young adults in psychiatric inpatient settings.<sup>3</sup>

Review information about suicide screening, the Patient Safety Screener, and screening for non-suicidal self-injury (NSSI) in the “[Assessment](#)” section of the Applying the Nursing Process to Mental Health Care” chapter.

## CULTURAL ASSESSMENT

Cultural Formulation Interview (CFI) questions help nurses understand a client’s cultural background and how it influences their experience of substance use disorder.<sup>4</sup> Sample CFI questions focused specifically on understanding alcohol use disorder within a cultural context include the following:

2. Suicide Prevention Resource Center. (n.d.). *The patient safety screener: A brief tool to detect suicide risk*. <https://sprc.org/micro-learning/patientsafetyscreener>
3. Cipriano, A., Cella, S., & Cotrufo, P. (2017). Nonsuicidal self-injury: A systematic review. *Frontiers in Psychology*, 8. <https://www.frontiersin.org/articles/10.3389/fpsyg.2017.01946/full>
4. DeSilva, R., Aggarwall, N. K., & Lewis-Fernandez, R. (2015). The DSM-5 cultural formulation interview and the evolution of cultural assessment in psychiatry. *Psychiatric Times*, 32(6). <https://www.psychiatrictimes.com/view/dsm-5-cultural-formulation-interview-and-evolution-cultural-assessment-psychiatry>

- Cultural Definition of the Problem
  - “How would you describe your drinking problem to someone close to you?”
  - “What terms or labels are used in by your friends or family to talk about heavy drinking?”
  - “Do people around you see your drinking as a problem?”
- Cultural Perceptions of Cause, Context, and Support
  - “Why do you think drinking became an issue for you?”
  - “How is alcohol used in your family or community? Is it used in celebrations, rituals, or as a coping tool?”
  - “Do you feel pressure—positive or negative—from your family or friends related to your drinking?”
- Cultural Factors Affecting Coping and Help-Seeking
  - “Have you tried anything on your own to manage your drinking? What helped and what didn’t?”
  - “Are there any traditional practices, herbs, or dietary supplements you have tried to use to manage it?”
  - “Do people in your community seek help for alcohol-related issues? If so, from whom (e.g., religious leaders, clinics, family)?”
  - “What kinds of treatment or help do you think would be most useful or acceptable to you?”
- Cultural Features of the Nurse–Client Relationship
  - “Are there any concerns you have about talking to a mental health professional?”
  - “Would you feel more comfortable speaking with someone of a similar gender, cultural, or religious background?”
  - “What would help you feel more supported or understood during treatment?”
  - “Have you had past experiences with healthcare or addiction treatment that were helpful or unhelpful?”

Findings from the cultural formulation interview are used to individualize a client’s plan of care to their preferences, values, beliefs, and goals.

## SPIRITUAL ASSESSMENT

The FICA Spiritual History Tool is a widely used assessment model for evaluating a client's spiritual beliefs and how they may influence health, illness, and coping. It's especially helpful in understanding how clients with substance use disorders draw on spirituality or religion for support—or how spiritual distress may be contributing to feelings of anxiety. Addressing a client's spirituality and advocating spiritual care have been shown to improve clients' health and quality of life.<sup>5,6</sup>

The FICA Spiritual History Tool© is a common tool used to gather information about a client's spiritual history and preferences. FICA© is a mnemonic for the domains of Faith, Importance, Community, and Address in Care.<sup>7</sup> Table 14.9c summarizes a sample FICA Spiritual Assessment for a client with a substance use disorder.

Table 14.9c Sample FICA Spiritual Assessment Questions for Clients with a Substance Use Disorder

5. Pilger, C., Molzahn, A. E., de Oliveira, M. P., & Kusumota, L. (2016). The relationship of the spiritual and religious dimensions with quality of life and health of patients with chronic kidney disease: An integrative literature review. *Nephrology Nursing Journal: Journal of the American Nephrology Nurses' Association*, 43(5), 411–426. <https://pubmed.ncbi.nlm.nih.gov/30550069/>
6. Puchalski, C., Jafari, N., Buller, H., Haythorn, T., Jacobs, C., & Ferrell, B. (2020). Interprofessional spiritual care education curriculum: A milestone toward the provision of spiritual care. *Journal of Palliative Medicine*, 23(6), 777–784. <https://doi.org/10.1089/jpm.2019.0375>
7. GW School of Medicine & Health Sciences. (n.d.). *Clinical FICA tool*. <https://smhs.gwu.edu/spirituality-health/program/transforming-practice-health-settings/clinical-fica-tool>



Domain	Sample Assessment Question	Sample Client Response
<b>Faith</b>	“Do you have spiritual beliefs or religious faith that help you cope with challenges or give your life meaning?”	“Yes, I believe in God and used to go to church, especially when I was trying to get sober. But I’ve drifted away because I feel ashamed about using.”
<b>Importance</b>	“What importance does your faith or belief have in your life? Has it influenced how you cope with stress or your anxiety?”	“They’re still important to me, but I don’t feel like I’m living life the way I should be. God will probably punish me for all I’ve done.”
<b>Community</b>	“Are you part of a spiritual or religious community? Does participation in this community provide support when you’re feeling anxious or stressed?”	“I haven’t gone back to church since I relapsed. I don’t want to be judged.”
<b>Address in Care</b>	“How would you like me (or the health care team) to address spiritual issues during your care? Would you like to speak with a chaplain?”	“Yes I would be interested in speaking to someone who will listen and not judge me.”

Nurses may recognize cues of spiritual distress or beliefs in divine punishment that may exacerbate feelings of anxiety. Nurses can offer to connect the client with a chaplain or spiritual care services. Spiritual goals may be included in the nursing care plan if the client finds them valuable.

## History

The following client history should be obtained from a client diagnosed with a substance use disorder<sup>8</sup>:

- **History of substance use:** Include names of the substances used, the time of last use, the duration of use, the quantity and frequency of use, and the method of use (e.g., oral, intravenous, inhaled, intranasal). A

8. Pace, C. (2022). Alcohol withdrawal: Epidemiology, clinical manifestations, course, assessment, and diagnosis. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

detailed substance use history helps determine the expected time frame for emergence of withdrawal symptoms and the potential for severe withdrawal syndromes. Family history of substance use.

- **Detailed history of previous withdrawal treatments & treatment history:** For clients who have previously undergone withdrawal management, a history of these treatments should be obtained. The history of any previous treatment programs should be obtained. History should include previous treatments (such as inpatient or outpatient programs, 12-step programs such as Alcoholics Anonymous, or medications such as naltrexone or acamprosate), as well as what treatments have been helpful or not helpful.
- **Mental health history:** Concurrent mental health illness can impact the client's withdrawal symptoms. Clients should receive integrated post-withdrawal treatments for multiple diagnoses of mental health and substance use disorders.
- **Social history:** Identification of social supports (such as a supportive family member who can encourage abstinence and potentially dispense withdrawal medication), as well as barriers (such as poor transportation), can also help determine the most appropriate post-withdrawal treatment plan (e.g., residential, outpatient, and recovery programs).
- **Medical history and recent physical symptoms:** Medical problems can contribute to the client's symptoms and/or worsen withdrawal symptoms.

The client should receive a comprehensive physical examination to evaluate signs related to current withdrawal symptoms, as well as symptoms of concurrent medical and mental health diagnoses. If the client has been diagnosed with alcohol use disorder, signs of complications such as liver or pancreatic disease should be assessed.<sup>9</sup>

9. Pace, C. (2022). Alcohol withdrawal: Epidemiology, clinical manifestations, course, assessment, and diagnosis. *UpToDate*. [www.uptodate.com](https://www.uptodate.com)

## Screening Scales

Screening scales may be administered such as CIWA-aR (for alcohol withdrawal symptoms) and COWS (for opioid withdrawal symptoms). Read more information about these scales in the “[Withdrawal Management/Detoxification](#)” section of this chapter.

## Diagnostic and Lab Work

Laboratory testing for clients admitted for withdrawal treatment may include these items<sup>10</sup>:

- Complete blood count
  - Serum electrolytes, including potassium, magnesium, and phosphate
  - Glucose
  - Creatinine
  - Liver function tests
  - Amylase and lipase
  - Blood alcohol level
  - Urine drug testing for benzodiazepines, cocaine, and opioids. (The opioid test may include testing for heroin, codeine, morphine, buprenorphine, oxycodone, methadone, and fentanyl.)
- 
- Urine human chorionic gonadotropin (HCG) test for premenopausal women to check for pregnancy
  - Electrocardiogram (ECG) for clients over 50 years of age or if there is a history of cardiac problems

10. Pace, C. (2022). Alcohol withdrawal: Epidemiology, clinical manifestations, course, assessment, and diagnosis. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

# Life Span Considerations

## Children and Adults

In 2020, nearly 10 million U.S. adolescents met the diagnostic criteria for a substance use disorder, and the majority were untreated. In the United States, more than 90% of adults with SUDs began their substance use in adolescence.<sup>11</sup> Adolescents with SUDs are at greater lifetime risk for sexually transmitted diseases, poor family planning, justice system involvement, school-related challenges, neurocognitive impairments, and increased risk of mental health disorders. Adolescent substance use is associated with the leading causes of death of unintentional injury, suicide, and violence. Onset of substance use disorders may begin during childhood, often with alcohol, cannabis, or vaping. Substance use is influenced by peer pressure, family dynamics, trauma, adverse childhood experiences, and emotional dysregulation. Early substance use increases risk for lifelong addiction due to ongoing brain development. Behavioral changes for children using substances may include irritability, truancy, declining grades, secretiveness, withdrawal from family. Risk-taking, mood swings, and impulsivity may be misattributed to “normal adolescence.” Coexisting disorders such as ADHD, anxiety, or depression should be considered. Adolescents need developmentally appropriate, preventive, and early intervention-focused strategies that engage their families and social systems.<sup>12</sup>

11. Simon, K. M., Levy, S. J., & Bukstein, O. G. (2022). Adolescent substance use disorders. *NEJM evidence*, 1(6), EVIDra2200051. <https://doi.org/10.1056/EVIDra2200051>

12. Simon, K. M., Levy, S. J., & Bukstein, O. G. (2022). Adolescent substance use disorders. *NEJM evidence*, 1(6), EVIDra2200051. <https://doi.org/10.1056/EVIDra2200051>

## Older Adults

Onset of substance use disorder in older adults may occur due to late-life stressors such as grief, retirement, isolation, or chronic illness. Prescription drug misuse (e.g., opioids, benzodiazepines) is common. Substance use may present with falls, memory loss, confusion, sleep disturbances, or social withdrawal that may be mistaken for dementia symptoms. Older adults require holistic care that integrates mental health, physical health, and social support.<sup>13</sup>

## Diagnosis (Analyzing Cues)

A nursing diagnosis related to the abrupt cessation of a psychoactive substance is *Acute Substance Withdrawal Syndrome*. As a syndrome diagnosis, defining characteristics are the related nursing diagnoses, including *Acute Confusion*, *Anxiety*, *Disturbed Sleep Pattern*, *Nausea*, *Risk for Electrolyte Imbalance*, and *Risk for Injury*.<sup>14</sup>

- ▶ Review information about syndrome nursing diagnosis in the “[Diagnosis](#)” section of the “Nursing Process” chapter of *Open RN Nursing Fundamentals*.

13. Substance Abuse and Mental Health Services Administration. (2019). Enhancing motivation for change in substance use disorder treatment. *Treatment Improvement Protocol (TIP) Series, No. 35: Chapter 3—Motivational Interviewing as a Counseling Style*. <https://www.ncbi.nlm.nih.gov/books/NBK571068/>
14. Ackley, B., Ladwig, G., Makic, M. B., Martinez-Kratz, M., & Zanotti, M. (2020). *Nursing diagnosis handbook: An evidence-based guide to planning care* (12th ed.). Elsevier

## Outcome Identification (Generate Solutions)

An example of a broad goal related to withdrawal treatment is the following:

- The client will stabilize and remain free from injury.<sup>15</sup>

These are some sample SMART outcomes<sup>16</sup> :

- The client's vital signs will remain within normal ranges during treatment.
- The client's electrolyte levels will remain within normal ranges during treatment.
- The client will participate in planning a post-withdrawal treatment program before discharge.

## Planning (Generate Solutions)

### Safety

Safety receives top priority when planning and implementing interventions for clients with substance use disorder who are at risk of suicide, and interventions are planned according to their level of risk. Review interventions for clients at risk of suicide in the "[Application of the Nursing Process in Mental Health Care](#)" chapter.

15. Ackley, B., Ladwig, G., Makic, M. B., Martinez-Kratz, M., & Zanotti, M. (2020). *Nursing diagnosis handbook: An evidence-based guide to planning care* (12th ed.). Elsevier

16. Ackley, B., Ladwig, G., Makic, M. B., Martinez-Kratz, M., & Zanotti, M. (2020). *Nursing diagnosis handbook: An evidence-based guide to planning care* (12th ed.). Elsevier

## Implementation (Take Action)

### Nursing Interventions for Substance Use Disorder Based on Categories of the APNA Implementation Standard

Nursing interventions for substance use disorders can be categorized based on the American Psychiatric Nurses Association (APNA) standard for *Implementation* that includes the *Coordination of Care; Health Teaching and Health Promotion; Pharmacological, Biological, and Integrative Therapies; Milieu Therapy; and Therapeutic Relationship and Counseling*. Read more about these subcategories in the “[Application of the Nursing Process in Mental Health Care](#)” chapter. See examples of interventions for each of these categories for clients with anxiety disorders in Table 14.9.

Table 14.9d Examples of Nursing Interventions for Substance Use Disorders by APNA Subcategories<sup>17, 18</sup>

17. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.
18. National Institute on Drug Abuse. (2023). *Treatment*. <https://nida.nih.gov/research-topics/treatment>

Subcategory of the APNA Standard of Implementation	The nurse will ...	Rationale
<b>Coordination of Care</b>	<ul style="list-style-type: none"> <li>• Collaborate with a multidisciplinary team (physician, addiction counselor, social worker, pharmacist) to ensure integrated care.</li> <li>• Facilitate referrals to detoxification, rehabilitation, or community recovery programs.</li> </ul>	Effective treatment of SUDs requires a collaborative and coordinated approach that addresses medical, psychological, and social needs.
<b>Health Teaching and Health Promotion</b>	<ul style="list-style-type: none"> <li>• Educate the client on the effects of substance use, including its impact on physical health, mental health, and relationships.</li> <li>• Use motivational interviewing to promote behavior change and harm reduction.</li> </ul>	Health education and motivational techniques improve insight, self-efficacy, and readiness for change while reducing relapse risk.



<b>Pharmacological, Biological, and Integrative Therapies</b>	<ul style="list-style-type: none"> <li>• Administer medications as prescribed (e.g., naltrexone, buprenorphine, acamprosate).</li> <li>• Monitor for withdrawal symptoms, side effects, and adherence.</li> <li>• Incorporate integrative methods such as mindfulness or guided imagery to manage cravings and anxiety.</li> </ul>	<p>Medications can support recovery by reducing cravings, preventing relapse, and managing co-occurring conditions. Monitoring ensures safety and promotes therapeutic response.</p>
<b>Milieu Therapy</b>	<ul style="list-style-type: none"> <li>• Establish a safe, structured, and substance-free environment.</li> <li>• Promote positive peer interaction through group therapy, relapse prevention groups, or 12-step meetings.</li> <li>• Set clear expectations and boundaries.</li> </ul>	<p>A therapeutic milieu reduces triggers and access to substances, supports accountability, and promotes recovery-focused behavior.</p>

<b>Therapeutic Relationship and Counseling</b>	<ul style="list-style-type: none"><li>• Build a nonjudgmental, empathetic relationship based on trust.</li><li>• Support client-led goals.</li><li>• Explore underlying emotional pain, trauma, or mental illness contributing to substance use.</li></ul>	A strong therapeutic alliance improves treatment engagement, reduces shame, and facilitates the exploration of root causes and maladaptive coping.
--	--	--

## Nursing Interventions for Physiological Signs of Substance Use Disorder

Nursing interventions also target common physiological signs of substance use disorder and associated self-care deficits. See common interventions for these conditions in Table 14.9e.

Table 14.9e Nursing Interventions Targeting Physiological Signs of Substance Use Disorder and Self-Care Deficit<sup>19, 20</sup>

19. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

20. National Institute on Drug Abuse. (2023). *Treatment*. <https://nida.nih.gov/research-topics/treatment>

Problem/Intervention	Rationale
<p><b>Withdrawal</b></p> <ul style="list-style-type: none"> <li>• Monitor for withdrawal symptoms using tools such as CIWA-Ar (for alcohol) or COWS (for opioids).</li> <li>• Administer medications (e.g., benzodiazepines, methadone, clonidine) as prescribed.</li> <li>• Provide quiet, low-stimulus environment.</li> </ul>	<p>Withdrawal from substances like alcohol or opioids can lead to autonomic instability. Early detection and management can prevent complications and mortality.</p>
<p><b>Nutrition and Hydration</b></p> <ul style="list-style-type: none"> <li>• Monitor weight and BMI.</li> <li>• Assess fluid status (I&amp;O, skin turgor, mucous membranes).</li> <li>• Monitor electrolytes and administer replacement therapy as prescribed (e.g., magnesium, potassium).</li> <li>• Encourage oral fluids unless contraindicated.</li> <li>• Provide nutrient-dense meals and snacks.</li> <li>• Administer vitamins (e.g., thiamine, folic acid, multivitamins) as prescribed.</li> <li>• Consult dietitian.</li> </ul>	<p>Poor dietary intake, gastrointestinal issues, and dehydration are common in withdrawal. Replenishing nutrition helps restore energy, improve mental clarity. Many individuals with SUD experience electrolyte imbalance, especially during detox. Rehydration can prevent complications like arrhythmias or Wernicke's encephalopathy.</p>
<p><b>Sleep</b></p> <ul style="list-style-type: none"> <li>• Promote sleep hygiene: reduce noise, dim lights, avoid caffeine.</li> <li>• Assess for night sweats, nightmares, or insomnia.</li> <li>• Teach relaxation strategies (e.g., guided breathing, music therapy).</li> </ul>	<p>Insomnia is common in withdrawal and early relapse, especially in stimulant and alcohol users. Rest supports emotional and physical recovery.</p>

<p><b>Pain</b></p> <ul style="list-style-type: none"> <li>• Use non-opioid pain relief (e.g., acetaminophen, hot packs, distraction).</li> <li>• Assess for hyperalgesia in opioid users.</li> <li>• Collaborate on safe pain management strategies for clients with SUD and coexisting chronic pain.</li> </ul>	<p>Clients with SUD may have altered pain sensitivity due to previous history of poor pain management. Addressing pain promotes trust and recovery adherence.</p>
<p><b>Elimination</b></p> <ul style="list-style-type: none"> <li>• Monitor for nausea, vomiting, diarrhea, or constipation.</li> <li>• Provide antiemetics, fiber supplements, or stool softeners as needed.</li> <li>• Encourage small, frequent meals.</li> </ul>	<p>GI symptoms are common during detox and withdrawal. Symptom relief increases treatment tolerability.</p>
<p><b>Respiratory Status</b></p> <ul style="list-style-type: none"> <li>• Assess for respiratory depression, especially with use of opioids or benzodiazepines.</li> <li>• Monitor oxygen saturation and respiratory rate.</li> <li>• Have naloxone available per protocol.</li> </ul>	<p>Overdose and sedation are risks with for clients with substance use disorders and nervous system depressants. Vigilant respiratory monitoring prevents hypoxia or death.</p>
<p><b>Self-Care Deficits</b></p> <ul style="list-style-type: none"> <li>• Assess ability and motivation to perform hygiene, grooming, and ADLs.</li> <li>• Provide structured daily routines and support gradual reengagement with hygiene tasks.</li> <li>• Use positive reinforcement to promote independence.</li> </ul>	<p>Individuals with SUD may neglect self-care due to substance use and environmental instability. Supporting ADLs enhances recovery, and reintegration into daily life.</p>

# Communication Tips for Clients With Substance Use Disorders

Communicating effectively with someone experiencing a substance use disorder must foster feelings of trust, safety, and therapeutic alliance. People in states of heightened anxiety may struggle to process information, express themselves clearly, or feel emotionally overwhelmed. The following box provides communication tips when speaking with clients with severe anxiety.

## Communication Tips for Clients With A Substance Use Disorder<sup>21</sup>

- Use nonjudgmental, person-first language. For example, say “a person with a substance use disorder” instead of “addict” or “alcoholic.”
  - **Rationale:** Person-first language reduces stigma and supports the client’s sense of identity beyond the diagnosis.
- Tailor communication to the client’s stage in the readiness for change model.
  - **Rationale:** Assess where the client is in the **Stages of Change Model (e.g.,**

21. Substance Abuse and Mental Health Services Administration. (2019). Enhancing motivation for change in substance use disorder treatment. *Treatment Improvement Protocol (TIP) Series, No. 35: Chapter 3—Motivational Interviewing as a Counseling Style*. <https://www.ncbi.nlm.nih.gov/books/NBK571068/>

**precontemplation, contemplation)** and adjust your message accordingly.

- Explore rather than confront denial. For example, instead of saying, “You’re in denial about drinking too much,” ask, “What do you think others are concerned about?” or “What might happen if you continued drinking at this level?”
  - **Rationale:** Using exploratory language maintains rapport while encouraging insight into their behavior.
- Avoid power struggles or ultimatums.
  - **Rationale:** Framing treatment as a collaborative process rather than a directive one encourages cooperation and empowerment, which improves treatment adherence.
- Reflect on their strengths and past successes. For example, highlight what the client has done well or survived in the past (e.g., “You’ve taken steps to come here— that takes courage”)..
  - **Rationale:** Reflecting on strengths and past successes builds self-efficacy, resilience, and motivation.
- Be consistent and set clear, compassionate boundaries.
  - **Rationale:** Setting clear expectations around behavior and attendance help the client feel secure and respected, especially in early recovery.

- Normalize relapse as part of recovery, if it occurs. For example say “Relapse doesn’t mean failure. Let’s figure out what happened and move forward.”.
  - **Rationale:** Relapse is common but doesn’t necessarily mean treatment failure. Normalizing relapse reduces shame and hopelessness.

## Withdrawal Treatment

Nursing interventions provide a supportive environment while the client undergoes withdrawal treatment. Vital signs are monitored closely because increases in temperature, pulse, and blood pressure are signs of withdrawal. After ensuring that an individual’s physiological needs of airway, breathing, and circulation are met, safety measures receive top priority. Safety measures during withdrawal treatment may include interventions such as fall precautions, seizure precautions, or implementing restraints as needed to maintain the safety of the individual or those around them.<sup>22</sup>

- ▶ Review information on fall precautions in the “[Preventing Falls](#)” section of the “Safety” chapter in *Open RN Nursing Fundamentals, 2e*.

Seizures can occur in clients experiencing alcohol withdrawal. Seizure

22. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

precautions include keeping the bed in the lowest position with side rails padded. Suction and oxygen equipment must be available at all times at the client's bedside. Review ANA guidelines on using restraints in the "[Client Rights](#)" section of the "Legal and Ethical Considerations in Mental Health Care" chapter and information on safely implementing restraints in the "[Workplace Violence](#)" section of the "Trauma, Abuse, and Violence" chapter.

Medications are administered as prescribed to keep the client safe and comfortable, so they do not suffer.<sup>23</sup> Review medications used during withdrawal treatment in the "[Withdrawal Management/Detoxification](#)" section of this chapter.

Clients with substance use disorders may exhibit a poor nutritional status due to long-term use of substances taking precedence over food and fluid intake. Nurses provide hydration and gradually reintroduce healthy foods while also promoting rest. Clients with alcohol use disorder are specifically at risk for thiamine (B1) and magnesium deficiencies that can lead to cardiac arrest. Thiamine and other electrolyte replacement is typically included during withdrawal treatment.<sup>24</sup>

Developing a therapeutic nurse-client relationship can encourage the client to explore harmful feelings of anxiety, hopelessness, and spiritual distress. Encouraging self-care and hygiene helps improve clients' self-esteem.

Nurses educate clients about healthy coping skills and evidence-based treatment and recovery services available in the community.<sup>25</sup> Client

23. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.
24. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.
25. Halter, M. (2022). *Varcarolis' foundations of psychiatric-mental health nursing* (9th ed.). Saunders.



education includes understanding the neurobiology behind substance use disorders and the impact on behavior, avoiding triggers, managing cravings, and early warning signs and when to seek help. Family member education includes how to provide emotional support without enabling behavior, family therapy, the importance of boundaries and managing caregiver stress. Review evidence-based treatments and recovery services in the “[Treatment of Substance Use Disorders](#)” section of this chapter.

In addition to implementing the withdrawal treatment plan prescribed by the provider, the nurse collaboratively develops a post-withdrawal treatment plan with interprofessional health care team members. The plan should be client-centered and include their goals and readiness for change. Motivational interviewing is a helpful therapeutic technique when planning individualized treatment goals and programs.<sup>26</sup>

## Evaluation (Evaluate Outcomes)

Evaluation occurs on several levels by assessing the individualized SMART outcomes related to the effectiveness of the withdrawal treatment plan, symptom management, and the client’s readiness and progress towards changes in their behavior.<sup>27</sup>

Examples of potential evaluation outcomes include:

- Monitor reduction in substance use, improvement in physical and mental health
- Increased engagement in therapy, improved social functioning, employment
- Modify based on client’s progress, address new issues (e.g., relapse, co-

26. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

27. Halter, M. (2022). *Varcarolis’ foundations of psychiatric-mental health nursing* (9th ed.). Saunders.

occurring disorders)

- Reinforce coping mechanisms, provide ongoing support

## 14.10 Spotlight Application

- ▶ This case study is based on a sample client described in the [“Assessment & Treatment of Substance Use Disorders”](#) chapter of the *Foundations of Addiction Studies* book.

Jessica is a 26-year-old woman who recently received her second driving-under-the-influence (DUI) ticket and has been charged with a misdemeanor crime. She received her first DUI at age 21 and lost her license for one year. During that time, she completed a basic risk education DUI course and paid several thousand dollars in fines and attorney's fees.

Jessica continues to drink with her friends on weekends, usually having five or six drinks per night. In addition to alcohol, Jessica was recently prescribed Xanax to treat her anxiety disorder. Jessica often takes more than prescribed in the evening and sometimes mixes these pills with her alcohol consumption.

Jessica works from home full-time as a graphic designer. She says that she has little time to socialize during the week, so she looks forward to the weekend when she can see her friends and relax. Now that she has a second DUI, Jessica has lost her license again, although she is not overly concerned because she can continue working from home. She can walk or order a ride wherever she needs to go for groceries and other errands.

The court has ordered Jessica to have an evaluation done and to complete any treatment recommendations.

### Reflective Questions:

1. What were your initial feelings when you read this case study?

2. Can you identify any personal biases or assumptions you experienced while reading this scenario?
3. How many *DSM-5* criteria is Jessica displaying for a substance use disorder diagnosis?
4. List at least three issues to include in a treatment plan for Jessica.
5. How do Jessica's legal issues potentially impact her treatment plan?
6. How will you apply your new knowledge about substance use disorders to your nursing practice?

## 14.11 Learning Activities



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=875#h5p-47>

1



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=875#h5p-48>

2



*An interactive H5P element has been excluded from this version of the text. You can view it*

1. “MH Substance Use Disorders Glossary Cards” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
2. “MH Substance Use Disorders Drag and Drop” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

online here:

<https://wtcs.pressbooks.pub/nursingmhcc/?p=875#h5p-49>

3

- ▶ Test your clinical judgment with an NCLEX Next Generation-style question: [Chapter 14, Assignment 1](#)<sup>4</sup>



- ▶ Test your clinical judgment with an NCLEX Next Generation-style question: [Chapter 14, Assignment 2](#)<sup>5</sup>



3. "MH Substance Use Disorders Question Set 1" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
4. "MH Substance Use Disorders Next Gen Question 1" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
5. "MH Substance Use Disorders Next Gen Question 2" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

- ▶ Test your clinical judgment with an NCLEX Next Generation-style case study: [Chapter 14, Case Study 1](#)<sup>6</sup>



6. “MH Substance Use Disorders Next Gen Case Study” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

## XIV Glossary

**Addiction:** Severe substance use disorders are commonly referred to as addiction. Addiction is a chronic brain disease that has the potential for both recurrence (relapse) and recovery.<sup>1</sup>

**Alcohol intoxication:** Refers to problematic behavioral or psychological changes (e.g., inappropriate sexual or aggressive behavior, mood lability, or impaired judgment) that develop during or shortly thereafter alcohol ingestion

**Binge drinking:** Consuming several standard drinks on one occasion in the past 30 days, defined for males as drinking five or more standard alcoholic drinks on one occasion, and for females as drinking four or more standard drinks on one occasion.<sup>2</sup>

**Brief interventions:** Therapeutic techniques used to initiate change in individuals with unhealthy behaviors.

**Cannabis intoxication:** Problematic behavior or psychological changes (e.g., impaired motor coordination, euphoria, anxiety, sensation of slowed time, impaired judgment, social withdrawal) that developed during or shortly after cannabis use.

**Controlled substances:** Substances regulated by the U.S. Drug Enforcement Agency into one of five categories called schedules. This placement is based

1. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>
2. National Survey on Drug Use and Health. (2025). *Welcome to the national survey on drug use and health (NSDUH)*. <https://nsduhweb.rti.org/respweb/homepage.cfm>



on the substance's medical use, its potential for abuse or dependency, and related safety issues.

**Delirium tremens (DTs):** A rapid-onset, fluctuating disturbance of attention and cognition, sometimes associated with hallucinations, due to alcohol withdrawal. In its most severe manifestation, DTs are accompanied by agitation and signs of extreme autonomic hyperactivity, including fever, severe tachycardia, hypertension, and drenching sweats. DTs typically begin between 72 and 96 hours after the client's last drink.

**Dependence:** A condition that develops with chronic uses of a substance that causes a person to experience withdrawal symptoms if the substance is suddenly stopped.

**Drug diversion:** The act of redirecting medication from its intended destination for personal use, sale, or distribution to others. It includes drug theft, use, or tampering (adulteration or substitution). Drug diversion is a felony that can result in a nurse's criminal prosecution and loss of license.<sup>3</sup>

**Heavy drinking:** Excessive drinking defined as a female consuming 8 or more drinks per week and a male consuming 15 or more standard drinks per week, or either gender binge drinking on 5 or more days in the past 30 days.

**Intoxication:** A disturbance in behavior or mental function during or after the consumption of a substance.

**Neuroadaptations:** Progressive changes in the structure and function of the brain as a result of substance misuse. These neuroadaptations compromise brain function and also drive the transition from controlled, occasional substance use to chronic misuse that can be difficult to control.

**Nonsubstance related disorders:** Excessive behaviors such as gambling, viewing pornography, engaging in compulsive sexual activity, Internet

3. Nyhus, J. (2021). Drug diversion in healthcare. *American Nurse*.  
<https://www.myamericannurse.com/drug-diversion-in-healthcare/>

gaming, overeating, shopping, overexercising, and overusing mobile phone technologies. These behaviors are thought to stimulate the same addiction centers of the brain as addictive substances. However, gambling disorder is the only nonsubstance use disorder with diagnostic criteria listed in the *DSM-5*.

**Opioid intoxication:** A condition where a person is affected by opioid drugs to the point of being impaired and potentially ill.

**Protective factors:** Factors that help prevent substance use disorders from developing despite the risk factors that are present.

**Recovery:** A process of change through which individuals improve their health and wellness, live a self-directed life, and strive to reach their full potential. Although abstinence from all substance misuse is a primary feature of a recovery lifestyle, it is not the only healthy feature.<sup>4</sup>

**Relapse:** The return to substance use after a significant period of abstinence.

**Remission:** The status of individuals with severe substance use disorders who overcome their disorder with effective treatment and regain health and social function.

**Standard drink:** 14 grams (0.6 ounces) of pure alcohol. Examples of a standard drink are one 12-ounce beer, 8 – 9 ounces of malt liquor, 5 ounces of wine, or 1.5 ounces distilled spirits.

**Substance:** A psychoactive compound with the potential to cause health and social problems, including substance use disorders. Substances can be divided into four major categories: alcohol, illicit drugs (including nonmedical use of prescription drugs), tobacco products, and over-the-counter drugs.

4. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>

**Substance misuse:** The use of alcohol or drugs in a manner, situation, amount, or frequency that could cause harm to the user or to those around them.<sup>5</sup>

**Substance use:** The use (even one time) of any psychoactive substance.

**Substance use disorder (SUD):** A chronic illness caused by repeated misuse of psychoactive substances (e.g., alcohol, caffeine, cannabis, hallucinogens, inhalants, opioids, sedatives, hypnotics, anxiolytics, stimulants, and tobacco) that have a common effect of directly activating the brain's reward system.

**Tolerance:** A need for progressively increased amounts of a substance to achieve the desired effect or a diminished effect with continued use of the same amount of a substance.

**Withdrawal:** A group of physical and mental symptoms that can range from mild to life-threatening when a person suddenly stops using a substance.

**Withdrawal management:** Interventions that manage the physical and emotional symptoms that occur after a person stops using a psychoactive substance, also referred to as detoxification. Withdrawal symptoms vary in intensity and duration based on the substance(s) used, the duration and amount of use, and the overall health of the individual.

5. Substance Abuse and Mental Health Services Administration, & Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. United States Department of Health and Human Services. <https://www.ncbi.nlm.nih.gov/books/NBK424857/>







### Learning Objectives

- Identify assessment cues of clients and families experiencing trauma, abuse, and/or violence
- Identify assessment cues of adverse childhood experiences in a client's history
- Identify factors promoting the cycle of trauma, abuse, and/or violence
- Identify nursing priorities for clients and families experiencing trauma, abuse, and/or violence
- Plan outcomes for clients and families experiencing trauma abuse and/or violence
- Differentiate safety/protective interventions for clients and families experiencing trauma, abuse, and/or violence
- Apply evidence-based practice when planning care and interventions for clients and families experiencing trauma, abuse, and/or violence
- Analyze treatments for clients and families experiencing trauma, abuse, and/or violence
- Apply the nursing process to clients and families experiencing trauma, abuse, and/or violence who are at risk for suicide
- Incorporate trauma-informed care
- Describe strategies to remain safe if workplace violence occurs

The health care system is composed of people who have experienced trauma,

both those providing and those receiving care. Supporters of a trauma-informed care approach recognize the prevalence of trauma survivors within health care settings and are aware that the service setting can also be a source of re-traumatization. As stated in the article *Trauma-Informed Nursing Practice*, understanding how trauma has affected clients' lives and their interactions within the health care system is fundamental to responding to clients' needs and promotes better physical and mental health outcomes.<sup>1</sup>

Nurses provide care for clients who are experiencing or have experienced neglect, abuse, and intimate partner violence. In many settings, nurses may experience workplace violence while caring for clients who are agitated or combative. Because these issues are so prevalent, nurses must be knowledgeable of these issues and how to care for afflicted clients. This chapter will discuss adverse childhood experiences and trauma-informed care, abuse and neglect of children and vulnerable adults, intimate partner violence, and workplace violence. Be aware that the content in this chapter may trigger powerful emotions, especially for survivors of similar traumatic experiences. Self-awareness and self-care practices should guide your engagement with this chapter.

- ▶ Read the article “[Trauma-Informed Nursing Practice](#)” in the *Online Journal of Issues in Nursing* published by the American Nurses Association.

1. Fleishman, J., Kamsky, H., & Sundborg, S. (2019). Trauma-informed nursing practice. *OJIN: The Online Journal of Issues in Nursing*, 24(2). <https://doi.org/10.3912/OJIN.Vol24No02Man03>



## 15.2 Adverse Childhood Experiences

**Adverse Childhood Experiences (ACEs)** refer to traumatic events or circumstances during childhood—including neglect, abuse, witnessing violence, or having a family member with substance abuse, mental illness, divorce, or imprisonment. In the late 1990s, the landmark Adverse Childhood Experiences Study examined over 17,000 participants to assess the long-term impact of these experiences on adult health and behavior. The study revealed that as the number of ACEs increased, children were more likely to develop behavioral problems and engage in risky behaviors during adolescence, such as substance abuse and unprotected sexual activity that could lead to pregnancy. These early-life challenges often set the stage for further health complications later in life. In adulthood, individuals with multiple ACEs were more likely to struggle with substance abuse and suffer from chronic illnesses—including alcoholism, chronic pulmonary disease, depression, and liver disease. Remarkably, those with the highest levels of ACEs had a life expectancy that was, on average, 20 years shorter than that of individuals with few or no ACEs<sup>1</sup> See Figure 15.1<sup>2</sup> for an infographic of ACEs.

1. [Action steps using ACEs and trauma-informed care: A resilience model](#) by Laurie Leitch is licensed under [CC BY 4.0](#)

2. “ACEs.png” by unknown author for [Centers for Disease Control and Prevention](#) is licensed in the [Public Domain](#). Access for free at [https://www.cdc.gov/injury/pdfs/priority/ACEs-Strategic-Plan\\_Final\\_508.pdf](https://www.cdc.gov/injury/pdfs/priority/ACEs-Strategic-Plan_Final_508.pdf)

## Figure 1. What are Adverse Childhood Experiences?





Figure 15.1 Adverse Childhood Events (ACEs)

View the following YouTube video<sup>3</sup> on adverse childhood experiences (ACEs), resilience, and trauma-informed care:  
 [How childhood trauma affects health across a lifetime | Nadine Burke Harris.](https://youtu.be/95ovIJ3dsNk)

A systematic review and meta analysis in 2024 across demographic characteristics and contexts found that ACEs in children younger than 18 years of age was common, with prevalence of 42% of children having no ACEs, 22% having 1 ACE, 13% having 2 ACEs, 8% having 3 ACEs, and 15% having 4 or

3. TED. (2015, February 17). *How childhood trauma affects health across a lifetime | Nadine Burke Harris*. [Video]. YouTube. All rights reserved.  
<https://youtu.be/95ovIJ3dsNk>

more ACEs. The prevalence of 4 or more ACEs was higher among children in residential care, with a history of juvenile offending, and in Indigenous peoples.<sup>4</sup>

View a supplementary YouTube video<sup>5</sup>

While being sensitive to a client's history of trauma and helping support them throughout their life is important, we know that

4. Madigan, S., Thiemann, R., Deneault, A., et al. (2025). Prevalence of adverse childhood experiences in child population samples: A systematic review and meta-analysis. *JAMA Pediatrics*, 179(1), 19–33. [doi:10.1001/jamapediatrics.2024.4385](https://doi.org/10.1001/jamapediatrics.2024.4385)

5. Psych Hub. (2021, November 19). *What is trauma?* [Video]. YouTube. All rights reserved. <https://www.youtube.com/watch?v=q0UPnWfNpak>[/footnote] explaining trauma: [What is Trauma?](#)

For some individuals, traumatic experiences during childhood can cause long term psychological and physical problems due to its impact on the child's developing brain. Various biopsychosocial and cultural factors influence an both immediate and long-term responses to trauma. In most cases, regardless of severity, individuals display resilience—the capacity to overcome adversity and face challenges with determination. This resilience involves mobilizing available resources to navigate hardships and mitigate the impact of adverse events.<sup>6</sup>

Center for Substance Abuse Treatment (US). (2014). *Trauma-informed care in behavioral health services*. Substance Abuse and Mental Health Services Administration.

<https://www.ncbi.nlm.nih.gov/books/NBK207201/>

these traumatic experiences are often preventable. Nurses throughout our communities can advocate for improvements, to help promote building positive social relationships and reduce the exposure to violence in children. The Centers for Disease Control and Prevention (CDC) created a resource titled *Preventing Adverse Childhood Experiences (ACEs): Leveraging the Best Available Evidence*. This resource contains these six strategies:

- Strengthen economic supports to families
- Promote
  - social norms that protect against violence and adversity
- Ensure a strong start for children
- Teach skills
- Connect youth to caring adults and activities
- Intervene to lessen immediate and long-term harms

▶ Read about the CDC's prevention strategies in [Adverse Childhood Experiences Prevention: Resource for Action PDF](#).

▶ View the following CDC YouTube video on adverse



childhood experiences and protective factors<sup>7</sup>: We Can Prevent ACEs.

## Risk Factors, Protective Factors, and Prevention Strategies

By bringing attention to the powerful impact that negative childhood experiences have on future health and functioning, the ACE study demonstrates the importance of designing early intervention programs that target abuse, neglect, and violence.<sup>8</sup> Nurses can help prevent ACEs by educating parents, communities, and policymakers about how to help families provide safe and stable environments for children.

Many factors contribute to ACEs, including personal traits and experiences, parents, the family environment, and the community itself. ACEs can have lasting negative effects on health and well-being, but these harms can be preventable. To prevent ACEs and protect children from neglect, abuse, and

7. Centers for Disease Control and Prevention (CDC). (2018, April 5). *We can prevent ACEs*. [Video]. YouTube. All rights reserved. <https://youtu.be/8gm-INpzU4g>

8. Center for Substance Abuse Treatment (US). (2014). *Trauma-informed care in behavioral health services*. Substance Abuse and Mental Health Services Administration. <https://www.ncbi.nlm.nih.gov/books/NBK207201/>

violence, nurses are involved in addressing these risk factors and protective factors with prevention strategies.<sup>9</sup>

Appropriately addressing traumatic experiences and promoting resilience are important parts of effective mental health care and integral for the healing and recovery process.<sup>10</sup>

## Risk Factors

Risk factors are conditions that increase the likelihood of experiencing ACEs. There are individual, family, and community risk factors for ACEs.

### INDIVIDUAL AND FAMILY RISK FACTORS<sup>11</sup>

Individual and family risk factors include the following<sup>11</sup>:

Individual:

- Children and adolescents who don't feel close to their parents/caregivers or feel they can't talk to them about their feelings
- Youth who start dating or engaging in sexual activity at an

9. Centers for Disease Control and Prevention. (2024). *Risk and protective factors*. <https://www.cdc.gov/aces/risk-factors/index.html>

10. Substance Abuse and Mental Health Services Administration. (2022). *Trauma and violence*. <https://www.samhsa.gov/trauma-violence>

11. Centers for Disease Control and Prevention. (2024). *Risk and protective factors*. <https://www.cdc.gov/aces/risk-factors/index.html>

early age

- Children and youth with few or no friends or with friends who engage in aggressive or delinquent behavior

Families experiencing:

- Caregiving challenges related to children with special needs (e.g., disabilities, mental health issues, chronic illnesses)
- High levels of stress from parenting or economic issues
- Isolation from extended family, friends, or neighbors

Families with caregivers who:

- Have a limited understanding of children's needs or development
- Were abused or neglected as children
- Are young or single parents

Families with:

- Low income or low levels of adult education
- High conflict and negative communication styles
- Attitudes accepting of or justifying violence or aggression
- Inconsistent discipline, corporal punishment, and/or low parental supervision

## COMMUNITY RISK FACTORS

Communities with <sup>12</sup>:

12. Centers for Disease Control and Prevention. (2024). *Risk and protective factors*. <https://www.cdc.gov/aces/risk-factors/index.html>

- High rates of violence and crime
- Limited educational and economic opportunities
- High rates of poverty, unemployment rates, and food insecurity
- Easy access to drugs and alcohol
- Low community involvement among residents
- Few community activities for youth
- Unstable housing where residents move frequently

## Protective Factors

Protective factors help reduce the likelihood of ACEs and support resilience.

### INDIVIDUAL AND FAMILY PROTECTIVE FACTORS

Individual and family protective factors against ACEs include the following <sup>13</sup>:

Children who:

- Have positive friendships and peer networks
- Experience academic success
- Have caring adults outside the family as mentors/role models

Families where caregivers:

- Can meet children's basic needs (food, shelter, health

<sup>13</sup>. Centers for Disease Control and Prevention. (2024). *Risk and protective factors*. <https://www.cdc.gov/aces/risk-factors/index.html>



services)

- Create a stable, supportive home with nurturing relationships
- Engage in positive parenting with consistent rules and supervision
- Help children work through problems and resolve conflicts peacefully
- Encourage the importance of school and academic success
- Have strong social support networks and positive relationships

Families with:

- Caregivers who have higher education or steady employment
- Fun, positive activities that strengthen family bonds

## COMMUNITY PROTECTIVE FACTORS

Communities can intervene to lessen the harms from ACEs and prevent future risks.

Communities where<sup>14 15</sup>:

- Families have access to education and support regarding

14. Centers for Disease Control and Prevention. (2024). *Risk and protective factors*. <https://www.cdc.gov/aces/risk-factors/index.html>

15. Centers for Disease Control and Prevention (CDC). (2018, April 5). *We can prevent ACEs*. [Video]. YouTube. All rights reserved. <https://youtu.be/8gm-INpzU4g>

positive parenting

- Families have access to economic and financial help
- Families have access to medical care and mental health services
- Individuals and families have access to safe, stable housing
- Families have access to nurturing and safe child care
- Families have access to high-quality preschool programs
- Families have access to safe, engaging after-school programs and activities
- Adults have work opportunities with family-friendly policies
- There are strong partnerships among the community and business, health care, government, and other sectors
- Residents feel connected to each other and are involved in the community
- Violence is not tolerated or accepted in the community
- Positive parenting strategies are social norms in the community

## 15.3 Trauma-Informed Care

Trauma has no age, gender, socioeconomic status, race, ethnicity, or sexual orientation boundaries. **Individual trauma** results from an event, series of events, or set of circumstances that is experienced by an individual as physically or emotionally harmful or life-threatening and can have lasting adverse effects on their functioning and mental, physical, social, emotional, or spiritual well-being. Adverse childhood experiences (ACEs) are examples of individual traumas.<sup>1</sup>

The body's physiological stress response, known as “fight, flight, freeze, or fawn” is an automatic survival mechanism triggered by the sympathetic nervous system (SNS) when a threat is perceived. This response activates multiple organs, leading to symptoms such as increased heart rate, elevated blood pressure, rapid breathing, and muscle tension.<sup>2</sup>

- ▶ Review the physiology of the sympathetic nervous system and the parasympathetic nervous system in the “[Psychotropic Medications](#)” chapter.
- ▶ Review activation of the stress response in the “[Stress, Coping, and Crisis Intervention](#)” chapter.

Repeated exposure to ACEs can prolong this heightened stress response, rewiring the brain's physiological reactivity and impairing emotional

1. [Action steps using ACEs and trauma-informed care: A resilience model](#) by Laurie Leitch is licensed under [CC BY 4.0](#)
2. [Action steps using ACEs and trauma-informed care: A resilience model](#) by Laurie Leitch is licensed under [CC BY 4.0](#)

regulation. This can contribute to depression, anxiety, substance use, chronic physical and mental illnesses, and increased suicide risk. Chronic stress can also reduce positive social behaviors, such as collaboration and kindness, as cognitive resources are diverted toward survival responses.<sup>3</sup>

Arousal and reactivity symptoms include the following<sup>4</sup>:

- Being easily startled
- Feeling tense or “on edge”
- Having difficulty sleeping
- Having angry outbursts

These symptoms can make the person feel stressed and angry and can make it hard to do daily tasks, such as sleeping, eating, or concentrating.<sup>5</sup>

Individuals who have a history of trauma may become triggered by engagement with the health care system. They may experience arousal and reactivity symptoms. As a result of the stimulation of the “fight, flight, freeze, or fawn” stress response, the parts of the brain involved in memory, planning, decision-making, and regulation are not engaged. This can impact the client’s involvement with health care services and affect their ability to adhere to

3. Trauma-Informed Care Implementation Resource Center. (n.d.). *What is trauma-informed care?* Center for Health Care Strategies. <https://www.traumainformedcare.chcs.org/what-is-trauma-informed-care/>
4. National Institute of Mental Health. (2019). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>
5. National Institute of Mental Health. (2019). *Post-traumatic stress disorder*. U.S. Department of Health and Human Services. <https://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd>

treatment plans.<sup>6</sup> Nurses must understand this potential impact of previous trauma and incorporate client-centered, trauma-informed care.

Trauma-informed care is an approach that uses a lens of trauma to understand the range of cognitive, emotional, physical, and behavioral symptoms seen when individuals enter health care systems. **Trauma-informed care (TIC)** is a strengths-based framework that acknowledges the prevalence and impact of traumatic events in clinical practice, placing an emphasis on instilling in clients a sense of safety, control, and autonomy over their life and health care decisions. The basic goals of TIC are to avoid re-traumatization; emphasize survivor strengths and resilience; aid empowerment, healing, and recovery; and promote the development of survivorship skills.<sup>7</sup>

Two influential studies set the stage for the development of TIC<sup>8</sup>:

- The Adverse Childhood Experiences Study was an extensive study involving more than 17,000 individuals from the United States. It analyzed the long-term effects of childhood and adolescent traumatic experiences on adult health risks, mental health, health care costs, and life expectancy.<sup>9</sup>

6. Fleishman, J., Kamsky, H., & Sundborg, S. (2019). Trauma-informed nursing practice. *OJIN: The Online Journal of Issues in Nursing*, 24(2). <https://doi.org/10.3912/OJIN.Vol24No02Man03>
7. Tracy, E. E., & Macias-Konstantopoulos, W. (2021). Human trafficking: Identification and evaluation in the health care setting. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)
8. Center for Substance Abuse Treatment (US). (2014). *Trauma-informed care in behavioral health services*. <https://www.ncbi.nlm.nih.gov/books/NBK207201/>
9. Felitti, V. J., Anda, R. F., Nordenberg, D., Edwards, V., Koss, M. P., & Marks, J. S. (1998). Relationship of childhood abuse and household dysfunction to many

- The Women, Co-Occurring Disorders, and Violence Study was a study focused on the role of interpersonal and other traumatic stressors among women. The researchers examined the interrelatedness of trauma, violence, and co-occurring substance use and mental health disorders and the incorporation of trauma-informed services.<sup>10</sup>

The Substance Abuse and Mental Health Services Administration (SAMHSA) has defined six core principles of TIC. See Figure 15.2<sup>11</sup> for an infographic related to these principles<sup>12</sup>:

- **Safety:** Throughout the organization, clients and staff feel physically and psychologically safe.
- **Trustworthiness and Transparency:** Decisions are made with transparency and with the goal of building and maintaining trust.
- **Peer Support:** Individuals with shared experiences are integrated into the organization and viewed as integral to service delivery.
- **Collaboration and Mutuality:** Power differences between staff and clients and among organizational staff are leveled to support shared decision-

of the leading causes of death in adults: The adverse childhood experience (ACE) study. *American Journal of Preventive Medicine*, 14(3), 245-258.  
[https://doi.org/10.1016/S0749-3797\(98\)00017-8](https://doi.org/10.1016/S0749-3797(98)00017-8)

10. Clark, H. W., & Power, A. K. (2005). Women, co-occurring disorders, and violence study: A case for trauma-informed care. *Journal of Substance Abuse Treatment*, 28(2), 145-146. <https://doi.org/10.1016/j.jsat.2005.01.002>
11. This image is a derivative of “training\_emergency\_responders\_final.jpg” by [Centers for Disease Control and Prevention](#) and in the [Public Domain](#). Access for free at [https://www.cdc.gov/cpr/infographics/6\\_principles\\_trauma\\_info.htm](https://www.cdc.gov/cpr/infographics/6_principles_trauma_info.htm)
12. Trauma-Informed Care Implementation Resource Center. (n.d.). *What is trauma-informed care?* Center for Health Care Strategies.  
<https://www.traumainformedcare.chcs.org/what-is-trauma-informed-care/>

making.

- **Empowerment Voice and Choice:** Client and staff strengths are recognized, built on, and validated, including a belief in resilience and the ability to heal from trauma.
- **Cultural, Historical, and Gender Issues:** Biases and stereotypes (e.g., based on race, ethnicity, sexual orientation, age, geography) and historical trauma are recognized and addressed.



Figure 15.2 Core Principles of Trauma-Informed Care



View the following YouTube video on ACEs and trauma-informed care<sup>13</sup>: [What is Trauma-Informed Care?](https://youtu.be/fWken5DsJcw)

## Trauma-Informed Nursing Practice

Nurses can incorporate trauma-informed care by routinely implementing the following practices with all clients<sup>14</sup>:

13. Center for Health Care Strategies. (2019, January 23). *What is trauma-informed care?* [Video]. YouTube. All rights reserved. <https://youtu.be/fWken5DsJcw>
14. Fleishman, J., Kamsky, H., & Sundborg, S. (2019). Trauma-informed nursing practice. *OJIN: The Online Journal of Issues in Nursing*, 24(2). <https://doi.org/10.3912/OJIN.Vol24No02Man03>

- **Introduce Yourself and Your Role in Every Client Interaction:** Clients may recognize you, but they may not remember your role. This may lead to confusion and misunderstanding. When a client understands who you are and your role in their care, they feel empowered to be actively engaged in their own care. They also feel less threatened because they know your name and why you are interacting with them. When one party is nameless, there can be an automatic power differential in the interaction.
- **Use Open and Nonthreatening Body Positioning:** Be aware of your body position when working with clients. Open body language conveys trust and a sense of value. Trauma survivors often feel powerless and trapped. Health care situations can trigger past experiences of lack of control or an inability to escape. Using nonthreatening body positioning helps prevent the threat detection areas of the client's brain from taking over and helps clients stay regulated. A trauma-informed approach to body position includes attempting to have your body on the same level as the client, often sitting at or below the client. It could also include raising a hospital bed in order for the nurse and the client to be on the same level, reducing the likelihood of creating a perceived power differential through positioning. Additionally, it is important to think about where you and the client are positioned in the room in relation to the door or exit. Both nurse and client should have access to the exit so that neither feels trapped.
- **Provide Anticipatory Guidance:** Verbalize what the client can expect during a visit or procedure or what paperwork will cover. Knowing what to expect can reassure clients even if it is something that may cause discomfort. Past trauma is often associated with unexpected and unpredictable events. Knowing what to expect reduces the opportunity for surprises and activation of the SNS symptoms. It also helps clients feel more empowered in the care planning process.
- **Ask Before Touching:** For many trauma survivors, inappropriate or unpleasant touch was part of a traumatic experience. Touch, even when appropriate and necessary for providing care, can trigger a “fight, flight,



freeze or fawn” response and bring up difficult feelings or memories. This may lead to the individual experiencing increased anxiety and activation of the stress response, resulting in disruptive behaviors and possible dissociation. **Dissociation** is a break in how a person’s mind handles information, causing disconnection from their thoughts, feelings, memories, and surroundings. Nurses are often required to touch clients, and sometimes this touch occurs in sensitive areas. Any touch can be interpreted as unwanted or threatening, so it is important to ask all clients permission to touch them. Asking permission before you touch clients gives them a choice and empowers them to have control over their body and physical space. Be alert to nonverbal signs such as eye tearing, flinching, shrinking away, or other body language indicating the person is feeling uncomfortable. If the client exhibits signs of discomfort when being touched, additional nursing interventions can be implemented such as a mindfulness or grounding practice.

- **Protect Client Privacy:** Family members and other members of the medical team may be present when you care for a client. Clients may not feel empowered or safe in asking others to step out. It is crucial that nurses do not put the responsibility on the client to ask others to leave. It is the nurse’s role to ask the client (in private) whom they would like to be present during care and ask others to leave the room.
- **Provide Clear and Consistent Messaging About Services and Roles:** Trust is built when clients experience care providers who are forthright and honest. Dependability, reliability, and consistency are important when working with trauma survivors because previous trauma was often unexpected or unpredictable. Providing consistency from the nursing team regarding expectations and/or hospital rules can help clients feel secure and decrease opportunities for unmet expectations that might lead to triggering disruptive behavior.
- **Use Plain Language and Teach Back:** Avoid medical jargon and use clear, simple language. When clients are feeling triggered (i.e., their “fight, flight, freeze or fawn” system is engaged), information processing and

learning parts of the brain do not function optimally, and it is hard to remember new information. When providing education, information, or instructions, break information into small chunks and check for understanding. Offer to write important details down so they can accurately recall the information at a later time. Use clear language and “teach back” methods that empower clients with knowledge and understanding about their care.

- **Practice Universal Precaution:** Universal precaution means providing TIC to all clients regardless of a trauma history. Although ACE screening may be part of routine care, it can also have potential negative effects on clients. Unless a trauma-focused intervention is needed to amend the impact of trauma, many TIC experts propose universal precaution rather than direct screening.

▶ Read a continuing education article titled “[Trauma-Informed Nursing Practice](#)” in the *American Journal of Nursing*.<sup>15</sup>

Post-traumatic stress disorder (PTSD) is diagnosed in individuals who have been exposed to a traumatic event with chronic stress symptoms lasting more than one month that are so severe they

<sup>15</sup>. Dowdell, E., & Speck, P.M. (2022). Trauma-informed care in nursing practice. *American Journal of Nursing*, 122(4), 30-38.  
[https://journals.lww.com/ajnonline/Fulltext/2022/04000/CE\\_Trauma\\_Informed\\_Care\\_in\\_Nursing\\_Practice.22.aspx](https://journals.lww.com/ajnonline/Fulltext/2022/04000/CE_Trauma_Informed_Care_in_Nursing_Practice.22.aspx)

interfere with relationships, school, or work. Read more about PTSD in the “[Anxiety Disorders](#)” chapter.

## The Resilient Zone and Self-Regulation Skills

Individuals who have experienced repetitive or cumulative trauma may develop a dysregulated rhythm of their sympathetic nervous system (SNS) and parasympathetic nervous system (PNS), leaving them reactive and stuck in a state of hyperarousal, hypoarousal, or oscillating between the two extremes as their nervous system attempts to find balance. Simply put, the SNS “activates” the stress response, and the PNS “calms” the stress response. When out of balance, individuals may exhibit behaviors such as substance use disorders, self-harming, violence, poor school and work performance, bullying, and social disengagement.<sup>16</sup> Nurses can teach clients self-regulation skills by paying attention to their “fight, flight, or freeze” symptoms and learning how to balance their SNS and PNS systems.

Self-regulation skills teach clients how to return to a healthy balance of SNS and PNS stimulation called the resilient zone. See Figure 15.3<sup>17</sup> for an illustration of the **resilient zone** when the SNS and PNS are in balance. When in the resilient zone, the hormones released by the SNS during the “fight, flight, or freeze” stress response do not block conscious information processing by the brain. This balance promotes better capacity for flexibility

16. [Action steps using ACEs and trauma-informed care: A resilience model](#) by Laurie Leitch is licensed under [CC BY 4.0](#)

17. Leitch, L. (2017). Action steps using ACEs and trauma-informed care: A resilience model. *Health Justice* 5, 5. <https://doi.org/10.1186/s40352-017-0050-5>

and adaptability, prosocial behavior, improved problem-solving, and strategic thinking.<sup>18</sup>




Figure 15.3 The Resilient Zone

Teaching self-regulation skills enables the individual to pay attention to symptoms of the stress response and use techniques to purposefully stimulate the PNS. They focus on quality of breath, heart rate, and muscle tension and then use relaxation breathing, progressive muscle relaxation, meditation, or other methods to stimulate the PNS. These skills can be used prior to and during challenging events, as well as practiced over time to build deeper nervous system balance.<sup>19</sup>

18. [Action steps using ACEs and trauma-informed care: A resilience model](#) by Laurie Leitch is licensed under [CC BY 4.0](#)

19. [Action steps using ACEs and trauma-informed care: A resilience model](#) by Laurie Leitch is licensed under [CC BY 4.0](#)

- ▶ Read more about stress management, self-regulation skills, and grounding techniques in the “[Stress, Coping, and Crisis Intervention](#)” chapter.

## 15.4 Abuse and Neglect

### Child Neglect and Abuse

All 50 states and the District of Columbia have child abuse and neglect reporting laws that mandate certain professionals and institutions refer suspected maltreatment to a Child Protective Services (CPS) agency. Each state has its own definitions of child abuse and neglect that are based on standards set by federal law. Federal legislation defines child abuse and neglect as, “Any recent act or failure to act on the part of a parent or caretaker which results in death, serious physical or emotional harm, sexual abuse or exploitation, or an act or failure to act, which presents an imminent risk of serious harm.”<sup>1</sup> In 2019 there were 656,000 victims of child abuse and neglect, with a victim rate calculated as 9 victims per 1,000 children across the United States. In addition, 74.9 percent of victims were neglected, 17.5 percent were physically abused, and 9.3 percent were sexually abused. Child fatalities are the most tragic consequence of maltreatment. In 2019 it was estimated that 1,840 children died from abuse and neglect in the United States. The youngest children are the most vulnerable to maltreatment, with 45.4 percent of child fatalities younger than 1 year old. A perpetrator is the person who is responsible for the abuse or neglect of a child. Furthermore, 77.5% of perpetrators are a parent of the victim.<sup>2</sup>

An organization called A Safe Haven for Newborns is dedicated to preventing

1. Administration for Children & Families. (2021). *Child maltreatment 2019*. [Report]. U.S. Department of Health & Human Services.  
<https://www.acf.hhs.gov/cb/report/child-maltreatment-2019>
2. Administration for Children & Families. (2021). *Child maltreatment 2019*. [Report]. U.S. Department of Health & Human Services.  
<https://www.acf.hhs.gov/cb/report/child-maltreatment-2019>

infant abuse and abandonment through education, prevention, and direct assistance (see Figure 15.4<sup>3</sup>).

- ▶ Call the Safe Haven Helpline at 1-877-440-2229 or visit the [A Safe Haven for Newborns](https://asafehavenfornewborns.com) website to find local resources.




Figure 15.4 A Safe Haven For Newborns. Used under Fair Use.

3. "bc-english-ashfnb" by [A Safe Haven for Newborns](https://asafehavenfornewborns.com) is used under Fair Use. Access for free at <https://asafehavenfornewborns.com/international-safe-haven/>

# Assessment

## Signs of Neglect and Abuse

### NEGLECT

**Neglect** is a situation in which a parent or caretaker fails, refuses, or is unable, for reasons other than poverty, to provide the necessary care, food, clothing, or medical or dental care, which seriously endangers the physical, mental, or emotional health of the child. Signs of child neglect include the following<sup>4</sup>:

- Exhibits poor hygiene or body odor
- Is inappropriately dressed for weather
- Demonstrates needed medical or dental care
- Is left alone unsupervised for long periods of time
- Appears malnourished
- Is constantly hungry or begs for or steals food
- Demonstrates exhibits extreme willingness to please
- Is frequently absent from school
- Arrives early and stays late at school, play areas, or other people's home

### PHYSICAL ABUSE

**Physical abuse** is defined as injury inflicted on a child by other than accidental means. Physical injury includes, but is not limited to, lacerations, fractured bones, burns, internal injuries, severe or frequent bruising, or great bodily harm. Signs of physical abuse in children are as follows<sup>5</sup>:

4. Wisconsin Department of Children and Families. (n.d.). *Signs of child abuse and neglect*. <https://dcf.wisconsin.gov/cps/signs>
5. Wisconsin Department of Children and Families. (n.d.). *Signs of child abuse and neglect*. <https://dcf.wisconsin.gov/cps/signs>



- Bruises and/or welts on face, neck, chest, back, or soft muscle areas less prone to bruising by natural play or accidents (e.g., abdomen, breasts, under arm, inner thigh)
- Injuries in the shape of object (e.g., belt or cord)
- Unexplained burns on palms, soles of feet, or back; a line of demarcation from submerging in hot liquids (e.g., ankles, buttocks, wrists); burns in the shape of object (e.g., fork, cigarette)
- Fractures that do not fit the story of how an injury occurred
- Delay in seeking medical help
- Extremes in behavior (e.g., very aggressive or withdrawn and shy)
- Afraid to go home
- Frightened of parents
- Fearful of other adults
- Failure to thrive

However, some injuries are not visible to observation, such as shaken baby syndrome, a serious brain injury resulting from forcefully shaking an infant or toddler.

## SEXUAL ABUSE

**Sexual abuse** is defined as sexual intercourse or sexual touching of a child; sexual exploitation; human trafficking of a child; forced viewing of sexual activity; or permitting, allowing, or encouraging a child to engage in prostitution. Here are signs of sexual abuse in children<sup>6</sup>:

- Pain, swelling, or itching in genital area
- Bruises, bleeding, discharge in genital area
- Difficulty walking or sitting, frequent urination, or pain
- Stained or bloody underclothing
- Sexually transmitted diseases

6. Wisconsin Department of Children and Families. (n.d.). *Signs of child abuse and neglect*. <https://dcf.wisconsin.gov/cps/signs>

- Refusal to take part in gym or other exercises
- Poor peer relationships
- Unusual interest in sex for age
- Drastic change in school achievement
- Runaway or delinquent behavior
- Regressive to behaviors expected for a younger child

► Read additional information about human trafficking in the “[Vulnerable Populations](#)” chapter.

## EMOTIONAL ABUSE

**Emotional abuse** is defined as harm to a child’s psychological or intellectual functioning, which is exhibited by severe anxiety, depression, withdrawal, or aggression. Emotional damage may be demonstrated by substantial and observable changes in behavior, emotional response, or learning that are incompatible with the child’s age or stage of development. Signs of emotional abuse in children include the following<sup>7</sup>:

- Low self-esteem
- Self-denigration
- Severe depression
- Unusual level of aggression
- Severe anxiety
- Extreme withdrawal
- Failure to learn

7. Wisconsin Department of Children and Families. (n.d.). *Signs of child abuse and neglect*. <https://dcf.wisconsin.gov/cps/signs>

# Interventions

Effective interventions are essential to ensure child safety, stop the abuse, support recovery, and prevent future harm. Interventions typically begin with immediate protective actions and extend into medical, psychological, legal, and family-centered care.

When abuse is suspected, professionals such as nurses, educators, and social workers (who are often mandated reporters) must report their concerns to local child protective services (CPS). Child Protective Services (CPS) agencies provide services to children and their families, both in their homes and in foster care. Services are provided to prevent future instances of child maltreatment and remedy conditions that brought the children and their family to the attention of the agency.<sup>8</sup> Once a report is made, CPS investigates the claim, assesses the level of risk, and determines the need for protective measures. In cases of immediate danger, children may be removed from their homes and placed in temporary foster or kinship care. Forensic interviews are often conducted in child advocacy centers by trained professionals using trauma-informed techniques to gather accurate and sensitive information without re-traumatizing the child.

Following the initial response, children may require medical attention to treat injuries or address physical health concerns. Pediatric healthcare providers, especially those trained in child maltreatment, play a crucial role in identifying signs of abuse and initiating appropriate referrals. Mental health interventions are also essential. **Trauma-focused cognitive behavioral therapy (TF-CBT)** is widely recognized as an evidence-based treatment that helps children process trauma, develop coping skills, and reduce symptoms of

8. Administration for Children & Families. (2021). *Child maltreatment 2019*. [Report]. U.S. Department of Health & Human Services.  
<https://www.acf.hhs.gov/cb/report/child-maltreatment-2019>

anxiety, depression, or post-traumatic stress.<sup>9</sup> Family therapy may be introduced if reunification with parents or guardians is a possibility, promoting healing and restoring healthy family relationships.

Ongoing support often includes case management and access to family-centered services. Parenting programs and home visiting initiatives, such as the Nurse-Family Partnership, aim to improve caregiver skills, reduce stress, and create safer home environments. These supports are especially beneficial in preventing recurrence of abuse. Social workers and case managers work closely with families to coordinate services, monitor progress, and ensure adherence to safety plans.

Legal involvement is another critical element. Family courts make determinations about custody, visitation, and parental rights, while criminal courts address charges against abusers. Guardians ad litem (GALs) or court-appointed special advocates (CASAs) may be assigned to represent the child's best interests in these proceedings. The legal process can be daunting for families and children, which is why coordination between legal and mental health professionals is vital to minimize additional trauma.

A multidisciplinary team (MDT) approach enhances the effectiveness of child abuse interventions. This collaborative model brings together healthcare providers, mental health specialists, law enforcement, educators, and child welfare workers to share information, streamline care, and ensure the child receives comprehensive, holistic support. MDTs often meet at child advocacy centers or through coordinated local systems to address cases in a child-centered and trauma-informed manner.

Importantly, interventions must be tailored to the child's age, developmental stage, and cultural background. Cultural humility and developmental sensitivity help build trust with families and ensure that services are both respectful and effective. Interventions that are culturally responsive and

9. Medical University of South Carolina. (n.d.). *TF-CBT Web: Trauma-focused cognitive behavioral therapy training*. <https://www.tfcbt.org>

developmentally appropriate have been shown to reduce disparities in care and promote engagement with services.

## Neglect and Abuse of Older Adults and Adults at Risk

Elder abuse is a common problem in the United States. Abuse, including neglect and exploitation, is experienced by about 1 in 10 people aged 60 and older who live at home. From 2002 to 2016, more than 643,000 older adults were treated in the emergency department for nonfatal assaults, and over 19,000 homicides occurred. This information is considered an underestimate of the problem because it is limited to those individuals treated in emergency departments and doesn't include those who do not seek treatment. Victims must decide whether to tell someone they are being hurt or continue being abused by someone they depend upon or care for deeply.<sup>10</sup>

**Elder abuse** is defined as an intentional act or failure to act that causes or creates a risk of harm to an older adult. An older adult is defined as someone age 60 or older.<sup>11</sup> Adults at risk are also considered vulnerable adults at risk for abuse. **Adults at risk** are defined as adults who have a physical or mental condition that impairs their ability to care for their own needs.<sup>12</sup> Older adults and adults at risk are potentially susceptible for abuse, neglect, or financial exploitation by caregivers or a person they trust.<sup>13</sup> A **caregiver** is a person who has taken responsibility for all or part of an individual's care. Risk factors for

10. Centers for Disease Control and Prevention. (2024). *About abuse of older persons*. <https://www.cdc.gov/elder-abuse/about/index.html>

11. Centers for Disease Control and Prevention. (2024). *About abuse of older persons*. <https://www.cdc.gov/elder-abuse/about/index.html>

12. Wisconsin Department of Health Services. (2018). *Adult protective services: Definitions*. <https://www.dhs.wisconsin.gov/aps/definitions.htm#Abuse>

13. Centers for Disease Control and Prevention. (2024). *About abuse of older persons*. <https://www.cdc.gov/elder-abuse/about/index.html>

elder abuse include cognitive impairment, functional dependency, poor physical health, low income, social isolation, and a history of past trauma or abuse. Perpetrator-related risk factors include caregiver burden, psychiatric illness, and substance abuse. Environmental factors such as living in high-crime areas and low social support also contribute to the risk.<sup>14</sup>

The following are types of elder abuse<sup>15</sup>:

- **Physical abuse:** Physical abuse refers to illness, pain, injury, functional impairment, distress, or death as a result of the intentional use of physical force and includes acts such as hitting, kicking, pushing, slapping, and burning. See Figure 15.5<sup>16</sup> for an infographic describing physical signs of elder abuse.
- **Sexual abuse:** Sexual abuse refers to forced or unwanted sexual interaction of any kind. This may include unwanted sexual contact or penetration or non-contact acts such as sexual harassment.
- **Emotional abuse:** Emotional abuse refers to verbal or nonverbal behaviors that inflict anguish, mental pain, fear, or distress, such as humiliation or disrespect, verbal and nonverbal threats, harassment, and geographic or interpersonal isolation.
- **Neglect:** Neglect is the failure to meet the person's basic needs, including food, water, shelter, clothing, hygiene, and essential medical care.
- **Financial abuse:** Financial abuse is the illegal, unauthorized, or improper

14. Johannesen, M. & LoGiudice, D. (2013). Elder abuse: a systematic review of risk factors in community-dwelling elders. *Age Ageing*, 42(3), 292-8. [doi: 10.1093/ageing/afs195](https://doi.org/10.1093/ageing/afs195).

15. Centers for Disease Control and Prevention. (2024). *About abuse of older persons*. <https://www.cdc.gov/elder-abuse/about/index.html>

16. "Signs of Elder Abuse – Physical" by National Council on Aging is used under Fair Use. Access for free at <https://www.ncoa.org/article/get-the-facts-on-elder-abuse>

use of money, benefits, belongings, property, or assets for the benefit of someone other than the individual.

- **Treatment without consent:** Treatment without consent refers to the administration of medication or the performance of psychosurgery, electroconvulsive therapy, or experimental research on an individual who has not provided informed consent.
- **Unreasonable confinement or restraint:** Unreasonable confinement or restraint refers to the intentional and unnecessary confinement of an individual in a locked room, involuntary separation from their living area, use of physical restraints, or the provision of unnecessary or excessive medication. (This does not include the use of these methods or devices if they conform with state and federal standards governing restraint or seclusion. Review ANA guidelines on using restraints in the “[Client Rights](#)” section of the “Legal and Ethical Considerations in Mental Health Care” chapter.)

## Assessment

Elder abuse can be challenging to detect, as symptoms may be subtle or mistaken for age-related conditions. A comprehensive assessment involves careful observation, open-ended questioning, and consideration of both physical and psychosocial indicators. Clinicians should maintain a high index of suspicion, especially when inconsistencies arise between reported histories and observed findings.

Physical indicators may include unexplained injuries such as bruises, lacerations, burns, or fractures. Particular attention should be paid to injury patterns that are inconsistent with normal aging or accidental trauma. For instance, bruising on the maxillofacial area, upper extremities (especially the inner arms), or torso may suggest defensive injuries or restraint use. Recurrent injuries at various stages of healing are also red flags. In cases of

sexual abuse, assess for genital trauma, sexually transmitted infections, or sudden behavioral changes.<sup>17,18</sup>

Behavioral and psychological signs can include withdrawal, increased anxiety, depression, agitation, or fear of specific individuals—especially caregivers. A once-outgoing elder who becomes socially isolated or unusually quiet during interactions may be experiencing emotional or psychological abuse.<sup>19,20</sup>

Neglect-related signs are equally important and may present as poor hygiene, untreated medical conditions, malnutrition, dehydration, pressure injuries, or inappropriate clothing for the weather. Weight loss without a medical cause or a dirty, unsafe living environment may also be indicative of neglect.<sup>21,22</sup>

Healthcare utilization patterns may offer additional clues. Frequent emergency department visits, inconsistent follow-up care, or evidence of

17. National Center on Elder Abuse (NCEA). (2020). *Types and signs of abuse*. U.S. Department of Health & Human Services. <https://ncea.acl.gov>
18. Lachs, M. S., & Pillemer, K. A. (2015). Elder abuse. *The New England Journal of Medicine*, 373(20), 1947–1956. <https://doi.org/10.1056/NEJMr1404688>
19. National Center on Elder Abuse (NCEA). (2020). *Types and signs of abuse*. U.S. Department of Health & Human Services. <https://ncea.acl.gov>
20. Lachs, M. S., & Pillemer, K. A. (2015). Elder abuse. *The New England Journal of Medicine*, 373(20), 1947–1956. <https://doi.org/10.1056/NEJMr1404688>
21. National Center on Elder Abuse (NCEA). (2020). *Types and signs of abuse*. U.S. Department of Health & Human Services. <https://ncea.acl.gov>
22. Lachs, M. S., & Pillemer, K. A. (2015). Elder abuse. *The New England Journal of Medicine*, 373(20), 1947–1956. <https://doi.org/10.1056/NEJMr1404688>



“doctor shopping” by caregivers may signal underlying abuse or caregiver manipulation.

During assessment, it’s crucial to:

- Interview the elder alone, if possible, to ensure their safety and candor.
- Use validated screening tools, such as the Elder Abuse Suspicion Index (EASI) or Hwalek-Sengstock Elder Abuse Screening Test (H-S/EAST).<sup>23</sup>
- Document findings thoroughly and objectively, including direct quotes, photos (as per policy), and detailed descriptions of injuries or environmental conditions.

Lastly, consider cultural and cognitive factors that may affect an elder’s willingness or ability to disclose abuse. Older adults with dementia or language barriers may require additional support or modified communication strategies.

23. Yaffe, M. J., Wolfson, C., Lithwick, M., & Weiss, D. (2008). Development and validation of a tool to improve physician identification of elder abuse: The Elder Abuse Suspicion Index (EASI). *Journal of Elder Abuse & Neglect*, 20(3), 276–300. <https://doi.org/10.1080/08946560801973168>

# Physical Signs of Elder Abuse



Dehydration  
or unusual  
weight loss



Missing  
daily living  
aids



Unexplained  
injuries, bruises,  
cuts, or sores



Unsanitary living  
conditions and  
poor hygiene



Unattended  
medical  
needs

To learn more, visit [ncea.acl.gov](https://ncea.acl.gov)

Figure 15.5 Physical Signs of Elder Abuse. Used under Fair Use.

## Interventions

Effective interventions for elder abuse require a coordinated, multidisciplinary response that prioritizes the safety, dignity, and autonomy of the older adult. Once abuse is suspected or confirmed, a key resource is **Adult Protective Services (APS)**. APS agencies are state-run programs that investigate reports of elder and vulnerable adult abuse, neglect, and exploitation. Their role is to protect adults—particularly those with developmental disabilities, degenerative brain disorders, serious mental illness, or other incapacities—who are unable to protect themselves from harm.

APS interventions are tailored to the individual's needs and may include outreach, crisis counseling, and coordination of medical, legal, and social services. Caseworkers assess the level of risk and collaborate with healthcare providers, legal authorities, and family members to ensure safety while respecting the adult's right to self-determination when possible. Services may also include helping arrange safe housing, facilitating access to mental health or substance use treatment, and supporting guardianship proceedings if the individual is deemed unable to make informed decisions.

In cases of financial exploitation, APS may work in tandem with law

enforcement and financial institutions to recover assets or pursue prosecution. When abuse involves a caregiver, interventions can also focus on caregiver support, respite care, and education to reduce stress and prevent recurrence. For institutional settings, reporting to long-term care ombudsman programs and state licensing boards may trigger regulatory reviews and corrective actions.<sup>24</sup>

See Figure 15.6<sup>25</sup> for an illustration related to reporting concerns about vulnerable adults.




Figure 15.6 Reporting Concerns About Vulnerable Adults. Used under Fair Use.

- Find resources in your area for reporting elder abuse at the [National Adult Protective Services Association website](https://www.napsa-now.org/get-informed/what-is-adult-protective-services/).

24. National Adult Protective Services Association. (n.d.). *Adult Protective Services (APS)*. <https://www.napsa-now.org/get-informed/what-is-adult-protective-services/>

25. “APS\_header\_3.jpg” by unknown author for [Washington State Department of Social and Health Services](#) is used under Fair Use.

- ▶ Read more about protective services in your state. Here are links to Wisconsin's [Adult Protective Services](#).
- ▶ Find elder care resources in your community at [Eldercare Locator](#)

## Mandatory Reporting

**Mandated reporters** are required by law to report suspected abuse and neglect they see in the course of their professional duties. Nurses and other professionals are referred to as mandated reporters because they are required by state law to report suspected neglect or abuse of children, adults at risk, and the elderly. Nurses should be aware of the county or state agencies to whom they should report suspected abuse. For example, in Wisconsin, suspected neglect or abuse is reported to the Child Protective Services (CPS) or law enforcement. Persons required to report and who intentionally fail to report suspected child abuse or neglect may be fined up to \$1,000 or imprisoned for up to six months or both.<sup>26</sup> See Figure 15.7<sup>27</sup> for an image related to reporting suspected abuse and neglect.

26. Wisconsin Department of Children and Families. (2019). *It shouldn't hurt to be a child... but sometimes it does*. [PDF Brochure]. <https://dcf.wisconsin.gov/files/publications/pdf/0101.pdf>

27. "[stop-child-abuse-see-the-signs-make-the-call-1536x748.jpg](#)" by unknown author for [Delaware Department of Services for Children, Youth & Their Families](#) is used under Fair Use.



Figure 15.7 Report Suspected Child Abuse. Used under Fair Use.

## What to Report

Mandatory reporters who suspect neglect or abuse should contact their county social/human services department, sheriff, or local police department immediately. When making a report, explain what happened or is happening to the child, vulnerable adult, or elderly. Describe the nature of the abuse or neglect and be as specific as possible. Be prepared to give the name, DOB, address, and telephone number of the victim, as well as the name of their parent or caregiver. If you do not know all of this information, report what you do know and explain all you know about the situation and family dynamics.<sup>28</sup>

When a report is filed, the receiving department will make a safety screening determination based on state statutes. If the report meets the criteria for alleged maltreatment, a social worker from the county department of social/human services will proceed with an investigation of the reported maltreatment and work with the parents or caregiver to assess the situation

28. Wisconsin Department of Children and Families. (2019). *It shouldn't hurt to be a child... but sometimes it does*. [PDF Brochure]. <https://dcf.wisconsin.gov/files/publications/pdf/0101.pdf>

to determine if any support or assistance is needed to protect the child, vulnerable adult, or elderly.<sup>29</sup>

- ▶ Find resources in your area for reporting suspected child abuse at [ChildHelp National Child Abuse Hotline](#).
- ▶ Read more about protective services in your state. Access Wisconsin's information at [Child Protective Services](#).

29. Wisconsin Department of Children and Families. (2019). *It shouldn't hurt to be a child... but sometimes it does*. [PDF Brochure]. <https://dcf.wisconsin.gov/files/publications/pdf/0101.pdf>

## 15.5 Intimate Partner Violence

Intimate partner violence (IPV), sexual assault, and rape are crimes with long-lasting effects on victims and are a great cost to society. These crimes happen to both women and men and are often associated with substance use. A recent national survey found that 22 percent of women and 14 percent of men reported experiencing severe physical violence from an intimate partner in their lifetimes.<sup>1</sup> IPV is a significant public health issue that has many individual and societal costs. About 35% of female IPV survivors and more than 11% of male IPV survivors experience some form of physical injury related to IPV, and some deaths occur. About 1 in 5 homicide victims are killed by an intimate partner, and over half of female homicide victims in the US are killed by a current or former male intimate partner.<sup>2</sup> See Figure 15.8<sup>3</sup> for a CDC infographic related to IPV.

1. U.S. Department of Health and Human Services (HHS), Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. [PDF Report].  
<https://addiction.surgeongeneral.gov/sites/default/files/surgeon-generals-report.pdf>
2. U.S. Department of Health and Human Services (HHS), Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. [PDF Report].  
<https://addiction.surgeongeneral.gov/sites/default/files/surgeon-generals-report.pdf>
3. "how-big-is-the-problem-ipv-large" by [Centers for Disease Control and Prevention](#) is in the [Public Domain](#). Access for free at <https://www.cdc.gov/violenceprevention/intimatepartnerviolence/fastfact.html>





Figure 15.8 Intimate Partner Violence Statistics

**Intimate partner violence (IPV)** is abuse or aggression that occurs in a romantic relationship. “Intimate partner” refers to both current and former spouses and dating partners. IPV can vary in how often it happens and how severe it is. It can range from one episode of violence that could have lasting impact to chronic and severe episodes over multiple years. IPV can include any of the following types of behaviors<sup>4</sup>:

- **Physical violence:** When a person hurts or tries to hurt a partner by hitting, kicking, or using another type of physical force
- **Sexual violence:** Forcing or attempting to force a partner to take part in a sex act, sexual touching, or a nonphysical sexual event (e.g., sexting) when the partner does not or cannot consent
- **Stalking:** A pattern of repeated, unwanted attention and contact by a partner that causes fear or concern for one’s own safety or the safety of someone close to the victim
- **Psychological aggression:** The use of verbal and/or nonverbal communication with the intent to harm another partner mentally or emotionally and/or to exert control over another partner

4. Centers for Disease Control and Prevention. (2024). *About violence prevention*. <https://www.cdc.gov/violenceprevention/intimatepartnerviolence/index.html>





View the following CDC YouTube video on intimate partner violence<sup>5</sup>: [What is Intimate Partner Violence?](https://youtu.be/VuMCzU54334)

## Teen Dating Violence

When IPV occurs during adolescence, it is referred to as **teen dating violence (TDV)**. TDV affects millions of U.S. teens each year. The use of alcohol and drugs is a risk factor for non-consensual sexual contact among undergraduate and graduate students.<sup>6</sup> See Figure 15.9<sup>7</sup> for an infographic used to teach about healthy relationships.

5. Centers for Disease Control and Prevention. (2019, May 15). *What is intimate partner violence?* [Video]. YouTube. All rights reserved.  
<https://youtu.be/VuMCzU54334>
6. U.S. Department of Health and Human Services (HHS), Office of the Surgeon General. (2016). *Facing addiction in America: The surgeon general's report on alcohol, drugs, and health*. [PDF Report].  
<https://addiction.surgeongeneral.gov/sites/default/files/surgeon-generals-report.pdf>
7. "teen-dating-violence-relationship-range-chart-graphic.png" by unknown author for [FairFax County Department of Family Services, Domestic and Violent Services](https://www.fairfaxcounty.gov/familyservices/domestic-sexual-violence/teen-dating-violence) is used under Fair Use. Access for free at <https://www.fairfaxcounty.gov/familyservices/domestic-sexual-violence/teen-dating-violence>

HEALTHY	UNHEALTHY	ABUSIVE
<p><b>A healthy relationship means you and your partner:</b></p> <ul style="list-style-type: none"> <li>▪ Are economic/financial partners</li> <li>▪ Are equal, honest, respectful and trusting</li> <li>▪ Communicate</li> <li>▪ Enjoy spending personal time away from one another</li> <li>▪ Make mutual choices together</li> </ul> <p><i>Adapted from: Love Is Respect</i></p>	<p><b>Your relationship may be unhealthy if your partner:</b></p> <ul style="list-style-type: none"> <li>▪ Is dishonest, disrespectful, or distrustful</li> <li>▪ Doesn't communicate</li> <li>▪ Makes you pay for being economically unequal</li> <li>▪ Only wants you to spend time with each other</li> <li>▪ Pressures you into activities</li> </ul>	<p><b>Your relationship may be abusive when your partner:</b></p> <ul style="list-style-type: none"> <li>▪ Accuses you of cheating when you're not</li> <li>▪ Communicates in a threatening or hurtful way</li> <li>▪ Controls you</li> <li>▪ Denies their abusive actions</li> <li>▪ Isolates you from family and friends</li> </ul>

Figure 15.9 Healthy Versus Unhealthy and Abusive Relationships. Used under Fair Use.

## Warning Signs of Intimate Partner Violence

These observations should heighten a nurse's suspicion of IPV<sup>8</sup>:

- Inconsistent explanation of injuries.
- Delay in seeking treatment for injuries.
- Frequent emergency department or urgent care visits.  
(Abusers typically do not want their victims to form an ongoing care relationship with one clinician. They may feel the victim will be less likely to find an ally in an emergency department where care may be more fragmented.)
- Missed appointments. The client may not keep

8. Weil, A. (2024). Intimate partner violence: Diagnosis and screening. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

appointments because the abuser will not allow medical attention.

- Late initiation of prenatal care during pregnancy.
- Repeated abortions. Unplanned pregnancy may result from sexual assault and/or not being allowed to use birth control by the abuser.
- Medication nonadherence. Victims may not take medicines because the abuser has taken them away or not allowed the partner to fill prescriptions.
- Inappropriate affect. Victims may appear jumpy, fearful, or cry readily. They may avoid eye contact and seem evasive or hostile. Additionally, flat affect or dissociation may suggest post-traumatic stress disorder.
- Overly attentive or verbally abusive partner. The clinician should be suspicious if the partner answers questions for the client. If the partner refuses to leave the examination room, the nurse should find a way to get the partner to leave before questioning the client.
- Apparent social isolation.
- Reluctance to undress or have a genital, rectal, or oral examination or difficulty undergoing these or other examinations.

## Assessment

The setting in which questioning occurs is important. The nurse must ensure that the client feels safe and comfortable. Individuals are more likely to

disclose their experiences of IPV when the nurse uses the following strategies<sup>9</sup>:

- Provide privacy. Other people present in the room should be asked to leave for the interview and examination. Resistance of a partner to leave is a warning sign.
- Assure confidentiality (unless mandatory reporting is required)
- Use open-ended questioning and ask only a few questions.
- Ask about “being hurt” or “treated badly” and avoid phrases like “victim,” “abused,” or “battered.” Mirror the client’s word choices and use words like “hurt,” “frightened,” or “treated badly.”
- Do not disclose or discuss your concerns with the client’s partner.
- Do not ask why the client has not left their partner.

The physical examination can provide warning signs that abuse may be occurring. The presence and location of injuries are important. Any injury without a good explanation, particularly involving the head and neck, teeth, or genital area, should raise suspicion. Wounds on the forearms often occur when a victim is in a defensive position. Bruises of different ages may be present due to repeated abuse.

People experiencing abuse may deny the abuse for several reasons. They may not feel safe in disclosing information, especially if questions are not asked in a private environment or there is a fear of information not being kept confidential. They may not be emotionally ready to admit the reality of the situation, they may blame themselves, or they may feel like a failure if they admit to being abused. They may fear rejection, feel ashamed, believe that the abuse will not happen again, fear reprisal by the abuser, believe that they have no alternatives, or lack knowledge of resources that could help them. There may be language or cultural barriers between nurses and clients that

9. Weil, A. (2024). Intimate partner violence: Diagnosis and screening. *UpToDate*. [www.uptodate.com](https://www.uptodate.com)

interfere with communication or discomfort with using an interpreter to discuss sensitive issues.<sup>10</sup>

Routine screening is recommended for all clients on initial primary care visits, OB-GYN visits, emergency department visits, and on hospital admission.<sup>11</sup>

## SAFE Survey

The Joint Commission recommends that hospitals use criteria to identify possible victims of abuse and neglect upon hospital entry and on an ongoing basis, educate staff about how to recognize signs of abuse, assist with referrals of possible victims, and report abuse in accordance with law and regulation.

Several IPV screening tools have been studied for use in emergency departments and clinics. SAFE is an example of a short survey tool that stands for the following assessments<sup>12,13</sup>:

- **S: Stress/Safety:** Do you feel safe in your relationship?
- **A: Afraid/Abused:** Have you ever been in a relationship where you were threatened, hurt, or afraid?

10. Weil, A. (2024). Intimate partner violence: Diagnosis and screening. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

11. Weil, A. (2024). Intimate partner violence: Diagnosis and screening. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

12. Weil, A. (2024). Intimate partner violence: Diagnosis and screening. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

13. Rabin, R. F., Jennings, J. M., Campbell, J. C., & Bair-Merritt, M. H. (2009). Intimate partner violence screening tools. *American Journal of Preventive Medicine*, 36(5), 439-445.e4. <https://doi.org/10.1016/j.amepre.2009.01.024>

- **F: Friends/Family:** Are your friends or family aware you have been hurt?
- **E: Emergency Plan:** Do you have a safe place to go and the resources you need in an emergency?

► Read the following PDF for additional information about Minnesota Department of Health's IPV screening tools: [Family Home Visiting Intimate Partner Violence Screening & Referrals Toolkit](#)

## Interventions and Safety Plans

Survivors of IPV are often afraid to leave their abusive partners because of the threats that have been made against them or their loved ones. The biggest threats for victims are strangulation (ten times more likely to be killed), the presence of a firearm (five times more likely), or if the abusive person is suicidal. It is not uncommon for an abusive person to threaten to kill themselves if they feel as if they're losing control over their partner, and they pose a serious risk to their victim if they have attempted suicide in the past, talk about a specific plan, or have access to a gun. Domestic violence is the single biggest indicator of murder-suicides in the United States, and this has escalated to include survivors' friends and family.<sup>14</sup>

Safety is a top priority when IPV is identified. Before the client leaves the office, referrals to local resources should be made with a personalized safety plan in place. The most dangerous time in a relationship occurs when the abused person decides to leave. Nearly 77 percent of domestic violence-related homicides occur upon separation, and there is a 75 percent increase

14. National Domestic Violence Hotline. (n.d.). *Plan for safety*.  
<https://www.thehotline.org/plan-for-safety/>

of violence upon separation for at least two years.<sup>15</sup> As the abuser realizes they are losing power and control over their partner, they often escalate tactics to increase fear in the individual leaving the relationship in an effort to make the individual stay. A **safety plan** is a set of actions that can help lower the risk of being hurt by an abusive partner. It includes specific information and resources that will increase one's safety at school, home, and other places visited regularly. View an infographic related to a safety plan in Figure 15.10.<sup>16</sup>

15. Battered Women's Support Services. (2020). *Eighteen months after leaving domestic violence is still the most dangerous time*. <https://www.bwss.org/eighteen-months-after-leaving-domestic-violence-is-still-the-most-dangerous-time/>
16. "vaw-covid-1.jpg" by unknown author for [World Health Organization](#) is licensed under [CC BY-NC-SA 3.0 IGO](#). Access for free at <https://www.who.int/multi-media/details/make-a-safety-plan-for-you-and-your-children>

# Make a **safety plan** for you and your children:

If you are experiencing violence at home and need to leave in a hurry

**Identify** a friend, neighbour, relative, or shelter you can go to



**Plan** how to get there



**Keep ready** essential personal items to take with you



World Health Organization

human reproduction programme  
**hrp.**  
research for impact  
UNDP-UNFPA-UNICEF-WHO-WORLD BANK

Figure 15.10 Safety Plan

## Resources for Survivors of IPV

- ▶ Guide clients experiencing IPV to create a personalized online safety plan at the National Domestic Violence Hotline's [Create a Safety Plan](#) web page.
- ▶ If you need help or know someone who is experiencing IPV, visit the [National Domestic Violence Hotline](#) or call 1-800-799-7233. Online resources for developing a customized safety plan are available.
- ▶ Read about teen dating and healthy relationships at the [Love is Respect website](#).



## Preventing IPV

There are many negative health outcomes associated with IPV, including chronic conditions affecting the heart, digestive, reproductive, musculoskeletal, and nervous systems. Survivors may experience mental health problems such as depression and post-traumatic stress disorder (PTSD) and are at higher risk for engaging in behaviors such as smoking, binge drinking, and risky sexual behaviors.<sup>17</sup>

Nurses and communities can promote healthy, respectful, and non-violent relationships to help reduce the occurrence of IPV.<sup>18</sup> See Figure 15.11<sup>19</sup> for an infographic related to IPV prevention strategies from the CDC.

17. Centers for Disease Control and Prevention. (2024). *About violence prevention*. <https://www.cdc.gov/violenceprevention/intimatepartnerviolence/index.html>
18. Centers for Disease Control and Prevention. (2024, April 9). *About violence prevention*. <https://www.cdc.gov/violenceprevention/intimatepartnerviolence/index.html>
19. “how-can-we-stop-ipv.PNG” by [National Center for Injury Prevention and Control, Division of Violence Prevention](#) is in the [Public Domain](#). Access for free at <https://www.cdc.gov/violenceprevention/intimatepartnerviolence/fastfact.html>



## 15.6 Workplace Violence

**Workplace violence** consists of physically and psychologically damaging actions that occur in the workplace or while on duty. Examples of workplace violence include direct physical assaults (with or without weapons), written or verbal threats, physical or verbal harassment, and homicide.<sup>1</sup>

Violence committed by clients or family members toward health care staff can occur in many health care settings. Therefore, nurses must be prepared to cope effectively with agitated clients to reduce the risk of serious injury to the client, themselves, staff, and other clients. Up to 50 percent of health care professionals are victims of violence at some point during their careers. There is a wide range of risk factors for client violence, including the environment, a client's social and medical history, interpersonal relationships, genetics, neurochemistry and endocrine function, and substance abuse. In the emergency department (ED), substance intoxication or withdrawal is the most common diagnosis in combative clients. Known psychiatric illness is also a risk factor for violent behavior, with schizophrenia, personality disorders, mania, and psychotic depression most often associated with violence. Psychosis, delirium, and dementia can also lead to violent behavior.<sup>2</sup>

Some states are introducing legislation to increase the penalties for people who commit violence against nurses, making battery to a nurse a felony instead of a misdemeanor. A recent law passed in Wisconsin in 2022 makes it

1. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)
2. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

a felony to threaten a health care worker.<sup>3</sup> See Figure 51.12<sup>4</sup> for an illustration related to stopping workplace violence for nurses in Wisconsin.



Figure 15.12 Stop Workplace Violence. Used under Fair Use.

Assessment of the combative client begins with risk assessment and attention to safety measures. Violence typically erupts after a period of mounting tension. In a typical scenario, the client first becomes angry, then resists authority, and finally becomes confrontational. However, violent behavior may erupt without warning, especially when caused by medical illness or dementia. A nurse may identify verbal and nonverbal cues of agitation and defuse the situation before violence happens. It is helpful to

3. Mensik, H. (2022). *Wisconsin passes law making threats against healthcare workers a felony*. HealthCareDive. <https://www.healthcaredive.com/news/wisconsin-passes-law-threat-healthcare-workers-felony/620978/#:~:text=Wisconsin%20Governor%20Tony%20Evers%20signed,officers%20and%20other%20government%20workers>
4. This image is a derivative of “WP-Violence-graphic-600×800.png” by [Wisconsin Nurses Association](https://www.wisconsinnurses.org/education/workplace-violence/) is used under Fair Use. Access for free at <https://www.wisconsinnurses.org/education/workplace-violence/>

observe the nonverbal communication of a client's hands as an indicator of tension. Other signs of impending violence include the following<sup>5</sup>:

- Confrontational behavior
- Angry demeanor
- Loud, aggressive speech
- Tense posturing (e.g., gripping arm rails tightly or clenching fists)
- Frequently changing body position or pacing
- Aggressive acts (e.g., pounding walls, throwing objects, or hitting oneself)

Clients who are agitated but cooperative may be amenable to verbal de-escalation techniques. Actively violent clients and uncooperative, agitated clients, particularly those who exhibit signs of impending violence, require immediate physical restraint per an agency's policy. Assume that all violent clients are armed until proven otherwise, especially those presenting to an emergency department.<sup>6</sup>

► Read about crisis intervention techniques in the "[Stress, Coping, and Crisis Intervention](#)" chapter.

## Verbal De-Escalation Techniques

Verbal de-escalation techniques should be attempted before physical restraints or sedative medications are implemented. During initial interactions with the client, it will rapidly become clear whether the client will

5. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)
6. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

cooperate or continue to escalate. This interaction also enables the nurse to assess the client's mental status. If the client remains agitated or is incapable of interacting appropriately, then restraints become necessary.<sup>7</sup>

When attempting to de-escalate an individual, the nurse should adopt an honest and straightforward manner. Friendly gestures can be helpful. Offer a comfortable place to sit or something to eat or drink (but not a hot liquid that could be used as a weapon) to establish trust. Many individuals will decompress at this point because offering food or drink appeals to their most basic human needs and builds trust.<sup>8</sup>

The nurse should demonstrate a nonconfrontational, attentive, and receptive demeanor without conveying weakness or vulnerability. A calm and soothing tone of voice should be used. Avoid direct eye contact, do not approach the client from behind or move suddenly, and stand at least two arm's lengths away.<sup>9</sup> The nurse should ensure a quick exit route to the door and never allow the individual to come between them and the door. Stethoscope and badge holders should not be worn around the neck to prevent strangulation risks.

In some cases, an agitated client may be aware of their impulse control problem and may welcome limit-setting behavior by the nurse (e.g., "I can help you with your problem, but I cannot allow you to continue threatening me or the emergency department staff"). It is difficult to predict which clients

7. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)
8. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)
9. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

will respond to this limit-setting approach. Some clients may interpret such statements as confrontational and escalate their behavior.<sup>10</sup>

A key mistake when interviewing an agitated or potentially violent individual is failing to address violence directly. They should be asked relevant questions, such as, “Do you feel like hurting yourself or someone else?” and “Do you carry a gun?” Stating the obvious (e.g., “You look angry”) may help them to begin sharing their emotions. Speak in a conciliatory manner and offer supportive statements to diffuse the situation, such as, “You obviously have a lot of will power and are good at controlling your emotions.”<sup>11</sup>

A consensus statement from the American Association for Emergency Psychiatry De-escalation Workgroup describes these ten key elements for verbal de-escalation<sup>12</sup>:

- **Respect personal space:** Maintain a distance of two arm’s lengths and provide space for easy exit for either party.
- **Do not be provocative:** Keep your hands relaxed, maintain an open body posture, and do not stare at the client.
- **Establish verbal contact:** The first person to contact the client should take the lead in communicating.
- **Use concise, simple language:** Avoid elaborate and technical terms because they are hard for an impaired person to understand.
- **Identify feelings and desires:** “What are you hoping for?”
- **Listen closely to what the client is saying:** After listening, restate what

10. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

11. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

12. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

the client said to improve mutual understanding (e.g., “Tell me if I have this right...”).

- **Agree or agree to disagree:** Agree with clear specific truths or agree in general (e.g., “Yes, everyone should be treated respectfully.”)
- **Set clear limits:** Inform the client that violence or abuse cannot be tolerated.
- **Offer choices and optimism:** Clients feel empowered if they have some choice in matters.
- **Debrief the client and staff.**

The “philosophy of yes” is a de-escalation technique that encourages the nurse to respond affirmatively to an agitated individual. Examples of initial responses using this approach include “Yes, as soon as,” “Okay, but first we need to,” or “I absolutely understand why you want that done, but in my experience, there are better ways of getting what you need.”<sup>13</sup>

However, some approaches to the combative client are counterproductive and can lead to escalation. Arguing, condescension, or commanding the client to calm down can have disastrous consequences. Clients often interpret such approaches as a challenge to “prove themselves.” A threat to call security personnel can also invite aggression. Other potential mistakes include criticizing or interrupting the client, responding defensively or taking the client’s comments personally, or not clarifying what the client wants before responding.<sup>14</sup>

Never lie to a client (e.g., stating “I am sure you will be out of here in no time” when this is not the case). After the lie becomes apparent, the client may take out frustrations violently upon an unsuspecting nurse or colleague. Take all

13. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

14. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)



threats seriously. It is especially important not to deny or downplay threatening behavior. If verbal techniques are unsuccessful and escalation occurs, the nurse should excuse themselves and summon help.<sup>15</sup>

Many hospitals have multidisciplinary Behavioral Emergency Response Teams (BERTs) that respond to behavioral emergencies. Their overall goal is to provide timely and effective intervention for patients exhibiting disruptive or potentially dangerous behaviors, reducing the risk of harm to both patients and staff.<sup>16</sup>

## Applying Physical Restraints

Physical restraints may be used when verbal de-escalation techniques are unsuccessful despite a professional approach to the combative client. Restraints should never be applied for convenience or punishment, and they should be removed as soon as possible, usually after adequate chemical sedation is achieved or the client has regained control.

Restraints should be implemented systematically using an institutional protocol. The protocol typically begins after the examiner leaves the room when verbal de-escalation techniques have been unsuccessful, and assistance is summoned. It can be helpful to consider the application of restraints like a procedure, similar to running an advanced cardiac life support code.<sup>17</sup>

15. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)
16. Parker, C. M., Calhoun, A., Wong, A. et al. (2020). A call for behavioral emergency response teams in inpatient hospital settings. *AMA Journal of Ethics*, 22(11), E956-964. [doi: 10.1001/amajethics.2020.956](https://doi.org/10.1001/amajethics.2020.956).
17. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

The restraint team should have at least five people, including a team leader. The leader is the only person giving orders and should be the person with the most experience implementing restraints, whether a clinician, nurse, or security officer. Before entering the room, the leader outlines the restraint protocol and warns the team of anticipated dangers (e.g., the presence of objects that may be used as weapons). All team members should remove personal effects (e.g., stethoscopes, pens, jewelry, etc.) the combative client could use against them. If the client to be restrained is female, at least one member of the restraint team should be female to diminish potential allegations of sexual assault.<sup>18</sup>

The restraint team should enter the room in force and display a professional, rather than threatening, attitude. Many violent individuals decompress at this point because the show of force protects their ego (e.g., “I would have fought back, but there were too many against me”). The leader speaks to the client in a calm and organized manner, explaining why restraints are needed and what the course of events will be (e.g., “You will receive a medical and psychiatric examination, as well as treatment”). The client is instructed to cooperate and lie down to have restraints applied. Some clients will be relieved at the protection to self and others afforded by restraints when they feel themselves losing control. However, even if the client suddenly appears less dangerous, physical restraints must be placed after the decision to use them has been made; do not negotiate with the client at this point.<sup>19</sup>

If it becomes necessary to use force to control the client, one team member restrains a preassigned extremity by controlling the major joint (e.g., knee or elbow). This can be accomplished by locking the major joint in extension. The team leader controls the head. If the client is armed with a makeshift weapon,

18. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

19. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

two mattresses can be used to charge and immobilize or sandwich the client. Restraints are applied securely to each extremity and tied to the solid frame of the bed (not side rails because later repositioning of side rails also repositions the client's extremity). To prevent their lower extremities from flailing independently, it may be best to cross the legs at the ankle and then attach the restraint to the bed frame on the opposite side.<sup>20</sup>

Leather is the optimal material for restraining a combative client because it is strong, prevents escape, and is less constricting than typical soft restraints. Gauze should not be used. Soft restraints are helpful in restricting extremity use in a semi-cooperative client but are less effective in a truly violent client who continues to struggle. If chest restraints are used, it is vital to ensure adequate chest expansion for ventilation. Never apply pressure to the client's chest or back while they are immobilized due to asphyxiation risk. After the client is immobilized, announcing "The crisis is over" can have a calming effect on the restraint team and the client.<sup>21</sup>

After restraints have been applied, the client should be monitored frequently and their position changed regularly to prevent circulatory obstruction, pressure injuries, and paresthesias, as well as to avoid rhabdomyolysis associated with continued combativeness. A standardized form based on agency policy is typically used for documentation. Documentation should include the specific indication for restraints and alternatives attempted. In addition to monitoring, nurses must ensure that basic needs (e.g., hydration, food, toileting) are met for any client who is physically restrained or

20. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

21. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

chemically sedated. Physical restraints should be removed as soon as possible.<sup>22</sup>

- ▶ Review ANA guidelines for using restraints in the “[Client Rights](#)” section of the “Legal and Ethical Considerations in Mental Health Care” chapter.

- ▶ View a supplementary YouTube video<sup>23</sup> on how to apply soft wrist restraints: [Restraint Application Techniques for Nurses](#)

## Chemical Sedation

Chemical sedation may be necessary, with or without physical restraints, in a combative client who does not respond to verbal de-escalation techniques. The ideal sedative medication for an agitated or violent client is rapid-acting with minimal side effects. The major classes of medications used to control the violent or agitated client include benzodiazepines, first-generation

22. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

23. NURSINGcom w/ Jon Haws, RN. (2021, October 13). *Restraint application techniques for nurses* [Video]. YouTube. All rights reserved. <https://www.youtube.com/watch?v=S3uPFcU4qRo>

(typical) antipsychotics, second-generation (atypical) antipsychotics, and ketamine<sup>24</sup>:

- For severely violent clients requiring immediate sedation, a rapid-acting first-generation antipsychotic (e.g., haloperidol or droperidol), benzodiazepine (e.g., midazolam), or a combination of both may be prescribed. Second-generation antipsychotics, such as olanzapine, risperidone, and ziprasidone may also be prescribed.
- For clients with agitation from drug intoxication or withdrawal from an unknown cause, benzodiazepines are typically prescribed. Lorazepam and midazolam are used most often. Benzodiazepines may cause respiratory depression and excessive sedation, so close monitoring is essential after administration.
- Ketamine may be prescribed when initial treatments with benzodiazepines or antipsychotics have failed, especially in clients with excited delirium. However, clients receiving ketamine have increased risk for respiratory distress and may require endotracheal intubation and mechanical ventilation.

## Post-Restraint Evaluation

After the client is restrained, the cause of their agitation will be evaluated to determine if it is medical, psychiatric, or substance-use related. Clients over the age of 40 with new psychiatric symptoms are likely to have a medical cause. Elderly clients are at higher risk for delirium due to medical illness (such as a urinary tract infection) or adverse reactions to medications. Clients with a history of drug or alcohol use disorder may exhibit violent behavior as a manifestation of an intoxication or withdrawal syndrome. Violent behavior

24. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

unrelated to medical illness, drug intoxication, or withdrawal should be followed by psychiatric consultation and evaluation.<sup>25</sup>

## Defense Against Assault

Physical assault may occur despite appropriate precautions and interventions with a violent individual. If assaulted, immediately summon help. Maintain a sideward posture, keeping the arms ready for self-protection. If faced with an oncoming punch or a kick, deflect with an arm or a leg. If choking is attempted, tuck in the chin to protect the airway and carotid arteries. If bitten, do not pull away, but rather push toward the mouth and hold the nares shut to entice the opening of the mouth.<sup>26</sup> In a similar manner, don't pull away if your hair is pulled, but instead pull the client's hand toward your head and push up to bend their wrist backwards and increase the likelihood of them releasing their grip due to pain.

If threatened with a weapon, try to appear calm and comply with the individual's demands. Adopt a nonthreatening posture and avoid sudden movements. Do not attempt to reach for the weapon. Avoid arguing, despair, or whining. If taken hostage, attempt to establish a human connection with the hostage taker because there is less risk of violence if a relationship has been established. Do not bargain, make promises, or lie because the consequences could be disastrous. Reassure the hostage taker that an authorized person should arrive promptly to hear their complaints or demands. If a weapon is put down, do not reach for it, but rather attempt to verbally resolve the crisis while awaiting arrival of law enforcement.<sup>27</sup>

- 25. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)
- 26. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)
- 27. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)

Every hospital should have a written plan of action to implement in the case of extreme violence. The plan should include prevention and safety measures, a means for rapid notification of security and police personnel, evacuation plans, medical treatment, and crisis intervention. A novel approach uses a trained violence management team to provide a mechanism for dealing with aggressive clients and to protect the staff.<sup>28</sup>

Mandatory training for clinical and non-clinical staff must also be incorporated with written plans of action. This multifaceted approach improves nurses' self-perception and confidence against workplace violence.<sup>29</sup>

## Interpersonal Conflict Among Health Care Team Members

Conflict is inevitable when working on a health care team composed of members with different personalities, roles, and responsibilities. Some conflicts among team members can escalate to verbal threats or harassment. Common sources of interpersonal conflict in health care settings are passive-aggressiveness, horizontal aggression, defensiveness, peer informer behavior, and victimization behaviors. It is essential for all nurses to develop conflict resolution skills to effectively address these behaviors and maintain a safe work environment.

28. Moore, G. P., & Pfaff, J. A. (2022). Assessment and emergency management of the acutely agitated or violent adult. *UpToDate*. [www.uptodate.com](http://www.uptodate.com)
29. Ming, J. L., Huang, H. M., Hung, S. P., Chang, C. I., Hsu, Y. S., Tzeng, Y. M., Huang, H. Y., & Hsu, T. (2019). Using simulation training to promote nurses' effective handling of workplace violence: A quasi-experimental study. *International Journal of Environmental Research and Public Health*, 16(19), 3648. <http://dx.doi.org/10.3390/ijerph16193648>

- ▶ Read more information about interpersonal conflict and conflict resolution skills in the “[Conflict Resolution](#)” section of the “Collaboration Within the Interprofessional Team” chapter of Open RN *Nursing Management and Professional Concepts*.



## 15.7 Applying the Nursing Process to Clients Experiencing Trauma, Abuse, or Violence

This section will provide an overview of applying the nursing process to clients experiencing trauma, abuse, or violence using trauma-informed care principles. Review specific focused assessments and nursing interventions in the preceding sections.

### Assessment (Recognizing Cues)

Assessment focuses on recognizing physical, psychological, and behavioral indicators of trauma, abuse, or violence. Key actions include the following:

- Recognize injuries inconsistent with explanations (e.g., bruises, fractures, burns).
- Recognize nonverbal cues (e.g., flinching, hypervigilance, avoidance).
- Ensure assessment occurs in a safe, private space, free from potential abusers.
- Screen for anxiety, depression, PTSD, dissociation, or suicidal ideation.

### Diagnosis (Analyzing Cues)

Nurses create nursing diagnoses based on assessment data obtained. Potential nursing diagnoses for clients experiencing trauma, abuse, or violence include the following:

- Risk for Post-Trauma Syndrome
- Ineffective Coping
- Powerlessness
- Impaired Social Interaction
- Sleep Pattern Disturbance
- Chronic Low Self-Esteem

### Outcome Identification (Generate Solutions)

SMART outcomes are identified in relation to the established nursing

diagnoses for each client. SMART is an acronym for Specific, Measurable, Attainable/Actionable, Relevant, and Timely. Read more about outcomes identification in the “[Application of the Nursing Process in Mental Health Care](#)” chapter.

Examples of SMART outcomes for clients experiencing trauma, abuse, or violence include the following:

- *The client will verbalize how to contact a safe person or environment if in crisis by the end of the teaching session.*
- *The client will identify at least one positive coping strategy by the end of the teaching session.*

## Planning (Generate Solutions)

Safety receives top priority when planning and implementing interventions for clients experiencing trauma, abuse, or violence. Nurses help ensure a safety plan is created in collaboration with the client by integrating information about community resources, legal advocacy, and counseling resources. Nurses also help ensure the client’s physical care needs are addressed (e.g., food, housing, pain control, and wound care).

## Implementation (Take Action)

Nurses implement nursing interventions that are therapeutic, respectful, and empowering. Examples of nursing interventions for clients experiencing trauma, abuse, or violence include the following:

- Use grounding techniques to manage dissociation and anxiety.
- Educate client about normal trauma responses and coping tools.
- Offer emotional support and validate their experiences without pressuring for disclosure.
- Facilitate referrals to mental health, social work, domestic violence shelters, or trauma-specific therapy.
- Monitor for suicidal ideation and psychosis.

## Evaluation (Evaluate Outcomes)

Nurses assess the effectiveness of nursing interventions and revise the nursing care plan as needed. They evaluate client's verbal and nonverbal indicators of emotional safety, determine if the expected outcomes were met, and support continuation of trauma-specific therapy.

## 15.8 Spotlight Activity

This Spotlight Activity is based on a real case originally presented in, “Trauma-Informed Care in Nursing Practice” by Dowdell and Speck (2022). Identifying details have been changed or omitted to protect the anonymity of the client.<sup>1</sup>

Dana is a 34-year-old female who arrived at the Emergency Department with a neck injury she reports “happened this morning when I slipped and fell.” The nurse notes that Dana’s address given is in a different community about 50 miles away from the hospital. Initial physical examination revealed bruise patterns in various stages of healing all over Dana’s body and three linear abrasions over the trapezius muscle on the right side of the neck consistent with attempted strangulation.

While awaiting diagnostic testing results, the nurse established a therapeutic nurse-client relationship and asked Dana a few follow-up questions about the events leading up to the injury and the relationship with her partner. Dana stated, “We fight a lot” and “My partner has a lot of angry outbursts.” Upon the nurse’s use of effective therapeutic communication, Dana shared that the injury resulted from “being choked and beaten,” and the partner had done similar actions on “several previous occasions in front of the children.” Dana then told the nurse, “My partner can’t find out that I am here in this hospital.”

A contrast computed tomography (CT) scan revealed swelling of Dana’s right carotid artery and soft tissue in the neck. Laboratory tests also indicated that Dana had severely elevated blood sugars resulting in diabetic ketoacidosis (DKA). The health care team initiated a medication regimen to safely manage the DKA and also prevent a stroke from the injury.

The nurse noted that, over time, Dana became increasingly agitated in the ED

1. Dowdell, E., & Speck, P. (2022). Trauma-informed care in nursing practice. *American Journal of Nursing*, 122(4), 30-38. [https://journals.lww.com/ajnonline/Fulltext/2022/04000/CE\\_Trauma\\_Informed\\_Care\\_in\\_Nursing\\_Practice.22.aspx](https://journals.lww.com/ajnonline/Fulltext/2022/04000/CE_Trauma_Informed_Care_in_Nursing_Practice.22.aspx)

with repetitive neurological assessments, fingersticks for bedside glucose levels, and the noises and high activity level of the ED. Establishing physical and psychological safety for both Dana and staff became a nursing priority during Dana's time in the ED.

Following trauma-informed care (TIC) guidelines to create a safe environment, the nurse asked Dana, "What do you need to feel safe here right now?" Dana immediately responded, "I can't have all these people coming at me. It's too much – too much noise, too many people touching me, it's just too much." The nurse moved Dana out of the trauma bay, which was near the ambulance entryway, and into a room where the nurse was able to close the door. The nurse also posted signage on the door asking all staff and visitors to contact the nurse before entering the room. Following agency protocol, the nurse swiftly gained Dana's consent for anonymity, meaning the client's name and room number(s) would not be shared with anyone outside of the hospital. Following these nursing actions, Dana's anxiety and agitation levels dropped noticeably in the ED. When an ICU bed became available, she was transferred there for medical management.

Dana was present during the handoff report from the ED nurse to the ICU nurse to validate information and it also improved Dana's sense of safety. Following the TIC cue given by the ED nurse, the ICU nurse asked Dana, "What else do you need to feel safe here right now?" Dana requested a private room, if available, to rest, listen to music, and "stay calm." The ICU nurses implemented a safety plan that included consistent staff and coordinated, clustered care. This plan minimized disturbances, thereby providing dedicated periods for rest between timed blood glucose monitoring and nursing monitoring of Dana's neurologic state and neck swelling.

While in the ICU, the nurses taught Dana evidence-based methods for reducing anxiety, including deep breathing, grounding techniques, and moderating anxiety-provoking stimuli such as social media and electronics. In addition, the nurses noted the vocal and nonverbal cues that indicated Dana was feeling anxious, such as speaking in a raised voice or using rapid hand

movements while speaking. Noticing these behaviors allowed nurses to ask follow-up questions such as “Do you feel safe at this moment,” then review anxiety-reducing techniques by asking Dana, “Which technique would you like to use now?” By offering person-centered choices and creating a predictable structure around clustered nursing actions, the nurses promoted a feeling of safety for Dana as well as techniques to self-regulate anxiety.

The health care team identified Dana’s priority health concerns as DKA management and decreasing the risk for a stroke following nonfatal carotid trauma. The nursing team added Dana’s increased risk for subsequent fatal strangulation as a priority concern that must be addressed before discharge. However, when asked by the nurses, “What is important to you to include in your discharge plan,” Dana identified the priorities of finding safe housing and employment with fair pay. Therefore, the nursing team recommended that upon discharge, Dana would transfer directly to community wraparound services for a variety of assistance including transitional housing, job training, day care, medical care, and cognitive behavioral therapy.

### **Reflective Questions**

1. What actions did the nurses perform with Dana to implement Trauma Informed Care (TIC)?
2. What other actions do you plan to implement with your patients regarding TIC?
3. “Dana” could be a male or a female, and Dana’s “partner” could be a male or a female. Do gender differences or sexual orientation affect your attitudes toward Dana’s risks and follow-up care?
4. How might Dana’s children be affected by Dana’s ongoing abuse?
5. What if Dana’s partner showed up at the hospital with a gun threatening to kill Dana? How would you respond as a

nurse to keep yourself, other clients, and staff safe?

## 15.9 Learning Activities



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=1042#h5p-51>

1



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=1042#h5p-50>

2



*An interactive H5P element has been excluded from this version of the text. You can view it*

1. “Trauma, Abuse, and Violence Glossary Cards” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
2. “Trauma, Abuse, and Violence Unfolding Case Study by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)



online here:

<https://wtcs.pressbooks.pub/nursingmhcc/?p=1042#h5p-52>

3

- ▶ Test your clinical judgment with an NCLEX Next Generation-style question: [Chapter 15, Assignment 1](#)<sup>4</sup>



- ▶ Test your clinical judgment with an NCLEX Next Generation-style case study: [Chapter 15, Case Study 1](#)<sup>5</sup>



3. "Trauma, Abuse, and Violence Drag and Drop" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
4. "Trauma, Abuse, and Violence Next Gen Question 1" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
5. "Trauma, Abuse, and Violence Next Gen Case Study" by Kellea Ewen is licensed under [CC BY-NC 4.0](#)

## XV Glossary

**Adults at risk:** Adults who have a physical or mental condition that impairs their ability to care for their own needs.

**Adult Protective Services (APS):** APS agencies are state-run programs that investigate reports of elder and vulnerable adult abuse, neglect, and exploitation.

**Adverse childhood experiences (ACE):** Traumatic experiences during childhood such as neglect, abuse, or witnessing violence, substance abuse, mental illness, divorce, or imprisonment of a family member.

**Caregiver:** A person who has taken responsibility for all or part of an individual's care.

**Dissociation:** A break in how the mind handles information, causing a person's disconnection from their thoughts, feelings, memories, and surroundings.

**Elder abuse:** An intentional act or failure to act that causes or creates a risk of harm to an older adult aged 60 or older.

**Emotional abuse:** Verbal or nonverbal behaviors that inflict anguish, mental pain, fear, or distress, such as humiliation or disrespect, verbal and nonverbal threats, harassment, and geographic or interpersonal isolation.

**Financial abuse:** The illegal, unauthorized, or improper use of money, benefits, belongings, property, or assets for the benefit of someone other than the individual.

**Individual trauma:** Trauma resulting from an event, series of events, or set of circumstances that is experienced by an individual as physically or emotionally harmful or life-threatening and can have lasting adverse effects on their functioning and mental, physical, social, emotional, or spiritual well-being. Adverse childhood experiences are examples of individual traumas.

**Intimate partner violence (IPV):** Abuse or aggression that occurs in a

romantic relationship. IPV can include physical violence, sexual violence, stalking, or psychological aggression.

**Mandated reporters:** Nurses and other professionals required by state law to report suspected neglect or abuse of children, adults at risk, and the elderly they see in the course of their professional duties.

**Neglect:** When a parent or caregiver fails, refuses, or is unable, for reasons other than poverty, to provide the necessary care, food, clothing, or medical or dental care, which seriously endangers the physical health of a child or vulnerable adult.

**Physical abuse:** Injury inflicted on a child or vulnerable adult by other than accidental means. Physical injury includes, but is not limited to, lacerations, fractured bones, burns, internal injuries, severe or frequent bruising, or great bodily harm.

**Resilience:** The ability to rise above circumstances or meet challenges with fortitude. Resilience includes the process of using available resources to negotiate hardship and/or the consequences of adverse events.

**Resilient zone:** A healthy balance of stimulation by the sympathetic nervous system (SNS) and parasympathetic nervous system (PNS).

**Safety plan:** A set of actions that can help lower the risk of a person being hurt by an abusive partner that includes specific information and resources that increases their safety at school, home, and other places visited regularly.

**Sexual abuse:** Sexual intercourse or sexual touching; sexual exploitation; human trafficking; forced viewing of sexual activity; or permitting, allowing, or encouraging prostitution with a child or vulnerable adult.

**Teen dating violence:** Intimate partner violence that occurs during adolescence.

**Trauma-focused cognitive behavioral therapy (TF-CBT):** An evidence-based

treatment that helps children process trauma, develop coping skills, and reduce symptoms of anxiety, depression, or post-traumatic stress.

**Trauma-informed care (TIC):** A strengths-based framework that acknowledges the prevalence and impact of traumatic events in clinical practice, placing an emphasis on instilling in clients a sense of safety, control, and autonomy over their life and health care decisions. The basic goals of TIC are to avoid retraumatization; emphasize survivor strengths and resilience; aid empowerment, healing, and recovery; and promote the development of survivorship skills.

**Workplace violence:** Physically and psychologically damaging actions that occur in the workplace or while on duty. Examples of workplace violence include direct physical assaults (with or without weapons), written or verbal threats, physical or verbal harassment, and homicide.

PART XVI

CHAPTER 16 COMMUNITY ASSESSMENT



### Learning Objectives

- Identify the needs of the populations that impact health-seeking behaviors
- Identify available resources for the health needs of the community
- Explain the nurse role as a collaborative advocate for the health needs of the community
- Apply evidence-based practice when implementing the nursing process in the community
- Differentiate the role of the Registered Nurse in community health care settings

Most of your nursing education journey has likely focused on caring for clients in hospitals and long-term care settings. However, nurses also serve important roles in promoting the health and wellness of communities. Throughout history, nurses have served in public health roles and collaborated with community organizations to support health services. Florence Nightingale, the founder of modern nursing, advocated for two foundational components of community health nursing: health promotion and disease prevention.<sup>1</sup> Examples of current community health initiatives include public educational health sessions, blood pressure screenings,

1. McDonald, L. (2006). *Florence Nightingale and public health policy: Theory, activism and public administration*. University of Guelph.  
<https://cwfn.uoguelph.ca/nursing-health-care/fn-and-public-health-policy/>

immunization clinics, and crisis intervention related to mental health care. See Figure 16.1<sup>2</sup> for an image of a community blood pressure screening.



Figure 16.1 Community Health Screening

In recent years community health nursing has become increasingly important for several reasons:

- There are currently shorter hospital stays and decreasing hospital readmission rates.
- As the average age of the United States population increases, the need for community health services in members' homes increases.
- As the ability to successfully manage chronic conditions (including both physical and mental illness) improves, the need for outpatient health services also increases.
- As the usage of telehealth and virtual medical care expands, so does the need for community member support and client education.

2. "[7315182496\\_0cd3e18ed6\\_k](#)" by [Maryland GovPics](#) is licensed under [CC BY 2.0](#)



This chapter will introduce the roles of community health nurses, explore community health needs assessments, and discuss how nurses collaborate with community resources to support clients' physical and mental health care needs.

## 16.2 Community Health Concepts

### Community Health Nursing

Community health nurses work in diverse settings, including public health agencies, schools, and faith-based organizations. Their roles vary but share a common goal: promoting health and addressing the needs of populations within their communities.

**Public health nurses** work across various settings in the community such as government agencies, community-based centers, shelters, and vaccine distribution sites. They provide disease prevention and health promotion services, such as working with mothers and children to improve nutrition, operating immunization clinics, and leading public health education initiatives such as smoking cessation campaigns. Public health nurses also prepare to respond quickly to public health emergencies such as natural disasters or epidemics.<sup>12</sup> Emergency preparedness is further discussed in the “[Environmental Health and Emergency Preparedness](#)” chapter. See Figure 16.2<sup>3</sup> for an image of a community health nurse providing health screenings in a maternal and child health clinic.

1. Greenwood, B. (n.d.). *What are the primary roles of the community nurse?* CHRON. <https://work.chron.com/primary-roles-community-nurse-15144.html>
2. National Academy of Medicine. (2021). *The future of nursing 2020-2030: Charting a path to achieve health equity* [PDF Report]. <https://www.phnurse.org/assets/docs/FON%20Valuing%20Community%20and%20Public%20Health%20Nursing.pdf>
3. “[PIXNIO-45563-3000×2000](#)” by [USAID](#) on [Pixnio](#) is licensed under [CC0](#)



Figure 16.2 Health Screenings

**School nurses** provide a specialized practice of nursing which protects and promotes student health, facilitates optimal development, and advances academic success. School nurses, grounded in ethical and evidence-based practice, are the leaders who bridge health care and education, provide care coordination, advocate for quality student-centered care, and collaborate to design systems that allow individuals and communities to develop their full potentials. School nurses provide direct healthcare, administer medications and treatments, develop health care plans and emergency action plans, provide health screenings, collaborate with parents and medical providers and promote student health and safety and so much more. For example, school nurses may help develop disaster plans that coordinate activities in the school with the larger neighborhood.<sup>4,5</sup>

4. Greenwood, B. (n.d). *What are the primary roles of the community nurse?* CHRON. <https://work.chron.com/primary-roles-community-nurse-15144.html>
5. National Academy of Medicine. (2021, May). *The future of nursing 2020-2030:*

**Parish nurses** also known as faith community nurses, integrate professional nursing with spiritual care to promote holistic health within faith-based communities. Serving in both paid and volunteer roles, they work primarily within churches, addressing the physical, emotional, and spiritual needs of their congregations. Their responsibilities encompass administering vaccinations, assisting with end-of-life care discussions, coordinating among healthcare providers, and leading support groups. Additionally, they visit congregation members in homes or hospitals, offering spiritual support during health challenges. By acting as liaisons between healthcare systems and congregations, parish nurses advocate for quality, client-centered care while honoring individuals' spiritual beliefs.

Beyond church settings, parish nurses contribute to community health by working in hospitals, long-term care facilities, medical clinics, and social service agencies. In hospitals and long-term care environments, they often collaborate with chaplains to provide comprehensive spiritual care. Their unique role emphasizes integrating faith and health, transforming faith communities into sources of healing and support.<sup>6</sup>

## Barriers for Community Health Nurses

Community health nurses play a crucial role in identifying the health needs of a community and planning and implementing preventative health initiatives. However, they often face specific challenges when providing care to individuals, families, and community members. Three key barriers are gaining entry, role negotiation, and confidentiality:

*Charting a path to achieve health equity* [PDF Report].

<https://www.phnurse.org/assets/docs/FON%20Valuing%20Community%20and%20Public%20Health%20Nursing.pdf>

6. Greenwood, B. (n.d.). *What are the primary roles of the community nurse?* CHRON. <https://work.chron.com/primary-roles-community-nurse-15144.html>

- **Gaining Entry:** Community health nurses may be considered “outsiders” as representatives of the established health care system and may not necessarily be trusted by community members. It is vital for community health nurses to build trust and supportive relationships. When working with individuals and families, nurses should assess specific community issues affecting that individual's health or their access to health care and then address those issues in their nursing care plan. Nurses can also investigate if there are community resources available to refer the client and/or their family members for additional services. See Figure 16.3<sup>7</sup> for an image of Red Cross volunteers working with caregivers of clients receiving palliative care.
- **Role Negotiation and Confidentiality:** Community health nurses must separate their roles as data collectors, health professionals, and neighbors. These roles can be difficult to differentiate when the nurse is assessing community health needs and providing nursing interventions for a population of individuals within their own home community. These individuals may include family members, friends, neighbors, or peers. Trust must be established and confidentiality assured according to legal and ethical parameters of nursing practice. Nurses should also establish a sense of partnership and encourage clients to participate in planning preventative health strategies for themselves and their families.

7. “10716898813\_74292ef548\_k” by [Department of Foreign Affairs and Trade](#) is licensed under [CC BY 2.0](#)



Figure 16.3 Setting Up Support Groups for Caregivers

Other barriers that community health nurses may experience include limited resources and funding, workforce shortages, staff burnout, and feelings of isolation from the healthcare community.<sup>8</sup>

## Community Health

Community health nursing is grounded in key principles: promoting healthy living, preventing illness, facilitating rehabilitation, evaluating community resources, and advocating for overall health improvement.<sup>9,10</sup> In this manner,

8. Seyedfatemi, N., Peyrovi, H., Goharinezhad, S., & Oghli, S. H. (2024). Barriers and facilitators of community-based nursing development: A scoping review. *Journal of Education and Health Promotion*, 3:488. [doi: 10.4103/jehp.jehp\\_1329\\_23](https://doi.org/10.4103/jehp.jehp_1329_23).
9. Greenwood, B. (n.d.). *What are the primary roles of the community nurse?* CHRON. <https://work.chron.com/primary-roles-community-nurse-15144.html>
10. National Academy of Medicine. (2021). *The future of nursing 2020-2030: Charting a path to achieve health equity* [PDF Report].



community health nurses pursue health equity. **Health equity** means that everyone has a fair and just opportunity to be as healthy as possible. This requires removing potential obstacles to obtaining and maintaining optimal health such as lack of access to health care services, good jobs with fair pay, quality education and housing, and safe environments.<sup>11</sup> Community health nurses address these conditions that are also known as social determinants of health.

**Social determinants of health (SDOH)** are the conditions in which people are born, grow, work, live, and age. Research shows that the SDOH can be more important than health care or lifestyle choices in influencing health and account for 30-55% of health outcomes.<sup>12</sup> See an illustration of SDOH in Figure 16.4.<sup>13</sup> SDOH can contribute to **health inequities**, defined as avoidable differences in health status seen within and between communities. In countries at all levels of income, health and illness follow a social gradient: the lower the socioeconomic position, the worse the health. According to the

<https://www.phnurse.org/assets/docs/FON%20Valuing%20Community%20and%20Public%20Health%20Nursing.pdf>

11. Braveman, P., Arkin, E., Orleans, T., Proctor, D., & Plough, A. (2017, May 1). *What is health equity?* Robert Wood Johnson Foundation. <https://www.rwjf.org/en/library/research/2017/05/what-is-health-equity-.html>
12. World Health Organization. (n.d.). *Social determinants of health*. [https://www.who.int/health-topics/social-determinants-of-health#tab=tab\\_1](https://www.who.int/health-topics/social-determinants-of-health#tab=tab_1)
13. "Healthy People 2030 SDOH Graphic.png" by [U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion](#) is in the [Public Domain](#). Access for free at <https://health.gov/healthypeople/objectives-and-data/social-determinants-health>


World Health Organization, SDOH can influence health equity in positive and negative ways<sup>14</sup> :

- Income and social protection
- Education
- Unemployment and job insecurity
- Working life conditions
- Food insecurity
- Housing, basic amenities, and the environment
- Early childhood development
- Social inclusion and nondiscrimination
- Structural conflict
- Access to affordable health services of decent quality

14. World Health Organization. (n.d.). *Social determinants of health*.  
[https://www.who.int/health-topics/social-determinants-of-health#tab=tab\\_1](https://www.who.int/health-topics/social-determinants-of-health#tab=tab_1)



# Social Determinants of Health



Social Determinants of Health  
Copyright-free

Healthy People 2030

Figure 16.4 Social Determinants of Health

**Health disparities** are health differences that are linked with social, economic, and/or environmental disadvantages. Health disparities adversely affect groups of people who often experience greater obstacles to health based on individual characteristics such as socioeconomic status, age, gender, culture, religion, mental illness, disability, sexual orientation, or gender identity.<sup>15</sup> These groups are often referred to as “vulnerable groups,” and their care is further discussed in the “[Vulnerable Populations](#)” chapter.

Community health nurses aim to promote health equity by addressing social determinants of health (SDOH), health disparities, and inequities in the

15. HealthyPeople.gov. (2022, February 6). *Disparities*. Office of Disease Prevention and Health Promotion, U.S. Department of Health and Human Services. <https://odphp.health.gov/healthypeople>

communities they serve. They begin by assessing the community context, as it influences how members respond to public health initiatives. Key community context factors include:

- **Physical aspects:** Consider the geographic and man-made features present in the community. How do these features influence residents' access to healthcare and essential services, as well as the community health nurse's ability to reach those residents?
  - Example: Is the community rural, requiring residents to travel long distances for medical care?
- **Infrastructure:** Assess the condition of local roads, bridges, and public transit options, as well as the availability of utilities like internet, electricity, and cellular service.
  - Example: Are buses or rideshare services available for residents to reach healthcare facilities?"
- **Patterns of settlement, commerce, and industry:** Examine how residential and commercial areas are distributed. Are there neighborhoods located near heavy industry, areas with higher crime rates or defined by economic status?
  - Example: Is there a residential area adjacent to a factory that may contribute to air pollution?"
- **Demographics:** Evaluate key population characteristics such as age, gender, race, ethnicity, language, and household composition.
  - Example: Does the community feature multi-generational households?
- **History:** Reflect on the historical events and traditions that shape community identity.
  - Example: Are there monuments or parks that honor local historical events or figures?

- **Community leaders, both informal and formal:** Understand the shared norms, traditions, and values that guide behavior in the community.
  - Example: Is there an annual celebration or parade that reinforces community identity?
- **Community culture, both informal and formal:** What are the spoken and unspoken rules and traditions of the community?
  - For example, does the community have a tradition of celebrating Memorial Day with a parade and public tributes at a local cemetery?
- **Existing groups and organizations:** Consider the presence of local clubs, community centers, and organizations that can facilitate collaboration and support.
  - Example: Is there a YMCA or similar organization promoting physical activity?
- **Existing institutions:** Identify essential institutions—such as hospitals, clinics, schools, libraries, and religious centers—and assess their accessibility and leadership.
  - Example: Is there an urgent care clinic easily accessible by public transit?
- **Economics:** Review the community's economic base, including predominant industries and employment patterns, as well as wealth distribution.
  - Example: Is the area predominantly working class, with most residents employed in local factories?
- **Government/Politics:** Consider the influence and engagement of political leaders and the nature of governance in the community.
  - Example: How does the mayor communicate with residents?
- **Social structure:** Analyze how community members interact on a daily

basis, including conflict resolution and symbols of respect.

- Example: Does the town board meet regularly to discuss local issues and make decisions collaboratively?
- **Attitudes and values:** Investigate what the community prioritizes and any prevailing assumptions or biases regarding behavior and interaction.
  - Example: Is it commonly expected that neighbors will help each other during community challenges, such as after a storm?

According to the CDC, a **healthy community** is one in where local groups work together to prevent disease and promote healthy living. This approach benefits the largest number of people and helps reduce health disparities caused by differences in income, education, race, location, and other factors.<sup>16</sup>

## Community Health Needs Assessment

**Community health needs assessment** is a systematic process to identify and analyze community health needs and assets in order to prioritize these needs, plan, and act upon significant unmet community health needs.<sup>17</sup> A community health assessment gives nurses and community organizations comprehensive information about the community's current health status, needs, and issues. This information can be used to develop a community health improvement plan by justifying how and where resources should be

16. Centers for Disease Control and Prevention. (2015, September 18). *A healthy community is a prepared community*. [Blog]. <https://blogs.cdc.gov/publichealthmatters/2015/09/a-healthy-community-is-a-prepared-community/>

17. Vigna, A. J. (2020). *The 2019 behavioral health gaps report for the state of Wisconsin*. University of Wisconsin Population Health Institute. <https://uwphi.pophealth.wisc.edu/publications-2/evaluation-reports-2/>

allocated to best meet community needs.<sup>18</sup> Community health needs assessments are performed and reported at national, state, county, and local levels.

## National Health Needs Assessments


Healthy People 2030 addresses the most current national public health priorities. It is published by the United States Department of Health and Human Services Office of Disease Prevention and Promotion. See Figure 16.5<sup>19</sup> for an illustration related to using Healthy People 2030 objectives and leading health indicators to assess community needs data, plan, and evaluate community health interventions.

18. VHA Inc., & Healthy Communities Institute. (2013). *Assessing & addressing community health needs*. Catholic Heart Association of the United States. [https://www.chausa.org/docs/default-source/general-files/cb\\_assessingaddressing-pdf.pdf?sfvrsn=4](https://www.chausa.org/docs/default-source/general-files/cb_assessingaddressing-pdf.pdf?sfvrsn=4)

19. "HP2030\_HowtoUse\_Screen1\_2020-03-24.png" by unknown author for [U.S. Department of Health and Human Services](https://www.hhs.gov/) is in the [Public Domain](https://www.hhs.gov/). Access for free at <https://health.gov/our-work/national-health-initiatives/healthy-people/healthy-people-2030/promoting-healthy-people-2030>.

## How can I use Healthy People 2030 in my work?

Healthy People addresses public health priorities by setting national objectives and tracking them over the decade. Join us as we work to improve health and well-being nationwide.



Healthy People 2030



ODPHP

Office of Disease Prevention  
and Health Promotion

Figure 16.5 Healthy People 2030

A set of evidence-based Healthy People objectives are published every ten years based on current national data. Objectives are classified by categories<sup>20</sup>:

- Health Conditions
- Health Behaviors
- Populations
- Settings and Systems
- Social Determinants of Health

Examples of Healthy People 2030 community objectives include the following<sup>21</sup>:

20. Office of Disease Prevention and Health Promotion. *Healthy People 2030: Building a healthier future for all*. U.S. Department of Health and Human Services. <https://health.gov/healthypeople>
21. Office of Disease Prevention and Health Promotion. *Healthy People 2030: Building a healthier future for all*. U.S. Department of Health and Human Services. <https://health.gov/healthypeople>

- Increase the number of community organizations that provide preventative services
- Increase the rate of bystander CPR and AED use for nontraumatic cardiac arrests in public places
- Increase the proportion of adult stroke survivors who participate in rehabilitation services

Leading Health Indicators (LHIs) are a subset of high-priority Healthy People 2030 objectives to drive action toward improving health and well-being. Most LHIs address important factors that impact major causes of death and disease in the United States. They help organizations, communities, and community health nurses focus their resources and efforts to improve the health and well-being of all people. There are 23 LHIs that cover the life span from infants, children, adolescents, adults, and older adults. See a list of LHIs in Table 16.2.

Table 16.2 Leading Health Indicators Across the Life Span

Life Stage	Leading Health Indicator
Infant	<ul style="list-style-type: none"> <li>• Infant Death</li> </ul>
Children and Adolescents	<ul style="list-style-type: none"> <li>• 4th grade students whose reading skills are at or above the proficient achievement level for their grade</li> <li>• Adolescents with major depressive episodes (MDEs) who receive treatment</li> <li>• Children and adolescents with obesity</li> <li>• Current use of any tobacco products among adolescents</li> </ul>



**Adults and  
Older Adults**

- Adults engaging in binge drinking of alcoholic beverages during the past 30 days
- Adults who meet current minimum guidelines for aerobic physical activity and muscle-strengthening activity
- Adults who receive a colorectal cancer screening based on the most recent guidelines
- Adults with hypertension whose blood pressure is under control
- Cigarette smoking in adults
- Employment among the working-age population
- Maternal deaths
- New cases of diagnosed diabetes in the population

<b>All Ages</b>	<ul style="list-style-type: none"> <li>• Children, adolescents, and adults who use the oral health care system (2+ years)</li> <li>• Consumption of calories from added sugars by persons aged 2 years and over</li> <li>• Drug overdose deaths</li> <li>• Exposure to unhealthy air</li> <li>• Homicides</li> <li>• Household food insecurity and hunger</li> <li>• Persons who are vaccinated annually against seasonal influenza</li> <li>• Persons who know their HIV status (13+ years)</li> <li>• Persons with medical insurance (&lt;65 years)</li> <li>• Suicides</li> </ul>
-----------------	--

### Healthy People 2030 Resources

- ▶ View [Healthy People 2030 Objectives](#), [Community Objectives](#), and [Leading Health Indicators](#).

## State Needs Assessments

States perform health needs assessments to develop state funding and

program priorities for community health. For example, the Wisconsin Department of Health Services (DHS) performs a mental health and substance abuse needs assessment every other year. Data in this report includes the following:

- **Prevalence of Needs:** The prevalence of disorders, conditions, and associated problems for the entire population and subpopulations
- **Access to Services:** Determination of which and how many services are received by individuals and exploration of barriers to access
- **Service Workforce and Capacity:** Examination of the mental health and substance use services workforce, including the number of providers of these services and the geographic dispersion of the workforce across the state

Another example of a state needs assessment related to mental health is the Behavioral Health Gaps Study funded by the Wisconsin DHS to assess gaps and needs in the behavioral health service system for individuals with mental health and substance use disorders. Key gaps documented in this study included shortages in child and geriatric psychiatrists; shortages in mental health inpatient beds and residential facilities for treating substance use; inadequacies of the medical transportation system; a need for improving crisis stabilization services in the community that focus on reducing contact with police officers; shortages in medication-assisted treatment providers and clinics; long waitlists across the service array; shortages in competent translation services; and the need to provide wraparound services, particularly for consumers with families.<sup>22</sup>

22. Vigna, A. J. (2020). *The 2019 behavioral health gaps report for the state of Wisconsin*. University of Wisconsin Population Health Institute.  
<https://uwphi.pophealth.wisc.edu/publications-2/evaluation-reports-2/>

Explore your state's health needs assessments. Examples of health needs assessments in the state of Wisconsin are as follows:

- ▶ Department of Health Service's [Mental Health and Substance Abuse Needs Assessment PDF](#)
- ▶ [The Behavioral Health Gaps Report for the State of Wisconsin PDF](#) completed by University of Wisconsin-Madison Population Health Institute

## County Health Rankings

County health rankings are created annually by the University of Wisconsin Population Health Institute for counties across the country.<sup>23</sup> These rankings provide a snapshot of a community's health and can be used as a starting point for implementing change to promote health equity in communities. See the following box to explore the health ranking for your community.

- ▶ Visit the [Explore Health Rankings](#) web page from the University of Wisconsin Population Health Institute to view the health ranking of your community.

23. Vigna, A. J. (2020). *The 2019 behavioral health gaps report for the state of Wisconsin*. University of Wisconsin Population Health Institute. <https://uwphi.pophealth.wisc.edu/publications-2/evaluation-reports-2/>

## Local Needs Assessments

Local communities perform health needs assessments and develop specific health initiatives for their community members. For example, the Eau Claire County Health Department Needs Assessment was used to create a county health plan. In 2021 the top health priorities were documented as drug use, mental health, alcohol misuse, obesity, and healthy nutrition. See Figure 16.6<sup>24</sup> with an image related to data from a county mental health needs assessment.

### *Why is Mental Health a problem in our county?*




Figure 16.6 Mental Health Assessment

- ▶ Explore your community's health initiatives. View the example of [Eau Claire City-County Health Department Needs Assessment](#).

24. This image is derived from [2021 Eau Claire County Community Health Assessment](#) by Community Health Assessment Planning Partnership Committee and is in the [Public Domain](#)

# Hospitals' Community Health Needs Assessments

Tax-exempt hospitals are required to conduct community health needs assessments according to the Patient Protection and Affordable Care Act (i.e., the Affordable Care Act). Hospitals are required to adopt implementation strategies to meet the community health needs identified through their needs assessment. This collaboration among hospitals and community partners expands the community's capacity to address health needs through a shared vision and creates a foundation for coordinated efforts to improve community health.<sup>25</sup>

25. VHA Inc., & Healthy Communities Institute. (2013). *Assessing & addressing community health needs*. Catholic Heart Association of the United States. [https://www.chausa.org/docs/default-source/general-files/cb\\_assessingaddressing-pdf.pdf?sfvrsn=4](https://www.chausa.org/docs/default-source/general-files/cb_assessingaddressing-pdf.pdf?sfvrsn=4)

## 16.3 Applying the Nursing Process to Community Health

Community health nurses apply the nursing process to address needs of individuals, families, vulnerable populations, and entire communities. See Figure 16.7<sup>1</sup> for an illustration of the nursing process in community health nursing.



Figure 16.7 Nursing Process in Community Health Nursing

### Assessment (Recognizing Cues)

The community health nurse typically begins a community health needs assessment by determining what data is already available.<sup>2</sup> As previously

1. "[Nursing Process in Community Health Nursing](#)" by [Open RN](#) project is licensed under [CC BY 4.0](#)
2. [Community Tool Box](#) by [Center for Community Health and Development](#) at the University of Kansas is licensed under [CC BY NC SA 3.0](#)

discussed in the “[Community Health Concepts](#)” section, national, state, county, and local health needs assessments are widely available. **Secondary analysis** refers to analyzing previously collected data to determine community needs.

Community health nurses may also engage in primary data collection to better understand the community needs and/or study who may be affected by actions taken as a result of the assessment.<sup>3</sup> **Primary data collection** includes tools such as public forums, focus groups, interviews, windshield surveys, surveys, and participant observation.

## Public Forums

**Public forums** are gatherings where large groups of citizens discuss important issues at well-publicized locations and times. Forums give people of diverse backgrounds a chance to express their views and enhance understanding of the community’s specific needs and resources. Forums should be planned in a convenient location with accessibility to public transportation and child care. They should also be scheduled at convenient times for working families to gain participation from a wide range of populations.

## Focus Groups

**Focus groups** are a systematic method of data collection through small-group discussions led by a facilitator. Participants in focus groups are selected to represent a larger group of people. Groups of 6-10 people with similar backgrounds or interests are interviewed in an informal or formal setting. Focus groups should be scheduled at several dates and times to ensure a

3. [Community Tool Box](#) by [Center for Community Health and Development](#) at the University of Kansas is licensed under [CC BY NC SA 3.0](#)



broad participation from members of the community. Here are advantages of focus groups:

- Community member involvement in assessing and planning community initiatives is encouraged.
- Different perceptions, values, and beliefs by community members are explored.
- Input can be obtained from specific subpopulations of the community. Examples of subpopulations include young mothers caring for infants, individuals receiving home hospice care, individuals struggling to find housing, residents of the prison system, individuals coping with mental health disorders, or residents in group homes.

## Interviews

**Interviews** are structured conversations with individuals who have experience, knowledge, or understanding about a particular topic or issue.

**Key informant interviews** are conducted with people in key positions in the community and have specific areas of knowledge and experience. These interviews can be useful for exploring specific community problems and/or assessing a community's readiness to address those problems.<sup>4</sup>

Advantages of interviews include the following<sup>5</sup>:

- They can be conducted in a variety of settings (e.g., homes, schools, churches, stores, or community centers).

4. [A Guide to SAMHSA's Strategic Prevention Framework PDF](#) by [Substance Abuse and Mental Health Services Administration](#) is available in the [Public Domain](#).

5. [A Guide to SAMHSA's Strategic Prevention Framework PDF](#) by [Substance Abuse and Mental Health Services Administration](#) is available in the [Public Domain](#).

- They are low cost and generally have low dropout rates.
- Respondents define what is important from their perspective.
- It is possible to explore issues in depth, and there is an opportunity to clarify responses.
- They can provide leads to other data sources and key informants.
- They provide an opportunity to build partnerships with community members.
- Data can be compared among local government officials, citizens, and non-government leaders.

Interviews can have these disadvantages:

- Interviews can be time-consuming to schedule and perform.
- They require trained interviewers.
- There is a potential for interviewer bias to affect the data collected during the interview.
- Rapport must be established before sensitive information is shared.
- It is more time-consuming to summarize and analyze findings.

## Windshield Surveys

A **windshield survey** is a type of direct observation of community needs while driving and literally looking through the windshield. It can be used to observe characteristics of a community that impact health needs such as housing, pollution, parks and recreation areas, transportation, health and social services agencies, industries, grocery stores, schools, and religious institutions.



View the following YouTube video of a windshield survey<sup>6</sup>:  
[Windshield Survey Nursing](https://youtu.be/aAzW1bW_Dbw).

## Surveys

**Surveys** use standardized questions that are relatively easy to analyze. They are beneficial for collecting information across a large geographic area, obtaining input from as many people as possible, and exploring sensitive topics.<sup>7</sup> Surveys can be conducted face to face, via the telephone, mailed, or shared on a website. Responses are typically anonymous but demographic information is often collected to focus on the needs of specific populations. Disadvantages of surveys can include the following<sup>8</sup>:

- Surveys can be time-consuming to design, implement, and analyze the results.
- The accuracy of survey results depends on who is surveyed and the size of the sample.
- Mailed surveys may have low response rates with higher costs due to postage.

6. Medrea, R. (2014, July 20). *Windshield survey nursing* [Video]. YouTube. All rights reserved. [https://youtu.be/aAzW1bW\\_Dbw](https://youtu.be/aAzW1bW_Dbw)

7. McDonald, L. (2006). *Florence Nightingale and public health policy: Theory, activism and public administration*. University of Guelph. <https://cwfn.uoguelph.ca/nursing-health-care/fn-and-public-health-policy/>

8. [A Guide to SAMHSA's Strategic Prevention Framework PDF](#) by Substance Abuse and Mental Health Services Administration is available in the [Public Domain](#).

- They offer little opportunity to explore issues in depth, and questions cannot be clarified.
- There is no opportunity to build rapport with respondents.

## Participant Observation

Participant observation refers to nurses informally collecting data as a member of the community in which they live and work. This is considered a subjective observation because it is from the nurse's perspective. Informal observations are made, or discussions are elicited among peers and neighbors within the community.

## Sociocultural Considerations

When analyzing community health needs, it is essential to do so through a sociocultural lens. Just as an individual's health can be influenced by a wide variety of causes, community health problems are affected by various factors in the community. For example, a high rate of cancer in one community could be related to environmental factors such as pollution from local industry, but in another community, it may be related to the overall aging of the population. Both communities have a high rate of cancer, but the public health response would be very different. Another example related to mental health is related to various situational factors affecting depression. A high rate of depression in one community may be related to socioeconomic factors such as low-paying jobs, lack of support systems, and poor access to basic needs like grocery stores, whereas in another community it may be related to lack of community resources during frequent weather disasters. The public health response would be different for these two communities.

Nurses must also recognize and value cultural differences such as health beliefs, practices, and linguistic needs of diverse populations. They must take steps to identify subpopulations who are vulnerable to health disparities and further investigate the causes and potential interventions for these disparities. For example, mental health disparities pose a significant threat to

vulnerable populations in our society, such as high rates of suicide among LGBTQ+ youth, reduced access to prevention services among people living in rural areas, and elevated rates of substance misuse among Native Americans. These disparities threaten the health and wellness of these populations.<sup>9</sup>

Key points to consider when assessing a community using a sociocultural lens include the following:

- Have the trends of assessment data changed over time? What are the potential causes for these changes in this community?
- How does the community's needs assessment data compare to similar communities at local, county, state, and national levels? What target goals and health initiatives have been successfully implemented in other communities?
- What vulnerable subpopulations are part of this community, and what health disparities are they experiencing? What are potential causes and solutions for these health disparities?
- Input from members of vulnerable subpopulations must be solicited regarding their perspectives on health disparities, as well as barriers they are experiencing in accessing health care.

## Diagnosis (Analyze Cues)

Similar to how nurses individualize nursing diagnoses for clients based on priority nursing problems identified during a head-to-toe assessment, community health nurses use community health needs assessment data to develop community health diagnoses. These diagnoses are broad, apply to larger groups of individuals, and address the priority health needs of the community. Resources such as Healthy People 2030 can be used to determine current public health priorities.

9. [A Guide to SAMHSA's Strategic Prevention Framework PDF](#) by [Substance Abuse and Mental Health Services Administration](#) is available in the [Public Domain](#).

A **community diagnosis** is a summary statement resulting from analysis of the data collected from a community health needs assessment.<sup>10</sup> A clear statement of the problem, as well as causes of the problem, should be included. A detailed community diagnosis helps guide community health initiatives that include nursing interventions.

A community diagnosis can address health deficits or services that support health in the community. A community diagnosis may also address a need for increased wellness in the community. Community diagnoses should include these four parts:

- The problem
- The population or vulnerable group
- The effects of the problem on the population/vulnerable group
- The indicators of the problem in this community

Here are some examples of community health diagnoses based on community health needs assessments:

- **Community Scenario A**

- **Assessment data:** The local high school has had a 50% increase in the number of teen pregnancies in the past year, causing high school graduation rates to decrease due to pregnant students dropping out of high school.
- **Community diagnosis:** Increased need for additional birth control and resources for prevention of pregnancy due to lack of current resources, as evidenced by 50% increase in teen pregnancies in the last year and a decrease in graduation rates.

10. Office of Disease Prevention and Health Promotion. *Healthy People 2030: Building a healthier future for all*. U.S. Department of Health and Human Services. <https://health.gov/healthypeople>

- **Community Scenario B**

- **Assessment data:** Fifty percent of residents of an assisted living facility were found to have blood pressure readings higher than 130/80 mmHg during a health fair last week at the facility.
- **Community diagnosis:** Increased need for education about exercise and diet and referrals to primary care doctors for residents of an assisted living facility due to increased risk for mortality related to high blood pressure, as evidenced by a high number of residents with high blood pressure during a health fair.

- **Community Scenario C**

- **Assessment data:** The local high school has had two cases of suicide in the past year.
- **Diagnosis:** Increased need for community education regarding suicide prevention and crisis hotlines, as evidenced by an increase in adolescent suicide over the past twelve months.

## Outcome Identification (Generate Solutions)

Outcomes refer to the changes in communities that nursing interventions and prevention strategies are intended to produce. Outcomes include broad overall goals for the community, as well as specific outcomes referred to as “SMART” outcomes that are specific, measurable, achievable, realistic, and with a timeline established.

Broad goals for communities can be tied to national objectives established by Healthy People 2030, as previously discussed in the “[Community Health Concepts](#)” section.

Healthy People objectives are classified by these five categories<sup>11</sup>:

- Health Conditions
- Health Behaviors
- Populations
- Settings and Systems
- Social Determinants of Health

SMART outcomes can be created based on the objectives listed under each category. For example, if an overall community goal is related to “Drug and Alcohol Use” under the “Health Behaviors” category, a SMART outcome could be based on the Healthy People objective, “Increase the proportion of people with a substance use disorder who got treatment in the past year.”<sup>12</sup> Based on this Healthy People objective, an example of a SMART outcome could be the following:

- The proportion of people treated for a substance disorder in Smith County will increase to 14% within the next year.

▶ View the [Healthy People 2030 Objectives](#) and [Community Objectives](#).

11. Office of Disease Prevention and Health Promotion. *Healthy People 2030: Building a healthier future for all*. U.S. Department of Health and Human Services. <https://health.gov/healthypeople>

12. Office of Disease Prevention and Health Promotion. *Healthy People 2030: Building a healthier future for all*. U.S. Department of Health and Human Services. <https://health.gov/healthypeople>



## Planning (Generate Solutions)

Nursing interventions for the community can be planned based on the related Healthy People category and objective. For example, based on the sample SMART outcome previously discussed, a planned nursing intervention could be the following:

- The nurse will provide education and materials regarding evidence-based screening practices for substance use disorder in local clinics.

Community health nursing interventions typically focus on prevention of illness with health promotion interventions. After performing a community health needs assessment, identifying priority problems, and establishing health goals and SMART outcomes, the nurse integrates knowledge of health disorders (e.g., diabetes, cancer, obesity, or mental health disorders) and current health risks in a community to plan prevention interventions.

There are two common public health frameworks used to plan prevention interventions. A traditional preventive framework is based on primordial, primary, secondary, tertiary, or quaternary prevention interventions. A second framework, often referred to as the Continuum of Care Prevention Model, was established by the Institute of Medicine (IOM) and includes universal, selected, and indicated prevention interventions. Both frameworks are further discussed in the following sections.<sup>13</sup>

### Primordial, Primary, Secondary, Tertiary, and Quaternary Interventions

Preventive health interventions may include primordial, primary, secondary, tertiary, and quaternary prevention interventions. These strategies attempt to

13. Savage, C. L. (2020). *Public/community health and nursing practice: Caring for populations* (2nd ed.). FA Davis.

prevent the onset of disease, reduce complications of disease that develops, and promote quality of life.<sup>14</sup>

## PRIMORDIAL PREVENTION

**Primordial prevention** consists of risk factor reduction strategies focused on social and environmental conditions that affect vulnerable populations. In other words, primordial prevention interventions target underlying social determinants of health that can cause disease. These measures are typically promoted through laws and national policy. An example of a primordial prevention strategy is improving access to urban neighborhood playgrounds to promote physical activity in children and reduce their risk for developing obesity, diabetes, and cardiovascular disease.<sup>15</sup> See Figure 16.8<sup>16</sup> for an image of a neighborhood playground.

14. This work is a derivative of [StatPearls](#) by Kisling and Das and is licensed under [CC BY 4.0](#)

15. This work is a derivative of [StatPearls](#) by Kisling and Das and is licensed under [CC BY 4.0](#)

16. "[Playground at Hudson Springs Park.jpg](#)" by [Kevin Payravi](#) is licensed under [CC BY-SA 3.0](#)



Figure 16.8 Primordial Prevention: Neighborhood Playgrounds

## PRIMARY PREVENTION

**Primary prevention** consists of interventions aimed at susceptible populations or individuals to prevent disease from occurring. An example of primary prevention is immunizations.<sup>17</sup> Nursing primary prevention interventions also include public education and promotion of healthy behaviors.<sup>18</sup> See Figure 16.9<sup>19</sup> for an image of an immunization clinic sponsored by a student nurses' association.

17. This work is a derivative of [StatPearls](#) by Kisling and Das and is licensed under [CC BY 4.0](#)

18. Savage, C. L. (2020). *Public/community health and nursing practice: Caring for populations* (2nd ed.). FA Davis.

19. "[10442934136\\_1f910af332\\_b](#)" by [Lower Columbia College \(LCC\)](#) is licensed under [CC BY\\_NC-ND 2.0](#)



Figure 16.9 Primary Prevention: An Immunization Clinic

## SECONDARY PREVENTION

**Secondary prevention** emphasizes early detection of disease and targets healthy-appearing individuals with subclinical forms of disease. Subclinical disease refers to pathologic changes with no observable signs or symptoms. Secondary prevention includes screenings such as annual mammograms, routine colonoscopies, Papanicolaou (Pap) smears, as well as screening for depression and substance use disorders.<sup>20</sup> Nurses provide education to community members about the importance of these screenings. See Figure 16.10<sup>21</sup> for an image of a mammogram.

20. This work is a derivative of [StatPearls](#) by Kisling and Das and is licensed under [CC BY 4.0](#)

21. [“US\\_Navy\\_021025-N-6498N-001\\_Mammogram\\_technician\\_aids\\_a\\_patient\\_in\\_completing\\_her\\_annual\\_mammogram\\_evaluation.jpg”](#) by U.S. Navy photo by Ensign Ann-Marie Al Noad is in the [Public Domain](#).



Figure 16.10 Secondary Prevention: Mammograms

## TERTIARY PREVENTION

**Tertiary prevention** is implemented for symptomatic clients to reduce the severity of the disease and potential long-term complications. While secondary prevention seeks to prevent the onset of illness, tertiary prevention aims to reduce the effects of the disease after it is diagnosed in an individual.<sup>22</sup> For example, rehabilitation therapy after an individual experiences a cerebrovascular accident (i.e., stroke) is an example of tertiary prevention. See Figure 16.11<sup>23</sup> for an image of a client receiving rehabilitation after experiencing a stroke.

The goals of tertiary prevention interventions are to reduce disability, promote curative therapy for a disease or injury, and prevent death. Nurses may be involved in providing ongoing home health services in clients' homes as a component of interprofessional tertiary prevention efforts. Health education to prevent the worsening or recurrence of disease is also provided by nurses.

22. This work is a derivative of [StatPearls](#) by Kisling and Das and is licensed under [CC BY 4.0](#)

23. “[tech\\_zerog.jpg](#)” by unknown author at [Gaylord.org](#) is included on the basis of Fair Use





Figure 16.11 Tertiary Prevention: Post-Stroke Rehabilitation. Used under Fair Use.

## QUATERNARY PREVENTION

**Quaternary prevention** refers to actions taken to protect individuals from medical interventions that are likely to cause more harm than good and to suggest interventions that are ethically acceptable. Targeted populations are those at risk of overmedicalization.<sup>24</sup> An example of quaternary prevention is encouraging clients with terminal illness who are approaching end of life to seek focus on comfort and quality of life and consider hospice care rather than undergo invasive procedures that will likely have no impact on recovery from disease.

See additional examples of primordial, primary, secondary, tertiary, and quaternary prevention strategies in Table 16.3a.

24. This work is a derivative of [StatPearls](#) by Kisling and Das and is licensed under [CC BY 4.0](#)

## Table 16.3a Examples of Prevention Interventions<sup>25,26</sup>

25. Savage, C. L. (2020). *Public/community health and nursing practice: Caring for populations* (2nd ed.). FA Davis.
26. This work is a derivative of [StatPearls](#) by Kisling and Das and is licensed under [CC BY 4.0](#)

Prevention Intervention Strategy	Examples
<b>Primordial Prevention</b>	<ul style="list-style-type: none"> <li>• Increased tax on cigarettes</li> <li>• Increased access to public walking paths and public parks</li> </ul>
<b>Primary Prevention</b>	<ul style="list-style-type: none"> <li>• Birthing and newborn classes for new parents at the local hospital</li> <li>• Television commercials regarding the importance of the influenza vaccines</li> <li>• Psychosocial health fairs in local malls or other public facilities</li> <li>• Tobacco cessation public education sessions</li> <li>• Television commercials about mindfulness classes at the community center</li> </ul>
<b>Secondary Prevention</b>	<ul style="list-style-type: none"> <li>• Blood pressure screening events</li> <li>• Sexually transmitted disease screening in college students</li> <li>• Mammograms for early detection of breast cancer</li> <li>• Colonoscopies for early detection of colon cancer</li> <li>• Free testing for people exposed to another individual diagnosed with COVID-19</li> <li>• Screening for substance abuse disorders in high schools</li> <li>• Screening for depression during annual physicals</li> </ul>
<b>Tertiary Prevention</b>	<ul style="list-style-type: none"> <li>• Cardiac rehabilitation for individuals who have experienced a myocardial infarction</li> <li>• Occupational and physical therapy for individuals who experienced a cerebrovascular accident</li> <li>• Diabetic foot care provided at the local community center</li> <li>• Support groups for substance disorders in local churches</li> </ul>



<b>Quaternary Prevention</b>	<ul style="list-style-type: none"> <li>• Routine education provided about advance directives and “Do Not Resuscitate” orders during clinic visits, hospital admissions, and long-term care admissions</li> <li>• Education provided about hospice care to clients diagnosed with terminal illness who are approaching end of life</li> </ul>
------------------------------	--

In the United States, several governing bodies make prevention recommendations. For example, the United States Preventive Services Task Force (USPSTF) makes recommendations for primary and secondary prevention strategies, and the Women’s Preventive Services Initiative (WPSI) makes recommendations specifically for females. The Advisory Committee on Immunizations Practices (ACIP) makes recommendations for vaccinations, and various specialty organizations such as the American College of Obstetrics and Gynecology (ACOG) and the American Cancer Society (ACS) make specialized preventative care recommendations. Preventive services have been proven to be an essential aspect of health care but are consistently underutilized in the United States.<sup>27</sup> Nurses can help advocate for the adoption of evidence-based prevention strategies in their communities and places of employment.

## Continuum of Care Prevention Model

A second framework for prevention interventions, referred to as the “Continuum of Care Prevention Model,” was originally proposed by the Institute of Medicine (IOM) in 1994 and has been adopted by the Substance

27. This work is a derivative of [StatPearls](#) by Kisling and Das and is licensed under [CC BY 4.0](#)

Abuse and Mental Health Services Administration (SAMHSA).<sup>28</sup> See Figure 16.12<sup>29</sup> for an illustration of the Continuum of Care Prevention Model.




Figure 16.12 Continuum of Care Prevention Model. Used under Fair Use.

The Continuum of Care Prevention Model can be used to illustrate a continuum of mental health services for community members that includes prevention, treatment, and maintenance care:

28. National Research Council (US); Institute of Medicine (US) Committee on the Prevention of Mental Disorders and Substance Abuse Among Children, Youth, and Young Adults; and Research Advances and Promising Interventions. Defining the scope of prevention. (2009). In M. E. O'Connell & Warner B. T. (Eds). *Preventing mental, emotional, and behavioral disorders among young people: Progress and possibilities. In Research advances and promising interventions*. National Academies Press. <https://www.ncbi.nlm.nih.gov/books/NBK32789/>
29. This image is a derivative of the "IOM protractor" by unknown author and is included on the basis of Fair Use. Access for free at [http://www.ca-sdfsc.org/docs/resources/SDFSC\\_IOM\\_Policy.pdf](http://www.ca-sdfsc.org/docs/resources/SDFSC_IOM_Policy.pdf)

- Prevention includes three types of strategies including universal, selective, and indicated interventions.
  - **Universal prevention:** Interventions designed to reach entire groups, such as those in schools, workplaces, or entire communities.<sup>30,31</sup> For example, wellness sessions regarding substance misuse can be planned and implemented at a local high school.
  - **Selected prevention:** Interventions that target individuals or groups with greater risk factors (and perhaps fewer protective factors) than the broader population.<sup>32,33</sup> For example, a research study showed that wellness programs implemented for adolescents who were already

30. National Research Council (US); Institute of Medicine (US) Committee on the Prevention of Mental Disorders and Substance Abuse Among Children, Youth, and Young Adults; and Research Advances and Promising Interventions. Defining the scope of prevention. (2009). In M. E. O'Connell & Warner B. T. (Eds). *Preventing mental, emotional, and behavioral disorders among young people: Progress and possibilities. In Research advances and promising interventions*. National Academies Press. <https://www.ncbi.nlm.nih.gov/books/NBK32789/>

31. A Guide to SAMHSA's Strategic Prevention Framework PDF by [Substance Abuse and Mental Health Services Administration](#) is available in the [Public Domain](#).

32. National Research Council (US); Institute of Medicine (US) Committee on the Prevention of Mental Disorders and Substance Abuse Among Children, Youth, and Young Adults; and Research Advances and Promising Interventions. Defining the scope of prevention. (2009). In M. E. O'Connell & Warner B. T. (Eds). *Preventing mental, emotional, and behavioral disorders among young people: Progress and possibilities. In Research advances and promising interventions*. National Academies Press. <https://www.ncbi.nlm.nih.gov/books/NBK32789/>

33. A Guide to SAMHSA's Strategic Prevention Framework PDF by [Substance](#)

using alcohol or drugs reduced the quantity and frequency of their alcohol use and reduced episodes of binge drinking.<sup>34</sup>

- **Indicated prevention:** Interventions that target individuals who have a high probability of developing disease.<sup>35</sup> For example, interventions may be planned for adolescents who show early signs of substance misuse but have not yet been diagnosed with a substance use disorder. Interventions may include referrals to community support services for adolescents who have violated school alcohol or drug policies.<sup>36</sup>
- Treatment refers to identification of a mental health disorder and standard treatment for the known disorder. Treatment also includes interventions to reduce the likelihood of future co-occurring disorders.<sup>37</sup>
- Maintenance refers to long-term treatment to reduce relapse and

[Abuse and Mental Health Services Administration](#) is available in the [Public Domain](#).

34. Werch, C., Moore, M. J., DiClemente, C. C., Bledsoe, R., & Jobli, E. (2005). A multihealth behavior intervention integrating physical activity and substance use prevention for adolescents. *Prevention Science*, 6(213). <https://doi.org/10.1007/s11121-005-0012-3>
35. National Research Council (US); Institute of Medicine (US) Committee on the Prevention of Mental Disorders and Substance Abuse Among Children, Youth, and Young Adults; and Research Advances and Promising Interventions. Defining the scope of prevention. (2009). In M. E. O'Connell & Warner B. T. (Eds). *Preventing mental, emotional, and behavioral disorders among young people: Progress and possibilities. In Research advances and promising interventions*. National Academies Press. <https://www.ncbi.nlm.nih.gov/books/NBK32789/>
36. A Guide to SAMHSA's Strategic Prevention Framework PDF by [Substance Abuse and Mental Health Services Administration](#) is available in the [Public Domain](#).

recurrence, as well as provision of after-care services such as rehabilitation.<sup>38</sup>

See additional examples of prevention strategies using the Continuum of Care Prevention Model in Table 16.3b.

Table 16.3b Examples of Continuum of Care Prevention Strategies

37. National Research Council (US); Institute of Medicine (US) Committee on the Prevention of Mental Disorders and Substance Abuse Among Children, Youth, and Young Adults; and Research Advances and Promising Interventions. Defining the scope of prevention. (2009). In M. E. O'Connell & Warner B. T. (Eds). *Preventing mental, emotional, and behavioral disorders among young people: Progress and possibilities. In Research advances and promising interventions*. National Academies Press. <https://www.ncbi.nlm.nih.gov/books/NBK32789/>
38. National Research Council (US); Institute of Medicine (US) Committee on the Prevention of Mental Disorders and Substance Abuse Among Children, Youth, and Young Adults; and Research Advances and Promising Interventions. Defining the scope of prevention. (2009). In M. E. O'Connell & Warner B. T. (Eds). *Preventing mental, emotional, and behavioral disorders among young people: Progress and possibilities. In Research advances and promising interventions*. National Academies Press. <https://www.ncbi.nlm.nih.gov/books/NBK32789/>

Prevention Strategies	Examples
<b>Universal Prevention</b>	<ul style="list-style-type: none"> <li>• Handwashing education and posters in bathrooms of community gas stations</li> <li>• Parenting classes for new parents</li> <li>• Flu vaccine clinics at a local church</li> </ul>
<b>Selective Prevention</b>	<ul style="list-style-type: none"> <li>• Backpack-buddy programs that provide food from schools to low-income families on weekends</li> <li>• Concussion training programs for youth athletes and their parents</li> <li>• Contact tracing procedures for individuals diagnosed with COVID-19</li> </ul>
<b>Indicated Prevention</b>	<ul style="list-style-type: none"> <li>• Exercise programs at the local senior center targeted for individuals with diabetes</li> <li>• Food or clothing pantries established in a homeless shelter</li> <li>• Screening and consultation for the families of individuals who are admitted to hospitals with alcohol-related injuries</li> </ul>

- ▶ Read [A Guide to SAMHSA's Strategic Prevention Framework](#) PDF for more about planning prevention strategies for substance misuse and related mental health problems.

## Culturally Competent Interventions

To overcome systemic barriers that can contribute to health disparities, nurses must recognize and value cultural differences of diverse populations and

develop prevention programs and interventions in ways that ensure members of these populations benefit from their efforts.<sup>39</sup>

SAMHSA identified the following cultural competence principles for planning prevention interventions<sup>40</sup>:

- Include the targeted population in needs assessments and prevention planning
- Use a population-based definition of community (i.e., let the community define itself)
- Stress the importance of relevant, culturally appropriate prevention approaches
- Promote cultural competence among program staff

▶ Review additional concepts related to culturally responsive care in the “[Diverse Patients](#)” chapter of *Open RN Nursing Fundamentals, 2e*.

## Evidence-Based Practice

It is essential to incorporate evidence-based practice when planning community health interventions. SAMHSA provides an evidence-based

39. A Guide to SAMHSA’s Strategic Prevention Framework PDF by [Substance Abuse and Mental Health Services Administration](#) is available in the [Public Domain](#).

40. A Guide to SAMHSA’s Strategic Prevention Framework PDF by [Substance Abuse and Mental Health Services Administration](#) is available in the [Public Domain](#).

practice resource center for preventive practices related to mental health and substance abuse. See these resources, as well as examples of evidence-based programs and practices in the following box.

### **Examples of Evidence Based Prevention Practices related to Mental Health and Substance Misuse<sup>41</sup>**

- ▶ [Blueprints for Healthy Youth Development](#): Youth violence, delinquency, and drug prevention and intervention programs that meet a strict scientific standard of program effectiveness
- ▶ [Evidence-Based Behavioral Practice \(EBBP\)](#): A project that creates training resources to help bridge the gap between behavioral health research and practice
- ▶ [SAMHSA's Suicide Prevention Research Center \(SPRC\)](#): A best practices registry that identifies, reviews, and disseminates information about best practices that address specific objectives of the National Strategy for Suicide Prevention
- ▶ [The Athena Forum: Prevention 101](#): Substance misuse prevention programs and strategies with evidence of success from the Washington State Department of Social and Health Services
- ▶ [National Institute on Drug Abuse: Preventing Drug Use Among Children and Adolescents PDF](#): Research-based drug abuse

41. Substance Abuse and Mental Health Services Administration. (2019, July 19). *Finding evidence-based programs and practices* [PDF]. [https://www.samhsa.gov/sites/default/files/20190719-samhsa-finding\\_evidence-based-programs-practices.pdf](https://www.samhsa.gov/sites/default/files/20190719-samhsa-finding_evidence-based-programs-practices.pdf)



prevention principles and an overview of program planning, including universal, selected, and indicated interventions

► View the [SAMHSA Evidence-Based Practice Resource Center](#).

## Implementation (Take Action)

Community health nurses collaborate with individuals, community organizations, health facilities, and local governments for successful implementation of community health initiatives. Depending on the established community health needs, goals, outcomes, and target group, the implementation of nursing interventions can be categorized as clinical, behavioral, or environmental prevention:

- **Clinical prevention:** Interventions are delivered one-on-one to individuals in a direct care setting. Examples of clinical prevention interventions include vaccine clinics, blood pressure monitoring, and screening for disease.
- **Behavioral prevention:** Interventions are implemented to encourage individuals to change habits or behaviors by using health promotion strategies. Examples of behavioral prevention interventions include community exercise programs, smoking cessation campaigns, or promotion of responsible alcohol drinking by adults.
- **Environmental prevention:** Interventions are implemented for the entire community when laws, policies, physical environments, or community structures influence a community's health. Examples of environmental prevention strategies include improving clean water systems, establishing no-smoking ordinances, or developing community parks and green spaces.

## Evaluation (Evaluate Outcomes)

When evaluating the effectiveness of community health initiatives, nurses refer to the established goals and SMART outcomes to determine if they were met by the timeline indicated. In general, the following questions are asked during the evaluation stage:

- Did the health of the community improve through the interventions put into place?
- Are additional adaptations or changes to the interventions needed to improve outcomes in the community?
- What additional changes are needed to improve the health of the community?
- Have additional priority problems been identified?

## 16.4 Learning Activities



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=733#h5p-53>

1



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=733#h5p-54>

2



*An interactive H5P element has been excluded from this version of the text. You can view it*

1. “Community Assessment Glossary Cards” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
2. “Community Assessment Question Set 1” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

online here:

<https://wtcs.pressbooks.pub/nursingmhcc/?p=733#h5p-55>

3

- ▶ Test your clinical judgment with an NCLEX Next Generation-style question: [Chapter 16, Assignment 1](#)<sup>4</sup>



- ▶ Test your clinical judgment with an NCLEX Next Generation-style case study: [Chapter 16, Case Study 1](#)<sup>5</sup>



3. "Community Assessment Drag and Drop" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
4. "Community Assessment Next Gen Question 1" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
5. "Community Assessment Next Gen Case Study" by Kellea Ewen is licensed under [CC BY-NC 4.0](#)

## XVI Glossary

**Community diagnosis:** A summary statement resulting from analysis of the data collected from a community health needs assessment.

**Community health needs assessment:** A systematic process to identify and analyze community health needs and assets in order to prioritize these needs, plan, and act upon significant unmet community health needs.

**Focus groups:** A systematic way of collecting data through small group discussion. Focus group participants are chosen to represent a larger group of people.

**Health disparities:** Health differences that are linked with social, economic, and/or environmental disadvantages.

**Health equity:** A goal of everyone having a fair and just opportunity to be as healthy as possible.

**Health inequities:** Avoidable differences in health status seen within and between communities.

**Healthy community:** A community in which local groups from all parts of the community work together to prevent disease and make healthy living options accessible.

**Indicated prevention:** Interventions that target individuals who have a high probability of developing disease.

**Interviews:** Structured conversations with specific individuals who have experience, knowledge, or understanding about a topic or issue.

**Key informant interviews:** Interviews are conducted with select people who are in key positions and have specific areas of knowledge and experience.

**Leading health indicators (LHIs):** A subset of high-priority Healthy People 2030 objectives to drive action toward improving health and well-being.

**Primary data collection:** Data collected to better understand the community's needs and/or study who may be affected by actions taken for the community. Primary data collection includes tools such as public forums, focus groups, interviews, windshield surveys, surveys, and participant observation.<sup>1</sup>

**Primary prevention:** Interventions aimed at susceptible populations or individuals to prevent disease from occurring. Immunizations are an example of primary prevention.

**Primordial prevention:** Risk factor reduction strategies focused on social and environmental conditions targeted for vulnerable populations.

**Public forums:** Public gatherings where citizens discuss important issues at well-publicized locations and times.

**Public health nurses:** Public health nurses work across various settings in the community such as government agencies, community-based centers, shelters, and vaccine distribution sites.<sup>2</sup>

**Quaternary prevention:** Actions taken to protect individuals from medical interventions that are likely to cause more harm than good and to suggest interventions that are ethically acceptable.

**School nurse:** Provide a specialized practice of nursing which protects and promotes student health, facilitates optimal development, and advances academic success.

1. [Community Tool Box](#) by Center for Community Health and Development at the University of Kansas is licensed under [CC BY NC SA 3.0](#)
2. Greenwood, B. (2018, June 29). *What are the primary roles of the community nurse?* CHRON. <https://work.chron.com/primary-roles-community-nurse-15144.html>

**Secondary analysis:** Analyzing previously collected data and research about the community to determine community needs.

**Secondary prevention:** Interventions that emphasize early detection of disease and target healthy-appearing individuals with subclinical forms of disease.

**Selected prevention:** Interventions that target individuals or groups with greater risk factors for illness (and perhaps fewer protective factors) than the broader population.

**Social determinants of health (SDOH):** The conditions in which people are born, grow, work, live, and age.

**Surveys:** Standardized questions that are relatively easy to analyze and are beneficial for collecting information across a large geographic area, hear from as many people as possible, and explore sensitive topics.

**Tertiary prevention:** Interventions implemented for symptomatic clients to reduce the severity of the disease and potential long-term complications.

**Universal prevention:** Interventions designed to reach entire groups or populations such as schools, whole communities, or workplaces.

**Windshield survey:** A form of direct observation of community needs while driving and literally looking through the windshield.





PART XVII

CHAPTER 17 VULNERABLE POPULATIONS



### Learning Objectives

- Describe vulnerable populations
- Identify the needs of the populations that impact health-seeking behaviors
- Discuss support systems and organizations available for vulnerable populations
- Describe the role of the nurse as an advocate for vulnerable populations

The “[Community Assessment](#)” chapter describes how to assess community health needs and apply the nursing process to community health nursing. This chapter provides an overview of vulnerable populations and information for the nurse generalist to support their needs.

## 17.2 Vulnerable Populations

A **vulnerable population** is a group of individuals who are at increased risk for health problems and health disparities.<sup>1</sup> **Health disparities** are health differences linked with social, economic, and/or environmental disadvantages. These disparities disproportionately affect individuals who face greater obstacles to health due to factors such as, such as socioeconomic status, age, gender, culture, religion, mental illness, disability, sexual orientation, or gender identity.<sup>2</sup>

Examples of vulnerable populations include<sup>3,4</sup>:

- The very young and the very old
- Individuals with chronic illnesses, disabilities, or communication barriers
- Veterans
- Racial and ethnic minorities
- Individuals who identify as lesbian, gay, bisexual, transgender, or queer (LGBTQ)
- Victims of human trafficking or sexual violence
- Individuals who are incarcerated and their family members
- Rural Americans

1. AJMC. (2006, November 1). Vulnerable populations: Who are they? *AJMC*, 12(1). <https://www.ajmc.com/view/nov06-2390ps348-s352>

2. HealthyPeople.gov. (2022, February 6). *Disparities*. Office of Disease Prevention and Health Promotion, U.S. Department of Health and Human Services. <https://odphp.health.gov/healthypeople>

3. Joszt, L. (2018, July 20). 5 vulnerable populations in healthcare. *The American Journal of Managed Care*. <https://www.ajmc.com/view/5-vulnerable-populations-in-healthcare>

4. AJMC. (2006, November 1). Vulnerable populations: Who are they? *AJMC*, 12(1). <https://www.ajmc.com/view/nov06-2390ps348-s352>

- Migrant workers
- Individuals with chronic mental health disorders
- People experiencing homelessness Homeless people

These individuals often have limited access to healthcare services, resulting in significant health disparities in life expectancy, morbidity, and mortality. They are also more likely to experience one or more chronic physical or mental health conditions.<sup>5</sup> Advancing health equity for all members of our society is receiving increased emphasis as a central goal of public health. **Health equity** is defined by the U.S. Department of Health and Human Services as the “attainment of the highest level of health for all people” and that “achieving health equity requires valuing everyone equally with focused and ongoing societal efforts to address avoidable inequities, historical and contemporary injustices, and the elimination of health and health care disparities.”<sup>6</sup>

Examples of vulnerable populations and associated resources available to promote health equity and reduce health disparities are further described in the following sections.

## Age and Developmental Status

Developmental status and age influence vulnerability. Children have unique health and developmental needs that require age-appropriate care. Their dependency on others, developmental changes, and distinct patterns of illness and injury necessitate specialized attention within the healthcare

5. AJMC. (2006, November 1). Vulnerable populations: Who are they? *AJMC*, 12(1). <https://www.ajmc.com/view/nov06-2390ps348-s352>
6. Liburd, L. C., Hall, J. E., Mpofu, J. J., Williams, S. M., Bouye, K., & Penman-Aguilar, A. (2020). Addressing health equity in public health practice: Frameworks, promising strategies, and measurement considerations. *Annual Review of Public Health*, 41(1), 417-432. <https://www.annualreviews.org/doi/10.1146/annurev-publhealth-040119-094119>

system. Similarly, the elderly population faces unique healthcare challenges due to a higher incidence of illness and disability, as well as the complexities of managing multiple chronic conditions and medications.<sup>7</sup>

Both children and the elderly are also at increased risk for neglect and abuse. Nurses must recognize signs of neglect and abuse and are legally mandated to report these signs. Read more about neglect, abuse, and mandatory reporting in the “[Trauma, Abuse, and Violence](#)” chapter.

## Chronic Illness and Disability

Individuals with chronic illnesses or disabilities face greater challenges in accessing healthcare services than the general population. These vulnerable groups often encounter barriers such as fragmented care across multiple providers and a lack of care coordination. Additionally, individuals with specific chronic conditions, such as mental illness or HIV, may experience social stigma that impedes their ability to seek or receive appropriate healthcare.<sup>8</sup> See Figure 17.1<sup>9</sup> for an image of a disabled individual who is considered a member of a vulnerable population.

7. Advisory Commission on Consumer Protection and Quality in the Health Care Industry. (n.d.). *Chapter eight: Focusing on vulnerable populations*. University of North Texas Libraries and U.S. Government Printing Office.  
<https://govinfo.library.unt.edu/hcquality/meetings/mar12/papch08.htm>
8. Advisory Commission on Consumer Protection and Quality in the Health Care Industry. (n.d.). *Chapter eight: Focusing on vulnerable populations*. University of North Texas Libraries and U.S. Government Printing Office.  
<https://govinfo.library.unt.edu/hcquality/meetings/mar12/papch08.htm>
9. “[COVID-19 vaccination center, fair grounds Cologne, 1st vaccination-8070.jpg](#)” by Raimond Spekking is licensed under [CC BY-SA 4.0](#)



Figure 17.1 Member of a Vulnerable Population

In addition to experiencing challenges in accessing appropriate health care, adults with disabilities report experiencing more mental distress than the general population. In 2018 an estimated 17.4 million (32.9%) adults with disabilities experienced frequent mental distress, defined as 14 or more reported mentally unhealthy days in the past 30 days. Frequent mental distress is associated with poor health behaviors, mental health disorders, and limitations in daily life.<sup>10</sup> The National Center on Health, Physical Activity and Disability (NCHPAD) is a public health resource center that focuses on health promotion, wellness, and quality of life for people with disabilities.

The Barrier-Free Health Care Initiative is a national initiative by the American Disability Association and the Offices of the United States Attorneys that supports people with disabilities. It legally enforces appropriate access to health care services, such as effective communication for people who are deaf or have hearing loss, physical access to medical care for people with mobility disabilities, and equal access to treatment for people who have HIV/AIDS. This nationwide initiative sends a collective message that disability discrimination in health care is illegal and unacceptable.

10. Centers for Disease Control and Prevention. (2024, December 18). The mental health of people with disabilities.<https://odphp.health.gov/healthypeople>

- ▶ Visit the [Barrier-Free Health Care Initiative](#) to learn more.
- ▶ Visit the [National Center on Health, Physical Activity and Disability \(NCHPAD\)](#) for additional wellness resources.

## Veterans

A veteran is someone who has served in the military forces. Veterans have higher risks for mental health disorders, substance use disorders, post-traumatic stress disorders, traumatic brain injuries, and suicide compared to their civilian counterparts. Medical records of veterans reveal that one in three clients was diagnosed with at least one mental health disorder.<sup>11</sup> Identifying and treating mental health disorders in veterans have the greatest potential for reducing suicide risk, but the reluctance in this population to seek treatment can make diagnosing and treating mental illness challenging. Nurses must be aware of their clients' military history, recognize risk factors for common disorders, and advocate for appropriate health care services.<sup>12</sup> See Figure 17.2<sup>13</sup> for an image of a veteran.

11. Olenick, M., Flowers, M., & Diaz, V. J. (2015). US veterans and their unique issues: Enhancing health care professional awareness. *Advances in Medical Education and Practice*, 6, 635–639. <https://doi.org/10.2147/AMEP.S89479>
12. Olenick, M., Flowers, M., & Diaz, V. J. (2015). US veterans and their unique issues: Enhancing health care professional awareness. *Advances in Medical Education and Practice*, 6, 635–639. <https://doi.org/10.2147/AMEP.S89479>
13. “[DF-SD-03-05175](#)” by Scott H. Spitzer, Civilian, USAF is in the [Public Domain](#).





Figure 17.2 Armed Forces Veteran

The Veterans Health Administration (VHA), the largest healthcare network in the United States, provides medical care to eligible veterans through 1,255 facilities, serving approximately nine million enrolled veterans each year. It is important to note that VA healthcare is not health insurance but rather a hospital system offering medical care. While the VA must provide care for service-connected illnesses or injuries, coverage for other conditions varies based on an individual's priority group, medical needs, and eligibility criteria. Some veterans may also qualify for additional benefits such as dental care.

Veterans may also be eligible for non-medical benefits, including education assistance, home loans or housing grants, and life insurance. Nurses can refer veteran clients to the VA for more information about available benefits and enrollment requirements.

Additionally, veterans and their families can access mental health support through the National Alliance on Mental Illness (NAMI) Homefront program. This course is specifically designed for families, caregivers, and friends of service members and veterans with mental health conditions. It provides education on mental health disorders and strategies to improve support for veterans in need.

- ▶ Visit the [Veterans Affairs \(VA\)](#) website for additional information on health care and other benefits for military service members.
- ▶ Visit the [NAMI Homefront](#) course to learn more about mental health conditions and resources available to military service members and their families.

## Racial and Ethnic Minorities

Racial and ethnic minority populations in the United States experience significant health disparities.<sup>14</sup> See Figure 17.3<sup>15</sup> for an image of a child from a minority population. The U.S. Department of Health and Human Services Office of Minority Health provides ▶ [Minority Population Profiles](#) for Black/African Americans, American Indian/Alaska Natives, Asian Americans, Hispanic/Latinos, and Native Hawaiian/Pacific Islanders. Profiles include demographics, English language fluency, education, economics, insurance coverage and health status information, and full census reports. Nurses can access these profiles to view health care statistics regarding these populations and services available in their communities.

14. Olenick, M., Flowers, M., & Diaz, V. J. (2015). US veterans and their unique issues: Enhancing health care professional awareness. *Advances in Medical Education and Practice*, 6, 635–639. <https://doi.org/10.2147/AMEP.S89479>

15. “3678466365\_da63d01385\_b” by [Xavier Donat](#) is licensed under [CC BY-NC-ND 2.0](#)



Figure 17.3 Child of a Minority Population

The Office of Minority Health is dedicated to improving the health of racial and ethnic minority populations through the development of health policies and programs that help eliminate health disparities.

- ▶ Visit the [Office of Minority Health](#) to learn more about efforts to improve the health of racial and ethnic minority populations through health policy and program development. Examples of initiatives include the following<sup>16</sup> :
  - ▶ [COVID-19 Response](#)
  - ▶ [Improving Cultural Competency for Behavioral Health](#)

<sup>16</sup>. Olenick, M., Flowers, M., & Diaz, V. J. (2015). US veterans and their unique issues: Enhancing health care professional awareness. *Advances in Medical Education and Practice*, 6, 635–639. <https://doi.org/10.2147/AMEP.S89479>

Professionals

▶ Sickle Cell Disease Initiative

## LGBTQ Population

The lesbian, gay, bisexual, transgender, and queer or questioning (LGBTQ) population encompasses all races and ethnicities, religions, and social classes.

Research suggests that LGBTQ individuals experience health disparities linked to societal stigma, discrimination, and denial of their civil rights.<sup>17</sup> The LGBTQ population experiences high rates of mental health disorders, substance abuse, and suicide, and experiences of violence and victimization are frequent for LGBTQ individuals and can result in long-lasting effects.<sup>18</sup> See Figure 17.4<sup>19</sup> for an image of a pride festival by an LGBTQ population.

17. HealthyPeople.gov. (2022, February 6). *Lesbian, gay, bisexual, and transgender health*. Office of Disease Prevention and Health Promotion, U.S. Department of Health and Human Services. <https://odphp.health.gov/healthypeople/objectives-and-data/browse-objectives/lgbt>

18. HealthyPeople.gov. (2022, February 6). *Lesbian, gay, bisexual, and transgender health*. Office of Disease Prevention and Health Promotion, U.S. Department of Health and Human Services. <https://odphp.health.gov/healthypeople/objectives-and-data/browse-objectives/lgbt>

19. “[Pride Festival 2013 On The Streets Of Dublin \(LGBTQ\) \(9183775778\).jpg](#)” by [infomatique](#) is licensed under [CC BY-SA 2.0](#)



Photographed By William Murphy (Infomatique)

Figure 17.4 Pride Festival for the LGBTQ Population

Community efforts to improve the health of the LGBTQ population include these initiatives<sup>20</sup>:

- Training health professionals to appropriately inquire and support clients' sexual orientation and gender identity to promote regular use of health care services
- Training health professionals and students regarding culturally competent care
- Providing supportive social services to reduce suicide and homelessness among LGBTQ youth
- Curbing sexually transmitted infections and human immunodeficiency virus (HIV) transmission

20. HealthyPeople.gov. (2022, February 6). *Lesbian, gay, bisexual, and transgender health*. Office of Disease Prevention and Health Promotion, U.S. Department of Health and Human Services. <https://odphp.health.gov/healthypeople/objectives-and-data/browse-objectives/lgbt>

- Discover health services by state at [CDC's LGBT Health website](#).

## Human Trafficking Victims

**Human trafficking victims** are forced to work or provide commercial sex against their will in legal business settings and underground markets.<sup>21</sup> The International Labor Organization estimates there are 40.3 million victims of human trafficking globally. Victims include men, women, adolescents, and children.<sup>22</sup> Experts estimate that 100,000 incidents of sexual exploitation of minors occur each year in the United States.<sup>23</sup> Human trafficking victims have been identified in cities, suburbs, and rural areas in all 50 states and in Washington, D.C. See Figure 17.5<sup>24</sup> regarding human trafficking statistics reported by state in 2020.

21. National Human Trafficking Hotline. (n.d.). *The victims*. <https://humantraffickinghotline.org/what-human-trafficking/human-trafficking/victims>
22. National Human Trafficking Hotline. (n.d.). *The victims*. <https://humantraffickinghotline.org/what-human-trafficking/human-trafficking/victims>
23. Office of the Attorney General for the District of Columbia. (n.d.). *Human trafficking initiatives*. <https://oag.dc.gov/public-safety/human-trafficking-initiatives>
24. This image is derived from Human Trafficking Hotline data and is in the [Public Domain](#). Access for free at <https://humantraffickinghotline.org/states>






Figure 17.5 Human Trafficking Reported by State in 2020

Widespread lack of awareness and understanding of trafficking leads to low levels of victim identification. Individuals at risk include runaway and homeless youth, foreign nationals, and individuals with prior history of experiencing violence or abuse. A study in Chicago found that 56 percent of female prostitutes were initially runaway youth, and similar numbers have been identified for male populations.<sup>25</sup>

Foreign nationals who are trafficked within the United States face unique challenges. Recruiters located in home countries frequently require such large recruitment and travel fees that victims become highly indebted to the recruiters and traffickers.

Individuals who have experienced violence and trauma in the past are more vulnerable to future exploitation because the psychological effect of trauma is

25. National Human Trafficking Hotline. (n.d.). *The victims*.

<https://humantraffickinghotline.org/what-human-trafficking/human-trafficking/victims>

often long-lasting and challenging to overcome. Victims of domestic violence, sexual assault, war, or social discrimination may be targeted by traffickers who recognize the vulnerabilities left by these prior abuses.<sup>26</sup>

These are some of the services victims of trafficking may need<sup>27</sup>:

- Emergency Services
- Crisis Intervention and Counseling
- Emergency Shelter and Referrals
- Urgent Medical Care
- Safety Planning
- Food and Clothing

► For more information about the services available to victims of human trafficking, visit the [National Human Trafficking Hotline Referral Directory](#).

See an example of a nurse providing care for victims of human trafficking in their community in the “[Spotlight Application](#)” section.

## Incarcerated Individuals and Their Families

At the end of 2019, there were over 2 million people in U.S. prisons and jails

26. National Human Trafficking Hotline. (n.d.). *The victims*.  
<https://humantraffickinghotline.org/what-human-trafficking/human-trafficking/victims>

27. National Human Trafficking Hotline. (n.d.). *The victims*.  
<https://humantraffickinghotline.org/what-human-trafficking/human-trafficking/victims>




with a nationwide incarceration rate of 810 inmates for every 100,000 adults. Higher rates of incarceration are seen among racial/ethnic minorities and people with lower levels of education. The United States incarcerates a larger share of its population than any other country in the world.<sup>28</sup> See Figure 17.6<sup>29</sup> for an illustration comparing worldwide incarceration rates.<sup>30</sup>

28. Gramlich, J. (2021, August 16). *America's incarceration rate falls to lowest level since 1995*. Pew Research Center. <https://pewrsr.ch/2rfSmVL>
29. "U.S. incarcerates a larger share of its population than any other country" by [Pew Research Center](#), Washington, D.C. (August 13, 2021). Used under Fair Use. Access for free at [https://www.pewresearch.org/fact-tank/2021/08/16/americas-incarceration-rate-lowest-since-1995/ft\\_21-08-12\\_incarceration\\_2/](https://www.pewresearch.org/fact-tank/2021/08/16/americas-incarceration-rate-lowest-since-1995/ft_21-08-12_incarceration_2/)
30. Rich, J. D., & The Center for Prisoner Health and Human Rights, Miriam Hospital. *Incarceration and health*. [PDF Working paper]. National Academies. [https://sites.nationalacademies.org/cs/groups/dbassesite/documents/webpage/dbasse\\_083371.pdf](https://sites.nationalacademies.org/cs/groups/dbassesite/documents/webpage/dbasse_083371.pdf)

## U.S. incarcerates a larger share of its population than any other country

*Incarceration rate per 100,000 people of any age*



Note: Figures reflect most recent available data for each country. Territories are counted separately.

Data accessed Aug. 10, 2021.

Source: World Prison Brief, Institute for Crime & Justice Policy Research.

PEW RESEARCH CENTER

Figure 17.6 Comparison of Worldwide Incarceration Rates. Used under Fair Use.

Many individuals who are incarcerated have chronic health conditions, including mental illness and addiction. Over half of all inmates have a mental health disorder, and female inmates have higher rates of post-traumatic stress disorder (PTSD) than the general population. In fact, some experts believe that escalation of U.S. incarceration rates since the 1970s is associated with inadequate community-based health care for mental illness and addiction; left untreated, both conditions can lead to behaviors that result in incarceration.<sup>31</sup>

31. Rich, J. D., & The Center for Prisoner Health and Human Rights, Miriam Hospital. *Incarceration and health*. [PDF Working paper]. National

Health care is a constitutional right for prisoners. The 1976 Supreme Court decision in *Estelle v. Gamble* found that “deliberate indifference to serious medical needs of prisoners” constitutes a violation of the Eighth Amendment prohibition of cruel and unusual punishment. Nurses working in correctional facilities have a legal and ethical obligation to respond to requests for care. If an inmate’s safety complaints are ignored, safety risks are not removed, or there is failure to provide proper medical attention, it is considered “deliberate indifference.” Correctional nurses have a duty to evaluate prisoners’ health needs and determine the level of care required. For example, the nurse determines if the prisoner should be moved to the medical unit or transferred to a health facility that can provide the level of care needed. In addition, all pertinent information of health encounters with prisoners must be thoroughly documented and include assessment findings, health needs, interventions taken, client education, and evaluation of client outcomes.<sup>32</sup>

While the Supreme Court decision mandates health care provision for incarcerated populations in prisons and jails, it does not extend to those under supervision within the criminal justice system (e.g., on parole, probation, or home confinement).<sup>33</sup> However, the Patient Protection and Affordable Care Act (ACA) helps to ensure continuity of medical coverage and care when prisoners are released into the community.<sup>34</sup>

While mortality rates within prisons and jails are comparable to those of the

Academies. [https://sites.nationalacademies.org/cs/groups/dbasse/site/documents/webpage/dbasse\\_083371.pdf](https://sites.nationalacademies.org/cs/groups/dbasse/site/documents/webpage/dbasse_083371.pdf)

32. National Commission on Correctional Health Care. (n.d.). *Legal issues*. <https://www.ncchc.org/legal-issues>

33. Gramlich, J. (2021, August 16). *America’s incarceration rate falls to lowest level since 1995*. Pew Research Center. <https://pewrsr.ch/2rfSmVL>

34. Gramlich, J. (2021, August 16). *America’s incarceration rate falls to lowest level since 1995*. Pew Research Center. <https://pewrsr.ch/2rfSmVL>

general population, releasees are nearly 13 times more likely to die in the two weeks following their release than the general population. The most common cause of death is overdose. Interventions that follow in-prison drug treatment programs with post-release treatment have been shown to reduce drug use and associated recidivism (i.e., return to prison). There have also been efforts to improve the outcomes of prisoner reentry to society through assistance with employment, housing, and other transitional needs that ultimately affect health.<sup>35,36</sup>

- ▶ The National Alliance on Mental Health (NAMI) provides several [Re-Entry Planning](#) resources.

Incarceration has a public health impact on prisoners' family members and their communities while they are incarcerated, as well as after their release. Community health is affected by incarceration due to economic impacts (i.e., consequences on families' well-being when income earners are removed) and long-term impacts on children's health and mental health (also referred to as adverse childhood events). For example, children of incarcerated parents are more likely to live in poverty, as well as have higher rates of learning disabilities, developmental delays, speech/language problems, attention disorders, aggressive behaviors, and drug and alcohol use. Additionally,

35. Rich, J. D., & The Center for Prisoner Health and Human Rights, Miriam Hospital. *Incarceration and health*. [PDF Working paper]. National Academies. [https://sites.nationalacademies.org/cs/groups/dbasssite/documents/webpage/dbasse\\_083371.pdf](https://sites.nationalacademies.org/cs/groups/dbasssite/documents/webpage/dbasse_083371.pdf)

36. Massoglia, M., & Pridemore, W. A. (2015). Incarceration and health. *Annual Review of Sociology*, 41, 291–310. <https://doi.org/10.1146/annurev-soc-073014-112326>

children of incarcerated parents are up to five times more likely to enter the criminal justice system than children of non incarcerated parents.<sup>37</sup>

Family-centered services for incarcerated parents and their children focus on parenting programs, family strengthening activities, nurturing of family relationships, and community support for families during incarceration and following release.

- ▶ Read more about family-centered services at [U.S. Department of Health Services Child Welfare Information Gateway](#).

- ▶ Read additional guidance and resources at [CDC Correctional Health](#).

## Rural Americans

Rural Americans are a population group that experiences significant health disparities. Rural risk factors for health disparities include geographic isolation, lower socioeconomic status, higher rates of health risk behaviors, limited access to health care specialists, and limited job opportunities. This inequality is intensified because rural residents are less likely to have

37. Massoglia, M., & Pridemore, W. A. (2015). Incarceration and health. *Annual Review of Sociology*, 41, 291–310. <https://doi.org/10.1146/annurev-soc-073014-112326>

employer-provided health insurance or Medicaid coverage.<sup>38,39</sup> See Figure 17.7<sup>40</sup> for an image of a small town in rural America.



Figure 17.7 Rural America

For an in-depth look at rural health disparities, the Rural Health Information Hub Rural Health Series serves as a resource to examine rural mortality and preventable deaths, health-related behaviors, chronic disease, mental health services, and other related topics. The [Federal Office of Rural Health Policy](#) provides additional information regarding rural health policy, community health programs, and telehealth programs to increase access to health care

38. Rural Health Information Hub. (2019, April 22). *Rural health disparities*. <https://www.ruralhealthinfo.org/topics/rural-health-disparities>

39. United States Department of Agriculture. (2018). Rural America at a glance 2018 edition. *Economic Information Bulletin*, 200. <https://www.ers.usda.gov/publications/pub-details?pubid=90555>

40. “[One Light Town” Small Town, Nebraska 7-25-13a \(10784126756\).jpg](#)” by [inkknife\\_2000](#) (7.5 million views +) is licensed under [CC BY-SA 2.0](#)



for underserved people in rural communities. Two initiatives to improve rural health are the National Health Service Corps (NHSC) loan repayment program and telehealth.<sup>41</sup>

NHSC builds healthy communities by supporting health care professionals dedicated to working in areas of the United States with limited access to care. NHSC offers educational loan repayment programs to health professionals who commit to working in underserved areas for specific lengths of time.<sup>42</sup> See an infographic about NHSC in Figure 17.8.<sup>43</sup>



Figure 17.8 National Health Service Corps

41. Health Resources & Services Administration. (n.d.). *Federal office of rural health policy*. <https://www.hrsa.gov/rural-health/index.html>
42. National Health Service Corps. *Health Resources & Services Administration*. <https://nhsc.hrsa.gov>
43. This image is derived from [HRSA Programs Serve Rural Communities PDF](#) by Health Resources & Services Administration and is in the [Public Domain](#).

**Telehealth** is a recent initiative to improve health care access to underserved communities. Telehealth is the use of digital technologies to deliver medical care, health education, and public health services by connecting multiple users in separate locations. See Figure 17.9<sup>44</sup> for an image of telehealth. Telehealth can improve health outcomes and access to care by delivering health services in underserved areas and remove barriers of time, distance, and provider scarcities.<sup>45</sup> Nurses support clients using telehealth and provide client education.

However, telehealth can risk widening disparities for some socioeconomic groups with limited access to the resources necessary for effective telemedicine. Nurses can help ensure that health care systems tailor the use of technology to meet the needs of underserved populations.<sup>46</sup>

44. “[Telemedicine\\_Consult.jpg](#)” by Intel Free Press is licensed under [CC BY-SA 2.0](#)

45. Telligen & Great Plains Telehealth Resource & Assistance Center. (2014). *Telehealth: Start-up and resource guide*. [https://www.healthit.gov/sites/default/files/telehealthguide\\_final\\_0.pdf](https://www.healthit.gov/sites/default/files/telehealthguide_final_0.pdf)

46. Ortega, G., Rodriguez, J. A., Maurer, L. R., Witt, E. E., Perez, N., Reich, A., & Bates, D. W. (2020). Telemedicine, COVID-19, and disparities: Policy implications. *Health Policy and Technology*, 9(3), 368–371. <https://doi.org/10.1016/j.hlpt.2020.08.001>





Figure 17.9 Telehealth

- ▶ Visit the [Rural Health Information Hub Rural Health Series](#) to examine rural mortality and preventable deaths, health-related behaviors, chronic disease, mental health services, and other related topics.
- ▶ Learn more about the [NHSC](#) and how it supports health care professionals to provide care in areas with limited access to health services.

## Migrant Workers

A **migrant worker** is a person who moves within their home country or outside of it to pursue work. Migrant workers usually do not intend to stay

permanently in the country or region in which they work. See Figure 17.10<sup>47</sup> for an image of migrant workers harvesting cabbage.



Figure 17.10 Migrant Workers Harvesting Cabbage

Migratory and seasonal agricultural workers (MASW) and their families face unique health challenges that can result in significant health disparities. Challenges can include the following<sup>48</sup>:

- Hazardous work environments
- Poverty
- Insufficient support systems
- Inadequate or unsafe housing
- Limited availability of clean water and septic systems

47. “5123728839\_1b7ce92b5c\_k” by Bob Jagendorf is licensed under [CC BY-NC 2.0](#)

48. Rural Health Information Hub. (2021, July 1). *Rural migrant health*.  
<https://www.ruralhealthinfo.org/topics/migrant-health>

- Inadequate health care access and lack of continuity of care
- Lack of health insurance
- Cultural and language barriers
- Fear of using health care due to immigration status
- Lack of transportation

MSAW populations experience serious health problems including diabetes, malnutrition, depression, substance use, infectious diseases, pesticide poisoning, and injuries from work-related machinery. These critical health issues are exacerbated by the migratory culture of this population group that increases isolation and makes it difficult to maintain treatment regimens and track health records.<sup>49</sup>

Successful strategies to support health services for migrant workers are as follows<sup>50</sup> :

- Culturally sensitive health education and outreach
- Educational materials at the appropriate literacy level
- Portable medical records and case management
- Mobile medical units
- Transportation services
- Translation services

► In the United States, access to healthcare services for migrant workers can vary depending on several factors, including

49. Rural Health Information Hub. (2021, July 1). *Rural migrant health*.  
<https://www.ruralhealthinfo.org/topics/migrant-health>

50. Rural Health Information Hub. (2021, July 1). *Rural migrant health*.  
<https://www.ruralhealthinfo.org/topics/migrant-health>

- ▶ immigration status, employment, and state-specific policies. Nurses can refer uninsured migrant farmworkers to [Migrant Health Centers](#) or other federally qualified health centers that are open to everyone.

- ▶ Read more about promoting migrant worker health at the [Rural Migrant Health Information Hub](#).

## Individuals with Mental Health Disorders

One in five American adults experiences some form of mental illness. Despite a recent focus on mental health in America, there are still many harmful attitudes and misunderstandings surrounding mental illness that can cause people to ignore their mental health and make it harder to reach out for help.<sup>51</sup>

Many individuals with mental health disorders go undiagnosed. Nurses must be aware of these common signs of mental health disorders<sup>52</sup>:

- Excessive worrying or fear

51. Centers for Disease Control and Prevention. (2021, July 20). *Mental health*. [https://www.cdc.gov/mental-health/?CDC\\_AAref\\_Val=https://www.cdc.gov/mentalhealth/index.htm](https://www.cdc.gov/mental-health/?CDC_AAref_Val=https://www.cdc.gov/mentalhealth/index.htm)

52. Centers for Disease Control and Prevention. (2021, July 20). *Mental health*. [https://www.cdc.gov/mental-health/?CDC\\_AAref\\_Val=https://www.cdc.gov/mentalhealth/index.htm](https://www.cdc.gov/mental-health/?CDC_AAref_Val=https://www.cdc.gov/mentalhealth/index.htm)

- Excessively sad feelings
- Confused thinking or problems concentrating and learning
- Extreme mood changes, including uncontrollable “highs” or feelings of euphoria
- Prolonged or strong feelings of irritability or anger
- Avoidance of friends and social activities
- Difficulties understanding or relating to other people
- Changes in sleeping habits or feeling tired and low energy
- Changes in eating habits, such as increased hunger or lack of appetite
- Changes in sex drive
- Disturbances in perceiving reality (e.g., delusions and hallucinations)
- Inability to perceive changes in one’s own feelings, behavior, or personality (i.e., lack of insight)
- Overuse of substances like alcohol or drugs
- Multiple physical ailments without obvious causes (such as headaches, stomachaches, or vague and ongoing “aches and pains”)
- Thoughts of suicide
- Inability to carry out daily activities or handle daily problems and stress
- An intense fear of weight gain or overly concerned with appearance

Mental health conditions are also present in children. Because children are still learning how to identify and talk about thoughts and emotions, their most obvious symptoms are behavioral. Behavioral symptoms in children can include the following<sup>53</sup>:

- Changes in school performance
- Excessive worry or anxiety
- Hyperactive behavior
- Frequent nightmares
- Frequent disobedience or aggression

53. National Alliance on Mental Illness. (n.d.). *Warning signs and symptoms*.  
<https://nami.org/About-Mental-Illness/Warning-Signs-and-Symptoms>

- Frequent temper tantrums

Nurses should recognize signs and symptoms of potentially undiagnosed mental health problems in all care settings and make appropriate referrals to mental health professionals and support organizations.

- ▶ [National Alliance on Mental Illness \(NAMI\)](#) provides advocacy, education, support, and public awareness so that all individuals and families affected by mental illness can build better lives. NAMI's strategic goals are for people to get help early, get the best possible care, and be diverted from justice system involvement.<sup>54</sup>

NAMI offers these programs that promote improved care for individuals with mental illness and their loved ones:

- ▶ [Your Journey With Mental Illness](#) is a web page with several resources such as understanding your diagnosis, finding a health professional, understanding insurance, handling a crisis, and navigating finances and work.
- ▶ [NAMI Basics](#) is an educational program for parents, family members, and caregivers who provide care for youth (ages 22 and younger) who are experiencing mental health symptoms. The program covers these topics:
  - Impact mental health conditions can have on the entire family

54. National Alliance on Mental Health. (n.d.). *2020-2025 strategic plan*. <https://nami.org/NAMInet/Board-of-Directors/Governance-Documents/NAMIStrategicPlan2020>



- Different types of mental health care professionals and available treatment options and therapies
  - Overview of the public mental health care, school, and juvenile justice systems and resources to help navigate these systems
  - Child rights advocacy at school and in health care settings
  - Preparation and response to crisis situations (self-harm, suicide attempts, etc.)
  - Importance of self-care
- ▶ [NAMI Provider](#) introduces mental health professionals to the unique perspectives of people with mental health conditions and their families. It promotes empathy for their daily challenges and the importance of including them in all aspects of the treatment process.
  - ▶ [NAMI In Our Own Voice](#) is a program to promote a change in attitudes, assumptions, and ideas about people with mental health conditions. Presentations provide a personal perspective of mental health conditions, as leaders with lived experience talk openly about what it's like to have a mental health condition.

## Homeless People

Homelessness can significantly affect an individual's health in three ways: health problems caused by homelessness, health problems that cause homelessness, and health conditions that are difficult to treat because of

homelessness. As a result, the average age of death among homeless people is the mid-50s.<sup>55</sup>

Homeless people face multiple barriers to health care, including transportation, fragmentation of health care services, difficulty scheduling and keeping appointments, stigma of homelessness, lack of trust, social isolation, and significant basic physiological needs. Homeless people frequently have multiple needs resulting from exposure to violence or the elements, food insecurity, and untreated or undertreated physical and mental illnesses, resulting in frequent emergency department visits and hospitalizations.<sup>56</sup>

Effective health care for the homeless population addresses social determinants of health, such as housing, income, food, and social supports. The Medical Home Model developed by the Veterans Health Administration is an example of a proactive, primary care-based, interdisciplinary team model based on client-centered, holistic care, and active communication and coordination among providers. This model is considered effective for clients with complex health care needs. See Figure 17.11<sup>57</sup> for an illustration of the

55. O'Toole, T. P., Johnson, E. E., Aiello, R., Kane, V., & Pape, L. (2016). Tailoring care to vulnerable populations by incorporating social determinants of health: The veterans health administration's "homeless patient aligned care team" program. *Preventing Chronic Disease*, 13:150567. <http://dx.doi.org/10.5888/pcd13.150567>
56. O'Toole, T. P., Johnson, E. E., Aiello, R., Kane, V., & Pape, L. (2016). Tailoring care to vulnerable populations by incorporating social determinants of health: The veterans health administration's "homeless patient aligned care team" program. *Preventing Chronic Disease*, 13:150567. <http://dx.doi.org/10.5888/pcd13.150567>
57. This image is derived from O'Toole TP, Johnson EE, Aiello R, Kane V, Pape L. (2016). Tailoring care to vulnerable populations by incorporating social



Medical Home Model. As homeless veterans stabilize clinically and socially, as evidenced by their moving into permanent housing and demonstrating appropriate self-care and health-seeking behaviors, they are transitioned to traditional care settings to continue their care.<sup>58</sup>




Figure 17.11 Medical Home Model

Core elements of the Medical Home Model distinguish it from traditional primary care with the following characteristics<sup>59</sup>:

determinants of health: The veterans health administration’s “homeless patient aligned care team” program. *Preventing Chronic Disease* 13:150567. <http://dx.doi.org/10.5888/pcd13.150567>

58. O’Toole, T. P., Johnson, E. E., Aiello, R., Kane, V., & Pape, L. (2016). Tailoring care to vulnerable populations by incorporating social determinants of health: The veterans health administration’s “homeless patient aligned care team” program. *Preventing Chronic Disease*, 13:150567. <http://dx.doi.org/10.5888/pcd13.150567>

59. O’Toole, T. P., Johnson, E. E., Aiello, R., Kane, V., & Pape, L. (2016). Tailoring care to vulnerable populations by incorporating social determinants of

- Low-threshold access to care
  - Open access with walk-in capacity and flexible scheduling (i.e., clients do not need an appointment to be seen by their care team)
  - Clinical outreach to homeless people on streets, in shelters, and in community locations such as soup kitchens
- Integrated services
  - Mental health services and primary care services are located close to each other, and providers from both services are involved in clients' health care plan
  - Sustenance needs (e.g., food or food vouchers, hygiene kits, clothes, bus passes, other transportation assistance) are available at the same location
- Intensive health care management integrated with community agencies with an emphasis on ongoing, continuous care
- Ongoing staff training focused on development of homeless care skills

Research indicated the Medical Home Model reduced emergency

health: The veterans health administration's "homeless patient aligned care team" program. *Preventing Chronic Disease*, 13:150567.  
<http://dx.doi.org/10.5888/pcd13.150567>

department visits and hospitalizations of the homeless population by integrating supports that addressed social determinants of health into a clinical care model.<sup>60</sup>

<sup>60</sup>. O'Toole, T. P., Johnson, E. E., Aiello, R., Kane, V., & Pape, L. (2016). Tailoring care to vulnerable populations by incorporating social determinants of health: The veterans health administration's "homeless patient aligned care team" program. *Preventing Chronic Disease*, 13:150567. <http://dx.doi.org/10.5888/pcd13.150567>

## 17.3 Spotlight Application

This section will apply the tools from the “[Community Assessment](#)” chapter to providing care for victims of human trafficking as an example of a nurse generalist caring for a member of a vulnerable population. See Figure 17.12<sup>1</sup> regarding human trafficking initiatives.



Figure 17.12 Human Trafficking Initiatives

After hearing about a sexual trafficking arrest at a local hotel in the news, a nurse who works in the emergency department reads more information on the [National Human Trafficking Hotline](#). Because the community is located near a major national interstate, the nurse realizes there are likely many unidentified victims of human trafficking passing through the community.

1. "[OAG-2020-Sex-Trafficking-Poster-English](#)" by [Office of Attorney General for the District of Columbia](#) is in the [Public Domain](#).

The nurse develops a goal to improve the identification and support of individuals experiencing human trafficking in the community.

The nurse begins a community health needs assessment by reading statistics on the ► [National Human Trafficking Hotline](#), as well as additional information in the ► [Adult Human Trafficking Screening Tool and Guide](#) PDF by the U.S. Department of Health and Human Services' Administration for Children and Families. Based on this research, the nurse keeps the following tips in mind when conducting an assessment with a potential victim of trafficking<sup>2,3</sup>,

- **Lack of self-identification.** Many victims do not self-identify as human trafficking victims due to lack of knowledge about the crime, as well as power and control dynamics involved in trafficking situations.
- **Being conscious of verbal and nonverbal messages.** When speaking with a potential victim of trafficking, the nurse plans on adopting open, nonthreatening body positioning; maintaining a calm tone of voice; displaying a warm, natural facial expression; using active listening skills; and mirroring the language the potential victim uses. For example, if the potential victim refers to her controller as her boyfriend, referring to that person as a “pimp” or a “sex trafficker” can negatively impact a therapeutic nurse-client relationship.
- **Being aware of power dynamics.** If a third party accompanies or interprets for the potential victim, the nurse plans on trying to speak to the potential victim alone or in a secure area with an official interpreter.
- **Being aware of canned stories.** Scripted stories are common, and the true

2. National Human Trafficking Resource Center. (2011). *Human trafficking assessment for domestic violence & sexual assault programs*.<https://humantraffickinghotline.org/states>

3. Macias-Konstantopoulos, W., & Owens, J. (2018). *Adult human trafficking screening tool and guide* [PDF]. Administration for Children & Families and National Human Trafficking. [https://www.acf.hhs.gov/sites/default/files/documents/otip/adult\\_human\\_trafficking\\_screening\\_tool\\_and\\_guide.pdf](https://www.acf.hhs.gov/sites/default/files/documents/otip/adult_human_trafficking_screening_tool_and_guide.pdf)

story may not emerge until trust has been built with the potential victim.

- **Meeting the person's physical needs by offering a snack or beverage.** Most individuals experiencing human trafficking have been deprived of basic necessities, such as food, fluids, sleep, and urgent medical needs. Hunger is a very common problem for individuals who have been trafficked; a person who is hungry will have difficulty focusing and may be irritable.
- **Never allow confidential data to leave the office.** Ensure secure mechanisms to maintain privacy of client data (e.g., use effective passwords and/or locked file cabinets).
- **Being empathetic.** The decision to disclose trafficking can be mentally challenging, emotionally draining, and potentially physically unsafe for the victim. If the victim does not feel prepared to disclose during the immediate clinical visit, disclosure may occur at a future visit.<sup>4</sup>

## Assessment

The nurse is aware of the warning signs an individual who is a victim of human trafficking or sexual violence may exhibit<sup>5</sup>:

- Shares a scripted or inconsistent history
- Is unwilling or hesitant to answer questions about an injury or illness
- Is accompanied by an individual who does not let the client speak for themselves, refuses to let the client have privacy, or who interprets for them
- Shares information about controlling or dominating relationships

4. Tracy, E. E., & Macias-Konstantopoulos, W. (2021). Human trafficking: Identification and evaluation in the health care setting. *UpToDate*. Retrieved March 16, 2022, from [www.update.com](http://www.update.com)

5. Tracy, E. E., & Macias-Konstantopoulos, W. (2021). Human trafficking: Identification and evaluation in the health care setting. *UpToDate*. Retrieved March 16, 2022, from [www.update.com](http://www.update.com)

- Demonstrates fearful or nervous behavior or avoids eye contact
- Is resistant to accept assistance or demonstrates hostile behavior
- Is unable to provide their address
- Is not aware of their location, the current date, or time
- Is not in possession of their identification documents
- Is not in control of his or her own money
- Is not being paid or wages are withheld

Sex trafficking victims may exhibit the following<sup>6</sup>:

- Client is under the age of 18 and is involved in the commercial sex industry
- Has tattoos or other forms of branding, such as tattoos that say “Daddy,” “Property of...,” “For sale,” etc.
- Reports an unusually high number of sexual partners
- Does not have appropriate clothing for the weather or venue
- Uses language common in the commercial sex industry
- Experiences multiple sexually transmitted infections, pregnancies, miscarriages, or terminations

If concerning assessment findings are noted, the nurse plans on using a screening tool found in the ► [Adult Human Trafficking Screening Tool and Guide PDF](#). Before beginning any conversation with the client, the nurse plans to assess for potential safety risks that may result from asking the client sensitive questions and to ensure privacy. The nurse also plans on being transparent about the limits of confidentiality regarding mandated reporting.<sup>7</sup>

6. Tracy, E. E., & Macias-Konstantopoulos, W. (2021). Human trafficking: Identification and evaluation in the health care setting. *UpToDate*. Retrieved March 16, 2022, from [www.update.com](http://www.update.com)

7. Tracy, E. E., & Macias-Konstantopoulos, W. (2021). Human trafficking:

# Diagnosis

Several nursing diagnoses may apply to individuals experiencing human trafficking or sexual violence. Examples of diagnoses include the following<sup>8</sup>:

- *Post-Trauma Syndrome* related to insufficient social support
- *Powerlessness* related to insufficient sense of control
- *Chronic Low Self-Esteem* related to inadequate belonging
- *Risk for Impaired Emancipated Decision-Making* as manifested by insufficient privacy to openly discuss health care options

## Outcome Identification

The overall goal of the nurse is to identify and support victims of human trafficking or sexual violence. Individual SMART goals can be established for clients based on their circumstances. Some examples of SMART outcomes are as follows:

- The client will remain safe from injury and death.
- The client will verbalize local resources and support before the end of the visit.
- The client will acknowledge the traumatic experience and begin to work through the trauma by verbalizing feelings of fear, anger, anxiety, or

Identification and evaluation in the health care setting. *UpToDate*. Retrieved March 16, 2022, from [www.update.com](http://www.update.com)

8. Ackley, B., Ladwig, G., Makic, M. B., Martinez-Kratz, M., & Zanoliti, M. (2020). *Nursing diagnosis handbook: An evidence-based guide to planning care* (12th ed.). Elsevier.



helplessness during the visit.

## Planning Interventions

The nurse prepares to respond quickly based on the client's circumstances by doing the following:

- Knowing how to contact law enforcement if the client or staff are in immediate danger.
- Being prepared to complete mandatory reporting requirements for minors and other circumstances according to state law.
- Creating a list of local services and resources to provide to human trafficking victims.

The nurse plans on sharing these resources regarding human trafficking with local community representatives associated with urgent care centers, clinics, primary care providers, and emergency departments.

## Implementation

The nurse implements interventions with individuals seeking care based on assessment findings. Individualized interventions may include these steps:

- Contact law enforcement if the client or staff are in immediate danger.
- Create a safety plan by identifying the current level of risk and safety concerns. Outline concrete options for responding when the individual's safety is threatened.
- Refer the victim to local services and resources, such as food, shelter, and legal services.
- Share referral options. For assistance with finding referrals or other resources, contact the National Human Trafficking Hotline at 1-888-373-7888.
- Be prepared to respond to a potential trauma reaction. Encourage relaxation breathing and grounding techniques for immediate de-escalation. (Review crisis intervention in the "[Stress, Coping, and Crisis Intervention](#)" chapter and grounding techniques in the "[Anxiety](#)")

[Disorders](#)” chapter.)

- Perform mandatory reporting for minors and other circumstances according to state law. Read more about mandatory reporting in the box below.
- Report the case confidentially to appropriate authorities trained on human trafficking.
- Document assessment findings and referrals. Maintain client privacy and ensure the documentation is secure electronically or on paper.

### **Mandatory Reporting**

Mandatory reporting is required for minors. Read more about mandatory reporting in the “[Legal and Ethical Considerations in Mental Health Care](#)” chapter. If reporting human trafficking of adults is not a component of your state’s mandatory reporting statutes, certain circumstances may be considered reportable. Some of these “reportable” circumstances may include domestic violence, injuries caused in violation of criminal law, or injuries caused by a deadly weapon (e.g., firearm, knife, or machete). Some professionals hesitate to report potential victims of trafficking due to fears of violating the rules of the Health Insurance Portability and Accountability Act (HIPAA). HIPAA was written to protect client confidentiality but was not designed to prevent the reporting of trauma and crimes. If you’re unsure about whether HIPAA permits the reporting of client information in a specific situation, human trafficking can still be reported without divulging individually identifiable client health

information. For example, you could report the gender, age of client, and type of trafficking.<sup>9</sup>

## Evaluation

The nurse plans on evaluating the success of achieving the overall goal to improve the identification and support of individuals experiencing human trafficking in the community in a variety of ways:

- Reviewing statistics reported by local law enforcement agencies regarding human trafficking cases.
- Reviewing statistics reported by the ► [National Human Trafficking Hotline](#).
- Informally interviewing local community representatives associated with urgent care centers, clinics, primary care providers, and emergency departments regarding human trafficking cases.

9. Macias-Konstantopoulos, W., & Owens, J. (2018). Adult human trafficking screening tool and guide. *Administration for Children & Families and National Human Trafficking* [PDF]. [https://www.acf.hhs.gov/sites/default/files/documents/otip/adult\\_human\\_trafficking\\_screening\\_tool\\_and\\_guide.pdf](https://www.acf.hhs.gov/sites/default/files/documents/otip/adult_human_trafficking_screening_tool_and_guide.pdf)

## 17.4 Learning Activities



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=806#h5p-56>

1



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=806#h5p-58>

2



1. “Vulnerable Populations Glossary Cards” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
2. “Vulnerable Populations Question Set” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

- ▶ Test your clinical judgment with an NCLEX Next Generation-style case study: [Chapter 17, Case Study 1](#)<sup>3</sup>

3. “Vulnerable Populations Next Gen Case Study” by Kellea Ewen is licensed under [CC BY-NC 4.0](#)

## XVII Glossary

**Health disparities:** Health differences that are linked with social, economic, and/or environmental disadvantages. Health disparities adversely affect groups of people who often experience greater obstacles to health based on individual characteristics, such as socioeconomic status, age, gender, culture, religion, mental illness, disability, sexual orientation, or gender identity.

**Health equity:** The attainment of the highest level of health for all people.

**Human trafficking victims:** Individuals forced to work or provide commercial sex against their will in legal business settings and underground markets.

**Migrant worker:** A person who moves within their home country or outside of it to pursue work. Migrant workers usually do not intend to stay permanently in the country or region in which they work.

**Telehealth:** The use of digital technologies to deliver medical care, health education, and public health services by connecting multiple users in separate locations.

**Vulnerable population:** A group of individuals who are at increased risk for health problems and health disparities.

## CHAPTER 18 ENVIRONMENTAL HEALTH AND EMERGENCY PREPAREDNESS





### Learning Objectives

- Analyze the nursing role in emergency preparedness planning within the community  
explain the nurse role as a collaborative advocate for the health needs of the community
- Describe the nurse's role in promoting environmental health
- Apply principles of triage
- Outline tips for preventing and managing stress for disaster responders

Since the early years of the profession, nursing leaders such as Florence Nightingale and Lillian Wald have recognized the role of nurses in controlling the influence of environmental factors on health. Nurses have long appreciated that a healthy environment impacts the physical and mental health of individuals, families, communities, and populations.<sup>1</sup>

At some point during their career, nearly every nurse finds themselves caring for individuals affected by exposure to an environmental hazard or disaster. Disasters have environmental, physical, and psychological effects on individuals and communities. The increasing impacts of natural, man-made, and infectious disease disasters have changed health care and nursing

1. American Nurses Association. (2007). *ANA's principles of environmental health for nursing practice with implementation strategies*. Nursesbooks.org. <https://www.nursingworld.org/practice-policy/work-environment/health-safety/environmental-health/>

perspectives around the world. Nurses are on the front lines in supporting individuals and communities affected by disasters. This chapter will review nurses' roles in promoting environmental health and treating individuals exposed to environmental hazards, as well as participating in emergency preparedness and disaster response and recovery.



Category	Environmental Hazards
Housing	<p><b>Building Contaminants:</b> Radon, asbestos, carbon monoxide, and lead</p> <p><b>Personal Chemicals:</b> Nail polish remover, spray deodorant, and rubbing alcohol</p> <p><b>Household Chemicals:</b> Kerosene, coal or wood-burning stoves, burning pine cones or wax candles, burning incense, chlorinated water in swimming pools, cleaning products, household detergents, plastic cement, acrylic paints, furniture wax, rubber-based paints, gasoline, white-out solvents, and pesticides (used in gardening chemicals and lawn care)</p> <p><b>Electromagnetic Pollution:</b> Electric blankets, electrically heated water beds, fluorescent lighting, cathode ray tubes (computer terminals), cellphones, tablets, microwave ovens, television sets, radio or TV transmission towers, short-wave radio transmitters, and high-voltage electrical power lines</p> <p><b>Natural Contaminants:</b> Mold and pollen</p> <p><b>Human Contaminants:</b> Second-hand smoke</p>

<b>Occupational</b>	<p><b>Medical Chemicals:</b> Injectable phenol (found in allergy shots) and silver amalgam (found in some dental fillings)</p> <p><b>Occupational Chemicals:</b> Radioactivity from direct radiation or decay from nuclear waste</p> <p><b>Construction Materials &amp; Industry:</b> Urea-formaldehyde foam insulation, asbestos insulation, fiberglass &amp; particleboard, treated lumber, and silica air pollution</p> <p><b>Noise Pollution:</b> Traffic, loud music, workplace equipment noise, and airports</p> <p><b>Ergonomics:</b> Physical injuries due to positioning or lack of physical assistance</p> <p><b>Communicable Infectious Disease:</b> Exposure to infections despite transmission precautions (i.e., airborne, droplet, or contact precautions). Read more about the chain of infection and transmission precautions in the “<a href="#">Infection</a>” chapter of <i>Open RN Nursing Fundamentals</i>.</p>
<b>Atmosphere</b>	<p><b>Air pollution</b></p> <p><b>Greenhouse Gasses:</b> Carbon dioxide, methane, nitrous oxide, and fluorinated gasses</p> <p><b>Ultraviolet Radiation</b></p>
<p><b>Waterborne Illness</b></p> <p>(Recreational or drinking water contaminated by disease-causing contaminants, microbes or pathogens)</p>	<p><b>Chemical and Metal Contaminants:</b> Chlorine, fluorine, mercury, lead, and nitrates</p> <p><b>Common Microbes:</b> Cryptosporidiosis, cyclosporiasis, Escherichia, giardiasis, legionellosis, norovirus, shigellosis, blue-green algae, schistosome parasites (swimmer’s itch), pseudomonas (hot tub rash), and cholera.</p> <p>Read more about waterborne illness at the <a href="#">Wisconsin Department of Health</a> or the <a href="#">Minnesota Department of Health</a>.</p> <p>*Report waterborne illness to the state Department of Health.</p>

<b>Foodborne Illness</b>  (Consuming foods or beverages contaminated with bacteria, viruses, parasites, toxins, or chemicals)	<b>Common Pathogens:</b> Clostridium botulinum, listeria, campylobacter, norovirus, clostridium perfringens, cryptosporidiosis, cyclosporiasis, Escherichia, giardiasis, salmonellosis, shigellosis, toxoplasmosis, yersiniosis, and hepatitis A  Read more about foodborne illness at the <a href="#">Wisconsin Department of Health</a> or <a href="#">Minnesota Department of Health</a> .  *Report foodborne illness to the state Department of Health.
---	---

Recognizing environmental hazards can help community health nurses provide information to community members and implement prevention strategies that promote positive outcomes. See Figure 18.1<sup>2</sup> for an example of prevention strategies related to the environmental hazard of lead poisoning.

2. “preventable.jpg” by unknown author for [National Center for Environmental Health, Division of Environmental Health Science and Practice](#) is in the [Public Domain](#).



Figure 18.1 Lead Poisoning

The American Nurses Association (ANA) established an *Environmental Health* standard defined as, “The registered nurse practices in a manner that advances environmental safety and health.”<sup>3</sup> Review the competencies for this standard in the following box.

3. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.

## ANA's Environmental Health Competencies<sup>4</sup>

The registered nurse:

- Creates a safe and healthy workplace and professional practice environment
- Fosters a professional environment that does not tolerate abusive, destructive, and oppressive behaviors
- Promotes evidence-based practices to create a psychologically and physically safe environment
- Assesses the environment to identify and address the impact of social determinants of health on risk factors
- Reduces environmental health risks to self, colleagues, health care consumers, and the world
- Integrates environmental health concepts in practice
- Communicates information about environmental health risks and exposure risk strategies
- Advocates for the implementation of environmental health principles in communities in which they work and live
- Incorporates technologies to promote safe practice environments
- Uses products or treatments consistent with evidence-based practice to reduce environmental threats and hazards
- Examines how the health care consumer's biography affects their biology, resultant health issues, and the ecosystem

4. American Nurses Association. (2021). *Nursing: Scope and standards of practice* (4th ed.). American Nurses Association.



- Analyzes the impacts of social, political, and economic influences on the human health experience and global environment
- Advances environmental concerns and complaints through advocacy and appropriate reporting mechanisms
- Promotes sustainable global environmental health policies and conditions that focus on prevention of hazards to people and the natural environment

Caring for individuals exposed to environmental hazards will be discussed in the following “Emergency Preparedness, Response, and Recovery” section.

Read more information about promoting environmental health and safety in nursing in the following box.

- ▶ Read a free e-book titled [\*Environmental Health in Nursing\*](#) published by the Alliance of Nurses for Healthy Environments (ANHE).
- ▶ Review the “[Environmental Safety](#)” section of the “Safety” chapter in *Open RN Nursing Fundamentals*.

## 18.3 Emergency Preparedness, Response, and Recovery

A **disaster** is defined as a serious disruption of the functioning of a community or a society at any scale due to hazardous events interacting with conditions of exposure, vulnerability, and capacity that lead to human, material, economic, and environmental losses and impacts.<sup>1</sup> Every community must prepare to respond to disasters that include natural events (e.g., tornadoes, hurricanes, floods, wildfires, earthquakes, or disease outbreaks), man-made events (e.g., harmful chemical spills, mass shootings, or terrorist attacks), or infectious disease outbreaks. See Figure 18.2<sup>2</sup> for an image of the effects of the natural disaster Hurricane Katrina.



Figure 18.2 Effects of Hurricane Katrina

**Emergency preparedness** is the planning process focused on avoiding or reducing the risks and hazards resulting from a disaster to optimize

1. National Academies of Sciences, Engineering, and Medicine. (2021). *The future of nursing 2020-2030: Charting a path to achieve health equity*. National Academies Press. <https://doi.org/10.17226/25982>
2. “LA\_1603\_9thDistDam121.jpg” by Booher, Andrea, Photographer is in the Public Domain.

population health and safety. **Disaster management** refers to the integration of emergency response plans throughout the life cycle of a disaster event. Because disasters cause physical and psychological effects in a community, emergency preparedness and disaster management emphasize collaboration and cooperative aid among health care institutions and community agencies to ensure a coordinated and effective response.<sup>3</sup>

- ▶ Read the American Nurses Association resource regarding [Disaster Preparedness](#).

Emergency preparedness and disaster management are based on four key concepts: preparedness, mitigation, response, and recovery. This process guides decision-making when an emergency or disaster occurs in a community.<sup>4</sup> After the disaster event has concluded, evaluation of the effectiveness of the response occurs as part of planning emergency preparedness. See Figure 18.3<sup>5</sup> for a diagram that illustrates this theoretical framework for emergency preparedness. Each of these concepts is further discussed in the following subsections.

3. Savage, C. L. (2020). *Public/community health and nursing practice: Caring for populations* (2nd ed.). FA Davis.
4. Emergency Management Institute. (2013). *ISS-111.A: Livestock in disasters*. Federal Emergency Management Agency. <https://training.fema.gov/is/courseoverview.aspx?code=iss-111.a&lang=en>
5. “Environmental Health and Emergency Preparedness” by Dawn Barone for [Open RN](#) is licensed under [CC BY 4.0](#)




Figure 18.3 Key Concepts in Emergency Preparedness and Response

## Preparedness

**Preparedness** includes planning, training personnel, and providing educational activities regarding potential disastrous events. Planning includes evaluating environmental risks and social vulnerabilities of a community. Environmental risk refers to the probability and consequences of an unwanted accident in the environment in which community members live, work, or play. Risk assessment also includes assessing social vulnerabilities that affect community resilience.<sup>6</sup>

6. Flanagan, B. E., Hallisey, E. J., Adams, E., & Lavery, A. (2018). Measuring community vulnerability to natural and anthropogenic hazards: The Centers

**Social vulnerability** refers to the characteristics of a person or a community that affect their capacity to anticipate, confront, repair, and recover from the effects of a disaster.<sup>7</sup> Populations living in a disaster-stricken area are not affected equally. Many factors can weaken community members' ability to respond to disasters, including poverty, lack of access to transportation, and crowded housing. Evidence indicates that those living in poverty are more vulnerable at all stages of a catastrophic event, as are racial and ethnic minorities, children, elderly, and disabled people.<sup>8</sup> Socially vulnerable communities are more likely to experience higher rates of mortality, morbidity, and property destruction and are less likely to fully recover in the wake of a disaster compared to communities that are less socially vulnerable. Community health nurses must plan emergency responses to disasters that address these social vulnerabilities to decrease human suffering and financial loss.

The Centers for Disease Control and Prevention (CDC) and the Agency for Toxic Substances and Disease Registry (ATSDR) created a Social Vulnerability Index database and mapping tool designed to assist state, local, and tribal

for Disease Control and Prevention's social vulnerability index. *Journal of Environmental Health*, 80(10). 34–36. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7179070/>

7. Flanagan, B. E., Hallisey, E. J., Adams, E., & Lavery, A. (2018). Measuring community vulnerability to natural and anthropogenic hazards: The Centers for Disease Control and Prevention's social vulnerability index. *Journal of Environmental Health*, 80(10). 34–36. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7179070/>
8. Flanagan, B. E., Hallisey, E. J., Adams, E., & Lavery, A. (2018). Measuring community vulnerability to natural and anthropogenic hazards: The Centers for Disease Control and Prevention's social vulnerability index. *Journal of Environmental Health*, 80(10). 34–36. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7179070/>



disaster management officials in identifying the locations of their most socially vulnerable populations. Geographic patterns of social vulnerabilities can be used in all phases of emergency preparedness and disaster management. See Figure 18.4<sup>9</sup> for an image of social vulnerability mapping.




Figure 18.4 Social Vulnerability Mapping Waiting on permission

9. This image is derivative of Flanagan, B. E., Hallisey, E. J., Adams, E., & Lavery, A. (2018). Measuring community vulnerability to natural and anthropogenic hazards: The Centers for Disease Control and Prevention's social vulnerability index. *Journal of Environmental Health*, 80(10), 34–36. Access the report at <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7179070/>

► View the [CDC/ATSDR Social Vulnerability Index Interactive Map](#).

## Mitigation

**Mitigation** refers to actions taken to prevent or reduce the cause, impact, and consequences of disasters. Health care institutions and community health agencies plan using the three C's to mitigate the effects of a disaster:

- **Communication:** An emergency communication plan identifies tools, resources, teams, and strategies to ensure effective actions during emergencies.
- **Coordination:** Coordination plays a crucial role in efficiency and effectiveness of disaster management by providing a big picture of an emergency and reducing uncertainty levels among responders.
- **Collaboration:** Collaboration allows responders to act together smoothly and helps reduce impact of the disaster.

## Response

The **response** phase occurs in the immediate aftermath of a disaster. When a disaster occurs, actions are taken to save lives, treat injuries, and minimize the effect of the disaster. Immediate needs are addressed, such as medical treatment, shelter, food, and water, as well as psychological support of survivors. Personal safety and well-being in an emergency and the duration of the response phase depend on the level of a community's preparedness. Examples of response activities include implementing disaster response plans, conducting search and rescue missions, and taking actions to protect oneself, family members, pets, and other community members.<sup>10</sup>

10. Emergency Management Institute. (2013). *ISS-111.A: Livestock in disasters*.

While the immediate actions of responding to a disaster are treating physical injuries, psychological effects must be addressed as well. To minimize psychological effects, nurses and first responders can provide support to victims of the disaster by following these tips from the Substance Abuse and Mental Health Services Administration (SAMHSA)<sup>11</sup>:

- **Promote safety:** Ensure basic needs are met and provide simple instructions about how to receive these basic needs.
- **Promote calm:** Listen to people express their feelings and provide empathy and compassion even if they are angry, upset, or acting out. Offer objective information about the situation and efforts being made to help those affected by the disaster.
- **Promote connectedness:** Help people connect with friends, family members, and other loved ones. Keep families and family units together as best as possible, especially by keeping children with those whom they feel safe.
- **Promote self-efficacy:** Give suggestions about how people can help themselves and guide them toward the resources available. Encourage families and individuals to help meet their own needs.
- **Promote help and hope:** Know what services are available and direct people to those services and continue to update people about what is being done. When people are worried or scared, remind them that help is on the way.

Federal Emergency Management Agency. <https://training.fema.gov/is/courseoverview.aspx?code=is-111.a&lang=en>

11. Substance Abuse and Mental Health Services Administration. (2005). *Psychological first aid for first responders*. [Handout]. U.S. Department of Health & Human Services. <https://store.samhsa.gov/product/Psychological-First-Aid-for-First-Responders/NMH05-0210>



# Disaster Response Protocols

When thinking about responding to a disaster, first responders and emergency personnel come to mind such as law enforcement, fire departments, and emergency medical technicians (EMTs). However, nurses are also called upon to assist in emergencies or disasters and must be competent in responding. Nurses may be involved in triaging individuals for treatment.


To respond effectively when a disaster occurs, emergency responders perform **triage** by prioritizing treatment for individuals affected by the disaster or emergency. **Field triage** sorts victims affected by the event and ranks victims based on the severity of their symptoms. **Disaster triage** determines the severity of injuries suffered by victims and then systematically distributes them to local health care facilities based on their severity.

**Simple Triage and Rapid Treatment (START)** is an example of a triage system established by the U.S. Department of Health and Human Services that prioritizes treatment of victims by using standard colors indicating the severity of symptoms and prognosis. See Figure 18.5<sup>12</sup> for the START algorithm. The following colors indicate severity of injury and prognosis:

- **RED:** Emergent needs
  - Life-threatening needs, such as alterations in airway, breathing, and circulation; impairment in neurological systems; or severe, life-threatening injuries.
  - They may have less than 60 minutes to survive.
  - These patients will be seen first or immediately.
- **YELLOW:** Urgent, but delayed needs

12. "[StartAdultTriageAlgorithm.png](#)" by unknown author at [CHEMM](#) is in the [Public Domain](#).

- Life-threatening needs; status is not anticipated to change quickly or significantly in the next hours, so transport can be delayed.
- **GREEN:** Non-urgent needs, often referred to as the “walking wounded”
  - Minor injuries; status is not likely to deteriorate over the next several days.
  - Many individuals can assist with obtaining their own care.
- **BLACK:** The person has died or is expected to die soon
  - This person is unlikely to survive given the severity of their injuries, level of available care, or both.
  - Palliative care and pain relief should be provided.



Adopted from <http://www.start-triage.com>

Figure 18.5 START Adult Triage

# Providing Care for Those Exposed to Environmental Hazards

Nurses may be involved in caring for clients who have been exposed to chemicals or other environmental hazards. See Table 18.3 for assessment findings and interventions for a variety of exposures. **Chelation therapy** is a treatment indicated for heavy metal poisoning such as mercury, arsenic, and lead. Chelators are medications that bind to the metals in the bloodstream to increase urinary excretion of the substance.

Some chemical exposures require decontamination to treat the individual, as well as to protect others around them, including first responders, nurses, and other clients. **Decontamination** is any process that removes or neutralizes a chemical hazard on or in the client to prevent or mitigate adverse health effects to the client; protect emergency first responders, health care facility first receivers, and other clients from secondary contamination; and reduce the potential for secondary contamination of response and health care infrastructure. For example, if a farmer enters a rural hospital's emergency department after chemical exposure to an insecticide spray, decontamination may be required. See Figure 18.6<sup>13</sup> for an image of decontamination.

13. "[decontamination26.jpg](#)" by Benjamin Crossley CDP/[FEMA](#) is in the [Public Domain](#).



Figure 18.6 Decontamination

The decision to decontaminate an individual should take into account a combination of these key indicators<sup>14</sup> :

- Signs and symptoms of exposure displayed by the client
- Visible evidence of contamination on the client's skin or clothing
- Proximity of the client to the location of the chemical release
- Contamination detected on the client using appropriate detection technology
- The chemical and its properties
- Request by the client for decontamination, even if contamination is

14. U.S. Department of Homeland Security, & U.S. Department of Health & Human Services. (2014). *Patient decontamination in a mass chemical exposure incident: National planning guidance for communities* [PDF]. <http://www.phe.gov/Preparedness/responders/Documents/patient-decont-natl-plng-guide.pdf>

unlikely

## Table 18.3 Assessment Findings and Interventions for Exposure to Various Environmental Hazards



Chemical or Hazard	Assessment Findings	Interventions
<p><b>Carbon Monoxide (CO) Poisoning</b></p> <p>(Auto exhaust and improperly vented or malfunctioning furnaces or fuel-burning devices)<sup>15</sup></p>	<p>Primarily decreased mental status from confusion to coma</p> <p>May have cherry-red appearance of the lips and skin</p> <p>*Note: Pulse oximetry does not reflect accurate oxygenation levels because CO binds to hemoglobin.</p>	<ul style="list-style-type: none"> <li>• Remove from source of exposure</li> <li>• Provide fresh air</li> <li>• Administer high-flow oxygen by face mask; may require intubation and mechanical ventilation</li> <li>• Obtain arterial blood gas (ABG) sample of carboxyhemoglobin</li> <li>• Administer hyperbaric oxygen, if necessary</li> </ul>
<p><b>Lead Poisoning</b></p> <p>(Lead-contaminated paint dust, water, or food and bullets in wild game)<sup>16, 17</sup></p>	<p>Abdominal pain, constipation, fatigue, joint pain, muscle pain, headache, anemia, memory deficits, psychiatric symptoms, elevated blood pressure, decreased kidney function, decreased sperm count, increased mortality</p> <p>*Note: Some symptoms may be irreversible.</p>	<ul style="list-style-type: none"> <li>• Remove source of exposure</li> <li>• Monitor blood lead level (toxic effects occur if &gt; 10 mcg)</li> <li>• Monitor hemoglobin and hematocrit for anemia</li> <li>• Monitor urinary output</li> <li>• Administer chelation therapy with ethylenediaminetetraacetic acid (EDTA)</li> <li>• Educate about lead-related health risks and removal of lead-based paint and other sources of exposure using safe OSHA practices</li> </ul>

15. Clardy, P. F., & Manaker, S. (2021, June 17). Carbon monoxide poisoning. *UpToDate*. Accessed April 3, 2022, from [www.update.com](http://www.update.com)
16. World Health Organization. (2021, October 11). *Lead poisoning*. <https://www.who.int/news-room/fact-sheets/detail/lead-poisoning-and-health>
17. Goldman, R. H., & Hu, H. (2021, November 4). Lead exposure and poisoning in adults. *UpToDate*. Accessed April 3, 2022, from [www.update.com](http://www.update.com)



<p><b>Formaldehyde Poisoning</b></p> <p>(Construction and agriculture products and disinfectants)<sup>18</sup></p>	<p>Eye and skin irritation, abdominal pain, bronchospasm, shortness of breath, decreased respiratory rate, acute kidney failure</p>	<ul style="list-style-type: none"> <li>• Remove from source of exposure</li> <li>• Provide fresh air (fumes are toxic)</li> <li>• Decontaminate</li> <li>• Flush eyes with water</li> <li>• Shower to cleanse skin</li> <li>• Administer oxygen as needed</li> <li>• If swallowed, don't induce vomiting</li> <li>• Administer activated charcoal or gastric lavage</li> </ul>
<p><b>Arsenic Poisoning</b></p> <p>(Contaminated groundwater, tobacco smoke, hide tanning, and pressure treated wood)<sup>19,20</sup></p>	<p>Nausea/vomiting, abdominal pain, diarrhea, paresthesias, muscle cramping, skin pigmentation changes, skin lesions and cancers, cardiac dysrhythmias, death</p>	<ul style="list-style-type: none"> <li>• Decontaminate skin and hair</li> <li>• Test urine for arsenic (it is rapidly cleared from blood)</li> <li>• Test hair follicles, fingernails, and skin for long-term exposure</li> <li>• Administer activated charcoal</li> <li>• Administer chelation therapy with dimercaprol or DMSA</li> <li>• Administer IV fluids to maintain urine output</li> <li>• Provide continuous cardiac monitoring</li> <li>• Identify and remove source of exposure</li> </ul>

18. Agency for Toxic Substances and Disease Registry. (2014, October 21). *Medical management guidelines for formaldehyde*. Centers for Disease Control and Prevention. <https://wwwn.cdc.gov/TSP/MMG/MMGDetails.aspx?mmgid=216&toxid=39>
19. World Health Organization. (2018, February 15). *Arsenic*. <https://www.who.int/news-room/fact-sheets/detail/arsenic>
20. Goldman, R. H. (2020, October 8). Arsenic exposure and poisoning. *UpToDate*. Accessed April 3, 2022, from [www.update.com](http://www.update.com)

<p><b>Mercury</b></p> <p>(Thermometers, sphygmomanometers, fluorescent light bulbs, amalgam tooth fillings, and contaminated fish)<sup>21</sup></p>	<p>Acute inhalation exposure in occupational settings may cause cough, dyspnea, chest pain, excessive salivation, inflammation of gums, severe nausea/vomiting, diarrhea, dermatitis</p>	<ul style="list-style-type: none"> <li>• Remove from source of exposure</li> <li>• Blood or urine mercury tests</li> <li>• If inhaled, oxygenation, bronchodilators, chelation treatment</li> <li>• If ingested, administer chelation therapy</li> <li>• Fish with 0.3 to 0.49 ppm of mercury can be safely consumed three times per month but those with greater than 0.5 ppm should be avoided</li> </ul>
<p><b>Radon Gas</b></p> <p>(Naturally occurring gas resulting from the decay of trace amounts of uranium found in the earth's crust)<sup>22</sup></p>	<p>Persistent cough, hoarseness, wheezing, shortness of breath, coughing up blood, chest pain, frequent respiratory infections like bronchitis and pneumonia, loss of appetite, weight loss, fatigue, lung cancer</p>	<ul style="list-style-type: none"> <li>• Measure levels of radon in the home (testing kits available)</li> <li>• Install radon mitigation system in home</li> <li>• Provide public education</li> </ul>

21. Beauchamp, G., Kusin, S., & Elinder, C. (2022, February 1). *Mercury toxicity*. UpToDate. Accessed April 3, 2022, from [www.update.com](http://www.update.com)

22. National Radon Defense. (n.d.). *Radon symptoms*. <https://www.nationalradondefense.com/radon-information/radon-symptoms.html>

<p><b>Infectious Disease</b></p> <p>(HIV, hepatitis, sexually transmitted diseases, and COVID-19)</p>	<p>Symptoms are based on disease process</p>	<ul style="list-style-type: none"> <li>• Blood tests for HIV, hepatitis, syphilis</li> <li>• Quarantine and contact trace as indicated by health department</li> <li>• Implement transmission-based precautions based on route of transmission</li> <li>• Provide public education about the disease and how it is spread</li> <li>• Report communicable disease based on state reporting requirements. Each state has requirements for reporting communicable disease to the Department of Health. For example, see <ul style="list-style-type: none"> <li>▶ <a href="#">Wisconsin's Department of Health Communicable Disease Reporting Requirements</a> or</li> <li>▶ <a href="#">Minnesota's Department of Health Communicable Disease Reporting Requirements</a>.</li> </ul> </li> </ul>
---	--	---

<b>Frostbite</b>  (Overexposure of skin to cold) <sup>23</sup>	White or grayish color of exposed skin, may be hard or waxy to touch; lack of sensitivity to touch or numbness and tingling; clear or blood-filled blisters after thawing; cyanosis after rewarming indicates necrosis	<ul style="list-style-type: none"> <li>• Remove wet clothing</li> <li>• Warm skin slowly in warm (not hot) water or with body heat</li> <li>• Do not rub area or allow walking on frostbitten feet because of potential tissue damage</li> <li>• May administer tPA for severe injury with thrombosis</li> </ul>
<b>Organophosphates</b>  (Insecticides and bioterrorism nerve agents) <sup>24</sup>	<p>Acute onset of symptoms related to cholinergic excess: bradycardia, increased salivation, tearing, urination, vomiting and diarrhea, diaphoresis, paralysis, respiratory failure, hypotension, seizures</p> <p>Intermediate syndrome: neck flexion weakness, cranial nerve abnormalities, muscle weakness</p>	<ul style="list-style-type: none"> <li>• Remove from source of exposure</li> <li>• Aggressively decontaminate</li> <li>• Administer 100% oxygen (may require endotracheal intubation)</li> <li>• Monitor RBC acetylcholinesterase</li> <li>• Administer atropine (prevents cholinergic activation)</li> <li>• Administer IV fluids for hypotension</li> <li>• Administer benzodiazepines for seizures</li> </ul>

23. Zafren, K., & Mechem, C. C. (2021, February 1). Frostbite: Emergency care and prevention. *UpToDate*. Accessed April 3, 2022, from [www.update.com](http://www.update.com)

<b>Bi terrorism</b>  (Anthrax, smallpox, nerve agents, and ricin) <sup>25</sup>	Symptoms are based on the agent	<ul style="list-style-type: none"> <li>• Activate 911</li> <li>• Decontaminate as indicated</li> <li>• Manage ABCs</li> <li>• Coordinate with emergency management officials</li> </ul>
---	---------------------------------	---

- ▶ Access up-to-date, evidence-based information for suspected poisoning at the [Poison Control Center](#) or call 1-800-222-1222.
- ▶ Read more the U.S. Department of Homeland Security and the U.S. Department of Health and Human Services’s [Patient Decontamination in a Mass Chemical Exposure Incident PDF](#).

## Recovery

During the recovery period, restoration efforts occur concurrently with regular operations and activities. The recovery period from a disaster can be prolonged. Examples of recovery activities include the following<sup>26</sup>:

24. Bird, S. (2021, September 23). Organophosphate and carbamate poisoning. *UpToDate*. Accessed April 3, 2022, from [www.update.com](http://www.update.com)
25. Adalja, A. A. (2022, January 10). Identifying and managing casualties of biological terrorism. *UpToDate*. Accessed April 3, 2022, from [www.update.com](http://www.update.com)
26. Emergency Management Institute. (2013). *ISS-111.A: Livestock in disasters*. Federal Emergency Management Agency. <https://training.fema.gov/is/courseoverview.aspx?code=iss-111.a&lang=en>

- Preventing or reducing stress-related illnesses and excessive financial burdens
- Rebuilding damaged structures
- Reducing vulnerability to future disasters

When people are affected by a disaster, they may respond in a variety of different ways. It is natural and expected to respond to a disaster with emotions such as fear, worry, sadness, anxiety, depression, and despair. Many people exhibit **resiliency**, the ability to cope with adversity and recover emotionally from a traumatic event.<sup>27</sup> However, the mental health of the population must be considered and monitored during recovery from any disastrous event. For example, some people may relive previous traumatic experiences or revert to using substances to cope. Behavioral health responses such as post-traumatic stress disorder (PTSD), substance use disorder, and increased risk for suicide should always be considered when assessing individuals' responses to a disaster.

Effects from trauma extend beyond the physical damages from any disaster. It may take time for individuals to recover physically and emotionally. Survivors of a community disaster should be encouraged to take steps to support each other to promote adaptive coping. Use the following box to read additional information in the “Tips for Survivors of a Traumatic Event” handout by the Substance Abuse and Mental Health Services Administration (SAMHSA).

27. Substance Abuse and Mental Health Services Administration. (2022, March 23). *Disaster preparedness, response, and recovery*. U.S. Department of Health & Human Services. <https://www.samhsa.gov/disaster-preparedness>

View a supplementary YouTube video<sup>28</sup> on disaster management and triage: [Disasters and Triage: Community Health – Fundamentals of Nursing | @LevelUpRN](#)

▶ Read the SAMHSA handout on “[Tips for Survivors of a Traumatic Event](#)” PDF.

▶ Review concepts related to loss and the stages of grief in the “[Grief and Loss](#)” chapter of *Open RN Nursing Fundamentals*.

## Agencies Providing Emergency Assistance

Many federal, state, and local agencies provide support to communities during disasters. The Federal Emergency Management Agency (FEMA) is the agency that promotes disaster mitigation and readiness and coordinates response and recovery following the declaration of a major disaster. FEMA defines a disaster as an event that results in large numbers of deaths and injuries; causes extensive damage or destruction of facilities that provide and sustain human needs; produces an overwhelming demand on state and local

28. Level Up RN. (2023, June 21). *Disasters and triage: community health – fundamentals of nursing* | @LevelUpRN [Video]. YouTube. All rights reserved. [https://www.youtube.com/watch?v=VA\\_W6OHLx5M](https://www.youtube.com/watch?v=VA_W6OHLx5M)



response resources and mechanisms; causes a severe long-term effect on general economic activity; and severely affects state, local, and private sector capabilities to begin and sustain response activities.<sup>29</sup> FEMA employees represent every U.S. state, local, tribal, and territorial area and are committed to serving our country before, during, and after disasters.

Disasters are declared using established guidelines and procedures. Because all disasters are local, they are initially declared at the local level. This declaration is typically made by the local mayor. When the mayor determines that capabilities of local resources have been or are expected to be exceeded, state assistance is requested. If the state chooses to respond to a disaster, the governor of the state will direct implementation of the state's emergency plan. If the governor determines that the resource capabilities of the state are exceeded, the governor can request that the president declare a major disaster in order to make federal resources and assistance available to qualified state and local governments. This ordered sequence is important to ensure appropriate financial assistance.<sup>30</sup>

A **state of emergency** is declared when public health or the economic stability of a community is threatened, and extraordinary measures of control may be needed. For example, an infectious disease outbreak like COVID-19 can cause the declaration of a state of emergency. A county or municipal agency is designated as the local emergency management agency, and local law specifies the chain of command in emergencies. Use the following box to access more information about federal and local agencies that provide emergency assistance.

29. Emergency Management Institute. (2013). *ISS-111.A: Livestock in disasters*. Federal Emergency Management Agency. <https://training.fema.gov/is/courseoverview.aspx?code=is-111.a&lang=en>

30. Emergency Management Institute. (2013). *ISS-111.A: Livestock in disasters*. Federal Emergency Management Agency. <https://training.fema.gov/is/courseoverview.aspx?code=is-111.a&lang=en>

## Examples of Organizations That Provide Emergency Assistance

### Federal

- ▶ [Federal Emergency Management Agency \(FEMA\)](#)
- ▶ [Strategic National Stockpile](#)
- ▶ [Cybersecurity and Infrastructure Security Agency \(CISA\)](#)

### Local

- ▶ [Local FEMA agencies \(each state\)](#)
- ▶ [American Red Cross](#)

Local county emergency management divisions

## 18.4 Nurse Roles in Emergency Response

Together with emergency personnel, community organizations, health care providers, and other members of the health care team, nurses play a key role in carrying out emergency response plans. Before, during, and after disasters, nurses provide education and health promotion interventions to protect the community and keep people safe from environmental hazards. During the emergency response, nurses assist in first aid, triage clients' needs, direct victims to resources and community support, and continuously assess and monitor clients' physical and psychological needs. Nurses also assist with community organization through logistical organization of response and prevention needs, such as coordination of blood drives, clothing drives, management of shelters, and other social needs that exist in the community. When disaster strikes, nurses are challenged to use their skills and scope of practice to the fullest to provide the best care for their community.<sup>1</sup>

► See an example [Disaster Nursing Timeline on page 250](#) of *The Future of Nursing 2020-2030 : Charting a Path to Achieve Health Equity* ebook.

The ability to care for and protect vulnerable citizens depends substantially on the preparedness of the nursing workforce. The nursing workforce available to participate in U.S. disaster and public health emergency response includes all licensed nurses (licensed practical/vocational nurses [LPNs/LVNs] and registered nurses [RNs]), civilian and uniformed services nurses at the federal and state levels, nurses who have recently

1. National Academies of Sciences, Engineering, and Medicine. (2021). *The future of nursing 2020-2030: Charting a path to achieve health equity*. National Academies Press. <https://doi.org/10.17226/25982>

retired, and nurses who volunteer in organizations (e.g., National Disaster Medical System, Medical Reserve Corps, National Voluntary Organizations Active in Disasters, and the American Red Cross). Each of these entities plays a critical role in the nation's ability to respond to and recover from disasters and large-scale public health emergencies such as the COVID-19 pandemic.<sup>2</sup> The increasing frequency of natural and environmental disasters, along with public health emergencies such as the COVID-19 pandemic, highlights the critical importance of having a national nursing workforce prepared with the knowledge, skills, and abilities to respond.<sup>3</sup>

## Nurse Roles in Pandemics and Other Infectious Disease Outbreaks

When infectious disease outbreaks occur, nurses are called to the front lines to care for clients very quickly. In the past few decades, infectious disease outbreaks have been occurring more frequently and with more intensity, and nurses, along with health care agencies, must be prepared to respond to such emergencies. Some of the ways nurses respond are by doing the following<sup>4</sup>:

- Tracking disease prevalence and epidemiology of diseases
- Assisting with screening and testing, along with vaccine distribution
- Providing direct care to affected individuals in hospitals and outpatient

2. National Academies of Sciences, Engineering, and Medicine. (2021). *The future of nursing 2020-2030: Charting a path to achieve health equity*. National Academies Press. <https://doi.org/10.17226/25982>

3. National Academies of Sciences, Engineering, and Medicine. (2021). *The future of nursing 2020-2030: Charting a path to achieve health equity*. National Academies Press. <https://doi.org/10.17226/25982>

4. National Academies of Sciences, Engineering, and Medicine. (2021). *The future of nursing 2020-2030: Charting a path to achieve health equity*. National Academies Press. <https://doi.org/10.17226/25982>

clinics

- Educating the public and community about disease prevention and treatment
- Providing psychological support to those fearful of the emergency

New infectious diseases require nurses to receive continuing education regarding infection prevention, diagnosis, treatment, and health promotion. This may include gaining new knowledge about medications, transmission precautions, personal protective equipment (PPE), and evidence-based nursing interventions and medical treatments. Nurses must use critical thinking and apply new evidence-based practices to provide quality care to individuals in their community affected by disease.

## The Impact of the COVID-19 Pandemic on Nurses

In December 2019 the novel coronavirus, known as COVID-19, was first detected in China. By March 2020 the World Health Organization declared a worldwide pandemic for what has been called the worst public health emergency in the past 100 years. By March 2022 almost 80 million cases were identified in the United States, with over 977,000 deaths. Worldwide, over 488 million cases were diagnosed, with over 6 million deaths.<sup>5,6</sup>

► View the CDC's [Covid Data Tracker](#) or [Our World in Data's Data Tracker](#) for up-to-date COVID information.

5. Centers for Disease Control and Prevention. (n.d.). *COVID data tracker*. <https://covid.cdc.gov/covid-data-tracker/#datatracker-home>

6. [Daily new confirmed COVID-19 feathers per million people](#) by [Our World in Data](#) is licensed under [CC BY 4.0](#)

The COVID-19 pandemic created a health care environment never before seen in modern times. Nurses were called upon to care for clients, often lacking proper personal protective equipment (PPE) and medical equipment, to properly support client health. They cared for clients above and beyond the acuity that they normally managed in their area of expertise and provided frequent end-of-life care. They communicated regularly by phone with loved ones of clients who were dying due to quarantine guidelines and visitor limitations. Staffing shortages became critical as illness spread to nurses and their family members.<sup>7</sup>

As a result, thousands of nurses were still stressed, frustrated, and overwhelmed two years into the pandemic. In October 2021 the American Nurses Association found the following in a mental health and wellness survey of nurses<sup>8</sup>:

- Over 34% rated their emotional health as not, or not at all, emotionally healthy.
- 42% reported having an extremely stressful, disturbing, or traumatic experience due to COVID-19.
- Among nurses who said they intend to leave their position in the next six months, 48% cited the top reason as work negatively affecting their health and well-being, followed closely by insufficient staffing (41%).

The American Nurses Foundation Executive Director, Kate Judge stated, “The

7. National Academies of Sciences, Engineering, and Medicine. (2021). *The future of nursing 2020-2030: Charting a path to achieve health equity*. National Academies Press. <https://doi.org/10.17226/25982>

8. American Nurses Foundation. (2021, October 26). *New survey data: Thousands of nurses are still stressed, frustrated, and overwhelmed almost 2 years into the pandemic*. [News release]. <https://www.nursingworld.org/news/news-releases/2021/american-nurses-foundation-releases-comprehensive-survey-about-nurses/>

COVID-19 pandemic is not over yet, and its impact will persist for a long time. Its challenges have left the nursing profession in a particularly vulnerable state, exacerbating nurse staffing shortages, and negatively impacting nurses' quality of life. Nurses are playing a pivotal role in efforts to end this pandemic, so we must ensure nurses are physically and psychologically safe and healthy to function optimally in caring first for themselves, their families, clients, and communities. A robust nursing workforce is essential to our nation's health and, therefore, nurses' well-being and mental health must be a top priority. The data collected from this survey overwhelmingly demonstrate the need to provide consistent and comprehensive support for our nation's nurses."<sup>9</sup>

- ▶ The American Psychiatric Nurses Association provides [COVID Resources](#) to support nurses as they address concerns and promote mental health as a result of the COVID-19 pandemic.

## Preventing and Managing Stress in Disaster Responders

Engaging in disaster and emergency response work can cause emotional distress for nurses and first responders. Depending on the nature of the event, sources of stress may include exposure to scenes of human suffering, risk for personal harm, life-and-death decision-making, intense workloads, limited resources, and separation from family members who may also be in

9. American Nurses Foundation. (2021, October 26). *New survey data: Thousands of nurses are still stressed, frustrated, and overwhelmed almost 2 years into the pandemic*. [News release]. <https://www.nursingworld.org/news/news-releases/2021/american-nurses-foundation-releases-comprehensive-survey-about-nurses/>

harm's way. Use the link in the following box to view a SAMSHA brochure outlining tips for coping with stress after a disaster or traumatic event.

- ▶ Read the [Disaster Distress Helpline](#) PDF brochure from the SAMHSA on tips for coping with stress after a disaster.

- ▶ Read more information about stress in the health care system and self-care in the “[Burnout & Self-Care](#)” chapter of *Open RN Nursing Management and Professional Concepts*.



## 18.5 Spotlight Application

Sam, a nurse who works at the local hospital, hears about a flooding disaster in a nearby community. Major roads are flooded, and grocery stores have been destroyed. Power lines were knocked down during the storm, and cell phone service is not working. Sam starts to wonder, “What if?”... “What if a disaster happens here and I’m not home with my family – how will I know if they are safe? How will I stay safe? What obligations will I have at work?”

As a compassionate nurse, Sam feels a need to respond to others in a crisis but also has a concern about ensuring the well-being of his family members. A Google search reveals the American Nurses Association document “[Who Will Be There?](#)” PDF. He realizes the discomfort he is feeling about responding to a potential future disaster event is due to the conflict of ethical principles regarding the nurse’s obligation to care for others, as well as the obligation to care for oneself.<sup>1</sup> He decides to respond to this ethical dilemma by preparing his personal response should a disaster occur.

Sam reads additional information called “Make a Plan” provided by the U.S. Department of Homeland Security’s website “[Ready.gov](#)”<sup>2</sup> Several questions come to mind as Sam reflects on his anticipated response to a disaster:

1. How will I receive alerts and warnings about a disaster in my community?
2. Are there any laws in my state that require my participation as a nurse in disaster response?
3. What is my employer’s emergency response plan? What is my expected

1. American Nurses Association. (2017). *Who Will Be There? Ethical, the law, and a nurse’s duty to respond in a disaster.*

[https://www.nursingworld.org/~4ad845/globalassets/docs/ana/who-will-be-there\\_disaster-preparedness\\_2017.pdf](https://www.nursingworld.org/~4ad845/globalassets/docs/ana/who-will-be-there_disaster-preparedness_2017.pdf)

2. U.S. Department of Homeland Security. (2022, May 18). *Make a plan.*

<https://www.ready.gov/plan>

role as an employed nurse?

4. What state and local disaster preparedness plans currently exist in my community?
5. How will I get to work safely if a disaster occurs?
6. How will I communicate with my family members and ensure their safety if a disaster occurs?

Sam reads additional information provided by the Federal Emergency Management Agency (FEMA) and starts to create his [!\[\]\(3dfb8d66e81160ad61421a3452093d1b\_img.jpg\) personal and family disaster plan PDF](#).<sup>3</sup>

### **Reflective Questions:**

1. Consider the questions Sam posed as he considered his potential disaster response. What additional questions do you have about your role in a disaster response? What are the answers to these questions in your community and at your place of employment?
2. Create your own personal and family disaster plan.

3. FEMA. (2018). *Create your family emergency communication plan* [PDF]. <https://www.ready.gov/sites/default/files/2021-04/family-emergency-communication-plan.pdf>

## 18.6 Learning Activities



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=1074#h5p-59>

1



*An interactive H5P element has been excluded from this version of the text. You can view it online here:*

<https://wtcs.pressbooks.pub/nursingmhcc/?p=1074#h5p-60>

2



1. “Environmental Health and Emergency Preparedness Crossword” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)
2. “Environmental Health and Emergency Preparedness Drag and Drop” by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

- ▶ Test your clinical judgment with a NCLEX Next Generation-style question: [Chapter 18, Assignment 1](#)<sup>3</sup>

- ▶ Test your clinical judgment with a NCLEX Next Generation-style case study: [Chapter 18, Case Study 1](#)<sup>4</sup>



3. "Environmental Health and Emergency Preparedness Next Gen Question 1" by [OpenRN](#) is licensed under [CC BY-NC 4.0](#)

4. "Environmental Health and Emergency Preparedness Next Gen Case Study 1" by Kellea Ewen is licensed under [CC BY-NC 4.0](#)

## XVIII Glossary

**Chelation therapy:** Treatment for heavy metal poisoning such as mercury, arsenic, and lead. Chelators are medications that bind to the metals in the bloodstream to increase urinary excretion of the substance.

**Decontamination:** Any process that removes or neutralizes a chemical hazard on or in the client in order to prevent or mitigate adverse health effects to the client; protect emergency first responders, health care facility first receivers, and other clients from secondary contamination; and reduce the potential for secondary contamination of response and health care infrastructure.

**Disaster:** A serious disruption of the functioning of a community or a society at any scale due to hazardous events interacting with conditions of exposure, vulnerability, and capacity that lead to human, material, economic, and environmental losses and impacts.

**Disaster management:** Response to a disaster with integration of emergency response plans throughout the life cycle of a disaster event.

**Emergency preparedness:** The planning process focused on avoiding or ameliorating the risks and hazards resulting from a disaster to optimize population health and safety. The process includes four key concepts: preparedness, mitigation, response, and recovery. Evaluation occurs after a disaster event as the planning process continues.

**Environmental health hazard:** A substance or pathogen that has the ability to cause an adverse health event in individuals or communities.

**Mitigation:** Actions taken to prevent or reduce the cause, impact, and consequences of disasters.

**Preparedness:** Planning, training personnel, and providing educational activities regarding potential disastrous events. Planning includes evaluating environmental risks and social vulnerabilities of a community.

**Recovery:** Restoration efforts occur concurrently with regular operations and

activities, such as preventing or reducing stress-related illnesses and excessive financial burdens.

**Resiliency:** The ability to cope with adversity and recover emotionally from a traumatic event.

**Response:** Actions taken in the immediate aftermath of a disaster, such as saving lives, treating injuries, and minimizing the effects of the disaster. Immediate needs are addressed, such as medical treatment, shelter, food, and water, as well as psychological support of survivors.

**Simple Triage and Rapid Treatment (START):** A triage system from the U.S. Department of Health and Human Services that prioritizes victims by color (red, yellow, green, and black) for efficient and effective treatment.

**Social vulnerability:** The characteristics of a person or a community that affect their capacity to anticipate, confront, repair, and recover from the effects of a disaster.

**State of emergency:** Status that is declared when the public health or the economic stability of a community is threatened, and extraordinary measures of control may be needed.

**Triage:** Prioritizing care for individuals affected by a disaster or emergency.

PART XIX

ANSWER KEY





## Chapter 1

Answers to interactive elements are given within the interactive element.

## Chapter 2

Answers to interactive elements are given within the interactive element.

## Chapter 3

Answers to interactive elements are given within the interactive element.

## Chapter 4

Answers to interactive elements are given within the interactive element.

## Chapter 5

Answers to interactive elements are given within the interactive element.

## Chapter 6

Answers to interactive elements are given within the interactive element.

## Chapter 7

Answers to interactive elements are given within the interactive element.

## Chapter 8

Answers to interactive elements are given within the interactive element.



## Chapter 9

Answers to interactive elements are given within the interactive element.

## Chapter 10

### Critical Thinking Questions

1. DSM-5 lists these as unique mental health illness diagnoses, but the symptoms of each may cross over. In obsessive-compulsive personality disorder, the client exhibits orderliness and perfectionism. However, in personality disorders, the behaviors do not come and go or fluctuate, and there is no insight that the behavior is abnormal. In fact, clients typically believe their behavior is normal and beneficial. Additionally, the client does not perform repetitive actions that are classic to OCD. In contrast, OCD is caused by anxiety, and the severity of the obsessions and compulsions vary according to the level of anxiety. Clients with OCD exhibit repetitive actions such as excessive handwashing or repeatedly checking the door is locked or counting items. Clients have insight into the behavior and view it as abnormal and distressing, so they are more likely to seek treatment.
2. Name that personality disorder:
  - a. Histrionic
  - b. Paranoid
  - c. Borderline
  - d. Schizoid
  - e. Dependent
  - f. Antisocial
  - g. Avoidant
  - h. Narcissistic
  - i. Obsessive-Compulsive
  - j. Schizotypal
3. Check your medication knowledge:
  - a. Low-dose antipsychotics
  - b. Mood stabilizers and low-dose antipsychotics
  - c. Mood stabilizers and omega-3
4. Compare normal adolescent development with trait similarities of personality disorders:

- a. Cluster A: May not trust others, prefers to be alone, and may have magical thinking (imagination)
- b. Cluster B: Disregards right from wrong, lies, gets into trouble, impulsive, overreactive, emotional, and easily influenced by others
- c. Cluster C: Social discomfort, shy, fearful, and lacks self-confidence

Answers to interactive elements are given within the interactive element.

## Chapter 11

Answers to interactive elements are given within the interactive element.

## Chapter 12

Answers to interactive elements are given within the interactive element.

## Chapter 13

### Questions:

1. What symptoms of an eating disorder is Tiffany demonstrating? *Tiffany is exhibiting symptoms of anorexia nervosa.*
2. What other assessment findings does the nurse anticipate? (See *Table 13.3a – Chapter 13*)
3. What laboratory tests will likely be ordered during this visit? *Laboratory tests may include a complete blood count, electrolyte levels, glucose level, thyroid function tests, erythrocyte sedimentation rate (ESR), and creatine phosphokinase (CPK).*
4. What type of psychotherapy would be helpful for Tiffany? *Cognitive behavioral therapy and family-based therapy would be useful psychotherapy treatments for Tiffany.*
5. What conditions would cause Tiffany to be hospitalized? *Tiffany would be hospitalized if her vital signs are unstable, laboratory findings present acute health risk, and she has rapidly worsening symptoms or suicidal ideation with a plan and is unable to contract for safety.*
6. Tiffany is hospitalized. Create a brief nursing care plan for Tiffany including a nursing diagnosis, SMART goal, and 3-5 nursing interventions.

*Diagnosis: Imbalanced Nutrition: Less Than Body Requirements*

*SMART Goal: Client will gain at least one pound weekly until reaching healthy weight classification according to her BMI.*

*Sample Interventions:*

- *The client will be weighed on Mondays, Wednesdays, and Fridays and progress determined based on a target weight established by the dietician.*
- *A pleasant, calm atmosphere will be provided at mealtimes.*
- *The client will be observed during meals to prevent hiding or throwing away food and at least one hour after eating to prevent purging.*

- *The client will be encouraged to make her own menu choices as she approaches her goal weight.*
- *Knowledge and skills gained from individual, family, and group therapy sessions will be reinforced with the client and her family.*

Answers to interactive elements are given within the interactive element.

## Chapter 14

Answers to interactive elements are given within the interactive element.



## Chapter 15

Answers to interactive elements are given within the interactive element.

## Chapter 16

Answers to interactive elements are given within the interactive element.

## Chapter 17

Answers to interactive activities are provided within the activity.

## Chapter 18

Answers to interactive elements are given within the interactive element.